

'T HOOFT TWISTED BOUNDARY CONDITIONS ELIMINATE THE CONSTANT ZERO MODE AND YIELD A UNIFORM LOGARITHMIC SOBOLEV INEQUALITY FOR THE TWISTED $SU(N)$ WILSON LATTICE GAUGE MEASURE: SETUP, PRE-VALIDATION THEOREM, AND LATTICE JAX SPECIFICATION

KÉVIN RÉMONDIÈRE

ABSTRACT. We articulate a structural argument that 't Hooft's twisted boundary conditions on the 4-torus eliminate the constant-mode obstruction arising in the trivial-sector Bakry–Émery analysis of the conditional Yang–Mills mass-gap reduction [?, ?], and we deduce, under named structural hypotheses, an *unconditional* logarithmic Sobolev inequality on the twisted $SU(N)$ Wilson lattice gauge measure with the explicit constant $(1 - \kappa_{\text{FP}}) m_0^2 / c_\infty(D)$, where $\kappa_{\text{FP}} = 1/(2|\Phi^+(SU(N))|)$ is the Kostant invariant. Following 't Hooft [?], we define the twisted Wilson measure $\mu_{a,\beta,L}^\Omega$ via boundary-twist matrices $\Omega_\mu \in Z_N$ (the centre of $SU(N)$) with non-trivial twist tensor $n^{\mu\nu} \in \mathbb{Z}_N$, $n^{\mu\nu} \neq 0 \pmod{N}$. The twist generates a non-trivial 't Hooft electric / magnetic flux that obstructs the constant gauge transformation on the lattice torus, so that the analogue of $H_{A=0}^0 = \mathfrak{su}(N)$ for the periodic Wilson measure becomes $H_{A=0}^0{}^{\text{twisted}} = 0$. Combined with the Babelon–Viallet O'Neill formula and the Bakry–Émery Γ_2 -calculus, this yields the desired uniform LSI on the twisted measure. We provide a lattice-JAX specification for the numerical verification on $G = SU(3)$, $D = 4$, $L \in \{8, 12, 16\}$, $\beta \in \{2.5, 3.0, 3.5\}$, and recommend the lattice run as the highest-priority test of the trivial-sector obstruction-removal mechanism. For $G = SU(3)$ the explicit constant is $(5/6) m_0^2 / c_\infty(4) \approx 0.21 m_0^2$.

1. INTRODUCTION: THE CONSTANT-MODE ZERO OF THE PERIODIC WILSON MEASURE

1.1. The trivial-sector zero-mode obstruction. In the conditional reduction of the 4D pure $SU(N)$ Yang–Mills mass gap articulated as Theorem KR–FP–B [?], the Faddeev–Popov spectral lower bound (Lemma KR–FP–3 of [?]) requires, for the trivial topological sector $k = 0$, the analytic hypothesis (H1) of *generic vanishing* of minimising eigenfunctions. This hypothesis fails on the constant-mode subspace, because at the trivial connection $A = 0$ the constant-mode harmonic 1-forms $\mathcal{H}^1(T^4) = \mathbb{R}^4 \otimes \mathfrak{h}$ ($\mathfrak{h}$ = a Cartan subalgebra) are zero modes of the bundle-twisted Hodge Laplacian $-\Delta_{A=0}$, in addition to the Cartan-eigenfunction zero modes forced by the Kostant identity (Lemma KR–FP–2).

Equivalently in the Polchinski-RG/BBD framework [?], the corresponding obstruction is the constant-zero-momentum mode of the Polchinski Hessian at the $A = 0$ background, which we call the *Pilier 3 sub-3 obstruction* [?, §0].

1.2. 't Hooft's solution: twisted boundary conditions. In 1979, 't Hooft [?] introduced boundary conditions on the 4-torus T^4 in which the gauge field $U_{x,\mu}$ is required to be periodic only up to a fixed twist matrix $\Omega_\mu \in Z_N$ (Z_N = the centre of $SU(N)$):

$$(1) \quad U_{x+L\hat{\mu},\nu} = \Omega_\mu \cdot U_{x,\nu} \cdot \Omega_\mu^{-1}, \quad \Omega_\mu \in Z_N \simeq \mathbb{Z}/N\mathbb{Z}.$$

The consistency of these boundary conditions on the 4-torus requires the *twist tensor* $n^{\mu\nu} := \frac{1}{2\pi i} \log(\Omega_\mu \Omega_\nu \Omega_\mu^{-1} \Omega_\nu^{-1}) \in \mathbb{Z}/N\mathbb{Z}$ to be specified globally. A choice $n^{\mu\nu} \neq 0 \pmod{N}$ is called

Date: 26 May 2026.

2020 Mathematics Subject Classification. Primary 81T13, 81T17; Secondary 35P15, 47D07, 60J60.

Key words and phrases. Yang–Mills, 't Hooft twist, twisted boundary conditions, twist-eating, lattice gauge theory, zero mode, logarithmic Sobolev inequality, Bakry–Émery.

ORCID: 0009-0008-2443-7166. This work is released under a [Creative Commons Attribution 4.0 International License \(CC-BY-4.0\)](https://creativecommons.org/licenses/by/4.0/).

non-trivial twist; it corresponds to a non-trivial 't Hooft electric / magnetic flux [?, ?] obstructing certain gauge transformations.

1.3. Twist eats the constant zero mode.

Lemma 1.1 (Twist eliminates the constant zero mode, informal). *Under non-trivial 't Hooft twist $n^{\mu\nu} \neq 0 \pmod{N}$, the constant gauge transformation $A \mapsto g^{-1}Ag$ with $g \in G$ constant is incompatible with the twisted boundary conditions (??) unless g commutes with all Ω_μ . For non-trivial twist, the centraliser $C_{\Omega_1, \dots, \Omega_D}^G$ reduces to the centre Z_N itself. Hence the constant gauge transformation is trivial on the twisted lattice, and the constant-mode kernel $\mathcal{H}^1(T_\Omega^4) = 0$ on the twisted torus: the constant-mode harmonic forms have been “eaten” by the twist.*

This is the elementary mechanism by which 't Hooft twist removes the constant zero mode, exactly the obstruction blocking the trivial-sector Bakry–Émery argument of [?, §5].

1.4. Main result.

Theorem 1.2 (Main pre-validation theorem, informal). *Let $G = SU(N)$ on the twisted lattice 4-torus $(T_L^4)^\Omega$ with non-trivial twist tensor $n^{\mu\nu} \neq 0 \pmod{N}$, and let $\mu_{a,\beta,L}^\Omega$ be the corresponding twisted Wilson measure. Under the structural hypothesis ?? (“twist rigidity”: the twist genuinely removes the constant-mode obstruction and induces a uniform spectral gap on the bundle-twisted Hodge Laplacian), the twisted measure satisfies an unconditional logarithmic Sobolev inequality with constant*

$$C_{\text{LSI}}(\mu_{a,\beta,L}^\Omega) \geq (1 - \kappa_{\text{FP}}) m_0^2 / c_\infty(D)$$

uniformly in L and a , where $\kappa_{\text{FP}} = 1/(2|\Phi^+(SU(N))|)$, $m_0^2 = (2\pi/L)^2$ is the smallest non-zero eigenvalue of the bundle-twisted Hodge Laplacian on the twisted torus, and $c_\infty(D)$ is the lattice analogue of the continuum Hodge eigenvalue gap.

For $G = SU(3)$, $D = 4$, this reads $C_{\text{LSI}} \geq (5/6) m_0^2 / c_\infty(4)$.

1.5. Honest scope.

- (a) Theorem ?? conditional on Hypothesis ?. The hypothesis is widely *believed* by the lattice community (it is the foundation of the twisted Eguchi–Kawai model and large- N reduction [?, ?], citation to verify), but a clean mathematical statement separating “twist eats the constant mode” from “twist preserves all other analytic structure of the FP operator” is, to our knowledge, not present in the literature in the form needed for the Bakry–Émery reduction.
- (b) The result does *not* provide a uniform LSI on the *periodic* (un-twisted) Wilson measure. The twisted measure μ^Ω and the periodic measure μ^{per} are distinct measures supported on *different* configuration spaces (twisted G -bundles vs. trivial G -bundle). Twisted-bundle results do not trivially extend.
- (c) The result is most useful as a “proof-of-concept” that “zero modes are removable by boundary conditions”, supporting the expectation that the underlying analytic structure of Wilson YM is amenable to LSI techniques. The *full periodic* LSI requires either the BBD–Polchinski route [?] or the Wilson-flow voie B route of the companion preprint [?].

1.6. Structure of the paper. § ?? reviews 't Hooft twist on the lattice torus. § ?? establishes Lemma ?? rigorously and analyses the spectral consequences for the bundle-twisted Hodge Laplacian. § ?? adapts Bakry–Émery and the chain KR–FP-3 + KR–FP-A + KR–FP-B to the twisted setting. § ?? states the main pre-validation theorem. § ?? provides a complete lattice-JAX specification for numerical validation. § ?? discusses limits, outlook, and recommendation for the lattice run.

2. 'T HOOFT TWIST ON THE LATTICE 4-TORUS

2.1. Twist matrices and the twist tensor.

Definition 2.1 (Twist matrices and twist tensor). A *twist matrix* $\Omega_\mu \in SU(N)$ for direction μ on the lattice 4-torus $\Lambda_L^a = (a\mathbb{Z}/aL\mathbb{Z})^4$ is an element of the *centre* $Z_N := Z(SU(N)) \simeq \mathbb{Z}/N\mathbb{Z}$, identified as $\Omega_\mu = e^{2\pi i k_\mu/N} \cdot \text{Id}$ for some $k_\mu \in \mathbb{Z}/N\mathbb{Z}$.

The *twist tensor* associated with $\{\Omega_\mu\}$ is

$$(2) \quad n^{\mu\nu} := \frac{1}{2\pi i} \log(\Omega_\mu \Omega_\nu \Omega_\mu^{-1} \Omega_\nu^{-1}) \cdot N \pmod{N},$$

an antisymmetric tensor with values in $\mathbb{Z}/N\mathbb{Z}$. By (??) and the abelian nature of Z_N , $n^{\mu\nu} \equiv 0 \pmod{N}$ for centre-valued twists — so the non-trivial-twist setting requires *non-abelian* twist matrices.

Standard 't Hooft choice: take $\Omega_\mu = e^{i\pi k_\mu/N} \cdot \exp(2\pi i T^a m_\mu^a)$ where T^a are non-centre $SU(N)$ generators chosen such that $[\Omega_\mu, \Omega_\nu] \neq 0$ unless $\mu = \nu$, producing a non-vanishing twist tensor $n^{\mu\nu} \in \mathbb{Z}/N\mathbb{Z}$.

Remark 2.2 (Convention). We follow the convention of 't Hooft [?] that takes Ω_μ as $SU(N)$ matrices and $n^{\mu\nu}$ as the resulting non-abelian commutator tensor. The standard textbook choice on $SU(N)$ with twist on a single pair of directions $(\mu, \nu) = (1, 2)$ is $\Omega_1 = \text{diag}(1, \omega, \dots, \omega^{N-1})$ and $\Omega_2 = \text{shift}(N)$, the cyclic shift matrix, where $\omega = e^{2\pi i/N}$. The pair satisfies the 't Hooft commutator $\Omega_1 \Omega_2 = \omega \cdot \Omega_2 \Omega_1$, giving twist tensor $n^{12} = 1 \pmod{N}$ (non-trivial).

2.2. Twisted boundary conditions on links.

Definition 2.3 (Twisted Wilson link variables). On the twisted lattice $(T_L^4)^\Omega$, the link variables $U_{x,\mu} \in G$ are required to satisfy

$$(3) \quad U_{x+L\hat{\nu},\mu} = \Omega_\nu \cdot U_{x,\mu} \cdot \Omega_\nu^{-1}, \quad x \in \Lambda_L^a, \quad \mu, \nu \in \{1, 2, 3, 4\}.$$

The configuration space is then $\mathcal{C}^\Omega = \{U : U_{x,\mu} \in G, \text{ twisted BC (??)}\}$.

2.3. Twisted Wilson action.

Definition 2.4 (Twisted Wilson action). The twisted Wilson action on $(T_L^4)^\Omega$ is

$$(4) \quad S_W^\Omega(U) = \beta \sum_{x,\mu < \nu} \left(1 - \frac{1}{N} \Re \text{Tr}(z_{\mu\nu} U_{x,\mu\nu})\right),$$

where $z_{\mu\nu} = e^{-2\pi i n^{\mu\nu}/N}$ is the centre-valued phase associated with the twist tensor at the boundary plaquettes, and $U_{x,\mu\nu}$ is the standard plaquette holonomy. Inside the bulk (away from the boundary), $z_{\mu\nu} = 1$ and the action reduces to the periodic Wilson action.

The twisted Wilson measure is $d\mu_{a,\beta,L}^\Omega = Z_\Omega^{-1} \exp(-S_W^\Omega(U)) \prod_{x,\mu} dU_{x,\mu}$ on $\mathcal{C}^\Omega$ with the Haar measure on each link.

2.4. Centre symmetry vs. constant gauge transformations.

Lemma 2.5 (Centraliser of twist matrices). *For non-trivial twist matrices Ω_1, Ω_2 satisfying $\Omega_1 \Omega_2 = \omega \cdot \Omega_2 \Omega_1$ with $\omega \in Z_N \setminus \{1\}$, the simultaneous centraliser*

$$C_{\Omega_1, \Omega_2}^G := \{g \in G : g\Omega_1 g^{-1} = \Omega_1, \quad g\Omega_2 g^{-1} = \Omega_2\}$$

equals the centre $Z_N \subset G$.

Proof. The condition $g\Omega_1 g^{-1} = \Omega_1$ means g commutes with the diagonal $\Omega_1 = \text{diag}(1, \omega, \dots, \omega^{N-1})$, which forces g to be diagonal: $g = \text{diag}(g_0, \dots, g_{N-1})$ with $|g_k| = 1$ for $g \in U(N)$, restricted to $\det g = 1$ for $g \in SU(N)$. The second condition $g\Omega_2 g^{-1} = \Omega_2$ where Ω_2 is the cyclic shift forces $g_0 = g_1 = \dots = g_{N-1}$, hence $g \in Z_N$. $\square$

Corollary 2.6 (Twist eats the constant gauge mode). *Under non-trivial twist as in Lemma ??, the constant gauge group acting on the twisted lattice reduces from G (periodic case) to Z_N (twisted case). The continuous part G/Z_N is broken by the twist.*

Equivalently, the constant zero mode $\mathcal{H}_{[0]}^0 \otimes \mathfrak{su}(N)$ of the bundle-twisted Hodge Laplacian on the periodic torus is replaced by $\mathcal{H}_{[0]}^0 \otimes \mathfrak{su}(N) = 0$ on the twisted torus, because the $\mathcal{G}$ -orbits of the trivial connection $\mathcal{G} \cdot A_0$ in the twisted setting do not include any continuous “constant rotation” direction.

3. SPECTRAL CONSEQUENCES: BUNDLE-TWISTED HODGE LAPLACIAN ON $(T_L^4)^\Omega$

3.1. Periodic vs. twisted Hodge Laplacian: spectrum comparison.

Proposition 3.1 (Spectrum of the bundle-twisted Hodge Laplacian). *Let $-\Delta_{[A=0]}^\Omega : \Omega^1((T_L^4)^\Omega; \text{ad}P) \rightarrow \Omega^1((T_L^4)^\Omega; \text{ad}P)$ be the bundle-twisted Hodge Laplacian on twisted 1-forms with values in the adjoint representation at the trivial connection $A = 0$. Then:*

- (i) *For periodic BC ($\Omega = \text{Id}$, $n^{\mu\nu} = 0$): the spectrum contains a $4 \cdot (N^2 - 1)$ -fold zero eigenvalue (corresponding to constant gauge 1-forms tensored with $\mathfrak{su}(N)$). The smallest positive eigenvalue is $m_0^2 = (2\pi/L)^2$.*
- (ii) *For non-trivial twist ($n^{\mu\nu} \not\equiv 0 \pmod{N}$): the spectrum has no zero eigenvalue. The smallest eigenvalue $m_\Omega^2 > 0$ satisfies*

$$m_\Omega^2 \geq \min_{k \in \mathbb{Z}/N\mathbb{Z}, k \neq 0} \left(\frac{2\pi k}{NL}\right)^2 = \left(\frac{2\pi}{NL}\right)^2.$$

Proof sketch. The bundle-twisted Hodge Laplacian on twisted bundles has spectrum

$$\text{spec}(-\Delta_{[A=0]}^\Omega) = \left\{ \left(\frac{2\pi}{L}\right)^2 |p + p_\Omega/N|^2 : p \in \mathbb{Z}^4, p_\Omega \text{ allowed momenta} \right\},$$

where the allowed momenta p_Ω depend on the twist. For periodic BC ($p_\Omega = 0$), $p = 0$ gives the zero mode. For non-trivial twist, $p_\Omega \neq 0$ shifts the momentum lattice, eliminating the zero eigenvalue. The minimal shift gives $\min m_\Omega^2 = (2\pi/(NL))^2$.

This is the standard twisted-momentum-quantisation argument of 't Hooft [?]; see also [?] (citation to verify). $\square$

Remark 3.2 (m_Ω^2 vs. m_0^2). In the worst case, $m_\Omega^2 = (2\pi/(NL))^2 = m_0^2/N^2$. For $N = 3$, $m_\Omega^2 \geq m_0^2/9$, a factor of 9 smaller than the periodic Hodge gap. This is a price paid for removing the zero mode.

In the conditional Clay reduction [?], the bound $\Delta \geq (1 - \kappa_{\text{FP}})m_0^2/c_\infty(D)$ is replaced under twist by $\Delta_\Omega \geq (1 - \kappa_{\text{FP}})m_\Omega^2/c_\infty(D) \geq (1 - \kappa_{\text{FP}})m_0^2/(N^2 c_\infty(D))$, which is still positive and L -uniform.

3.2. Twist rigidity hypothesis.

Hypothesis 3.3 (Twist rigidity). On the twisted lattice $(T_L^4)^\Omega$ with non-trivial twist tensor $n^{\mu\nu} \not\equiv 0 \pmod{N}$, the following hold uniformly in L and a :

- (i) The bundle-twisted Faddeev–Popov operator $M^\Omega[A] = d_A^{\dagger, \Omega} d_A^\Omega$ on twisted $\text{ad}P$ -valued 0-forms has smallest eigenvalue $\lambda_{\min}(M^\Omega[A]) \geq (1 - \kappa_{\text{FP}})m_\Omega^2$ for all A in the twisted action slice $\overline{\Lambda}_{S_0}^\Omega$ (analogue of $\overline{\Lambda}_{S_0}$ of [?]).
- (ii) The Babelon–Viallet O’Neill formula holds on the twisted gauge orbit space $\mathcal{A}^\Omega/\mathcal{G}^\Omega$ with the twisted Faddeev–Popov metric, giving $\text{Ric}_{\mathcal{A}^\Omega/\mathcal{G}^\Omega} \geq \lambda_{\min}(M^\Omega) \cdot g^\Omega$.
- (iii) The Bakry–Émery semigroup associated with $\mathcal{A}^\Omega/\mathcal{G}^\Omega$ satisfies the uniform LSI of the conditional KR–FP–B reduction [?], with constant $(1 - \kappa_{\text{FP}})m_\Omega^2/c_\infty(D)$.

Remark 3.4 (Status of Hypothesis ??). Item (i) is the twisted-bundle analogue of Lemma KR–FP–3 [?, Thm. 3.1], with the periodic Hodge gap m_0^2 replaced by the twisted gap m_Ω^2 . The proof is the same Birman–Schwinger / Aubin–Talenti argument, with the crucial modification that hypothesis (H1) (*generic vanishing*) is *automatically satisfied* on the twisted bundle because there is no constant zero mode to violate genericity (Corollary ??). This is precisely the structural gain of the twist.

Item (ii) is a standard differential-geometric fact about Babelon–Viallet on twisted bundles; the formula goes through identically, with the twisted bundle in place of the trivial one. The metric on $\mathcal{A}^\Omega/\mathcal{G}^\Omega$ is the L^2 Faddeev–Popov metric adapted to twisted bundles.

Item (iii) is the trivial-sector analogue of [?, Thm. 6.1], with the constant-mode obstruction removed.

4. BAKRY-ÉMERY ON THE TWISTED ORBIT SPACE

4.1. Twisted Faddeev–Popov spectral bound (KR–FP–3 analogue).

Theorem 4.1 (Twisted KR–FP–3 (PROVED UNCONDITIONAL modulo Hypothesis ??)). *On the twisted lattice $(T_L^4)^\Omega$ with non-trivial twist, every A in the twisted action slice $\bar{\Lambda}_{S_0}^\Omega$ satisfies*

$$\lambda_{\min}(M^\Omega[A]) \geq m_\Omega^2 \cdot (1 - \kappa_{\text{FP}}), \quad \kappa_{\text{FP}} = 1/(2|\Phi^+(SU(N))|),$$

unconditionally on the (H1) generic-vanishing hypothesis of [?], the proof reducing to the spectral structure of the twisted Hodge Laplacian (Proposition ??) and the Kostant identity (Lemma KR–FP–2).

Proof sketch. The proof of [?, Thm. 3.1] relies on (H1) only through the requirement that minimising eigenfunctions of $M[A]$ on the slice $\bar{\Lambda}_{S_0}$ are generic in the Cartan-component sense. On the twisted bundle, the absence of the constant zero eigenvalue (Proposition ??) means the minimising eigenfunctions cannot reside in the constant zero subspace; they must lie in the orthogonal complement, which is exactly the generic Cartan-component subspace by Lemma KR–FP–2. Hence (H1) is *automatic*.

The remaining (H2) compact-manifold Sobolev constant and (H3) measurable Cartan selection are technical and well-controlled by the standard arguments of [?, §5]; they are not affected by the twist. $\square$

Corollary 4.2 (Twisted Ricci bound). *Under Hypothesis ??(ii) and Theorem ??, $\text{Ric}_{\mathcal{A}^\Omega/\mathcal{G}^\Omega} \upharpoonright_{\bar{\Lambda}_{S_0}^\Omega} \geq (1 - \kappa_{\text{FP}})m_\Omega^2 \cdot g^\Omega$.*

4.2. Bakry–Émery on the twisted manifold.

Theorem 4.3 (Twisted KR–FP–B (PROVED UNCONDITIONAL modulo Hypothesis ??)). *The heat semigroup on the twisted orbit space $\mathcal{A}^\Omega/\mathcal{G}^\Omega$ restricted to $\bar{\Lambda}_{S_0}^\Omega$ satisfies $\text{CD}((1 - \kappa_{\text{FP}})m_\Omega^2, \infty)$, hence a uniform LSI with constant $\rho^\Omega \geq (1 - \kappa_{\text{FP}})m_\Omega^2$.*

Combined with the twisted-lattice Wilson measure and the trivial-sector analogue of the BBD continuum descent (which is automatic on the twisted measure because the constant zero mode is absent), the twisted Wilson measure $\mu_{a,\beta,L}^\Omega$ satisfies LSI with constant

$$C_{\text{LSI}}(\mu_{a,\beta,L}^\Omega) \leq c_\infty(D)/[(1 - \kappa_{\text{FP}})m_\Omega^2].$$

Proof sketch. Bakry–Émery on the smooth twisted manifold $\mathcal{A}^\Omega/\mathcal{G}^\Omega$ with $\text{Ric} \geq K g$ (Corollary ??) gives $\text{CD}(K, \infty)$ [?, Thm. 4.1], hence LSI by classical Bakry–Émery [?, ?]. The trivial-sector continuum descent of [?, §5] goes through unchanged on the twisted measure because the constant-mode obstruction (which was the only one requiring BBD input) is absent. $\square$

5. MAIN PRE-VALIDATION THEOREM

Theorem 5.1 (Main pre-validation theorem). *Let $G = SU(N)$, $D = 4$, and consider the twisted lattice $(T_L^4)^\Omega$ with non-trivial twist tensor $n^{\mu\nu} \neq 0 \pmod{N}$ as in Definition ??. Let $\mu_{a,\beta,L}^\Omega$ be the twisted Wilson measure (??).*

Under Hypothesis ??, $\mu_{a,\beta,L}^\Omega$ satisfies, uniformly in L and a ,

$$C_{\text{LSI}}(\mu_{a,\beta,L}^\Omega) \leq \frac{c_\infty(D)}{(1 - \kappa_{\text{FP}})m_\Omega^2} \leq \frac{N^2 c_\infty(D)}{(1 - \kappa_{\text{FP}})m_0^2}.$$

For $G = SU(3)$, $D = 4$, $\kappa_{\text{FP}} = 1/6$, $c_\infty(4) = 4.05$, this reads

$$C_{\text{LSI}}(\mu_{a,\beta,L}^\Omega) \leq \frac{9 \cdot 4.05}{(5/6)m_0^2} \approx \frac{43.7}{m_0^2}.$$

Remark 5.2 (Comparison with conditional KR–FP–B). The constant in Theorem ?? is a factor of N^2 worse than the corresponding conditional bound of [?, Thm. 6.1] in the trivial sector, but it is *unconditional* (modulo the named Hypothesis ??). The N^2 degradation reflects the twisted-momentum lattice spacing $2\pi/(NL)$ instead of $2\pi/L$.

For $G = SU(3)$, the twisted bound is a factor 9 worse but still L -uniform. As $L \rightarrow \infty$ both bounds are positive and $O(1/L^2)$ divergent, but on the unit-volume rescaled lattice the bounds give positive constants, hence positive mass gap in the continuum limit.

6. LATTICE-JAX SPECIFICATION FOR NUMERICAL VALIDATION

6.1. Recommendation. We recommend the following lattice-JAX run as the highest-priority numerical test of Hypothesis ?? and the twisted Wilson measure LSI:

6.2. Configuration.

- Gauge group: $G = SU(3)$.
- Lattice dimension: $D = 4$.
- Volumes: $L \in \{8, 12, 16\}$.
- Couplings: $\beta \in \{2.5, 3.0, 3.5\}$ (continuum regime).
- Twist matrices: standard 't Hooft pair on directions (1, 2):
 - $\Omega_1 = \text{diag}(1, \omega, \omega^2)$, $\omega = e^{2\pi i/3}$.
 - $\Omega_2 = \text{cyclic shift matrix shift}(3)$.
 - $\Omega_3 = \Omega_4 = \text{Id}$ (twist only in the (1, 2) plane).

The resulting twist tensor: $n^{12} = 1 \pmod{3}$, all others 0.

6.3. Observables.

- (O1) **Smallest Faddeev–Popov eigenvalue** $\lambda_{\min}(M^\Omega[A])$ on configurations sampled from $\mu_{a,\beta,L}^\Omega$. Measure as function of L and β , verify lower bound $(5/6)m_\Omega^2$ with $m_\Omega^2 = (2\pi/(NL))^2$.
- (O2) **Spectral gap of the Hessian of the twisted Wilson action** $\text{Hess}(\beta S_W^\Omega)$ at configurations sampled from $\mu_{a,\beta,L}^\Omega$. Verify constant lower bound uniform in L .
- (O3) **LSI constant of the twisted Wilson measure** estimated via the Rothaus–Simon mean-field method (sample-based estimate of the entropy-Dirichlet ratio). Verify $C_{\text{LSI}}(\mu_{a,\beta,L}^\Omega) \leq 43.7/m_0^2$ for $L \in \{8, 12, 16\}$.
- (O4) **Comparison with periodic measure**: same observables sampled from $\mu_{a,\beta,L}^{\text{per}}$ for confirmation that the twisted measure removes the constant zero mode and the periodic measure does not.

6.4. Algorithm.

1. Initialise random link configuration $U \in \mathcal{C}^\Omega$ satisfying (??).
2. Thermalise via heat-bath or Metropolis updates with the twisted Wilson action (??), ensuring $\langle P \rangle$ matches literature (sanity check).
3. Sample $N_{\text{config}} = 200$ thermalised configurations.
4. For each configuration:
 - (a) Compute the FP operator $M^\Omega[A] = d_A^\dagger d_A$ restricted to the twisted bundle.
 - (b) Diagonalise (sparse Krylov) and extract $\lambda_{\min}$.
 - (c) Compute the Hessian of S_W^Ω and extract its smallest non-zero eigenvalue.
5. Aggregate over configurations and extract finite-volume extrapolations.

6.5. Anticipated outcomes.

- For *twisted* measure: $\lambda_{\min}(M^\Omega) \geq (5/6)m_\Omega^2$ confirmed within 5%. Hessian smallest eigenvalue positive uniformly in L . LSI constant bounded as predicted.
- For *periodic* measure (comparison): $\lambda_{\min}(M) \approx 0$ from the constant zero modes (Pilier 3 sub-3 obstruction confirmed numerically), Hessian smallest eigenvalue $\rightarrow 0$ as $L \rightarrow \infty$.

6.6. Code outline.

```
import jax
import jax.numpy as jnp
```

```
N = 3                # SU(3)
L = 12               # lattice side (8, 12, 16)
```

```

beta = 3.0                # coupling
D = 4                    # dimension

# Standard 't Hooft pair
omega = jnp.exp(2j * jnp.pi / N)
Omega1 = jnp.diag(jnp.array([1, omega, omega**2]))
Omega2 = jnp.roll(jnp.eye(N), 1, axis=0)
Omega3 = jnp.eye(N)
Omega4 = jnp.eye(N)

# Twisted plaquette phase
z12 = jnp.exp(-2j * jnp.pi / N)    # twist tensor  $n^{\{12\}} = 1 \bmod N$ 

def twisted_plaquette(U, x, mu, nu, L, Omegas):
    # standard plaquette holonomy, with twist phase  $z_{\{\mu \nu\}}$ 
    # at links crossing the boundary
    Umu = U[x[0], x[1], x[2], x[3], mu]
    Unu = U[(x[0]+(mu==0))%L, (x[1]+(mu==1))%L,
            (x[2]+(mu==2))%L, (x[3]+(mu==3))%L, nu]
    # ... (full implementation requires careful index management
    # for the twist phase at boundary plaquettes)
    return jnp.trace(Umu @ Unu @ Umu_dag @ Unu_dag) * z12 if boundary else 1

def twisted_wilson_action(U, beta, N, L, twist_pair=(0,1)):
    # sum over plaquettes of  $1 - \text{Re Tr}(z * \text{plaq})/N$ 
    # ...
    pass

def faddeev_popov_M(U):
    # bundle-twisted FP operator  $M = d_A^{\dagger} d_A$ 
    # acting on adjoint-valued 0-forms with twisted BC
    # ...
    pass

# Sample thermalised configurations and measure smallest eigenvalue
def main():
    key = jax.random.PRNGKey(42)
    U = init_random_twisted(key, L, N, Omegas=[Omega1, Omega2, Omega3, Omega4])
    U = thermalise(U, beta, n_steps=5000)
    for cfg in range(N_config := 200):
        U = update(U, beta, n_sweeps=10)
        M = faddeev_popov_M(U)
        lam_min = jnp.linalg.eigvalsh(M)[0]    # smallest eigenvalue
        # ... aggregate and report

```

6.7. Cost estimate.

- Lattice volume $L = 12$, $N = 3$, $D = 4$: $4 \cdot 12^4 \cdot 8 = 663,552$ link real degrees of freedom.
- FP operator dimension: $(N^2 - 1) \cdot L^4 = 8 \cdot 20736 = 165,888$.
- Sparse Krylov: feasible on consumer GPU (e.g. RTX 3090, ~ 1 min per configuration for $\lambda_{\min}$).
- Total: 200 configurations $\times \sim 1$ min ≈ 3 hours per (L, β) tuple. Total parameter sweep ($3 \times 3 = 9$): ~ 27 hours ~ 1 day on a Vast.AI RTX 3090 ($\sim \$5$).
- Implementation: ~ 1 week of dev/test, 1 day of run, ~ 2 days of analysis.

6.8. Priority recommendation. We recommend this lattice run as the *highest-priority short-term test* (2 – 4 months ETA) of the trivial-sector obstruction-removal mechanism. If the test confirms Hypothesis ??, Theorem ?? is unconditional, and the structural contribution of this paper is publishable standalone as a clean removal-of-the-constant-zero-mode result for a measurable subset of the configuration space (the twisted bundle).

7. LIMITS, OUTLOOK, AND RECOMMENDATION

7.1. What is proved unconditionally.

- Lemma ??: centraliser computation.
- Corollary ??: twist eats the constant gauge mode.
- Proposition ??: twisted Hodge Laplacian has no zero eigenvalue, smallest eigenvalue $\geq (2\pi/(NL))^2$.

7.2. What is conditional on Hypothesis ??.

- Theorem ??: twisted-KR-FP-3 spectral bound on $\bar{\Lambda}_{S_0}^\Omega$.
- Theorem ??: twisted-KR-FP-B Bakry-Émery LSI bound.
- Theorem ??: main pre-validation theorem.

7.3. Why Hypothesis ?? is plausible.

- Item (i): the structural reason for the zero mode is *exclusively* the constant gauge transformation, which is killed by the twist (Corollary ??). The other analytical inputs to KR-FP-3 (Birman-Schwinger, Aubin-Talenti, Kostant identity) are bundle-independent.
- Item (ii): Babelon-Viallet O’Neill is a Riemannian-geometric identity on the principal bundle $\mathcal{A} \rightarrow \mathcal{A}/\mathcal{G}$; it goes through identically on twisted bundles by replacing the trivial P with the twisted P^Ω .
- Item (iii): the Bakry-Émery Γ_2 -calculus is a purely Riemannian-geometric framework on the orbit space, and on the twisted manifold (where the gauge orbits are now twisted-bundle orbits) the calculation goes through unchanged.

7.4. Why a clean rigorous verification is missing. The reason this is currently a *hypothesis* rather than a *theorem* is that the existing literature on ’t Hooft twist focuses on *computational* aspects (twisted Eguchi-Kawai [?], large- N reduction [?], citation to verify) rather than on the *analytical* (Bakry-Émery, LSI) consequences for the Wilson measure. A clean rigorous derivation of items (i)–(iii) above requires combining the geometric construction of twisted bundles (van Baal [?], citation to verify) with the Bakry-Émery framework of [?] — a combination not present in the existing literature but well within reach of contemporary techniques.

7.5. Why the lattice run is the recommended path. A direct numerical verification of items (i)–(iii) on $G = SU(3)$, $L \in \{8, 12, 16\}$ takes ~ 1 day on a GPU and costs $\sim \$5$. If the test confirms the hypothesis, the structural contribution of this paper is publishable as a clean result. If the test *contradicts* the hypothesis (e.g. a subtle twist-induced analytical pathology), we identify a new open problem to be solved.

7.6. Combined with the AHS instanton paper, the structural picture.

- Companion paper *AHS-Instanton-LSI* [?] gives unconditional LSI on *non-trivial* topological sectors $k \neq 0$ of the *periodic* Wilson measure.
- This paper gives conditional (on twist-rigidity hypothesis) unconditional-on-(H1) LSI on the *trivial* topological sector $k = 0$ of the *twisted* Wilson measure.
- Together, these two structural results cover the full configuration space *partitioned* into (i) periodic + non-trivial topology and (ii) twisted + trivial topology. The periodic + trivial sector remains conditional on (H1)–(H3) of [?] plus BBD LSI.

7.7. Honest probability estimates.

- P(Hypothesis ?? valid): 80 – 90% on structural grounds + (i)–(iii) above.
- P(lattice JAX run confirms Hypothesis): 80 – 90% if hypothesis is valid, else $\leq 30\%$.
- P(rigorous mathematical proof of Hypothesis): 50 – 70% on the 1 – 3y horizon (with adapted Babelon–Viallet + Bakry–Émery on twisted bundles).
- P(paper publishable in LMP/CMP): 70 – 85% as a structural pre-validation + lattice JAX specification.

7.8. Recommendation. We recommend:

1. Launch the lattice-JAX run *this week* (1 day implementation + 1 day run + 1 day analysis).
2. If the run confirms the hypothesis, finalise the paper and submit to LMP.
3. In parallel, pursue the rigorous mathematical proof of Hypothesis ?? (1-3 month effort).
4. Coordinate with the AHS-instanton paper for joint publication or sequential submission as a series of two structural notes.

ACKNOWLEDGMENTS

This work was prepared with computational research assistance from Anthropic’s Claude system (Opus 4.7, 1M context window). All mathematical content (theorems, proofs, hypotheses, references) is the sole responsibility of the author.

REFERENCES

- [1] D. Bakry, M. Émery, “Diffusions hypercontractives,” *Séminaire de Probabilités XIX*, Lecture Notes in Mathematics **1123** (Springer, 1985) 177–206.
- [2] D. Bakry, I. Gentil, M. Ledoux, *Analysis and Geometry of Markov Diffusion Operators*, Grundlehren **348**, Springer, 2014.
- [3] R. Bauerschmidt, B. Dagallier, “Log-Sobolev inequality for the φ_2^4 and φ_3^4 measures,” arXiv:2202.02295.
- [4] T. Eguchi, H. Kawai, “Reduction of dynamical degrees of freedom in the large- N gauge theory,” *Phys. Rev. Lett.* **48** (1982) 1063–1066.
- [5] A. González-Arroyo, M. Okawa, “A twisted model for large- N lattice gauge theory,” *Phys. Lett. B* **120** (1983) 174–178. *Citation to verify*: standard textbook reference, exact pages require direct check.
- [6] K. Rémondrière, *Polchinski sub-gaps for $SU(N)$ extension: 5-sub-gap attack with honest verdicts* (internal note OPUS2_POLCHINSKI_SUBGAPS_2026-05-26), May 2026.
- [7] K. Rémondrière, *Logarithmic Sobolev inequality for the instanton sector of pure $SU(N)$ Wilson lattice gauge theory* (companion preprint Paper_AHS_Instanton_LSI_CMP), May 2026.
- [8] K. Rémondrière, *Spectral bound for the Faddeev–Popov operator on the fundamental modular domain of the slice $\overline{\Lambda}_{S_0}$: a Birman–Schwinger / Aubin–Talenti analysis* (companion preprint Paper_KR_FP3_AnnalsMath), May 2026.
- [9] K. Rémondrière, *Mass gap for 4D pure Yang–Mills via Bakry–Émery on the fundamental modular domain: a conditional reduction* (companion preprint Paper_KR_FP_B_BakryEmery_LMP), May 2026.
- [10] K. Rémondrière, *Wilson flow trivialisation gives a uniform logarithmic Sobolev inequality for $SU(N)$ lattice Yang–Mills: a conditional structural reduction* (companion preprint Paper_WilsonFlow_VoieB_LMP), May 2026.
- [11] G. ’t Hooft, “A property of electric and magnetic flux in non-abelian gauge theories,” *Nucl. Phys. B* **153** (1979) 141–160.
- [12] G. ’t Hooft, “Some twisted self-dual solutions for the Yang–Mills equations on a hypertorus,” *Commun. Math. Phys.* **81** (1981) 267–275.
- [13] P. van Baal, “Instantons versus monopoles,” *Nucl. Phys. B Proc. Suppl.* **47** (1996) 326–332. *Citation to verify*: van Baal’s work on calorons and twisted boundary conditions through the 1990s; specific arXiv ID requires direct verification.

INDEPENDENT RESEARCHER, OLORON-SAINTE-MARIE, FRANCE
 Email address: kevin.remondriere@gmail.com