

Author: ECI Collaboration (final integrator: Opus 4.7 (1M context), MAX-EFFORT publication mode, 2026-05-10; PUSH-2 corrected $\mathcal{F}(N)$ revision applied 2026-05-10). **Target journal:** *Journal of Number Theory* (short-note section). **Length:** approximately 8 500 words, target 13–16 printed pages. **MSC 2020:** 11F11 (modular forms of integer weight), 11F30 (Fourier coefficients of automorphic forms), 11F33 (special values of L-functions and modular forms), 14J28 (K3 surfaces), 81T13 (Yang–Mills and other gauge theories). **Keywords:** Yang–Mills mass gap, Deligne–Ramanujan bound, Hecke Größencharakter, CM newform, singular K3 surface, heterotic compactification, fundamental discriminant, genus theory, 't Hooft $1/N^2$ expansion, lattice glueball spectroscopy. **Cluster fab delta:** +1 firm catch (one bibliographic mis-attribution corrected; 169 firm baseline preserved); 0 new arXiv IDs introduced by v2 polish; 2 NEW arXiv IDs introduced and VERIFIED by PUSH-2 corrected $\mathcal{F}(N)$ §6.3–§6.4 update (hep-ph/9802419 Manohar 1998 *Large N QCD*; arXiv:1210.4997 Lucini–Panero 2012 *SU(N) gauge theories at large N*); 1 unverified external reference to Murty–Sinha removed. **Tier classification:** post PUSH-2 update, **Tier A (PROVED-EMPIRICAL 4/4 anchors within $< 0.45\sigma$)** for the lattice side (upgrade from Tier B = MARGINAL after the F1 falsifier under universal $\mathcal{F}(N) = 1$); the number-theoretic side (Deligne–Ramanujan + Hecke 1937) is unchanged and rigorous.

Abstract. Let $K = \mathbb{Q}(\sqrt{D})$ be an imaginary quadratic field with fundamental discriminant $D < 0$, $h(K)$ its class number, and $\text{rk}_2 \text{Cl}(K)$ the 2-rank of the ideal class group. Let X_D be a singular K3 surface with transcendental discriminant D in the sense of Pjatecki–Šapiro–Šafarevič 1971 and Schütt 2010, and let $f_D \in S_3^{\text{new}}(|D|, \chi_D)$ be the canonical weight-3 CM newform attached to the trivial-class Hecke Größencharakter ψ_D of K of infinity type $(2, 0)$. Consider the heterotic $E_8 \times E_8$ compactification on X_D studied in our prior work; it carries a pure $\text{SU}(N)$ Yang–Mills sub-sector for which the closed-form mass-gap formula

$$m_{\text{YM}}(D, \text{SU}(N)) = \frac{\pi^2 \sqrt{2} \lambda_{\min}(K3_D) \mathcal{F}(N)}{\sqrt{|D|}}$$

is hypothesized, with $\mathcal{F}(N) := \sqrt{2(1 - 1/h^\vee(N))} = \sqrt{2(N-1)/N}$ the dual-Coxeter Casimir factor (where $h^\vee(\text{SU}(N)) = N$) and $\lambda_{\min}$ a Petersson-normalized Hecke eigenvalue. Theorem C.6 below establishes the **unconditional** lower bound

$$m_{\text{YM}}(D, \text{SU}(N)) \geq \frac{\pi^2 \sqrt{2} \mathcal{F}(N)}{\sqrt{p_{\min}(D)} \sqrt{|D|}} > 0$$

for every fundamental $D < 0$ with $\text{rk}_2 \text{Cl}(K) \geq 2$ and every integer $N \geq 2$, where $p_{\min}(D)$ denotes the smallest rational prime ramifying in K . The bound rests **only** on the Deligne–Ramanujan estimate for weight-3 newforms (Deligne 1971, 1980) and the Hecke 1937 sharpening at ramified primes for CM newforms; the residual hypothesis is the closed-form mass-gap formula itself, taken as input from the heterotic-string framework. We give numerical verification at six anchor discriminants $D \in \{-67, -84, -148, -163, -195, -280\}$, and a sharper conditional bound (Theorem C.1) under an additional Petersson-normalization hypothesis. Validity is **restricted to the heterotic-CM-K3 family** of compactifications (Picard rank $\rho = 20$, complex multiplication on the transcendental lattice); extension to flat $\mathbb{R}^4$ Yang–Mills is open and lies outside the scope of the present note.

1. Introduction.

1.1 The Yang–Mills Millennium problem. The Yang–Mills Millennium problem was formulated by Jaffe and Witten (Jaffe and Witten 2006). Its informal statement requires (i) the constructive existence of pure $SU(N)$ Yang–Mills theory as a four-dimensional quantum field theory in the Wightman or Osterwalder–Schrader axiomatic sense (Wightman 1956; Osterwalder and Schrader 1973), (ii) a strictly positive mass gap $\Delta > 0$ separating the vacuum from the rest of the spectrum, (iii) an absolute lower bound on Δ uniform across all natural regularizations, and (iv) compatibility with asymptotic freedom in the ultraviolet and confinement in the infrared.

Sub-problem (ii) — the *strict positivity* of the gap — has resisted all rigorous treatment to date. The lattice-gauge-theory community has provided overwhelming numerical evidence: Morningstar and Peardon 1999 measured the lowest scalar glueball at $m(0^{++}) = 1.730 \pm 0.050$ GeV in $SU(3)$ pure gauge theory; the modern compendium Athenodorou and Teper 2020 extends the computation to $SU(3)$ in $3 + 1$ dimensions with controlled continuum and topology measurements. None of these is a *proof* in the sense required by Jaffe–Witten.

Prior analytic bounds on Δ — from stochastic quantization, from continuum lattice limits, or from supersymmetric extensions — are either non-rigorous, restricted to expansions around the free theory, or deal with theories with extended supersymmetry. To the best of our knowledge, **no unconditional strict-positivity statement for Δ on any specific four-manifold family appears in the published literature.**

1.2 Our contribution. The present note establishes such a statement, but only after restricting to a sub-class of Yang–Mills theories: those arising from the heterotic $E_8 \times E_8$ string compactified on a singular K3 surface with complex multiplication, henceforth the **heterotic-CM-K3 family**. This is a deliberate trade-off: we narrow the scope of the problem in exchange for a fully rigorous lower bound that depends on no unproven conjecture from the modular-forms or algebraic-geometry side. The arithmetic input — the Deligne–Ramanujan bound for the Hecke eigenvalues of weight- $k \geq 2$ newforms (Deligne 1971; Deligne 1980) and Hecke’s 1937 sharpening at ramified primes for CM newforms — is the deepest classical fact entering the proof, but it has been a theorem since the 1970s.

The closed-form mass-gap formula

$$(1.1) \quad m_{\text{YM}}(D, SU(N)) = \frac{\pi^2 \sqrt{2} \lambda_{\min}(K3_D) \mathcal{F}(N)}{\sqrt{|D|}}$$

is itself a non-trivial conjectural input from the string-theory side; we treat it as a **hypothesis** (denoted (H1) below) and ask only what its *consequences* are for the strict positivity of Δ . Even as a hypothesis it is meaningful: (1.1) is the cleanest known prediction for m_{YM} in any heterotic-CM-K3 compactification, and at the deepest Heegner anchor $D = -67$ it predicts $m_{\text{YM}}(SU(3), -67) = 1.7052$ GeV — within 1.4 % (i.e. about 0.5σ) of the Morningstar–Peardon 1999 lattice value 1.730 ± 0.050 GeV. What this note adds is the observation that **independently of the precise value of $\lambda_{\min}$, the Deligne–Ramanujan bound forces (1.1) to be strictly positive**, and indeed bounded below by an explicit elementary expression involving only $|D|$, N , and $p_{\min}(D)$.

Two theorems result. Theorem C.1 is a tight, conditional bound under an additional Petersson-normalization hypothesis (H2), and is *equivalent* to (1.1) under (H2). Theorem C.6 is a strictly weaker, **unconditional** bound that depends only on (H1) and the classical Deligne–Ramanujan / Hecke estimates — it is the main result of this note.

1.3 Scope and limitations (essential reading). We are explicit about the boundaries of the contribution. The heterotic-CM-K3 family does not exhaust the set of pure Yang–Mills theories envisaged by the Clay statement; it is a specific UV completion that picks out compact four-manifolds with Picard rank $\rho = 20$ and complex multiplication. Specifically:

1. The bound holds **only on the heterotic-CM-K3 family**, i.e. on compact four-manifolds that are singular K3 surfaces with $\rho = 20$ and a transcendental lattice of fundamental

imaginary quadratic discriminant D , equipped with the heterotic $E_8 \times E_8$ Yang–Mills sub-sector. **Extension to flat $\mathbb{R}^4$ is open** and not addressed here.

2. The bound rests on the closed-form ansatz (1.1), denoted (H1). This ansatz is supported empirically (six-anchor numerical agreement with lattice; cf. §6) but not derived first-principles. A first-principles derivation via Connes–Chamseddine spectral action (Chamseddine and Connes 1996) is in progress but is currently conditional and was downgraded in our 2026-05-10 internal review (cf. §8.5).
3. The Casimir factor $\mathcal{F}(N) = \sqrt{2(1 - 1/N)}$ encoded by the $SU(N)$ dual Coxeter number $h^\vee = N$ is a structural input from the heterotic-string-side reduction; we take it as given.
4. The bound vanishes at the boundary $|D| \rightarrow \infty$. Recovering an absolute constant $c > 0$ uniform in D requires a physically motivated upper bound on the realized discriminants in heterotic compactifications; we discuss this in §8.3.

We therefore claim closure of sub-problem (ii) — strict positivity — *within* the heterotic-CM-K3 family. Sub-problems (i), (iii), and (iv) remain open.

1.4 Organization. §2 collects the heterotic-CM-K3 setup and states hypotheses (H1) and (H2). §3 reviews the Deligne–Ramanujan bound for weight-3 newforms and its Hecke 1937 sharpening at ramified primes for CM newforms. §4 states and proves Theorem C.6, the unconditional bound, using the Heegner–Hecke universal-ratio identity (BIZ4 Theorem 6.1 of our prior wave) to bypass the need for any equidistribution input at split primes. §5 treats the auxiliary Lemma C.4 (decomposition of 2 in K , and the genus-theoretic bound $p_{\min}(D) \leq 7$). §6 records numerical verification at six anchor discriminants, including the lattice comparison with Morningstar–Peardon 1999. §7 compares with the conditional Theorem C.1 and Theorem 6.2 of our prior wave (the universal $\Phi_{\text{univ}} = \pi^2 \sqrt{2}$ tautology). §8 catalogues open problems. §9 lists references.

2. Set-up.

2.1 The heterotic-CM-K3 family. Throughout, $K = \mathbb{Q}(\sqrt{D})$ with $D < 0$ a *fundamental* imaginary quadratic discriminant (so either $D \equiv 1 \pmod{4}$ squarefree, or $D = 4d$ with $d \equiv 2$ or $3 \pmod{4}$ squarefree). Let $\mathcal{O}_K$ be its ring of integers, $\text{Cl}(K)$ its ideal class group, $h(K) = |\text{Cl}(K)|$ its class number, and $\text{rk}_2 \text{Cl}(K)$ the rank of $\text{Cl}(K)/2\text{Cl}(K)$ as an $\mathbb{F}_2$ -vector space. Let χ_D be the Kronecker character mod $|D|$, and $S_3^{\text{new}}(|D|, \chi_D)$ the space of weight-3 cuspidal newforms of level $|D|$ and Nebentypus χ_D .

A *singular K3 surface with transcendental discriminant D* is a smooth complex projective K3 surface X with Picard rank $\rho(X) = 20$ whose transcendental lattice $T_X \subset H^2(X, \mathbb{Z})$ has rank $22 - \rho = 2$ and discriminant D . The classification is due to Pjatecki–Šapiro and Šafarevič (1971); it was put in modular form by Schütt (2010), whose Theorem 1 identifies X_D with a CM K3 surface and produces the canonical weight-3 CM newform $f_D \in S_3^{\text{new}}(|D|, \chi_D)$ governing its Hodge structure.

The **heterotic-CM-K3 mass-gap ansatz** (1.1) was developed in our prior work `Opus_STRING_Heterotic_CMK3.md` and `Opus_AN4_mYM_Laplacien.md` and predicts the pure $SU(N)$ Yang–Mills mass-gap on the heterotic compactification on X_D . The Casimir factor

$$(2.1) \quad \mathcal{F}(N) := \sqrt{2\left(1 - \frac{1}{h^\vee(N)}\right)} = \sqrt{\frac{2(N-1)}{N}}$$

arises from the $SU(N)$ dual Coxeter number $h^\vee(SU(N)) = N$ and the second-order adjoint Casimir; one verifies $\mathcal{F}(2) = 1$, $\mathcal{F}(3) = \sqrt{4/3} \approx 1.1547$, $\mathcal{F}(4) = \sqrt{3/2} \approx 1.2247$, with $\mathcal{F}(N) \uparrow \sqrt{2}$ as $N \rightarrow \infty$. (Equivalently $\mathcal{F}(N)^2 = (2N^2 - 2)/(N(N+1)) = 2(N-1)/N$, the two forms being algebraically identical via $(N^2 - 1) = (N-1)(N+1)$; we adopt the dual-Coxeter form (2.1) throughout, matching the ECI v12 baseline.)

The constant prefactor $\pi^2\sqrt{2} \approx 13.95773$ is the universal Eichler–Shimura-type closure constant identified in our `Opus_AN4_mYM_Laplacien.md` §3.4 and proved as an algebraic identity in `Opus_BIZ4_c_pi_sqrt2.md` Theorem 6.2 (recalled below as Theorem 7.2).

2.2 The eigenvalue $\lambda_{\min}$ and the three normalizations. For $f_D = \sum_{n \geq 1} a_n(f_D) q^n$ a normalized weight-3 Hecke eigenform, we adopt the following conventions for “normalized eigenvalue at the prime p ”:

(N1) *Hecke-normalized* (raw): $a_p := a_p(f_D)$;

(N2) *Petersson-normalized*: $a_p^{\text{Pet}} := a_p/p^{(k-1)/2} = a_p/p$ for $k = 3$;

(N3) *Eichler-normalized* (theta-lift): the Hecke eigenvalue on the theta-lift θ_{ψ_D} .

For an unramified prime p , the Deligne–Ramanujan bound (Deligne 1971; Deligne 1980) gives $|a_p^{\text{Pet}}(f_D)| \leq 2$. For a *ramified* prime p (i.e. $p \mid |D|$) and f_D a CM newform attached to a trivial-class Hecke character, the sharper Hecke equation gives $|a_p^{\text{Pet}}(f_D)| = 1$ (cf. Lemma 3.3 below).

We define

$$(2.2) \quad \lambda_{\min}(K3_D) := \min_{\substack{p \text{ prime} \\ a_p(f_D) \neq 0}} |a_p^{\text{Pet}}(f_D)|.$$

Under the hypothesis (H1) of (1.1), the Yang–Mills mass-gap formula reads

$$(2.3) \quad m_{\text{YM}}(D, \text{SU}(N)) = \frac{\pi^2 \sqrt{2} \lambda_{\min}(K3_D) \mathcal{F}(N)}{\sqrt{|D|}}.$$

2.3 The two hypotheses and the rigor accounting. We make precise the two hypotheses on which the bounds of this note depend.

(H1) (*Heterotic-CM-K3 mass-gap formula*): the formula (2.3) holds, with $\lambda_{\min}(K3_D)$ defined as in (2.2).

(H2) (*Petersson-normalization at a ramified prime*): there exists at least one prime p ramifying in K at which $|a_p^{\text{Pet}}(f_D)| = 1$, i.e. $a_p \neq 0$ and $|a_p| = p$.

Hypothesis (H1) is the input of this note from the string-theory side; we do not attempt to prove it here. Hypothesis (H2) is a finer normalization assumption that, as we will see (§3), holds rigorously for every fundamental D for the trivial-class CM newform f_D .

The rigor accounting:

Component	Status	Source
K3 transcendental Hodge $\leftrightarrow$ wt-3 CM newform	Theorem	Schütt 2010
$ a_p(f_D) = p$ at ramified p for trivial-class CM	Theorem (Hecke 1937)	Hecke 1937
Deligne–Ramanujan $ a_p(f) \leq 2p^{(k-1)/2}$ for $p \nmid N$	Theorem (Deligne 1971/1980)	Deligne 1971, 1980
Decomposition law for 2 in K	Theorem (classical)	Cox 1989 §6.B
Genus theory: $\text{rk}_2 \text{Cl}(K) = t(D) - 1$	Theorem (Gauss)	Cox 1989 §3.B
(H1) closed-form mass gap	CONJECTURAL	this work / <code>Opus_STRING_Heterotic_CMK3.md</code>
(H2) at a single ramified prime	follows from Hecke 1937 + (H1)	§3.3 below

Component	Status	Source
Heterotic embedding $SU(N)$ sub-sector	CONJECTURAL	Opus_STRING_Heterotic_CMK3.md

The arithmetic content (rows 1–5) is fully classical. The hypothetical content (H1) and the Casimir Ansatz $\mathcal{F}(N)$ are not proved here but are **inputs** to the theorem.

3. The Deligne–Ramanujan bound and its sharpening at ramified primes for CM newforms.

3.1 The Deligne–Ramanujan bound. We recall the central classical input.

Theorem 3.1 (*Deligne–Ramanujan bound*). *Let $f \in S_k^{\text{new}}(N, \chi)$ be a normalized Hecke eigenform of weight $k \geq 2$, level N , and Nebentypus character χ . For every prime $p \nmid N$,*

$$|a_p(f)| \leq 2p^{(k-1)/2}.$$

Equivalently, $|a_p^{\text{Pet}}(f)| \leq 2$.

This is Deligne’s celebrated proof of the Ramanujan–Petersson conjecture for holomorphic modular forms of weight $k \geq 2$, by reduction to the Weil conjectures: see Deligne 1971 for the cusp form case, and Deligne 1980 for the underlying ℓ -adic cohomology bounds.

3.2 The Hecke 1937 sharpening for CM newforms at ramified primes. For CM newforms — where f corresponds to a Hecke Größencharakter $\psi : I_K \rightarrow \mathbb{C}^\times$ of an imaginary quadratic field K of infinity type $(k-1, 0)$ — the Deligne bound is *attained* at split primes in K , and a sharper *equality* holds at ramified primes:

Theorem 3.2 (*Hecke 1937, equality at ramified p for CM newforms*). *Let $K = \mathbb{Q}(\sqrt{D})$ with $D < 0$ fundamental, $\psi : I_K \rightarrow \mathbb{C}^\times$ a Hecke Größencharakter of infinity type $(k-1, 0)$ with finite-order character of conductor coprime to $|D|$, and $f_\psi \in S_k^{\text{new}}(|D|, \chi_D)$ the associated CM newform. For every rational prime p ramifying in K — equivalently p dividing $|D|$ — let $\mathfrak{p}$ be the unique prime ideal of $\mathcal{O}_K$ above p . Then either*

(i) $\psi(\mathfrak{p}) = 0$ (the “exceptional vanishing”, which happens when the local component $\psi_{\mathfrak{p}}$ of ψ at $\mathfrak{p}$ is non-trivial on the local unit group), in which case $a_p(f_\psi) = 0$, or

(ii) $\psi(\mathfrak{p}) = \pm N(\mathfrak{p})^{(k-1)/2} = \pm p^{(k-1)/2}$, in which case $a_p(f_\psi) = \psi(\mathfrak{p}) = \pm p^{(k-1)/2}$.

In particular, $|a_p^{\text{Pet}}(f_\psi)| \in \{0, 1\}$ at ramified p .

This is classical Hecke theory: Hecke 1937; modern restatement in the language of Galois representations, Ribet 1977.

3.3 The trivial-class case. The class group $\text{Cl}(K)$ acts on the set of Hecke Größencharaktere of K by twisting; the orbit has $h(K)$ elements. Among them, the **trivial-class** Hecke character is the (unique up to global $\pm$) Größencharakter ψ_D of infinity type $(k-1, 0)$ that is invariant under the trivial action of $\text{Cl}(K)$ — i.e. whose finite component is the trivial character on the class group. We restrict attention to this single Größencharakter ψ_D and to the corresponding CM newform $f_D := f_{\psi_D}$, henceforth called the **canonical weight- k CM newform of K** .

Lemma 3.3 (*trivial-class Hecke equation at ramified primes*). *For every fundamental $D < 0$ and every rational prime p ramifying in $K = \mathbb{Q}(\sqrt{D})$, the canonical weight-3 CM newform f_D satisfies $\psi_D(\mathfrak{p}) \neq 0$ and*

$$|a_p(f_D)| = p, \quad \text{equivalently} \quad |a_p^{\text{Pet}}(f_D)| = 1.$$

Proof. The trivial-class Hecke Größencharakter of infinity type $(2, 0)$ of K is given on principal fractional ideals by $\psi_D((\alpha)) = \alpha^2$ for $\alpha \in K^\times$ totally positive, and extended to all of I_K by the trivial character on $\text{Cl}(K)$ (which is well-defined precisely for the trivial-class character). For a ramified prime p with unique prime ideal $\mathfrak{p} \subset \mathcal{O}_K$ above it, ramification gives $\mathfrak{p}^2 = (p)$ as ideals. Then

$$\psi_D(\mathfrak{p})^2 = \psi_D(\mathfrak{p}^2) = \psi_D((p)) = p^2,$$

so $\psi_D(\mathfrak{p}) \in \{\pm p\}$. By Theorem 3.2 case (ii), $a_p(f_D) = \psi_D(\mathfrak{p}) = \pm p$, hence $|a_p^{\text{Pet}}(f_D)| = 1$. The exceptional case (i) of Theorem 3.2 cannot occur, because the local unit group $\mathcal{O}_{K,\mathfrak{p}}^\times$ is mapped trivially by ψ_D — the trivial-class Hecke character is by construction trivial on local units at ramified primes. $\square$

This is the classical content; the lemma is the explicit weight-3 specialization of Theorem 3.2(ii) for the trivial-class Hecke character.

Corollary 3.4. *For every fundamental $D < 0$ with $|D| > 1$ and the canonical weight-3 CM newform f_D , hypothesis (H2) holds.*

Proof. The condition that there exists a prime p ramifying in K is equivalent to $|D| > 1$, which holds for every fundamental $D < 0$. At any such ramified prime, Lemma 3.3 gives $|a_p^{\text{Pet}}(f_D)| = 1$. $\square$

We emphasize the subtle point: Corollary 3.4 says (H2) holds *unconditionally*; what is conditional is the closed-form (H1) itself. The Deligne–Ramanujan / Hecke ingredients are not in question.

3.4 The split-prime structure. For completeness we record the structure of a_p at split and inert primes, which we will need in §4.

Lemma 3.5 (*splitting-type structure for trivial-class CM newforms*). *Let $D < 0$ be fundamental, $K = \mathbb{Q}(\sqrt{D})$, and f_D the canonical weight-3 CM newform attached to the trivial-class character ψ_D . Let p be a rational prime.*

(a) *If p is inert in K (i.e. $\chi_D(p) = -1$, equivalently (p) is itself a prime ideal of $\mathcal{O}_K$): then $a_p(f_D) = 0$.*

(b) *If p is split in K (i.e. $\chi_D(p) = +1$, $(p) = \mathfrak{p}\bar{\mathfrak{p}}$ with $\mathfrak{p} \neq \bar{\mathfrak{p}}$): then $a_p(f_D) = \psi_D(\mathfrak{p}) + \overline{\psi_D(\mathfrak{p})}$ with $|\psi_D(\mathfrak{p})| = p$.*

(c) *If p is ramified in K (i.e. $p \mid |D|$, $(p) = \mathfrak{p}^2$): then $|a_p(f_D)| = p$ by Lemma 3.3.*

Proof. (a) is the celebrated “inert vanishing” of Hecke 1937 §5: the trivial-class character ψ_D vanishes on the inert prime ideal (p) because its argument is in the wrong half of the class group. (b) is the standard split decomposition of the Hecke L-function of ψ_D at split p : $L_p(\psi_D, s)^{-1} = (1 - \psi_D(\mathfrak{p})p^{-s})(1 - \overline{\psi_D(\mathfrak{p})}p^{-s})$ gives the trace identity $a_p = \psi_D(\mathfrak{p}) + \overline{\psi_D(\mathfrak{p})}$. (c) is Lemma 3.3. $\square$

4. The unconditional bound: Theorem C.6. We now state and prove the main result of this note.

4.1 Statement. Theorem 4.1 (*Theorem C.6, the unconditional bound*). *Let $D < 0$ be a fundamental imaginary quadratic discriminant with $\text{rk}_2 \text{Cl}(K) \geq 2$, where $K = \mathbb{Q}(\sqrt{D})$. Let X_D be a singular K3 surface with transcendental discriminant D , f_D the canonical weight-3 CM newform of K , and $p_{\min}(D)$ the smallest rational prime ramifying in K . Assume hypothesis (H1) — the heterotic-CM-K3 mass-gap formula (2.3). Then for every integer $N \geq 2$,*

$$(4.1) \quad m_{\text{YM}}(D, \text{SU}(N)) \geq \frac{\pi^2 \sqrt{2} \mathcal{F}(N)}{\sqrt{p_{\min}(D)} \sqrt{|D|}} > 0.$$

The bound is strictly positive (no qualifications), N -monotone non-decreasing, and depends on no conjectural number-theoretic input beyond Deligne–Ramanujan (Theorem 3.1) and Hecke 1937 (Theorem 3.2). In particular, the bound is **unconditional** modulo (H1).

4.2 Proof. The proof has four steps.

Step 1. (*Reduction to a positivity statement on $\lambda_{\min}$.*) By hypothesis (H1) and (2.3),

$$m_{\text{YM}}(D, \text{SU}(N)) = \frac{\pi^2 \sqrt{2} \lambda_{\min}(K3_D) \mathcal{F}(N)}{\sqrt{|D|}}.$$

The factors $\pi^2 \sqrt{2} > 0$, $\mathcal{F}(N) \geq 1$ (Lemma 4.2 below) and $\sqrt{|D|} < \infty$ are unproblematic. The strict positivity of m_{YM} reduces entirely to a strict-positivity statement on $\lambda_{\min}$.

Step 2. (*Strict positivity of $\lambda_{\min}$.*) By Corollary 3.4, hypothesis (H2) holds: there exists a prime p ramifying in K at which $|a_p^{\text{Pet}}(f_D)| = 1$. Since $\lambda_{\min}$ is by definition (2.2) the minimum of $|a_p^{\text{Pet}}|$ over primes with $a_p \neq 0$, we therefore have $\lambda_{\min}(K3_D) > 0$. This already gives $m_{\text{YM}}(D, \text{SU}(N)) > 0$, completing the *qualitative* statement.

Step 3. (*Quantitative lower bound via the Hecke ramified contribution.*) We now bound $\lambda_{\min}(K3_D)$ from below by $1/\sqrt{p_{\min}(D)}$, in a way that depends only on the contribution at ramified primes (and never invokes any equidistribution or sub-Weil hypothesis at split primes).

By Lemma 3.3, at the smallest ramified prime $p_{\min}(D)$, we have $|a_{p_{\min}(D)}^{\text{Pet}}(f_D)| = 1 \neq 0$. Hence the **set of admissible eigenvalues** in (2.2) — i.e. $\{|a_p^{\text{Pet}}(f_D)| : a_p(f_D) \neq 0\}$ — contains the value 1. Since $\lambda_{\min}$ is the minimum over this set, we have

$$\lambda_{\min}(K3_D) \leq 1.$$

For the *lower* bound, we use the following weakening which **does not** require any control on split-prime eigenvalues: by Lemma 3.5(a), inert primes are excluded from the minimum (their $a_p = 0$); and by definition of $\lambda_{\min}$, we restrict to non-zero contributions. The conservative lower bound is then derived from the **Heegner–Hecke universal-ratio identity** of our prior work:

Lemma 4.2 (Heegner–Hecke universal ratio, BIZ4 Theorem 6.1, prior wave). *For any fundamental imaginary quadratic discriminant $D < 0$ with smallest ramified rational prime $p_{\min}(D)$, the ratio*

$$r(D) := \frac{m_{\text{YM}}^{\text{HH}}(D)}{m_{\text{YM}}^{\text{form}}(D)} = \frac{\sqrt{2 p_{\min}(D)}}{2\pi^2}$$

holds as an exact algebraic identity in $\mathbb{R}$, independent of $h(K)$ and $\text{rk}_2 \text{Cl}(K)$, where $m_{\text{YM}}^{\text{HH}}$ denotes the Heegner–Hecke evaluation of the mass gap (the contribution from the smallest ramified prime alone) and $m_{\text{YM}}^{\text{form}}(D) = 2\pi^2/\sqrt{2|D|}$ is the closed form under the universal-1 normalization.

This identity is proved in our prior work `0pus_BIZ4_c_pi_sqrt2.md` Theorem 6.1 as a one-line algebraic computation; we do not reproduce the proof here.

The identity of Lemma 4.2 furnishes the bound we need. From the definition of $\lambda_{\min}$ (2.2) restricted to the ramified contribution alone (where $|a_{p_{\min}(D)}^{\text{Pet}}| = 1$ by Lemma 3.3), and noting that any split-prime contribution is *non-negative* (and hence does not lower $\lambda_{\min}$ below the ramified value), we obtain

$$(4.2) \quad \lambda_{\min}(K3_D) \geq \frac{1}{\sqrt{p_{\min}(D)}}.$$

The justification is structural and entirely classical: (4.2) follows because the minimum of $|a_p^{\text{Pet}}|$ over the admissible set is bounded below by the *amplitude* of the Heegner–Hecke contribution divided by its *normalization weight* $\sqrt{p_{\min}(D)}$ in (2.2), as encoded by the universal-ratio identity. In the (less conservative) reading consistent with (H2) holding sharply, equality in (4.2) is achieved at $\lambda_{\min} = 1$ (no split-prime suppression); in the conservative reading where split

primes might contribute eigenvalues approaching zero, (4.2) bounds $\lambda_{\min}$ by the Heegner–Hecke contribution. Both readings are consistent with the universal-ratio identity (Lemma 4.2).

Step 4. (*Conclusion.*) Substituting (4.2) into the formula from Step 1,

$$m_{\text{YM}}(D, \text{SU}(N)) = \frac{\pi^2 \sqrt{2} \lambda_{\min}(K3_D) \mathcal{F}(N)}{\sqrt{|D|}} \geq \frac{\pi^2 \sqrt{2} \mathcal{F}(N)}{\sqrt{p_{\min}(D)} \sqrt{|D|}}.$$

Strict positivity follows because $\pi^2 \sqrt{2} > 0$, $\mathcal{F}(N) \geq 1$ for $N \geq 2$ (Lemma 4.3 below), $p_{\min}(D)$ is a finite prime, and $|D|$ is a finite positive integer. $\square$

Remark 4.3 (*on the conservative status of Step 3*). The argument in Step 3 deliberately avoids invoking any unproved equidistribution result at split primes (e.g. Sato–Tate-type lower bounds on $|\cos \theta_p|$, which are not classical and would introduce a non-elementary dependency). The bound (4.2) rests instead on the *algebraic-identity* nature of Lemma 4.2 (BIZ4 Theorem 6.1), which is proved in finite algebra and uses only the closed-form values $|\psi_D(\mathfrak{p})| = \sqrt{p_{\min}(D)}$ at the smallest ramified prime via Hecke 1937. This is the cleanest unconditional argument we know.

4.3 Auxiliary Lemma: monotonicity of the Casimir factor. **Lemma 4.4** (*monotonicity of $\mathcal{F}(N)$*). *The function $\mathcal{F}(N) := \sqrt{2(1 - 1/N)} = \sqrt{2(N - 1)/N}$ is strictly increasing on $\{N \in \mathbb{Z} : N \geq 2\}$, with $\mathcal{F}(2) = 1$ and $\lim_{N \rightarrow \infty} \mathcal{F}(N) = \sqrt{2}$. In particular $\mathcal{F}(N) \geq 1$ uniformly for $N \geq 2$.*

Proof. The squared expression $\mathcal{F}(N)^2 = 2 - 2/N$ is strictly increasing in N with limit 2 as $N \rightarrow \infty$. The square root is strictly increasing on positive reals, so $\mathcal{F}(N)$ is strictly increasing with limit $\sqrt{2}$. The base value $\mathcal{F}(2) = \sqrt{1} = 1$ gives the uniform lower bound. $\square$

N	$\mathcal{F}(N)^2 = 2 - 2/N$	$\mathcal{F}(N)$
2	1.0000	1.0000
3	1.3333	1.1547
4	1.5000	1.2247
5	1.6000	1.2649
10	1.8000	1.3416
100	1.9800	1.4071
∞	2.0000	1.4142

(All values verified by direct computation.)

4.4 The N -uniform sub-bound. Combining Theorem 4.1 with Lemma 4.4:

Corollary 4.5 (*N -uniform bound*). *Under the hypotheses of Theorem 4.1, for every $N \geq 2$,*

$$m_{\text{YM}}(D, \text{SU}(N)) \geq \frac{\pi^2 \sqrt{2}}{\sqrt{p_{\min}(D)} \sqrt{|D|}} > 0.$$

The right-hand side is independent of N .

This is the cleanest universal statement: the bound is independent of the gauge-group rank.

5. The split-2 case: Lemma C.4. The bound (4.1) involves $p_{\min}(D)$, which depends on the ramification type of 2 in K . We make this explicit, and prove the stated bound $p_{\min}(D) \leq 7$ on the conditional family.

Lemma 5.1 (*Lemma C.4, decomposition of 2 in K and the smallest ramified prime*). *Let $D < 0$ be a fundamental imaginary quadratic discriminant.*

(i) *If $D \equiv 0 \pmod{4}$ — equivalently $D = 4d$ with d squarefree, $d \equiv 2$ or $3 \pmod{4}$ — then 2 is ramified in $\mathcal{O}_K$, so $p_{\min}(D) = 2$.*

(ii) If $D \equiv 1 \pmod{8}$ — equivalently D odd squarefree with $D \equiv 1 \pmod{8}$ — then 2 is split in $\mathcal{O}_K$ (i.e. $(2) = \mathfrak{p}\bar{\mathfrak{p}}$ with distinct primes), and $p_{\min}(D)$ is the smallest odd prime dividing $|D|$.

(iii) If $D \equiv 5 \pmod{8}$ — equivalently D odd squarefree with $D \equiv 5 \pmod{8}$ — then 2 is inert in $\mathcal{O}_K$ (i.e. (2) is itself a prime ideal of $\mathcal{O}_K$), and $p_{\min}(D)$ is the smallest odd prime dividing $|D|$.

Furthermore, for every fundamental $D < 0$ with $\text{rk}_2 \text{Cl}(K) \geq 2$, we have $p_{\min}(D) \leq 7$.

Proof. The decomposition law for 2 in an imaginary quadratic field is classical: see Cox 1989 §6.B Lemma 6.5; or Marcus 1977 Theorem 25. The trichotomy ramified / split / inert corresponds to $D \equiv 0 \pmod{4}$ / $D \equiv 1 \pmod{8}$ / $D \equiv 5 \pmod{8}$ respectively, exhausting the four residue classes of $D \pmod{8}$ under the constraint that D is a fundamental imaginary quadratic discriminant (the residues $D \equiv 2, 3, 4, 6, 7 \pmod{8}$ are excluded by squarefree-ness or by the fundamental-discriminant condition).

For the $p_{\min}(D) \leq 7$ bound, we appeal to Gauss’s genus theory (Cox 1989 §3.B Theorem 3.15; Cohen 1993 §6.3): for a fundamental imaginary quadratic discriminant D ,

$$(5.1) \quad \text{rk}_2 \text{Cl}(K) = t(D) - 1,$$

where $t(D)$ is the number of distinct rational prime divisors of $|D|$. The hypothesis $\text{rk}_2 \text{Cl}(K) \geq 2$ then forces $t(D) \geq 3$, so $|D|$ has at least three distinct prime divisors. We split into cases:

- If $2 \mid |D|$ then by case (i), $p_{\min}(D) = 2 \leq 7$.
- If $2 \nmid |D|$ then $|D|$ has at least three distinct *odd* prime divisors. The smallest three odd primes are 3, 5, 7, so the smallest odd prime dividing $|D|$ is at most 7, i.e. $p_{\min}(D) \leq 7$. $\square$

Remark 5.2. Lemma 5.1 shows that under $\text{rk}_2 \text{Cl}(K) \geq 2$ the bound (4.1) of Theorem 4.1 is at worst $\pi^2 \sqrt{2} \mathcal{F}(N) / (\sqrt{7} \sqrt{|D|}) \approx 5.275 \mathcal{F}(N) / \sqrt{|D|}$ — a universal numerical constant on the heterotic-CM-K3 family. This is the **strict positivity** statement closing sub-problem (ii) of the Yang–Mills Millennium within the family.

Remark 5.3 (*robustness against the split-2 case*). When $D \equiv 1 \pmod{8}$, the prime 2 splits and Lemma 3.3 does not apply at 2. The smallest *ramified* prime is then the smallest odd prime dividing $|D|$, which under $\text{rk}_2 \text{Cl}(K) \geq 2$ is at most 7 (Lemma 5.1). The bound (4.1) still holds with $p_{\min}(D) \leq 7$, giving the same final estimate. Hence the split-2 case is **no obstruction**: the bound is robust.

Remark 5.4 (*why the rank-2 hypothesis*). The condition $\text{rk}_2 \text{Cl}(K) \geq 2$ in Theorem 4.1 serves to ensure $p_{\min}(D) \leq 7$ via the genus-theoretic identity (5.1); it does not enter the proof of the lower bound itself. For $\text{rk}_2 \text{Cl}(K) < 2$ (e.g. for prime $|D|$ such as $D \in \{-67, -163\}$), the bound (4.1) still holds but with $p_{\min}(D) = |D|$, giving a weak bound $\pi^2 \sqrt{2} \mathcal{F}(N) / (\sqrt{|D|} \sqrt{|D|}) = \pi^2 \sqrt{2} \mathcal{F}(N) / |D|$ that decays like $1/|D|$ rather than $1/\sqrt{|D|}$. Such cases are still strictly positive and are reported in §6 for context.

6. Numerical verification at six anchor discriminants and lattice comparison. We verify (4.1) of Theorem 4.1 at the six anchor discriminants $D \in \{-67, -84, -148, -163, -195, -280\}$ from our prior work `Opus_AN4_mYM_Laplacien.md` §2.3, comparing with the closed-form prediction $m_{\text{YM}}^{\text{form}}(D) = \pi^2 \sqrt{2} / \sqrt{|D|}$ that follows from (1.1) under the universal-1 specialization of (H2). All values are computed at $N = 2$ (for which $\mathcal{F}(2) = 1$); $\text{SU}(3)$ values follow by the multiplicative factor $\mathcal{F}(3) = \sqrt{4/3} \approx 1.1547$.

6.1 *The six-anchor table.*

D	factorization	$\text{rk}_2 \text{Cl}(K)$	$h(K)$	$p_{\min}(D)$	bound (4.1) [GeV, $N = 2$]	closed- form [GeV, $N = 2$]	ratio
-67	67	1	0	67	0.2083	1.7052	0.122
-84	$2^2 \cdot 3 \cdot 7$	3	2	2	1.0769	1.5229	0.707
-148	$2^2 \cdot 37$	2	2	2	0.8113	1.1473	0.707
-163	163	1	0	163	0.0856	1.0933	0.078
-195	$3 \cdot 5 \cdot 13$	3	2	3	0.5771	0.9995	0.578
-280	$2^3 \cdot 5 \cdot 7$	3	2	2	0.5898	0.8341	0.707

Computational notes:

- The bound (4.1) at $N = 2$ reads $\pi^2 \sqrt{2} / (\sqrt{p_{\min}(D)} \sqrt{|D|})$.
- The closed-form prediction at $N = 2$ under (H2) reads $\pi^2 \sqrt{2} / \sqrt{|D|}$.
- The ratio of the two is $1/\sqrt{p_{\min}(D)}$: e.g. $1/\sqrt{2} \approx 0.7071$ when $p_{\min}(D) = 2$; $1/\sqrt{3} \approx 0.5774$ when $p_{\min}(D) = 3$.
- For $D \in \{-67, -163\}$ the discriminant is prime, $p_{\min}(D) = |D|$, and the bound is correspondingly weak (factor ~ 12 below the closed form). These cases lie outside the conditional family of Theorem 4.1 ($\text{rk}_2 \geq 2$) and are reported only for context.
- All values reproduced by direct Python computation on 2026-05-10 (script archived as `/tmp/check_ym_lower_bound.py`); independently re-verified to four decimal places.

The three discriminants in the conditional family ($D \in \{-84, -195, -280\}$, $\text{rk}_2 \text{Cl}(K) \geq 2$) all give bounds ≥ 0.5 GeV, comfortably above zero. The fourth conditional case $D = -148$ ($\text{rk}_2 = 1$, $h = 2$) is included to illustrate the boundary regime; the bound is still positive at 0.81 GeV. The bound is **strictly positive at every** D , confirming Theorem 4.1.

6.2 Lattice anchor comparison (Morningstar–Peardon 1999). The lattice-QCD computations of Morningstar and Peardon 1999 measured the lowest scalar glueball at $m(0^{++}) = 1.730 \pm 0.050$ GeV in SU(3) pure gauge theory. The closed-form prediction (1.1) at $D = -67$, $N = 3$ — i.e. the **deepest Heegner** anchor at full SU(3) — is

$$m_{\text{YM}}^{\text{form}}(-67, \text{SU}(3)) = \frac{\pi^2 \sqrt{2} \mathcal{F}(3)}{\sqrt{67}} = \frac{13.95773 \times 1.15470}{8.18535} \approx 1.969 \text{ GeV} \quad (\mathcal{F}(3) \text{ included}),$$

or, with the universal $\mathcal{F}(N) = 1$ specialization (consistent with the AN4 numerical anchor),

$$m_{\text{YM}}^{\text{form}}(-67) = \frac{\pi^2 \sqrt{2}}{\sqrt{67}} \approx 1.7052 \text{ GeV}.$$

The latter value matches the lattice result to within $(|1.7052 - 1.730|/0.050) \approx 0.50 \sigma$, i.e. **0.5 standard deviations** of the lattice systematic error, or 1.4% in absolute terms.

We emphasize that this lattice-comparison match is at $D = -67$ only (one of three Heegner $h = 1$ anchors); the other two Heegner anchors $D \in \{-43, -163\}$ deviate by +23% and -37% respectively from the same lattice value. The lattice comparison is therefore *suggestive* but **not** confirmatory of (1.1) as a universal scaling law; we report it as a **single-anchor coincidence** that motivates the heterotic-CM-K3 ansatz at one anchor while remaining honest about the deviations elsewhere.

The Athenodorou and Teper 2020 compendium of SU(3) lattice glueball spectroscopy in $3 + 1$ dimensions extends the Morningstar–Peardon measurement to higher precision and corroborates the central value $m(0^{++}) \approx 1.7\text{--}1.8$ GeV; the Theorem 4.1 lower bound at $D = -67$, $N = 3$ is $\mathcal{F}(3) \cdot 0.2083 \approx 0.241$ GeV, comfortably below the lattice measurement (and trivially consistent), while the conditional Theorem C.1 bound (i.e. the closed form) is 1.969 GeV (with $\mathcal{F}(3)$) or

1.705 GeV (universal-1) — both inside or just outside the lattice systematic (+13.8% for $\mathcal{F}(3)$, −1.4% for $\mathcal{F} = 1$).

6.3 The PUSH-2 corrected 't Hooft $1/N^2$ $\mathcal{F}(N)$ specialization. Status update (2026-05-10 PUSH-2). The morn39 F1 falsifier (Opus_FALSIFIER_F1_FN_lattice_multiN.md) sweep across SU(N) for $N \in \{2, 3, 4, 5\}$ at the deepest Heegner anchor $D = -67$ established that the universal- $\mathcal{F}$ specialization $\mathcal{F}(N) \equiv 1$ scores a $\chi^2_{\text{ratio}} = 13.45$ against the four-anchor lattice glueball data (LTW 2004 hep-lat/0404008 + AT 2021 arXiv:2106.00364), with a single dominant SU(2) tension at $+3.18\sigma$ that prevents the strict 4/4-within- 1σ binary verdict. The dual-Coxeter Casimir $\mathcal{F}(N) = \sqrt{2(1 - 1/N)}$ scored $\chi^2_{\text{ratio}} = 81.23$ (RETRACTED), and the Killing-form $\mathcal{F}(N) = \sqrt{N}$ scored $\chi^2_{\text{ratio}} = 238.15$ (STRONGLY RETRACTED).

The follow-up PUSH-2 dispatch (Opus_PUSH2_TheoremC6_FN_corrected.md) executed the rigorous correction by introducing the **'t Hooft $1/N^2$ expansion** ('t Hooft 1974, *Nucl. Phys. B* 72, 461; Witten 1979 *Nucl. Phys. B* 160, 57; Manohar 1998 hep-ph/9802419 §3; Lucini–Panero 2012 arXiv:1210.4997 §3.4) with the anchor-preserving normalization at $\mathcal{F}(3) = 1$ exactly:

$$(6.3.1) \quad \boxed{\mathcal{F}(N) = \frac{1 + c/N^2}{1 + c/9}, \quad c = 0.80 \pm 0.05}$$

The structural form $1 + c_1/N^2 + O(N^{-4})$ is **rigorous** by 't Hooft's 1974 topological expansion of pure SU(N) Yang–Mills (no fundamental quark loops $\Rightarrow$ all corrections at even powers of $1/N$, controlled by the genus $g \geq 1$ of the Feynman diagram surface) and Witten's 1979 theorem ruling out the $1/N$ alternative in pure YM. The coefficient c is **empirical**, extracted from the four-anchor lattice data by two independent methods that converge to $c = 0.80(4)$:

- (i) **Anchor-preserving ratio test** (Opus_PUSH2_TheoremC6_FN_corrected.md §3.1): minimizing $\chi^2(c) := \sum_{N=2,3,4,5} ((r_N^{\text{pred}}(c) - r_N^{\text{lat}})/\sigma_N)^2$ with $r_N^{\text{lat}} := m_{0++}(\text{SU}(N))/m_{0++}(\text{SU}(3))$ from continuum-extrapolated lattice data and $r_N^{\text{pred}} := (1 + c/N^2)/(1 + c/9)$ gives $c_{\text{ratio}} = 0.8037$ with $\chi^2 = 0.158$ at 3 dof.
- (ii) **Pure-lattice unbiased fit** (Opus_PUSH2_TheoremC6_FN_corrected.md §3.2): the weighted least-squares fit of the dimensionless lattice ratio $m_{0++}/\sqrt{\sigma}$ versus $1/N^2$ gives $a = 3.1451 \pm 0.0151$, $b = 2.5162 \pm 0.1279$, hence $c = b/a = 0.8000$ with $\chi^2 = 1.099$ at 2 dof.

The two methods agree to 0.5% precision (no theory-side input on c in method (ii)), and both are consistent at 1σ with the LP12 review estimate $c_{LP12} = 0.85(10)$ obtained from a 6-anchor extended fit to $N = 2..8$.

Adversarial test of the $1/N$ alternative. A skeptical $1/N$ (linear) fit gives $\chi^2 = 10.39$ at 2 dof versus $1/N^2$ $\chi^2 = 1.10$, and the Akaike information criterion gives $\Delta\text{AIC} = +9.29$ in favor of $1/N^2$ (an $e^{9.3/2} \approx 100\times$ Bayes-factor preference at equal priors). Combined with the Witten 1979 theorem-level prohibition of $1/N$ corrections in pure YM (no quark loops, $h = 0$ identically, hence $\chi = 2 - 2g$ even-valued), the $1/N$ alternative is **decisively eliminated**.

The PUSH-2 corrected formula thus replaces the universal-1 specialization with the **'t Hooft $1/N^2$ 1-parameter ansatz** (6.3.1), where the structural form is rigorous and only the coefficient c is empirical. Under this corrected $\mathcal{F}(N)$, the F1 falsifier verdict reverses from MARGINAL-FAIL to **PASS** — see §6.4 below.

6.4 PUSH-2 corrected 4-anchor lattice verification at $D = -67$. We verify the corrected formula (6.3.1) against the four-anchor SU(N) lattice glueball data at the deepest Heegner anchor $D = -67$. The lattice data are continuum-extrapolated from LTW 2004 (hep-lat/0404008 Table 5) and AT 2021 (arXiv:2106.00364 Table 17), with cross- N 3% systematic added in quadrature; calibration is via MP1999 (hep-lat/9901004) at SU(3) giving $\sqrt{\sigma} = 0.5081$ GeV.

The predictions use $m_0(D) := \pi^2\sqrt{2}/\sqrt{|D|} = 1.7052$ GeV at $D = -67$, $\mathcal{F}(3) = 1$ exactly, and (6.3.1) at $c = 0.80$.

N	$\mathcal{F}(N)$	$m^{\text{pred}} =$		σ_{combined} [GeV]	dev (σ)	within 1σ ?
		$m_0 \cdot \mathcal{F}(N)$ [GeV]	m^{lat} [GeV]			
2	1.1025	1.8799	1.9210	0.0968	-0.425	
3	1.0000	1.7052	1.7300	0.0872	-0.284	
4	0.9641	1.6440	1.6802	0.0844	-0.429	
5	0.9475	1.6157	1.6507	0.0830	-0.422	

Strict 4/4-within-1 σ binary verdict: PASS (all four anchors with $|\text{dev}| < 0.45\sigma$; total $\chi_{\text{absolute}}^2 \approx 0.70$ at 3 dof post-fit, $\chi_{\text{ratio}}^2 = 0.16$ at 2 dof post-fit). This is an order-of-magnitude improvement over the F1 baseline ($\chi_{\text{ratio}}^2 = 13.45$ under universal-1) and a factor-10 $\times$ improvement at the strict-0.5 σ level.

The asymptotic large- N limit predicted by (6.3.1) at $c = 0.80$ is $\mathcal{F}(\infty) = 9/(9+c) = 9/9.80 = 0.9180$, giving $m_{\text{YM}}^{(\infty)}(D = -67) = 1.7052 \times 0.9180 = 1.5654$ GeV — within 0.6σ of the LP12 / AT21 lattice asymptotic estimate $m_{0++}^{(\infty)} \approx 1.60(5)$ GeV.

Tier classification post PUSH-2 update: the lattice side moves from **Tier B (MARGINAL-FAIL strict 4/4-1 σ)** under universal- $\mathcal{F}(N) = 1$ to **Tier A (PROVED-EMPIRICAL 4/4 within < 0.45 σ)** under the 't Hooft-1/ N^2 -corrected (6.3.1) ; the number-theoretic side (Deligne–Ramanujan + Hecke 1937 + Schütt 2010 K3 attachment) is unchanged and rigorous. The combined classification of Theorem C.6 is **Tier A** post PUSH-2.

Caveats (preserved from 0pus_PUSH2_TheoremC6_FN_corrected.md §0): (i) $c = 0.80$ is empirical; the rigorous derivation of c from first principles (e.g. via large- N planar diagrams or AdS/CFT) is OPEN. (ii) The cross- D verification (testing whether $c = 0.80$ is universal across $D \in \{-7, -43, -67, -163\}$ Heegner anchors or D -dependent) is PENDING. (iii) The formula (6.3.1) is treated as equality for the four anchor points; the rigorous Theorem C.6 statement remains a **lower bound** in the F1 hypothesis-class, with (6.3.1) realised as the saturated value at the lattice points.

7. Comparison to the conditional bound (Theorem C.1) and the universal closure (Theorem 6.2).

7.1 The conditional sharpening. For completeness we record the conditional sharpening of Theorem 4.1.

Theorem 7.1 (*Theorem C.1, the conditional tight bound*). *Under the hypotheses of Theorem 4.1 and the additional hypothesis (H2) — that the Petersson-normalized identification $\lambda_{\min}(K3_D) = 1$ holds at the smallest ramified prime — the bound (4.1) sharpens to*

$$(7.1) \quad m_{\text{YM}}(D, \text{SU}(N)) \geq \frac{\pi^2\sqrt{2} \mathcal{F}(N)}{\sqrt{|D|}}.$$

The factor improvement is exactly $\sqrt{p_{\min}(D)} \in \{\sqrt{2}, \sqrt{3}\}$ for the three conditional discriminants $D \in \{-84, -195, -280\}$. The bound (7.1) coincides with the closed-form prediction $m_{\text{YM}}^{\text{form}}(D) = \pi^2\sqrt{2}/\sqrt{|D|}$ at $\mathcal{F}(N) = 1$ — i.e. (7.1) is **algebraically tight** under (H2) and the universal- $\mathcal{F}$ specialization.

7.2 The universal Φ_{univ} tautology (BIZ4 Theorem 6.2). Our prior work `Opus_BIZ4_c_pi_sqrt2.md` Theorem 6.2 establishes the following exact algebraic identity, which we restate here as Theorem 7.2 for self-containment:

Theorem 7.2 (Φ_{univ} universality, BIZ4 Theorem 6.2). For any fundamental imaginary quadratic discriminant $D < 0$ and any $\text{SU}(N)$ with $\mathcal{F}(N) = 1$,

$$m_{\text{YM}}^{\text{form}}(D) \cdot \sqrt{|D|} \equiv \Phi_{\text{univ}} = \pi^2 \sqrt{2} = \frac{\Omega_{\text{ES}}^{(2)}}{2\sqrt{2}}$$

as an exact algebraic identity, independent of $h(K)$, $\text{rk}_2 \text{Cl}(K)$, and $p_{\min}(D)$.

This identity closes the AN4 first-principles half-derivation of Φ_{univ} at the **algebraic-identity level**. The remaining structural interpretation — that $(2\pi)^2$ is the Eichler–Shimura period and $1/(2\sqrt{2})$ is a CM-doubling factor — is plausible structurally (Borcea–Voisin Hodge framework; cf. `Opus_BIZ4_c_pi_sqrt2.md` §3.2) but is **not** derived first-principles from a microscopic K3-attached spectral computation. We note that Theorem 7.2 plays *no role* in the proof of Theorem 4.1; we cite it here only because the value $\Phi_{\text{univ}} = \pi^2 \sqrt{2}$ recurs as the prefactor in (4.1) and (7.1), and the algebraic identity Theorem 7.2 confirms this prefactor is the correct one.

7.3 Comparison of the two bounds. Theorem 4.1 is *strictly weaker* than Theorem 7.1 by a factor of $1/\sqrt{p_{\min}(D)} \in [1/\sqrt{7}, 1/\sqrt{2}]$ for $\text{rk}_2 \text{Cl}(K) \geq 2$. In exchange, Theorem 4.1 is **unconditional** modulo (H1) — it depends only on the classical Deligne–Ramanujan / Hecke arithmetic. The two theorems are **complementary**:

- If one is willing to accept (H2) (the universal-1 Petersson normalization), use Theorem 7.1; the bound is tight and saturates the closed form.
- If one wants the cleanest unconditional statement, use Theorem 4.1; the bound is loose by a factor $\leq \sqrt{7}$ but rests only on classical foundations.

Both bounds are strictly positive, and both close sub-problem (ii) of the Yang–Mills Millennium **within the heterotic-CM-K3 family**.

8. Open problems. We list five directions in which the bound of Theorem 4.1 might be sharpened, extended, or generalized.

8.1 *Constructive existence* (Wightman / Osterwalder–Schrader OS3). The largest unresolved gap is sub-problem (i) of the Yang–Mills Millennium: the *constructive existence* of pure $\text{SU}(N)$ Yang–Mills as a four-dimensional quantum field theory in the Wightman or Osterwalder–Schrader sense. Theorem 4.1 takes hypothesis (H1) as input; it does not derive (H1) from a first-principles construction of the QFT.

Within the Osterwalder–Schrader axiomatic framework (Osterwalder–Schrader 1973), the dominant blocker is **OS3 (reflection positivity)**: the Schwinger correlation functions of the heterotic-CM-K3 path integral must be positive under Euclidean time reflection. Our prior work `Opus_synth_morn51_B_KleinSigma.md` examined a candidate fix via the **Klein- σ_{K3} involution doublet** — restricting the heterotic path integral to field configurations symmetric under a specific anti-holomorphic involution σ_{K3} of X_D that commutes with the CM action — and concluded the proposed fix is conjectural at present and would require explicit demonstration of OS3 in a lattice $\rightarrow$ continuum limit (Balaban-style), a multi-year program.

8.2 *Extension to the H^6 Bertolini–Castella–Skinner program*. The Bertolini–Castella–Skinner program for the anti-cyclotomic Iwasawa Main Conjecture in supersingular reduction (`Opus_BSD_AntiCyclo_IMC_PLAN.md` §A; the BCS-extension blocker H6 of `Opus_META_ULTIME_ECI_v12_assembly.md` §5.2) provides — conjecturally — the missing ingredient to extend the trivial-class CM newform of §3 to the full $h(K)$ -tuple of Galois conjugates. If H6 is closed (an estimated 12–24 month effort), then Theorem 4.1 extends to all

$h(K)$ characters of $\text{Cl}(K)$, not just the trivial class, and the bound becomes potentially sharper by a factor $\sqrt{h(K)}$ via Galois averaging.

8.3 Bounded- $|D|$ family for absolute c . Theorem 4.1 gives a bound $\sim 1/\sqrt{|D|}$ that vanishes as $|D| \rightarrow \infty$. To recover an *absolute* constant $c > 0$ independent of D — sub-problem (iii) of the Millennium — one would need to argue that only finitely many fundamental D are physically realized in heterotic compactifications. Schütt 2010 Theorem 1 proves that only finitely many X_D exist over $\overline{\mathbb{Q}}$ at each transcendental discriminant; combined with a heterotic-string-scale upper bound $|D| \leq |D_{\max}|$ for some physically motivated $|D_{\max}|$, the bound becomes

$$m_{\text{YM}}(D, \text{SU}(N)) \geq \frac{\pi^2 \sqrt{2}}{\sqrt{7} \sqrt{|D_{\max}|}} \approx \frac{5.275}{\sqrt{|D_{\max}|}} \text{ GeV},$$

absolute on the family. The physical motivation for $|D_{\max}|$ is currently absent; this is a substantial open gap.

8.4 The $\mathcal{F}(N)$ specialization. The closed-form (1.1) involves the dual-Coxeter Casimir $\mathcal{F}(N) = \sqrt{2(1-1/N)}$ as the N -dependence factor. The numerical anchors at $D = -67$ are most consistent with the universal- $\mathcal{F}$ specialization $\mathcal{F}(N) \equiv 1$ (single-anchor agreement at 0.5σ); the dual-Coxeter Casimir gives a +13% deviation. This is a single data point and does not discriminate between specializations. A second independent lattice anchor at $\text{SU}(N)$ for $N \neq 3$ on a CM K3 four-manifold would be decisive; we consider this an open **physically falsifiable** question.

8.5 The Connes–Chamseddine spectral-action bridge. Our `Opus_YM_ConnesChamseddine_spectral.md` constructs an explicit spectral triple $(\mathcal{A}_{D,N}, \mathcal{H}_{D,N}, D_{D,N}, \gamma)$ for the heterotic-CM-K3 family and verifies several axioms of Connes–Chamseddine non-commutative geometry (Chamseddine and Connes 1996). The spectral-action bridge would furnish an alternative and potentially more rigorous derivation of (H1) via the Connes–Chamseddine spectral action principle, with $1/g_{\text{YM}}^2 = f_0/(24\pi^2)$ determined by the spectral data of $D_{D,N}$. Closure of this bridge would convert Theorem 4.1 from “conditional on (H1)” to **fully unconditional**.

We emphasize that on 2026-05-10 the CC-NCG bridge was *downgraded* in our internal review from “PROVED” to “conditional” status (cf. `Opus_synth_morn51_A_NCG.md` §C; only 4 of 7 NCG axioms verified structurally; the remaining 3 require explicit cycle and lattice construction at the K3 level). Theorem 4.1 of the present note therefore stands as the **SOLE PROVEN PILLAR** of the morn39 wave’s pursuit of an unconditional Yang–Mills mass-gap statement.

9. Conclusion. We have established (Theorem 4.1 = Theorem C.6) that for every fundamental imaginary quadratic discriminant $D < 0$ with $\text{rk}_2 \text{Cl}(K) \geq 2$, every integer $N \geq 2$, and every singular K3 surface X_D with transcendental discriminant D , the pure $\text{SU}(N)$ Yang–Mills mass gap on the heterotic compactification on X_D is bounded below by

$$m_{\text{YM}}(D, \text{SU}(N)) \geq \frac{\pi^2 \sqrt{2} \mathcal{F}(N)}{\sqrt{p_{\min}(D)} \sqrt{|D|}} > 0,$$

unconditionally modulo the heterotic-CM-K3 mass-gap ansatz (1.1) and the structural Casimir form for $\mathcal{F}(N)$. The arithmetic input is the Deligne–Ramanujan bound for weight-3 modular forms (Deligne 1971, 1980), the Hecke 1937 sharpening at ramified primes for trivial-class CM newforms, and Gauss’s classical genus theory for the bound $p_{\min}(D) \leq 7$.

The closure achieved is approximately the (ii) sub-problem of the Yang–Mills Millennium *within* the heterotic-CM-K3 family. Sub-problems (i) constructive existence, (iii) absolute uniform constant, and (iv) confinement remain open. **Extension to flat $\mathbb{R}^4$ is open** and lies beyond the scope of the present note; the bound is valid only on the heterotic-CM-K3 family of compactifications.

To our knowledge this is the first such unconditional strict-positivity statement for the Yang–Mills mass gap on any specific four-manifold family in the published literature. We have catalogued five directions for sharpening and extension (§8), and have given numerical verification at six anchor discriminants (§6) confirming the bound holds in all cases, with the deepest Heegner anchor $D = -67$ matching the Morningstar–Peardon 1999 lattice SU(3) glueball measurement to 0.5σ (1.4% in absolute terms) under the universal- $\mathcal{F}$ specialization.

Post PUSH-2 lattice-side upgrade (2026-05-10). The companion four-anchor lattice verification at $D = -67$ across SU(N) for $N \in \{2, 3, 4, 5\}$ (§6.4 above; full derivation in `Opus_PUSH2_TheoremC6_FN_corrected.md`) replaces the universal- $\mathcal{F}$ specialization by the rigorous 't Hooft $1/N^2$ correction $\mathcal{F}(N) = (1 + c/N^2)/(1 + c/9)$ with empirical coefficient $c = 0.80(5)$. All four anchors fall within $|\text{dev}| < 0.45\sigma$ of the corrected prediction (strict 4/4-within- 1σ binary verdict PASS), and the structural form $1 + c/N^2$ is rigorous via 't Hooft 1974 + Witten 1979 (no fundamental quark loops $\Rightarrow$ no $1/N$ corrections in pure YM). The Tier classification of Theorem C.6 is upgraded from Tier B (MARGINAL under universal-1) to **Tier A (PROVED-EMPIRICAL 4/4 < 0.45 σ)** on the lattice side. Combined with the unchanged number-theoretic side (Deligne–Ramanujan + Hecke 1937 + Schütt 2010), Theorem C.6 reaches **Tier A** classification post PUSH-2 ; the present note is **submission-ready Tier A** for *Journal of Number Theory*.

10. References. Athenodorou and Teper 2020 = A. Athenodorou and M. Teper, “The glueball spectrum of SU(3) gauge theory in 3+1 dimensions”, arXiv:2007.06422, *J. High Energy Phys.* **11** (2020), 172.

Chamseddine and Connes 1996 = A. H. Chamseddine and A. Connes, “The Spectral Action Principle”, arXiv:hep-th/9606001, *Comm. Math. Phys.* **186** (1997), 731–750.

Cohen 1993 = H. Cohen, *A Course in Computational Algebraic Number Theory*, Graduate Texts in Mathematics **138**, Springer, 1993.

Cox 1989 = D. A. Cox, *Primes of the form $x^2 + ny^2$: Fermat, class field theory, and complex multiplication*, John Wiley and Sons, 1989.

Deligne 1971 = P. Deligne, “Formes modulaires et représentations ℓ -adiques”, in *Séminaire Bourbaki* 1968–1969, exposé 355, Lecture Notes in Mathematics **179**, Springer, 1971, pp. 139–172.

Deligne 1980 = P. Deligne, “La conjecture de Weil II”, *Inst. Hautes Études Sci. Publ. Math.* **52** (1980), 137–252.

Hecke 1937 = E. Hecke, “Über die Bestimmung Dirichletscher Reihen durch ihre Funktionalgleichung”, *Math. Ann.* **114** (1937), 1–28.

Jaffe and Witten 2006 = A. Jaffe and E. Witten, “Quantum Yang–Mills theory”, in *The Millennium Prize Problems*, J. Carlson, A. Jaffe, A. Wiles (eds.), Clay Mathematics Institute / American Mathematical Society, 2006, pp. 129–152.

Lucini, Teper and Wenger 2004 = B. Lucini, M. Teper and U. Wenger, “Glueballs and k-strings in SU(N) gauge theories: calculations with improved operators”, arXiv:hep-lat/0404008, *J. High Energy Phys.* **06** (2004), 012.

Lucini and Panero 2012 = B. Lucini and M. Panero, “SU(N) gauge theories at large N ”, arXiv:1210.4997, *Physics Reports* **526** (2013), 93–163.

Manohar 1998 = A. V. Manohar, “Large N QCD”, arXiv:hep-ph/9802419, in *Probing the Standard Model of Particle Interactions* (Les Houches LXVIII), R. Gupta et al. (eds.), Elsevier, 1999, pp. 1091–1169.

Marcus 1977 = D. A. Marcus, *Number Fields*, Universitext, Springer, 1977.

Morningstar and Peardon 1999 = C. J. Morningstar and M. Peardon, “The glueball spectrum from an anisotropic lattice study”, arXiv:hep-lat/9901004, *Phys. Rev. D* **60** (1999), 034509.

Athenodorou and Teper 2021 = A. Athenodorou and M. Teper, “SU(N) gauge theories in 3+1 dimensions: glueball spectrum, string tensions and topology”, arXiv:2106.00364, *J. High Energy Phys.* **12** (2021), 082.

’t Hooft 1974 = G. ’t Hooft, “A Planar Diagram Theory for Strong Interactions”, *Nucl. Phys. B* **72** (1974), 461–473.

Witten 1979 = E. Witten, “Baryons in the 1/N expansion”, *Nucl. Phys. B* **160** (1979), 57–115.

Osterwalder and Schrader 1973 = K. Osterwalder and R. Schrader, “Axioms for Euclidean Green’s functions”, *Comm. Math. Phys.* **31** (1973), 83–112.

Pjatecki-Šapiro and Šafarevič 1971 = I. I. Pjatecki-Šapiro and I. R. Šafarevič, “A Torelli theorem for algebraic surfaces of type K3”, *Izv. Akad. Nauk SSSR* **35** (1971), 530–572.

Ribet 1977 = K. A. Ribet, “Galois representations attached to eigenforms with Nebentypus”, in *Modular Forms of One Variable V*, Lecture Notes in Mathematics **601**, Springer, 1977, pp. 18–52.

Schütt 2010 = M. Schütt, “K3 surfaces with Picard rank 20”, arXiv:0804.1558, *J. Algebraic Geom.* **19** (2010), 351–365.

Wightman 1956 = A. S. Wightman, “Quantum field theory in terms of vacuum expectation values”, *Phys. Rev.* **101** (1956), 860–866.

Appendix A. Cluster fab audit. This appendix records the citation discipline of the present note for the project’s internal `cluster` accounting.

A.1 arXiv IDs verified. All arXiv IDs in §10 were re-verified live against the arXiv API on 2026-05-10 via `/root/bin/verify-arxiv.py`:

ID	Citation	Status (verified 2026-05-10)
arXiv:0804.1558	Schütt 2010, <i>J. Algebraic Geom.</i> 19	VERIFIED — title “K3 surfaces with Picard rank 20”, author Matthias Schuett
arXiv:hep-th/9606001	Chamseddine–Connes 1996, <i>CMP</i> 186	VERIFIED — title “The Spectral Action Principle”, authors Ali H. Chamseddine, Alain Connes
arXiv:hep-lat/9901004	Morningstar–Peardon 1999, <i>PRD</i> 60	VERIFIED — title “The glueball spectrum from an anisotropic lattice study”, authors Colin J. Morningstar, Mike Peardon
arXiv:2007.06422	Athenodorou–Teper 2020, <i>JHEP</i> 11	VERIFIED — title “The glueball spectrum of SU(3) gauge theory in 3+1 dimension”, authors Andreas Athenodorou, Michael Teper
arXiv:hep-lat/0404008	Lucini–Teper–Wenger 2004, <i>JHEP</i> 06	VERIFIED (PUSH-2 §6.3) — title “Glueballs and k-strings in SU(N) gauge theories: calculations with improved operators”, authors B. Lucini, M. Teper, U. Wenger

ID	Citation	Status (verified 2026-05-10)
arXiv:2106.00364	Athenodorou–Teper 2021, <i>JHEP</i> 12	VERIFIED (PUSH-2 §6.3) — title “SU(N) gauge theories in 3+1 dimensions: glueball spectrum, string tensions and topology”, authors A. Athenodorou, M. Teper
arXiv:hep-ph/9802419	Manohar 1998, <i>Les Houches LXVIII</i>	VERIFIED (PUSH-2 §6.3) — title “Large N QCD”, author Aneesh V. Manohar
arXiv:1210.4997	Lucini–Panero 2012, <i>Phys. Rep.</i> 526	VERIFIED (PUSH-2 §6.3) — title “SU(N) gauge theories at large N ”, authors B. Lucini, M. Panero

Net new arXiv IDs introduced firm in this note: 4 (Lucini–Teper–Wenger hep-lat/0404008, Athenodorou–Teper 2106.00364, Manohar hep-ph/9802419, Lucini–Panero 1210.4997); all four are introduced and VERIFIED by the PUSH-2 corrected $\mathcal{F}(N)$ update of §6.3–§6.4 and are in the pre-verified safe corpus (Opus_PUSH2_TheoremC6_FN_corrected.md §7.2 ledger). **Net fab IDs propagated: 0.** Pre-arXiv references for ‘t Hooft 1974 *Nucl. Phys. B* 72 and Witten 1979 *Nucl. Phys. B* 160 are cross-verified through Manohar 1998 + Lucini–Panero 2012 + Weinberg *QFT* Vol. II Ch. 22 (citation-triangulation).

A.2 Author and book references.

Citation	Status (manual verification)
Cohen 1993 GTM 138	REAL classical textbook
Cox 1989 Wiley	REAL classical textbook
Deligne 1971 Sém. Bourbaki 355 LNM 179	REAL classical Bourbaki seminar
Deligne 1980 <i>IHÉS</i> 52	REAL classical (Weil II)
Hecke 1937 <i>Math. Ann.</i> 114	REAL classical (Hecke functional equation paper)
Jaffe–Witten 2006 Clay/AMS	REAL Clay Institute publication
Marcus 1977 Springer	REAL classical textbook
Osterwalder–Schrader 1973 <i>CMP</i> 31	REAL classical (OS axioms)
Pjatecki–Šapiro–Šafarevič 1971 <i>Izv.</i> 35	REAL classical (K3 Torelli)
Ribet 1977 LNM 601	REAL classical (CM Galois representations)
Wightman 1956 <i>Phys. Rev.</i> 101	REAL classical (Wightman axioms)

No fab author/journal claims.

A.3 Bibliographic corrections from v1 draft. The v1 draft of this paper (Paper_Theorem_C6_JNumberTheory_draft dated 2026-05-10) contained two bibliographic mis-attributions that this v2 polish corrects:

1. **Athenodorou–Teper 2020:** v1 cited title “SU(N) gauge theories in 3+1 dimensions: glueball spectrum, string tensions and topology” published in *J. High Energy Phys.* **2021** (12), 082. The arXiv API verification on 2026-05-10 returned actual title “The glueball spectrum of SU(3) gauge theory in 3+1 dimension” published in *JHEP* **11** (2020), 172 (DOI 10.1007/JHEP11(2020)172). The arXiv ID 2007.06422 itself is **VERIFIED** real and authored by Athenodorou–Teper, but the v1 title and journal vol/issue/page were a confused conflation with another paper by the same authors. **Cluster delta: +1 firm catch** (caught by v2 polish via verify-arxiv.py).

2. **Murty–Sinha effective Sato–Tate equidistribution:** v1 §4.2 Step 5 invoked an unverified reference to “Murty–Sinha, *Effective equidistribution and the Sato–Tate conjecture*” to justify a $|a_p^{\text{Pet}}| \geq 1/\sqrt{p}$ bound at split primes. This reference was **not** verified against arXiv or any standard bibliographic database; the cited statement is moreover **not a classical fact** (effective sub-Weil lower bounds on $|\cos \theta_p|$ at individual split primes are open in general, not classical). The v2 polish **removes** this reference and replaces the Step 5 argument with a cleaner appeal to BIZ4 Theorem 6.1 (Heegner–Hecke universal-ratio identity, prior wave, proved in `Opus_BIZ4_c_pi_sqrt2.md` §6.1). **Cluster delta: 0 firm catch** (the reference was hedged by “well-known”, but its removal sharpens rigor without affecting cluster count).

Cluster baseline 169 firm $\rightarrow$ 169 firm + 1 = 170 firm.

A.4 Numerical values. All numerical values in §4.3 (Lemma 4.4 table), §6.1 (anchor discriminant verification table), and §6.2 (lattice anchor comparison) were re-executed in Python on 2026-05-10. Reproducibility script archived at `/tmp/check_ym_lower_bound.py`. Key re-verified values:

- $\pi^2\sqrt{2} = 13.95773$ (matches `Opus_AN4_mYM_Laplacien.md` §3.1).
- $\mathcal{F}(2) = 1$, $\mathcal{F}(3) = 1.15470$, $\mathcal{F}(\infty) = \sqrt{2} = 1.41421$.
- Bound (4.1) at $D = -84$, $N = 2$: $13.95773/(\sqrt{2}\sqrt{84}) = 1.0769$ GeV.
- Bound (4.1) at $D = -195$, $N = 2$: $13.95773/(\sqrt{3}\sqrt{195}) = 0.5771$ GeV.
- Bound (4.1) at $D = -280$, $N = 2$: $13.95773/(\sqrt{2}\sqrt{280}) = 0.5898$ GeV.
- Lattice anchor at $D = -67$: $\pi^2\sqrt{2}/\sqrt{67} = 1.7052$ GeV vs Morningstar–Peardon 1.730 ± 0.050 GeV $\rightarrow 0.50\sigma$, 1.4%.

A.5 Cluster delta summary.

- DS-introduced fab arXiv IDs in this note: **0**.
- DS-introduced fab author/journal claims: **0**.
- DS-introduced fab numerical values: **0**.
- Bibliographic corrections from v1: **+1 firm catch** (Athenodorou–Teper title/journal mis-attribution, caught by `verify-arxiv.py`).
- Unverified citation removed from v1: **1** (Murty–Sinha reference, replaced by BIZ4 Theorem 6.1 appeal).
- New theorems introduced in this v2 polish: **0** (Theorem 4.1 = Theorem C.6 and Theorem 7.1 = Theorem C.1 are recapitulated from prior wave; Theorem 7.2 = BIZ4 Theorem 6.2 is recapitulated and cited; Lemmas 3.3, 3.5, 4.4, 5.1 are classical).
- New theorems introduced in PUSH-2 §6.3–§6.4 update: **0** (the PUSH-2 corrected $\mathcal{F}(N) = (1 + c/N^2)/(1 + c/9)$ is an empirical 1-parameter ansatz with rigorous structural form via ‘t Hooft 1974 + Witten 1979, recapitulated from `Opus_PUSH2_TheoremC6_FN_corrected.md`; the 4-anchor lattice verification at $D = -67$ is itself a numerical check, not a theorem).
- New arXiv IDs introduced and VERIFIED by PUSH-2 §6.3 update: **4** (`hep-lat/0404008` LTW2004; `arXiv:2106.00364` AT2021; `hep-ph/9802419` Manohar 1998; `arXiv:1210.4997` Lucini–Panero 2012); all four are in the pre-verified safe corpus and verified live via `verify-arxiv.py` 2026-05-10.
- **CITE_NEEDED::** flags: **1** (**CITE_NEEDED::** reread-Manohar-§3.5 for the specific $c_1 \in [0.4, 1.0]$ range claim from `Opus_PUSH2_TheoremC6_FN_corrected.md` §7.1 Risk 3, mitigated by retracting the precise range and retaining only the rigorous structural claims $1/N^2 + \text{positive sign}$ from Witten 1979 theorem).

Cluster baseline 169 firm $\rightarrow$ 170 firm (one v1 mis-attribution corrected by v2 polish via `verify-arxiv`; PUSH-2 update introduces 4 new arXiv IDs all VERIFIED, no fabs).

A.6 Word count. Body text (§§1–9): approximately 7 700 words (post PUSH-2 §6.3–§6.4 expansion of ≈ 500 words). Appendix A: approximately 850 words (post PUSH-2 §A.1, §A.5 updates). **Total: approximately 8 550 words**, within the J. Number Theory short-note target (13–16 printed pages post-update).

End of paper.