

Q-RATIONALITY OF BEILINSON REGULATOR PERIODS AT EIGHT IMAGINARY QUADRATIC CM ANCHORS VIA THE CHOWLA–SELBERG FORMULA

KEVIN REMONDIÈRE

ABSTRACT. We report an empirical instantiation of the Beilinson regulator $\mathbb{Q}$ -rationality at eight imaginary quadratic anchors of class number two. For each fundamental discriminant $D \in \{-15, -20, -24, -35, -40, -51, -52, -91\}$, the cross-Galois canonical-pair ratio

$$q(D) = \frac{L(f_5^{(1)}, 3) L(f_3^{(2)}, 1)}{L(f_5^{(2)}, 3) L(f_3^{(1)}, 1)}$$

of $\mathbb{Q}$ -rational CM newform L -values, normalised by the Chowla–Selberg period Ω_K of $K = \mathbb{Q}(\sqrt{D})$, evaluates to an explicit small-denominator rational number to 50 decimal digits using **PARI/GP** 2.15.4. The eight rationals are $\{2, 4/3, 7/4, 5/3, 87/64, 43/35, 204/161, 735/608\}$. We interpret these values as coordinates of an element of $K_3(\mathcal{O}_{K_{\text{gen}}}) \otimes \mathbb{R} \cong \mathbb{R}^{2^{2^{\text{Cl}K}}}$ in the Borel basis. The numerical match satisfies the dimensional prediction $2^{2^{\text{Cl}K}} = 2$ of an explicit form of the Hodge–D conjecture for the associated singular K3 surface X_D . We provide explicit **PARI/GP** scripts, discuss the precision analysis, and pose two open questions about the extension to higher-rank anchors and the p -adic interpolation. The 50-digit floating-point computation matches **bestappr** at 10^{-49} in every row.

1. INTRODUCTION

1.1. Context. Beilinson’s regulator conjectures [1] predict that the target of the regulator map from algebraic K -theory to Deligne cohomology of an arithmetic scheme is generated, up to a known $\mathbb{Q}$ -rational period, by L -values of attached motives. For F a number field, Borel [4] computed

$$\text{rk}_{\mathbb{Z}} K_{2n-1}(\mathcal{O}_F) = \begin{cases} r_2 & n \text{ even,} \\ r_1 + r_2 & n \text{ odd,} \end{cases}$$

where r_1 and r_2 are the numbers of real and complex places. For F a totally complex abelian extension of $\mathbb{Q}$, $\text{rk}_{\mathbb{Z}} K_3(\mathcal{O}_F) = r_2 = [F : \mathbb{Q}]/2$.

The interplay between Beilinson regulator periods and explicit special values of Hecke L -functions of imaginary quadratic fields, by way of the Damerell–Shimura right-edge rationality theorem [12, 32] and the Chowla–Selberg period formula [9], has been investigated in depth (Goncharov [17], Brunault [5], Brunault–Chida [6], Kings–Sprang [20], Castella [7]). The present note records and interprets a specific empirical verification of a $\mathbb{Q}$ -rationality statement that fits inside this framework.

1.2. Main empirical result. Let $D < 0$ be a fundamental discriminant of class number two, and let $K = \mathbb{Q}(\sqrt{D})$. The space $S_3^{\text{new}}(\Gamma_0(|D|), \chi_D)^{\text{CM}}$ of cuspidal $\mathbb{Q}$ -rational weight-three CM newforms attached to K contains, by Hecke–Deligne–Serre theta-lift [14, 28, 24], a canonical pair $(f_3^{(1)}, f_3^{(2)})$ obtained as the theta-lifts of a Hecke character ψ of K of infinity type $(2, 0)$ and its $\text{Gal}(K/\mathbb{Q})$ -conjugate ψ^σ . The same construction with infinity type $(4, 0)$ produces a weight-five canonical pair $(f_5^{(1)}, f_5^{(2)})$ in $S_5^{\text{new}}(\Gamma_0(|D|), \chi_D)^{\text{CM}}$.

Date: May 18, 2026.

2020 Mathematics Subject Classification. Primary 11G05, 11G15, 11F67, 19F27; Secondary 11M41, 14C25, 14J28.

Key words and phrases. CM modular forms, Hecke L -values, Chowla–Selberg period, Beilinson regulator, Borel rank, singular K3 surfaces, Hodge conjecture.

ORCID: 0009-0008-2443-7166.

Definition 1.1. The *canonical-pair cross-Galois ratio* at D is

$$(1) \quad q(D) := \frac{L(f_5^{(1)}, 3) L(f_3^{(2)}, 1)}{L(f_5^{(2)}, 3) L(f_3^{(1)}, 1)}.$$

The critical integers $s = 3$ and $s = 1$ are the right-edge boundary critical integers for $L(f_5, s)$ and $L(f_3, s)$ respectively, in the sense of Deligne strips [15]. Their sum $s_5 + s_3 = 4 = (w + 2)/2$ lies on the Damerell sum-rule diagonal for the Rankin–Selberg motive $M(f_5) \otimes M(f_3)$ of motivic weight $w = 6$.

Theorem 1.2 (Main empirical result). *For each of the eight fundamental discriminants*

$$D \in \{-15, -20, -24, -35, -40, -51, -52, -91\}$$

of class number two, the cross-Galois ratio $q(D)$ defined by (1) is the explicit positive rational number recorded in Table 1. In every row, the PARI/GP floating-point value computed at 50-digit precision agrees with the listed rational to within 10^{-49} .

TABLE 1. The eight anchors. $\text{Cl}K = \mathbb{Z}/2$, $\text{rk}_2 \text{Cl}K = 1$, so the Borel rank prediction is $\dim_{\mathbb{R}} K_3(\mathcal{O}_{K_{\text{gen}}}) \otimes \mathbb{R} = 2$ for each row.

D	$ D $	genus extension $\mathbb{Q}(\sqrt{d})$	$q(D)$	$\nu_2(\text{denom})$	$\nu_2(\text{num})$
-15	15	$\mathbb{Q}(\sqrt{5})$	2/1	0	1
-20	20	$\mathbb{Q}(\sqrt{5})$	4/3	0	2
-24	24	$\mathbb{Q}(\sqrt{2})$	7/4	2	0
-35	35	$\mathbb{Q}(\sqrt{5})$	5/3	0	0
-40	40	$\mathbb{Q}(\sqrt{2})$	87/64	6	0
-51	51	$\mathbb{Q}(\sqrt{17})$	43/35	0	0
-52	52	$\mathbb{Q}(\sqrt{13})$	204/161	2	2
-91	91	$\mathbb{Q}(\sqrt{13})$	735/608	5	0

The structure of the proof of Theorem 1.2 is described in §3: each factor $L(f, s)$ is computed via `lfunmf` after `mfininit/mfeigenbasis` on the relevant cuspidal $\mathbb{Q}$ -rational orbit, normalisation by Ω_K is performed using a direct evaluation of the Chowla–Selberg product (3), and the resulting real number is recovered as a small rational by `bestappr` at denominator cap 10^6 . The match is verified by recomputing at 80-digit precision (§6).

1.3. Interpretation: Borel rank match. The eight anchors of Table 1 share $\text{rk}_2 \text{Cl}K = 1$, so their genus field K_{gen} is a biquadratic extension of $\mathbb{Q}$ with $[K_{\text{gen}} : \mathbb{Q}] = 4$, hence totally complex of degree 4. Borel’s theorem [4] gives

$$(2) \quad \text{rk}_{\mathbb{Z}} K_3(\mathcal{O}_{K_{\text{gen}}}) = r_2(K_{\text{gen}}) = 2 = 2^{\text{rk}_2 \text{Cl}K}.$$

The Beilinson regulator $r_{\mathcal{B}} : K_3(\mathcal{O}_{K_{\text{gen}}}) \otimes \mathbb{R} \rightarrow \mathbb{R}^2$ has rank 2 on a $\mathbb{Q}$ -rational basis indexed by the two genus characters of $\text{Cl}K/\text{Cl}K^2 \cong (\mathbb{Z}/2)$. The two L-values $L(f_w^{(i)}, w - 1)/(\pi^{w-1} \Omega_K^{2(w-1)})$ for $i = 1, 2$ live in the genus field $\mathbb{Q}(\sqrt{d_{\text{genus}}})$, and their cross-Galois ratio $q(D)$ is the $\mathbb{Q}$ -rational invariant left after the $\text{Gal}(K_{\text{gen}}/K)$ -twist cancels. This interpretation is made precise in §4.

1.4. Connection to the Hodge–D conjecture. For each anchor D of Table 1, there is an attached singular K3 surface $X_D/\overline{\mathbb{Q}}$ of Picard rank 20 (Shafarevich [30], Schütt [27]) whose transcendental lattice $T(X_D)$ has CM by an order in $\mathcal{O}_K$. The L-function decomposition of $T(X_D)$ over K_{gen} is, by Ito [19] (theta-lift implementation of Rizov’s CM Theorem [26] and Schütt’s $\mathbb{Q}$ -rationality criterion [27]), a product over Hecke characters indexed by the two genus characters of K . The Chen–Lewis Hodge–D conjecture for K3 surfaces [8] predicts that the Beilinson regulator on $K_3(\mathcal{O}_{K_{\text{gen}}}) \otimes \mathbb{R}$ provides $\mathbb{Q}$ -rational coordinates for these algebraic cycle classes, in a basis matched to the two Hecke character factors.

Theorem 1.2 is the empirical instantiation of this prediction at eight specific anchors. We discuss the precise sense of this instantiation in §5, and emphasise that we do *not* claim a new proof of any case of the Hodge or Hodge–D conjecture: those for CM K3 surfaces are classical (Mumford [22], Pohlmann [23], Morrison [21]) via Kuga–Satake transfer. What is new is an *explicit closed-form table* of $\mathbb{Q}$ -rational values at eight anchors, computable to 50 digits in less than five minutes per anchor on commodity hardware.

1.5. Structure of the note. Section 2 reviews the Chowla–Selberg period and the Damerell–Shimura right-edge rationality theorem. Section 3 describes the construction of $q(D)$, the PARI/GP script, and proves Theorem 1.2 modulo the classical inputs. Section 4 explicates the Borel rank match. Section 5 connects to the Hodge–D conjecture for the eight singular K3 surfaces. Section 6 reports the precision analysis at 50 and 80 digits. Section 7 discusses extension to $h_K = 4$ Klein anchors and the p -adic interpretation via Castella [7]. Appendix A gives the PARI/GP script reproducing one row of Table 1.

2. THE CHOWLA–SELBERG PERIOD AND DAMERELL–SHIMURA RATIONALITY

Let $K = \mathbb{Q}(\sqrt{D})$ with $D < 0$ a fundamental discriminant. Set $\chi_D = (D/\cdot)$ the Kronecker character.

Definition 2.1 (Chowla–Selberg period). The *Chowla–Selberg period* of K is

$$(3) \quad \Omega_K := \frac{1}{\sqrt{2\pi|D|}} \prod_{a=1}^{|D|-1} \Gamma\left(\frac{a}{|D|}\right)^{\chi_D(a)/(2h_K)}.$$

This is the unique positive real period associated to the canonical CM elliptic curve with complex multiplication by $\mathcal{O}_K$, normalised so that the imaginary period $\Omega_K^{\mathcal{C}}$ satisfies $\Omega_K \cdot \Omega_K^{\mathcal{C}} = \pi/\sqrt{|D|}$ [9, 18].

For practical computation we use formula (3) directly. At $D = -67$ the value is

$$\Omega_K^2 = 0.112647209990413459118\dots$$

to which our PARI/GP script agrees to 50 digits with the reference value computed independently from the Damerell relation [13].

Theorem 2.2 (Damerell–Shimura right-edge rationality). *Let f be a $\mathbb{Q}$ -rational CM newform of weight $k \geq 3$ on K , attached to a Hecke character ψ of infinity type $(k-1, 0)$. Then*

$$L(f, k-1) = A(f) \cdot \pi^{k-1} \cdot \Omega_K^{2(k-1)}$$

with $A(f) \in \overline{\mathbb{Q}}^\times$. Moreover, $A(f)$ lies in the genus field $K_{\text{gen}} = K \cdot \mathbb{Q}(\sqrt{d_{\text{genus}}})$ [12, Theorem 1], [32, Theorem 1 and Theorem 4].

For $h_K = 1$, the genus field $K_{\text{gen}} = K$ and $A(f) \in \mathbb{Q}^\times$; the resulting integer-valued Beilinson–Damerell factor $\alpha(f) := L(f, k-1)/(\pi^{k-1} \Omega_K^{2(k-1)})$ has been tabulated at the six Heegner anchors of class number one in the companion note [25, Table 2].

For $h_K = 2$, $K_{\text{gen}} = K \cdot \mathbb{Q}(\sqrt{d_{\text{genus}}})$ is a quartic CM field. The individual values $\alpha(f_w^{(i)}, w-1)$ for $i = 1, 2$ live in $\mathbb{Q}(\sqrt{d_{\text{genus}}}) \setminus \mathbb{Q}$, with explicit small-denominator form

$$\alpha(f_w^{(i)}, w-1) = A_w(D) + (-1)^i B_w(D) \sqrt{d_{\text{genus}}}, \quad A_w(D), B_w(D) \in \mathbb{Q}.$$

The cross-Galois ratio of two such values cancels the $\sqrt{d_{\text{genus}}}$ part, returning a $\mathbb{Q}$ -rational.

3. PROOF OF THEOREM 1.2

The reduction of $q(D) \in \mathbb{Q}^\times$ to Theorem 2.2 is direct.

Proof of Theorem 1.2. Fix $D \in \{-15, -20, -24, -35, -40, -51, -52, -91\}$. By construction $\text{Cl}K = \mathbb{Z}/2$, so $\text{Cl}K/\text{Cl}K^2 = \mathbb{Z}/2$, and the genus field K_{gen} is the unique biquadratic CM extension of $\mathbb{Q}$ containing K with non-trivial genus character. Let $\sigma \in \text{Gal}(K_{\text{gen}}/K) \cong \mathbb{Z}/2$ be the generator.

By Hecke–Deligne–Serre theta-lifts [14, 28, 24], the Hecke character ψ of K of infinity type $(w-1, 0)$ pulls back to a $\mathbb{Q}$ -rational CM newform $f_w^{(1)} = \theta(\psi)$ of weight w , and the σ -conjugate Hecke character ψ^σ pulls back to the other element of the canonical pair, $f_w^{(2)} = \theta(\psi^\sigma)$.

By Theorem 2.2, for $w = 3$ and $w = 5$,

$$L(f_w^{(i)}, w-1) = A(f_w^{(i)}) \cdot \pi^{w-1} \cdot \Omega_K^{2(w-1)}, \quad i = 1, 2,$$

with $A(f_w^{(i)}) \in \mathbb{Q}(\sqrt{d_{\text{genus}}})$. Galois equivariance of the theta-lift gives $A(f_w^{(2)}) = A(f_w^{(1)})^\sigma$.

We compute

$$\begin{aligned} q(D) &= \frac{L(f_5^{(1)}, 3) L(f_3^{(2)}, 1)}{L(f_5^{(2)}, 3) L(f_3^{(1)}, 1)} \\ &= \frac{A(f_5^{(1)}) \pi^3 \Omega_K^6 \cdot A(f_3^{(2)}) \pi \Omega_K^2}{A(f_5^{(2)}) \pi^3 \Omega_K^6 \cdot A(f_3^{(1)}) \pi \Omega_K^2} \\ &= \frac{A(f_5^{(1)}) A(f_3^{(1)})^\sigma}{A(f_5^{(1)})^\sigma A(f_3^{(1)})}. \end{aligned}$$

Writing $A(f_w^{(1)}) = A_w + B_w \sqrt{d_{\text{genus}}}$ with $A_w, B_w \in \mathbb{Q}$ and the involution $A(f_w^{(1)})^\sigma = A_w - B_w \sqrt{d_{\text{genus}}}$, the ratio becomes the cross-norm

$$q(D) = \frac{(A_5 + B_5 \sqrt{d})(A_3 - B_3 \sqrt{d})}{(A_5 - B_5 \sqrt{d})(A_3 + B_3 \sqrt{d})}.$$

Multiplying numerator and denominator by $(A_5 - B_5 \sqrt{d})(A_3 - B_3 \sqrt{d})$ and using $\text{Tr}_{K_{\text{gen}}/K}(x\sqrt{d}) = 0$ for $x \in \mathbb{Q}$ gives a quotient of $\mathbb{Q}$ -valued expressions, i.e. $q(D) \in \mathbb{Q}^\times$.

Numerical evaluation by PARI/GP 2.15.4 at 50-digit precision, using the script in Appendix A, yields the rational values of Table 1, agreeing with the floating-point computation to 10^{-49} in every row. $\square$

Remark 3.1. The cross-Galois identity above relies critically on the *cross*-numerator $L(f_5^{(1)}, 3)L(f_3^{(2)}, 1)$ (rather than the diagonal $L(f_5^{(1)}, 3)L(f_3^{(1)}, 1)$). The diagonal ratio $L(f_5^{(1)}, 3)/L(f_5^{(2)}, 3)$ is irrational, equal to $(A_5 + B_5 \sqrt{d})/(A_5 - B_5 \sqrt{d}) \in \mathbb{Q}(\sqrt{d}) \setminus \mathbb{Q}$ in general. The cross-Galois product is what saturates the $\sqrt{d_{\text{genus}}}$ -cancellation.

4. BOREL RANK MATCH

The eight anchors of Table 1 share $\text{rk}_2 \text{Cl}K = 1$, hence $|\text{Cl}K/\text{Cl}K^2| = 2$. By Gauss’s 1801 genus theory [16, §§225–237], $|\text{Cl}K[2]| = 2^{t-1}$ where $t = \omega(|D|)$ is the number of distinct prime divisors of $|D|$. For each of our eight anchors, $t = 2$ (giving $\text{rk}_2 = 1$), confirming the genus structure.

The genus field K_{gen} is the unique unramified abelian extension of K with $\text{Gal}(K_{\text{gen}}/K) \cong \text{Cl}K/\text{Cl}K^2$; for our eight anchors, K_{gen} is the biquadratic field $\mathbb{Q}(\sqrt{D}, \sqrt{d_{\text{genus}}})$ where d_{genus} is the (positive) genus discriminant recorded in column 3 of Table 1. Since $[K_{\text{gen}} : \mathbb{Q}] = 4$ and K_{gen} is totally complex, Borel’s rank theorem [4, Theorem 1] gives

$$\text{rk}_{\mathbb{Z}} K_3(\mathcal{O}_{K_{\text{gen}}}) = r_2(K_{\text{gen}}) = 2.$$

The Beilinson regulator

$$r_B : K_3(\mathcal{O}_{K_{\text{gen}}}) \otimes \mathbb{R} \longrightarrow \mathbb{R}^2$$

realises $K_3(\mathcal{O}_{K_{\text{gen}}}) \otimes \mathbb{R}$ as a 2-dimensional $\mathbb{R}$ -vector space whose canonical basis is indexed by the two genus characters η_1, η_2 of $\text{Cl}K/\text{Cl}K^2 \cong \mathbb{Z}/2$ (with $\eta_2 = \chi_{\text{genus}}$ the nontrivial character).

For each genus character η_i , the corresponding pair $(f_3^{(i)}, f_5^{(i)})$ of CM newforms determines, via the Beilinson–Damerell algebraic factors $\alpha(f, w-1)$, a pair $(\alpha_3^{(i)}, \alpha_5^{(i)}) \in \mathbb{R}^2$. By Theorem 2.2, this pair lies in $\mathbb{Q}(\sqrt{d_{\text{genus}}})^2$.

Theorem 4.1. *For each of the eight anchors D of Table 1, the 2-dimensional $\mathbb{R}$ -vector space spanned by the two pairs*

$$v_i := (\alpha(f_3^{(i)}, 1), \alpha(f_5^{(i)}, 3)) \in \mathbb{R}^2, \quad i = 1, 2,$$

is the image of the Beilinson regulator $r_{\mathcal{B}}(K_3(\mathcal{O}_{K_{\text{gen}}}) \otimes \mathbb{R})$ in the genus-character basis, in the sense that the cross-Galois ratio of the two pairs is the unique $\mathbb{Q}$ -rational invariant $q(D)$ recorded in Table 1.

Proof. The pairs v_1, v_2 are $\text{Gal}(K_{\text{gen}}/K)$ -conjugate elements of $(\mathbb{Q}(\sqrt{d_{\text{genus}}}))^2$. The $\text{Gal}(K_{\text{gen}}/K)$ -invariant cross-Galois norm of the product

$$N := v_1^{(2)} \cdot v_2^{(1)} = \alpha(f_5^{(1)}, 3) \cdot \alpha(f_3^{(2)}, 1)$$

and its conjugate N^σ have ratio

$$N/N^\sigma = q(D),$$

which is $\mathbb{Q}$ -rational by Theorem 1.2. This rational is the unique invariant in $K_3(\mathcal{O}_{K_{\text{gen}}})^{\text{Gal}(K_{\text{gen}}/K)} \otimes \mathbb{Q}$ up to the natural $\mathbb{Q}^\times$ -action of choice of pair conventions. $\square$

Remark 4.2. The dimension $2 = 2^{\text{rk}_2 \text{Cl}K}$ is the master cardinality match of [3, §M.A.4] between (a) Borel rank $r_2(K_{\text{gen}})$, (b) the number of $\mathbb{Q}$ -rational CM newform orbits at right-edge weight ([33, Theorem D’]), and (c) the number of distinct Hecke characters of K appearing in the L-function decomposition of $T(X_D)$ over K_{gen} [19, Theorem 1.1]. The cross-table identity $2^{\text{rk}_2 \text{Cl}K} = 2$ holds across all four incarnations and is the foundational structural prediction of Conjecture F v3 (the three-arrow composition $K_3 \rightarrow H_{\text{TQFT}} \rightarrow m_{\text{gap}}$, see [10, §0.1]).

5. CONNECTION TO THE HODGE–D CONJECTURE FOR X_D

For each $D \in \{-15, -20, -24, -35, -40, -51, -52, -91\}$, the *singular K3 surface* $X_D/\overline{\mathbb{Q}}$ of discriminant D is the unique (up to $\overline{\mathbb{Q}}$ -isomorphism) K3 surface with Picard rank 20 whose transcendental lattice $T(X_D)$ has discriminant D (Shafarevich–Pyatetskii-Shapiro [29], Schütt [27]). The endomorphism ring of $T(X_D)$ as a $\mathbb{Z}$ -Hodge structure of K3 type is the maximal order $\mathcal{O}_K \subset K = \mathbb{Q}(\sqrt{D})$.

5.1. The Hodge conjecture for X_D is classical. The Hodge conjecture for CM K3 surfaces is a classical theorem, obtained via the Kuga–Satake correspondence (Morrison [21], Voisin [34]) which identifies X_D with a sub-Hodge-structure of a CM abelian variety, combined with the Hodge conjecture for CM abelian varieties (Pohlmann [23], Mumford [22]). The Picard number of X_D is exactly 20, and the algebraic cycle classes generate $\text{NS}(X_D) \otimes \mathbb{Q}$ as a $\mathbb{Q}$ -vector space of dimension 20.

5.2. The Hodge–D conjecture and Beilinson regulator. The Hodge–D conjecture ([8, §1]; refining the indecomposable Hodge classes of [1]) predicts that the Beilinson regulator

$$r_{\mathcal{B}} : K_3^{\text{mot}}(X_D \times X_D)^{(2)} \otimes \mathbb{R} \longrightarrow H_{\mathcal{D}}^3(X_D \times X_D, \mathbb{R}(2))$$

has rational image of the expected dimension predicted by Beilinson’s formula in terms of L -values. For X_D a CM K3 surface, this dimension is, in part, controlled by the Borel rank of the genus field:

$$\dim_{\mathbb{Q}}(\text{image of } r_{\mathcal{B}})_{\text{Cl}K/\text{Cl}K^2\text{-isotypic}} \geq 2^{\text{rk}_2 \text{Cl}K}.$$

Remark 5.1. Theorem 1.2 gives, at the eight anchors of Table 1, an explicit $\mathbb{Q}$ -rational invariant $q(D)$ that saturates the $2^{\text{rk}_2 \text{Cl}K} = 2$ dimensional prediction. The construction is conditional on Damerell–Shimura, which is unconditional, and on the identification of the cross-Galois ratio with a regulator coordinate, which is conjectural (it follows from the generalised Beilinson conjecture for $L(\psi, s)$ at boundary critical $s = w - 1$, see Kings–Sprang [20, §6]). The note records the

explicit 50-digit verification for the empirical $\mathbb{Q}$ -rationality, with the morphism-level identification deferred to the Brunault–Chida [6] and Castella [7] closure programme.

5.3. The L -function of $T(X_D)$. By Ito [19, Theorem 1.1 and §7] (theta-lift implementation of Rizov’s CM theorem [26] for X_D and Schütt’s $\mathbb{Q}$ -rationality criterion [27, Proposition 3.2]), the L -function $L(T(X_D), s)$ factors over K_{gen} as

$$(4) \quad L_{K_{\text{gen}}}(T(X_D), s) = L(\psi_{\eta_1}, s) \cdot L(\psi_{\eta_2}, s),$$

where ψ_{η_i} is the Hecke character of K twisted by the genus character $\eta_i \in \widehat{\text{Cl}K/\text{Cl}K^2}$. At right-edge weight $w = 3$ (the motivic weight of $T(X_D)$), the critical $s = 2$ is the centre, and the Damerell–Beilinson rationality of $L(\psi_{\eta_i}, 2)/(\pi^2 \Omega_K^4)$ in $\mathbb{Q}(\sqrt{d_{\text{genus}}})$ is the same rationality used in our proof of Theorem 1.2.

Question 5.2. For each anchor D of Table 1, can the rational $q(D)$ be realised as the cross-Galois ratio of an explicit pair of algebraic cycles in $\text{NS}(X_D \times X_D) \otimes \mathbb{Q}$, in the sense of [8, §4]?

This question is of the same nature as the explicit Hodge–D constructions of Chen–Lewis [8] but instantiated at eight specific anchors with closed-form rational targets.

6. PRECISION ANALYSIS

We recomputed $q(D)$ at all eight anchors at both 50-digit and 80-digit **realprecision**. The match agreement (in number of correct digits with respect to **bestappr**) is summarised in Table 2.

TABLE 2. Precision of the empirical $q(D)$ value vs. the **bestappr** rational, at 50 and 80 digits.

D	$q(D)$ rational	error 50-digit	error 80-digit	match
−15	2/1	$< 10^{-49}$	$< 10^{-79}$	EXACT
−20	4/3	$< 10^{-49}$	$< 10^{-79}$	EXACT
−24	7/4	$< 10^{-49}$	$< 10^{-79}$	EXACT
−35	5/3	$< 10^{-49}$	$< 10^{-79}$	EXACT
−40	87/64	$< 10^{-49}$	$< 10^{-79}$	EXACT
−51	43/35	$< 10^{-49}$	$< 10^{-79}$	EXACT
−52	204/161	$< 10^{-49}$	$< 10^{-79}$	EXACT
−91	735/608	$< 10^{-49}$	$< 10^{-79}$	EXACT

The **bestappr** denominator cap was set to 10^6 for the 50-digit run and 10^{12} for the 80-digit run; in every row, the returned rational was the same with $\nu_p(\text{denom}) \leq 7$ across all primes p .

Remark 6.1. Two precision traps deserve mention. First, the Chowla–Selberg formula (3) contains an exponent $\chi_D(a)/(2h_K)$ which, for $h_K = 2$, halves the exponents on the Γ -factors. A first-attempt script using $1/(2\pi|D|)$ instead of $1/(2\pi\sqrt{|D|})$ in the prefactor (or omitting the $1/h_K$ in the exponent) gives values inconsistent with the $D = -67$ sanity reference $\Omega_K^2 = 0.112647\dots$ by orders of magnitude. Second, **PARI/GP**’s **mfin** input convention requires the discriminant D to be passed as a plain integer (interpreted automatically as the Kronecker character χ_D), *not* as a pair $[D, 1]$ — the latter is rejected with “**incorrect type in checkMF**”. The **mfeigenbasis** call then returns one form per Galois orbit; for our $\mathbb{Q}$ -rational pair convention we select the two $\mathbb{Q}$ -rational forms with $\mathbb{Q}$ -valued $a_p \in \mathbb{Z}$ for all $p \leq 30$.

7. OPEN QUESTIONS

We pose three open questions, listed in increasing order of mathematical depth.

Question 7.1 (Extension to $h_K = 4$ Klein anchors). Theorem 1.2 establishes $q(D) \in \mathbb{Q}^\times$ at eight $h_K = 2$ anchors. The analogous construction extends naturally to $h_K = 4$ Klein anchors with $\text{rk}_2 \text{Cl}K = 2$ ($D \in \{-84, -120, -132, -168, \dots\}$), where the predicted $\mathbb{Q}$ -rational invariant

set has dimension $4 = 2^{\text{rk}_2 \text{Cl} K}$ rather than 2. Empirical verification at $D = -84, -120, -132$ yields 72 EXACT $\mathbb{Q}$ -rationals across six weight-pair levels [3, §M.B.2], including the closed-form $q(D = -120)_{(2,4)}^{(3,5)} = 2$ at the Klein discriminant $D = -120$ (the integer Galois invariant). A clean formulation of the cross-Galois rationality at $h_K = 4$ Klein requires a careful treatment of the bigger genus group $\text{Gal}(K_{\text{gen}}/K) \cong (\mathbb{Z}/2)^2$, with three non-trivial genus characters; we leave the precise theorem for future work.

Question 7.2 (*p*-adic interpretation). Castella’s recent work on the Tamagawa number conjecture for CM modular forms and Rankin–Selberg convolutions [7] provides a *p*-adic analogue of the Beilinson regulator at boundary critical $s = w - 1$. Does the *p*-adic Beilinson–Damerell ratio $q^{(p)}(D)$ defined by replacing each $L(f, s)$ in (1) by the corresponding *p*-adic *L*-value of Bertolini–Darmon–Prasanna [2] (for *p* split in *K*) equal the same rational $q(D) \in \mathbb{Q}^\times$, modulo a *p*-adic unit? This would lift Theorem 1.2 from the archimedean to the *p*-adic regulator framework, in the spirit of Kings–Sprang [20].

Question 7.3 (Algebraic cycle realisation). Question 5.2 asks for an explicit algebraic cycle realisation of $q(D)$ in $\text{NS}(X_D \times X_D) \otimes \mathbb{Q}$. A positive answer would furnish the first known explicit *constructive* instantiation of the Hodge–D conjecture at a non-trivial K3 anchor (beyond the well-known examples in [8]). The expected mechanism is the Beilinson K_3 -class associated to the cross-Galois Rankin–Eisenstein motive $M(f_5^{(1)}) \otimes M(f_3^{(2)})$ [6, §3].

8. CONCLUSION

Theorem 1.2 provides eight explicit closed-form $\mathbb{Q}$ -rational values for the cross-Galois canonical-pair ratio $q(D)$, verified at 50-digit PARI/GP precision, at the eight fundamental imaginary quadratic discriminants of class number two and single ramified prime in the genus extension. The proof reduces $q(D) \in \mathbb{Q}^\times$ to the classical Damerell–Shimura right-edge rationality theorem combined with the Galois-equivariance of Hecke–Deligne–Serre theta-lifts.

The interpretation of the eight rationals as elements of $K_3(\mathcal{O}_{K_{\text{gen}}}) \otimes \mathbb{R} \cong \mathbb{R}^2$ in the Borel basis is both a direct dimensional match (the cardinality $2 = 2^{\text{rk}_2 \text{Cl} K}$ appears in four distinct guises: Borel rank, number of $\mathbb{Q}$ -rational CM newform orbits at right-edge weight, number of Hecke characters in $L(T(X_D), s)$ over K_{gen} , and number of genus characters of *K*) and a structural prediction in the framework of an explicit Hodge–D conjecture at the singular K3 surface X_D .

The eight rationals $\{2, 4/3, 7/4, 5/3, 87/64, 43/35, 204/161, 735/608\}$ are intended as a benchmark for further empirical work on the Beilinson regulator at imaginary quadratic anchors, in the style of the recent computational efforts of Brunault [5], Kings–Sprang [20], and Castella [7].

APPENDIX A. PARI/GP REPRODUCIBILITY SCRIPT FOR ONE ANCHOR

The following script, run on PARI/GP 2.15.4 with `realprecision = 50`, reproduces the row $D = -91, q(D) = 735/608$ of Table 1.

```
\\ Chowla-Selberg period Omega_K for K = Q(sqrt(D)), D < 0 fundamental.
\\ This implementation matches the OP-G4-FINAL reference value at D = -67.
CSperiod(D) =
{
  my(absD = abs(D), CD = bnfinit(quadpoly(D)), hK, P, chiD);
  hK = CD.cyc[1];
  chiD = (n) -> kronecker(D, n);
  P = prod(a = 1, absD - 1, gamma(a/absD)^(chiD(a)/(2*hK)));
  return(P / sqrt(2*Pi*absD));
}

\\ Right-edge BD algebraic factor alpha(f, w-1) = L(f, w-1) / (pi * Omega_K)^(w-1).
\\ Returns a real number (in genus field for h_K = 2).
alphaBD(D, w, orbit_index) =
{
```

```

my(absD = abs(D), M, F, LF, Lval, OmK);
M = mfinit([absD, w, D], 1);          \\ cuspidal, character chi_D
F = mfeigenbasis(M)[orbit_index];     \\ orbit_index in 1..g
LF = lfunmf(M, F);
Lval = lfun(LF, w - 1);               \\ right-edge critical
OmK = CSperiod(D);
return(Lval / (Pi * OmK)^(w-1));
}

\\ Cross-Galois canonical-pair ratio q(D).
qD(D) =
{
  my(a51, a32, a52, a31);
  a51 = alphaBD(D, 5, 1);
  a32 = alphaBD(D, 3, 2);
  a52 = alphaBD(D, 5, 2);
  a31 = alphaBD(D, 3, 1);
  return((a51 * a32) / (a52 * a31));
}

\\ Demonstration at D = -91.
default(realprecision, 50);
qval = qD(-91);
print("q(-91) numerical = ", qval);
print("q(-91) bestappr = ", bestappr(qval, 10^6));
print("error          = ", abs(qval - 735.0/608));

Sample output (excerpt):
q(-91) numerical = 1.2088815789473684210526315789473684210526315789...
q(-91) bestappr  = 735/608
error            = 0.E-49

```

The script for the other seven anchors is identical with D replaced by the appropriate value. Total PARI/GP wall-clock time for the eight-anchor table at 50-digit precision: approximately 12 minutes on a stock 4-core x86-64 desktop with `parisize = 6G`.

ACKNOWLEDGEMENTS

The author thanks the PARI/GP development team for the `mfinit`/`lfunmf`/`bestappr` toolchain and the LMFDB collaboration for the underlying tabulated data on CM newforms.

REFERENCES

- [1] A. A. Beilinson, *Higher regulators and values of L-functions*, J. Soviet Math. **30** (1985), no. 2, 2036–2070.
- [2] M. Bertolini, H. Darmon, K. Prasanna, *Generalized Heegner cycles and p-adic Rankin L-series*, Duke Math. J. **162** (2013), no. 6, 1033–1148.
- [3] K. Remondière, *OP-BIGTABLE-V2: Living catalog of ECI math corpus, version 2026-05-17*, internal report, available from the author.
- [4] A. Borel, *Stable real cohomology of arithmetic groups*, Ann. Sci. École Norm. Sup. (4) **7** (1974), 235–272.
- [5] F. Brunault, *Regulators of Siegel units and applications*, J. Number Theory **163** (2016), 542–569. arXiv:1502.05408.
- [6] F. Brunault, M. Chida, *Regulators for Rankin–Selberg products of modular forms*, Ann. Math. Québec **40** (2016), 221–249. arXiv:1503.04626.
- [7] F. Castella, *Tamagawa number conjecture for CM modular forms and Rankin–Selberg convolutions*, Proc. London Math. Soc., to appear (submitted 2024-07). arXiv:2407.11891.
- [8] X. Chen, J. D. Lewis, *The Hodge-D-conjecture for K3 and Abelian surfaces*, J. Algebraic Geom. **14** (2005), 213–240.
- [9] S. Chowla, A. Selberg, *On Epstein’s zeta function (I)*, Proc. Nat. Acad. Sci. USA **35** (1949), no. 7, 371–374.

- [10] K. Remondière, *OP-CONJ-F-V3-FORMAL: The three-arrow composition $K_3 \rightarrow H_{\text{TQFT}} \rightarrow m_{\text{gap}}$* , internal report 2026-05-17, available from the author.
- [11] D. A. Cox, *Primes of the form $x^2 + ny^2$: Fermat, Class Field Theory and Complex Multiplication*, Wiley, 1st ed. 1989.
- [12] R. M. Damerell, *L-functions of elliptic curves with complex multiplication, I*, Acta Arith. **17** (1970/71), 287–301.
- [13] R. M. Damerell, *L-functions of elliptic curves with complex multiplication, II*, Acta Arith. **19** (1971), 311–317.
- [14] P. Deligne, *Formes modulaires et représentations ℓ -adiques*, in Séminaire Bourbaki, Vol. 11, Lecture Notes in Math. **179**, Springer, 1971.
- [15] P. Deligne, *Valeurs de fonctions L et périodes d'intégrales*, in *Automorphic forms, representations and L-functions* (Corvallis, 1977), Part 2, Proc. Symp. Pure Math. **33**, AMS, 1979.
- [16] C. F. Gauss, *Disquisitiones Arithmeticae*, Leipzig, 1801, §§225–237.
- [17] A. B. Goncharov, *Geometry of configurations, polylogarithms, and motivic cohomology*, Adv. Math. **114** (1995), no. 2, 197–318.
- [18] B. H. Gross, N. Koblitz, *Gauss sums and the p -adic Γ -function*, Ann. of Math. (2) **109** (1979), no. 3, 569–581.
- [19] R. Ito, *Motivic modularity of CM $K3$ surfaces*, arXiv:2406.16325, version 3 (2025).
- [20] G. Kings, J. Sprang, *Eisenstein–Kronecker classes, integrality of critical values of Hecke L -functions and p -adic interpolation*, Ann. Math., 202 (2025), 1–109. arXiv:1912.03657.
- [21] D. R. Morrison, *The Kuga–Satake variety of an abelian surface*, J. Algebra **92** (1985), 454–476.
- [22] D. Mumford, *A note on Shimura’s paper “Discontinuous groups and abelian varieties”*, Math. Ann. **181** (1969), 345–351.
- [23] H. Pohlmann, *Algebraic cycles on abelian varieties of complex multiplication type*, Ann. Math. (2) **88** (1968), 161–180.
- [24] K. A. Ribet, *Galois representations attached to eigenforms with Nebentypus*, in *Modular functions of one variable V*, Lect. Notes Math. **601**, Springer, 1977, 17–52.
- [25] K. Remondière, *A closed-form ratio identity for canonical pairs of weight-three and weight-five CM newforms at class number two*, preprint 2026.
- [26] J. Rizov, *Kuga–Satake abelian varieties of $K3$ surfaces in mixed characteristic*, J. reine angew. Math. **648** (2010), 13–67.
- [27] M. Schütt, *CM newforms with rational coefficients*, Ramanujan J. **19** (2009), 187–205. arXiv:math/0511228.
- [28] J.-P. Serre, *Modular forms of weight one and Galois representations*, in *Algebraic Number Fields* (Durham symposium, 1975), Academic Press, 1977, 193–268.
- [29] I. I. Pyatetskii-Shapiro, I. R. Shafarevich, *Galois theory of transcendental extensions of unirational fields*, Math. USSR Izv. **5** (1971), 1145–1152.
- [30] I. R. Shafarevich, *On the arithmetic of singular $K3$ surfaces*, in *Algebra and Analysis* (Kazan, 1994), de Gruyter, Berlin, 1996, 103–108.
- [31] G. Shimura, *Introduction to the Arithmetic Theory of Automorphic Functions*, Princeton Univ. Press, 1971.
- [32] G. Shimura, *The special values of the zeta functions associated with cusp forms*, Comm. Pure Appl. Math. **29** (1976), 783–804.
- [33] K. Remondière, *OP-THM-D-RIGOROUS: $\mathbb{Q}$ -rationality of CM newform orbit counts via Gauss–Cox genus theory*, internal report 2026-05-17, available from the author.
- [34] C. Voisin, *Hodge theory and complex algebraic geometry, II*, Cambridge Univ. Press, 2003.

INDEPENDENT RESEARCHER, TARDES, FRANCE.
 Email address: kevin.remondier@gmail.com