A modular Lyapunov bound for finite-rank type-II$_\infty$ crossed-product algebras

with a saturation conjecture and an analog-Hawking falsifier

Kévin Remondière

Independent researcher, Tarbes, France

Submitted to Letters in Mathematical Physics

Preprint: May 5, 2026 (v2)

Abstract

Let $\mathcal{M}_R \otimes B(\mathcal{H}_n)$ be a finite-rank truncation of the type-II$_\infty$ crossed-product von Neumann algebra associated with an observer's causal wedge at Bisognano–Wichmann KMS inverse temperature $\beta = 2\pi$, and let K_R be its modular Hamiltonian. We prove that the modular Krylov (Lanczos) complexity $\mathcal{C}_K^{\mathrm{mod}}(s)$ of any operator $\mathcal{O}$ under the modular flow $\sigma_s = e^{iK_R s} \cdot e^{-iK_R s}$ satisfies the bound

$$\lambda_K^{\mathrm{mod}} \;:=\; \limsup_{s\to\infty} \frac{1}{s} \log \mathcal{C}_K^{\mathrm{mod}}(s) \;\leq\; \frac{2\pi}{\beta} \;=\; 1,$$

matching the Maldacena–Shenker–Stanford chaos bound *kinematically* on the truncation. The proof combines the universal operator-growth inequality of Parker, Cao, Avdoshkin, Scaffidi, and Altman [**?**] with the Bisognano–Wichmann theorem [**?**] and uses no large-N, holography, or chaos hypothesis. The converse statement – that saturation $\lambda_K^{\mathrm{mod}} = 2\pi/\beta$ characterises type-II$_\infty$ chaos – is shown to be *false in general*: free-QFT and integrable counter-examples (Avdoshkin–Dymarsky [**?**], Camargo–Jahnke–Kim–Nishida [**?**], Sreeram–Kannan–Modak–Aravinda [**?**]) attain the same kinematic ceiling without underlying chaos. We therefore retain saturation only as a conjectural *sufficient condition* (Conjecture **??**). A concrete one-sided falsifier is then derived: for the Kolobov–Golubkov–Muñoz de Nova–Steinhauer [**?**] analog black-hole BEC at $T_H = 0.35 \pm 0.10$ nK (Muñoz de Nova et al. [**?**]), the bound predicts $\Gamma_{\mathrm{meas}} \leq 2\pi k_B T_H/\hbar = (288 \pm 82)\,\mathrm{s}^{-1}$ for any exterior-phonon temporal correlation rate. We compare with the concurrent boundary-area-operator construction of Vardian [**?**] (AdS/CFT) and note that our scope (de Sitter static patch / generic type-II$_\infty$ truncation) is complementary.

1 Introduction

The Maldacena–Shenker–Stanford (MSS) bound on chaos [**?**]

$$\lambda_L \;\leq\; \frac{2\pi k_B T}{\hbar} \;=\; \frac{2\pi}{\beta} \tag{1}$$

limits the OTOC Lyapunov exponent of any thermal large-N quantum system, with maximally chaotic systems (SYK, near-extremal black holes) saturating it. Parker, Cao, Avdoshkin, Scaffidi, and Altman [**?**] subsequently introduced the universal operator-growth hypothesis: in chaotic systems the Lanczos coefficients b_n of any few-body operator grow linearly, $b_n \sim \alpha n$, and the Krylov (Lanczos) complexity satisfies the rigorous inequality

$$\lambda_K \;\leq\; 2\alpha, \qquad \mathcal{C}_K(t) \;\leq\; A\cosh(2\alpha t), \tag{2}$$

sharper than (**??**) and free of large-N assumptions. For thermal correlators in d-dimensional CFT one has the universal slope $\alpha = \pi/\beta$, recovering the MSS rate as a kinematic upper envelope.

In parallel, Witten [**?**] and Chandrasekaran–Longo–Penington–Witten [**?**] showed that gravitational observable algebras with quantum reference frames (QRFs) are type-II_∞ crossed products $\mathcal{A}_{\mathrm{II}_\infty} = \mathcal{A}_{\mathrm{III}_1} \rtimes_{\sigma^\phi} \mathbb{R}$, restoring a semifinite trace and finite generalised entropy. De Vuyst, Eccles, Höhn, and Kirklin [**?**, **?**] established observer-dependent gravitational entropy in this framework, and Faulkner–Speranza [**?**] derived the generalised second law in the crossed-product setting.

Aim of this paper. We prove a clean kinematic statement: on *any* finite-rank truncation of a type-II_∞ crossed product at the canonical Bisognano–Wichmann temperature $\beta = 2\pi$, the modular Krylov complexity is bounded by the MSS rate (Theorem **??**, §**??**). The proof combines the Parker–Cao–Avdoshkin–Scaffidi–Altman inequality with the Bisognano–Wichmann KMS condition and is independent of any large-N or chaos assumption.

What this paper does *not* prove. We do *not* prove the converse: saturation of the bound does not characterise type-II_∞ chaos. Free-QFT lattice models [**?**], free and interacting bounded-power-spectrum scalar fields [**?**], and operator/initial-state-dependent counter-examples [**?**] all attain the same kinematic ceiling without exhibiting MSS-saturating chaos. The earlier ECI "modular shadow" draft (v1, 2026-05-04) presented saturation as the central object; this revision (v2) downgrades saturation to a conjecture (Conjecture **??**, §**??**) that requires additional dynamical input beyond the type-II_∞ structure.

Concurrent work. While preparing this manuscript we became aware of Vardian [**?**] (arXiv:2602.02675, February 2026), "Modular Krylov Complexity as a Boundary Probe of Area Operator and Entanglement Islands", which uses modular Lanczos coefficients to reconstruct the QES area operator from boundary dynamics in AdS/CFT. Vardian's analysis is complementary to ours: it operates in the holographic AdS/CFT setting and targets the area-operator spectrum, whereas we work intrinsically in the finite-rank type-II_∞ algebra at any KMS temperature, and aim at a generic Lyapunov bound applicable to de Sitter and BEC analog systems. We compare scopes in §**??**.

Organization. §**??** fixes the algebraic setup (type-II_∞ crossed product, finite-rank truncation, modular Krylov complexity). §**??** states and proves the bound theorem. §**??** downgrades the saturation step to a conjecture and reviews the free-QFT counter-examples. §**??** applies the bound to the Kolobov–Steinhauer 2021 BEC analog black hole, deriving the falsifier $\Gamma_{\mathrm{meas}} \leq (288 \pm 82)\,\mathrm{s}^{-1}$. §**??** compares with Vardian and discusses scope. §**??** lists open problems.

2 Setup: type-II$_\infty$ crossed product, finite-rank truncation, and modular Krylov complexity

2.1 The crossed-product algebra

Let $\mathcal{A}_{\mathrm{III}_1}$ be the type-III$_1$ von Neumann algebra of observables in a relativistic QFT restricted to a Rindler wedge $\mathcal{W}$, and let ϕ be the vacuum KMS state at the Bisognano–Wichmann inverse temperature $\beta_{\mathrm{BW}} = 2\pi$. Following Witten [**?**] and CLPW [**?**], the crossed product with the modular automorphism group

$$\mathcal{A}_{\mathrm{II}_\infty} \;=\; \mathcal{A}_{\mathrm{III}_1} \rtimes_{\sigma^\phi} \mathbb{R} \tag{3}$$

is type-II$_\infty$, admits a faithful semifinite trace τ, and hosts a well-defined dressed modular Hamiltonian K_ϕ generating the modular flow. The Bisognano–Wichmann theorem [**?**] guarantees $K_\phi = 2\pi K_{\mathrm{boost}}$ where K_{boost} is the boost generator of the wedge.

2.2 Finite-rank truncation

Working with the full type-II$_\infty$ algebra requires care: K_ϕ is unbounded and the modular flow is generally non-trace-preserving. For our bound theorem we work on the *finite-rank truncation*

$$\mathcal{M}_R \otimes B(\mathcal{H}_n) \;:=\; P_n\, \mathcal{A}_{\mathrm{II}_\infty}\, P_n \tag{4}$$

where P_n is the spectral projector of K_ϕ onto the modular energy band $|K_\phi| \le \Lambda_n$ with $\dim(\operatorname{ran} P_n) = n$. This truncation preserves the $\beta = 2\pi$ KMS condition (it is the restriction of the GNS representation to a finite-dimensional subspace on which K_ϕ acts as a bounded self-adjoint operator) and inherits the semifinite trace from the type-II$_\infty$ parent.

Remark 1. *The Bisognano–Wichmann property survives the truncation in the following sense: although $\mathcal{M}_R \otimes B(\mathcal{H}_n)$ is not itself a wedge algebra, its modular flow restricted to operators of energy $\le \Lambda_n$ agrees with the boost flow on those modes, because the Bisognano–Wichmann modular operator commutes with the spectral projection P_n. This is the only structural fact about the truncation that we will need.*

2.3 Modular Krylov complexity

For an operator $\mathcal{O} \in \mathcal{M}_R \otimes B(\mathcal{H}_n)$, its modular evolution is

$$\mathcal{O}(s) \;=\; e^{iK_\phi s}\, \mathcal{O}\, e^{-iK_\phi s}, \qquad s \in \mathbb{R}. \tag{5}$$

Equip the truncation with the KMS Gel'fand–Naimark–Segal inner product $(A, B)_\phi := \tau(A^* B)$ (well-defined because τ is finite on the truncation). The Lanczos algorithm applied to the modular Liouvillian $\mathcal{L} := [K_\phi, \cdot\,]$ produces an orthonormal Krylov basis $\{|\mathcal{O}_n\rangle\}$ with tridiagonal coefficients $\{b_n\}_{n\ge1}$, and the modular Krylov complexity is

$$\mathcal{C}_K^{\mathrm{mod}}(s) \;:=\; \sum_{n\ge0} n\, |\varphi_n(s)|^2, \qquad \mathcal{O}(s) \;=\; \sum_{n\ge0} \varphi_n(s)\, \mathcal{O}_n, \tag{6}$$

following the modular construction of Caputa–Magán–Patramanis–Tonni [**?**] who showed $\lambda_L^{\mathrm{mod}} = 2\pi$ in dimensionless modular time for 2D CFTs.

2.4 Modular Lyapunov exponent

The *modular Lyapunov exponent* of $\mathcal{O}$ is

$$\lambda_K^{\mathrm{mod}}(\mathcal{O}) \;:=\; \limsup_{s\to\infty} \frac{1}{s} \log \mathcal{C}_K^{\mathrm{mod}}(s), \tag{7}$$

when this limit is finite. Our central object is the supremum of $\lambda_K^{\mathrm{mod}}(\mathcal{O})$ over operators $\mathcal{O}$ in a dense $*$-subalgebra of the truncation; the bound theorem (next section) controls this supremum.

3 The modular Lyapunov bound: a provable theorem

Theorem 1 (Modular Lyapunov bound on finite-rank type-II$_\infty$ truncations)**.** *Let $\mathcal{M}_R \otimes B(\mathcal{H}_n)$ be the finite-rank truncation (**??**) of the type-II$_\infty$ crossed product (**??**) at Bisognano–Wichmann KMS inverse temperature $\beta = 2\pi$, equipped with the KMS–GNS inner product $(A, B)_\phi = \tau(A^*B)$ and the modular flow σ_s of (**??**). For every $\mathcal{O} \in \mathcal{M}_R \otimes B(\mathcal{H}_n)$ with finite modular two-point function, the Lanczos coefficients $b_n(\mathcal{O})$ produced by the Lanczos algorithm with respect to $(\,\cdot\,,\,\cdot\,)_\phi$ and Liouvillian $\mathcal{L} = [K_\phi, \,\cdot\,]$ satisfy*

$$b_n(\mathcal{O}) \;\le\; n \cdot \frac{\pi}{\beta} + o(n), \qquad n \to \infty, \tag{8}$$

and consequently the modular Krylov complexity satisfies

$$\mathcal{C}_K^{\mathrm{mod}}(s) \;\le\; A\,\cosh\!\big(2b_1(\mathcal{O})s\big), \qquad \lambda_K^{\mathrm{mod}}(\mathcal{O}) \;\le\; \frac{2\pi}{\beta} \;=\; 1, \tag{9}$$

where $A > 0$ depends on $\|\mathcal{O}\|_\phi$ but not on s.

Proof sketch (full proof in Appendix A of an extended version). The argument has three steps.

Step 1 (Lanczos–Liouville inequality on KMS–GNS). The Lanczos algorithm on the KMS–GNS inner product produces a tridiagonal representation of $\mathcal{L}$:

$$\mathcal{L}\,\mathcal{O}_n \;=\; b_{n+1}\,\mathcal{O}_{n+1} + b_n\,\mathcal{O}_{n-1}$$

(with $a_n = 0$ for hermitian $\mathcal{L}$ on the KMS state, by KMS reflection symmetry). Parker et al. [**?**, §II.B] prove that for any such tridiagonal evolution

$$\mathcal{C}_K(s) \;\le\; A\,\cosh(2\,b_{\max}\,s), \qquad b_{\max} := \sup_n \frac{b_n}{n}.$$

This is a purely algebraic inequality from operator norms on the tridiagonal matrix; nothing about chaos enters at this step.

Step 2 (Bisognano–Wichmann moment bound). The Lanczos coefficients b_n are determined by the moments $M_{2k} = (\mathcal{O}, \mathcal{L}^{2k}\mathcal{O})_\phi$ of the modular two-point function via the Hankel determinant formula

$$b_n^2 \;=\; \frac{\det H_n \cdot \det H_{n-2}}{(\det H_{n-1})^2}, \qquad (H_k)_{ij} = M_{2(i+j)}.$$

On the truncation, by Remark **??** the modular flow agrees with the boost flow on $\leq \Lambda_n$ modes, and the boost two-point function in the Bisognano–Wichmann KMS state has the universal hyperbolic decay

$$C(s) \;=\; (2\sinh(\pi s/\beta))^{-\Delta} \cdot (\text{spatial factors})$$

with $\beta = 2\pi$. The moments of this $\sinh^{-\Delta}$ kernel have the universal asymptotic form $M_{2k} \sim C_\Delta\,(2\pi k/\beta)^{2k}$ (standard hypergeometric expansion), which by the Hankel–Lanczos inverse map yields exactly $b_n \leq n \cdot \pi/\beta + o(n)$.

Step 3 (combine). Substituting Step 2 into Step 1 gives (**??**). The Lyapunov exponent reads off the cosh growth as $\lambda_K^{\mathrm{mod}} \leq 2 \cdot (\pi/\beta) = 2\pi/\beta = 1$ in $\beta = 2\pi$ units. □

Remark 2. *Theorem* **??** *is purely kinematic. It uses (i) the Bisognano–Wichmann KMS structure (algebraic input) and (ii) the Parker–Cao–Avdoshkin–Scaffidi–Altman tridiagonal inequality (operator-norm input). No chaos hypothesis, no holography, no large-N assumption is invoked.*

Remark 3. *For 2D CFT in the Bisognano–Wichmann KMS state, Caputa–Magán–Patramanis–Tonni [**?**] compute $b_n^{\mathrm{CFT}} = n \cdot \pi/\beta$ exactly, saturating* (**??**) *as an equality. Thus the bound is tight on the CFT class. The full type-II$_\infty$ saturation question (do all dressed type-II$_\infty$ algebras saturate?) is open – see §**??**.*

Corollary 2 (MSS bound recovery)**.** *For the truncation $\mathcal{M}_R \otimes B(\mathcal{H}_n)$ at any KMS inverse temperature β, the modular Lyapunov exponent satisfies $\lambda_K^{\mathrm{mod}} \leq 2\pi/\beta$, which is the MSS rate* (**??**)*.*

Proof. Replace $\beta = 2\pi$ by general β in the proof of Theorem **??**; the hyperbolic moment kernel becomes $(2\sinh(\pi s/\beta))^{-\Delta}$ and the universal Lanczos slope becomes π/β, giving the MSS rate. □

4 Saturation as a conjecture: free-QFT counter-examples to the converse

The original ECI "modular shadow" draft [**?**] proposed the stronger statement that saturation $\lambda_K^{\mathrm{mod}} = 2\pi/\beta$ is a structural fingerprint of type-II$_\infty$. On audit (A11, 2026-05-05), this converse fails: there exist non-chaotic systems that saturate the kinematic ceiling. We collect three counter-examples and then state what survives as a conjecture.

Counter-example 1 (free-scalar lattice; Avdoshkin–Dymarsky 2019). Avdoshkin and Dymarsky [**?**] show that operator growth under imaginary-time evolution in 1D lattice models can attain exponential Lanczos-coefficient growth even in integrable regimes, with a 1D superexponential frequency suppression ensuring that "one-dimensional systems are special". This means linear b_n and exponential $\mathcal{C}_K$ are *not* signatures of chaos; they appear in integrable lattice systems as well.

Counter-example 2 (free & interacting massive scalar QFT; Camargo et al. 2023). Camargo, Jahnke, Kim, and Nishida [**?**] compute Krylov complexity in free and weakly interacting massive scalar field theories and find non-chaotic regimes where the

Lanczos coefficients exhibit the same linear envelope as in maximally chaotic systems, with mass and UV cutoff modifying the prefactor but preserving the kinematic form. Their work refutes the reading that linear $b_n \to 2\pi/\beta$ implies type-II_∞ MSS-saturating chaos.

Counter-example 3 (initial-operator dependence; Sreeram et al. 2025). Sreeram, Kannan, Modak, and Aravinda [**?**] (PRE 112 L032203) prove that Krylov complexity depends monotonically on the inverse participation ratio (IPR) of the initial operator in the Hamiltonian eigenbasis, and that this dependence persists in fully chaotic regimes. Saturation of λ_K^{mod} is therefore not a state- or operator-independent invariant of the algebra: it varies even within a fixed maximally-chaotic dynamics.

What survives as a conjecture. The strong claim of v1, "saturation $\Leftrightarrow$ type-II_∞", is refuted by counter-examples 1–3. What remains plausible is the weaker direction: type-II_∞ at $\beta = 2\pi$ is *sufficient* for the equality $b_n = n\pi/\beta$, even if not necessary.

Conjecture 1 (Modular saturation for dressed type-II_∞ algebras)**.** [CONJECTURE; v2 downgraded.] *Let $\mathcal{A}_{\mathrm{II}_\infty}$ be a type-II_∞ crossed product (**??**) arising from a QRF-dressing of a wedge or static-patch algebra, with the canonical $\beta = 2\pi$ KMS state. Then on a sequence of finite-rank truncations $\mathcal{M}_R \otimes B(\mathcal{H}_n)$, $n \to \infty$, the Lanczos coefficients of any few-mode operator $\mathcal{O}$ in a dense subalgebra satisfy*

$$b_n(\mathcal{O}) \xrightarrow[n\to\infty]{} n \cdot \frac{\pi}{\beta},$$

*saturating the bound (**??**) and yielding $\lambda_K^{\mathrm{mod}}(\mathcal{O}) = 2\pi/\beta$ in the large-truncation limit.*

Remark 4. *Conjecture **??** is consistent with the 2D-CFT computation of Caputa et al. [**?**] (Remark **??**), with the SYK Krylov saturation of Parker et al. [**?**], and with the DSSYK / sine-dilaton holographic checks of Heller, Papalini, and Schuhmann [**?**]. We do not propose it as a theorem; the counter-examples above show that the converse fails, and even a one-sided proof of the sufficient direction is beyond current methods.*

Erratum: Faulkner–Speranza identification. The v1 draft of this work attributed to Faulkner and Speranza [**?**] the identification $S_{\mathrm{gen}} \leftrightarrow \mathcal{C}_K^{\mathrm{mod}}$. On rereading, FS24 derive the crossed-product generalised entropy and prove the GSL via Wall monotonicity; they do *not* explicitly identify S_{gen} with the modular Krylov complexity. The bridge $S_{\mathrm{gen}} \leftrightarrow \mathcal{C}_K^{\mathrm{mod}}$ is a working conjecture of the ECI programme, not a theorem of FS24. We thank the A11/A19 audit for catching this over-attribution.

5 Application: a falsifier on Kolobov 2021 BEC analog Hawking radiation

We now derive a concrete observable consequence of Theorem **??** for the analog black-hole BEC platform of Kolobov, Golubkov, Muñoz de Nova, and Steinhauer [**?**] (Nat. Phys. 17, 362, 2021).

5.1 Hawking temperature and the predicted bound

Muñoz de Nova et al. [?] (Nature 569, 688, 2019) measured the analog Hawking temperature in the BEC sonic-horizon platform as

$$T_H \;=\; 0.35 \pm 0.10\,\mathrm{nK}, \tag{10}$$

implying the Bisognano–Wichmann KMS rate

$$\boxed{\Gamma_{\mathrm{bound}} \;:=\; \frac{2\pi k_B T_H}{\hbar} \;=\; (288 \pm 82)\,\mathrm{s}^{-1}} \tag{11}$$

($k_B/\hbar = 1.31 \times 10^{11}\,\mathrm{rad\,s^{-1}\,K^{-1}}$).

By Theorem **??** applied to the (assumed) finite-rank type-II_∞ truncation of the exterior phonon algebra at the analog horizon, any temporal correlator of exterior phonons must have its envelope decaying with rate $\leq \Gamma_{\mathrm{bound}}$:

$$\Gamma_{\mathrm{meas}} \;\leq\; \Gamma_{\mathrm{bound}} \;=\; (288 \pm 82)\,\mathrm{s}^{-1}, \tag{12}$$

where Γ_{meas} is the exponential-decay rate of any of: (i) the second-order coherence $g^{(2)}(\tau)$; (ii) the two-point density correlator $G^{(2)}(z_1, z_2, t)$ envelope; (iii) the Hawking-flux ramp-up rate.

5.2 Why this is a clean falsifier

The Kolobov 2021 dataset [?] contains time-resolved density correlations at six temporal points across a 124-day, 97,000-repetition acquisition: the temporal envelope of $G^{(2)}(z_1, z_2, t)$ can be extracted directly from the existing data without new experimental runs.

Proposition 3 (BEC bound falsifier)**.** *If a re-analysis of Kolobov 2021 data yields* $\Gamma_{\mathrm{meas}} > 1.5 \times \Gamma_{\mathrm{bound}} = 432\,\mathrm{s}^{-1}$ *at* 5σ*, then the type-*II_∞ *finite-rank scaffolding underlying Theorem* **??** *is falsified for the analog-BEC exterior phonon algebra.*

Robustness to integrability counter-examples. Crucially, the bound (**??**) is the BOUND form, not the saturation form: it is therefore robust to the free-QFT / integrability counter-examples of §**??** [**?**, **?**, **?**]. Even if the BEC phonon system is effectively integrable on the relevant timescales, the bound still holds; only an observed *violation* would refute the type-II_∞ kinematic prefactor.

Cross-platform redundancy. Independent platforms can cross-check (**??**): polariton-condensate analog black-hole mergers (Solnyshkov–Paquelier–Balmisse–Malpuech 2026, arXiv:2603.01664 [**?**]) and water-tank vortex QNM decay rates [**?**] provide independent substrates. A UV-finite volume-law entanglement negativity (Chandran–Fischer 2026, arXiv:2604.02075 [**?**]) likewise yields an entanglement-growth rate bounded by (**??**) via Theorem **??**.

6 Discussion: comparison with Vardian (2026) and scope

Vardian's concurrent boundary-area-operator construction. Vardian [**?**] (arXiv:2602.02675, February 2026, single-author hep-th) reconstructs the QES area operator from boundary modular Lanczos coefficients in AdS/CFT, using operator-algebra quantum error correction. Specifically: from boundary modular dynamics, the spectrum of the boundary modular Hamiltonian is extracted via the Lanczos algorithm; its central component is identified with the area operator of the dual quantum extremal surface. Applications discussed include diagnosing island formation and the Page transition for evaporating black holes.

Comparison: complementary scopes. Vardian's work and ours overlap in their use of modular Lanczos coefficients on a type-II_∞-like algebra, but the scopes differ along three axes:

- **Holographic context.** Vardian works in AdS/CFT and uses holographic boundary–bulk duality explicitly; we work intrinsically in the finite-rank truncation without invoking AdS/CFT and our results apply to any wedge or static-patch algebra (de Sitter, Rindler, BEC analog horizon).
- **Object of study.** Vardian targets the area operator (a specific element of the modular Hamiltonian's spectrum); we target the kinematic Lyapunov ceiling on Krylov growth (an envelope statement on all operators).
- **Claim type.** Vardian provides a constructive boundary–bulk dictionary; we provide a one-sided inequality with concrete falsifier on a laboratory analog system.

The two works together suggest that modular Krylov techniques are becoming a standard tool for type-II_∞ algebras, with Vardian probing the holographic side and our bound theorem covering the non-holographic side (de Sitter, BEC, and generic truncations).

Scope of Theorem ??. Our theorem applies to any finite-rank truncation of a type-II_∞ crossed product at canonical KMS temperature. This covers:

1. Rindler wedge of a free QFT (where the bound is tight by Caputa et al. [**?**]);
2. BTZ exterior in AdS_3 (the holographic paradigm);
3. de Sitter static patch [**?**], with $\beta = 2\pi/H$;
4. BEC and other analog-Hawking systems with a sonic horizon (Kolobov 2021, Solnyshkov 2026);
5. the SYK / DSSYK / sine-dilaton family at large N [**?**, **?**].

It does *not* apply directly to:

1. Type-III_1 algebras at finite N without a QRF dressing (no semifinite trace, modular flow ergodic);
2. Cosmological algebras (Bianchi IX, FRW) where the type classification is open;
3. Type-I or II_1 models without boost geometry (no Bisognano–Wichmann input).

Modular vs physical Lyapunov exponents. The bound (**??**) is on the modular Lyapunov exponent λ_K^{mod}, defined via modular flow. Translating to the physical Lyapunov exponent $\lambda_L^{\mathrm{OTOC}}$ requires the physical Hamiltonian to align with the modular Hamiltonian under the Connes–Rovelli [**?**] thermal-time hypothesis. For boost-symmetric wedges this alignment is automatic (Bisognano–Wichmann); for general dynamical settings it is an additional input.

7 Outlook

1. **Lifting the truncation.** Theorem **??** is stated on the finite-rank truncation $\mathcal{M}_R \otimes B(\mathcal{H}_n)$. A genuine type-$\mathrm{II}_\infty$ statement requires a uniform-in-n control of b_n and a passage to the inductive-limit algebra. The Hankel-determinant moment estimate of Step 2 in our proof sketch is uniform in n on the BW class, but the operator $\mathcal{O}$ must lie in a dense $*$-subalgebra of the type-II_∞ parent. We expect the lifted statement to hold; a complete proof is left for future work.

2. **de Sitter static patch.** CLPW [**?**] establish the type-II_∞ structure of the dS static patch with $\beta = 2\pi/H$ (Gibbons–Hawking temperature). By Theorem **??**, the modular Lyapunov exponent of any few-mode operator on the dS static patch is bounded by H (the Hubble rate). A concrete observable (e.g. a primordial-tensor-mode decoherence rate) testing this bound would be of cosmological interest.

3. **Bianchi IX and homogeneous chaotic cosmologies.** The mixmaster Bianchi IX universe exhibits chaotic dynamics with a known modular structure on the kasner-segment wedge algebra. Whether this algebra is type-II_∞ and whether its modular Krylov exponent saturates (**??**) is open.

4. **Connes–Størmer entropy.** The Connes–Størmer entropy on type-II_1 reductions of our truncation provides an alternative diagnostic (see Connes–Rovelli [**?**]); whether it inherits the same kinematic ceiling is an open question.

5. **Bridge $S_{\mathrm{gen}} \leftrightarrow \mathcal{C}_K^{\mathrm{mod}}$.** The bridge between the FS24 generalised entropy and modular Krylov complexity remains a working conjecture of the ECI programme. A rigorous derivation would close the loop between Theorem **??** and the Faulkner–Speranza GSL.

6. **Relation to Vardian's area-operator reconstruction.** The boundary area operator of Vardian [**?**] sits in the central part of the modular Hamiltonian. Its Krylov-Lanczos signature should sit at a specific level of our truncation; making this precise would unify the two approaches.

Acknowledgments

This work was extracted and reworked from the ECI framework (v6.0.53, Zenodo record `10.5281/zenodo.20030684`). The split between the (provable) bound and the (conjectural) saturation, and the identification of the Avdoshkin–Dymarsky / Camargo et al. / Sreeram et al. counter-examples, owe to the A11 audit (2026-05-05). The discovery of Vardian [**?**] as concurrent independent work was made via the same audit, with arXiv

ID 2602.02675 live-verified. Erratum on the Faulkner–Speranza over-attribution (§**??**) follows the A11 cross-check. We thank [collaborators] for discussions.

References

[1] J. Maldacena, S. H. Shenker, and D. Stanford, "A bound on chaos," *JHEP* **08** (2016) 106, arXiv:1503.01409 [hep-th].

[2] D. E. Parker, X. Cao, A. Avdoshkin, T. Scaffidi, and E. Altman, "A Universal Operator Growth Hypothesis," *Phys. Rev. X* **9** (2019) 041017, arXiv:1812.08657 [cond-mat.str-el].

[3] P. Caputa, J. M. Magán, D. Patramanis, and E. Tonni, "Krylov complexity of modular Hamiltonian evolution," *Phys. Rev. D* **109** (2024) 086004, arXiv:2306.14732 [hep-th].

[4] E. Witten, "Gravity and the crossed product," *JHEP* **10** (2022) 008, arXiv:2112.12828 [hep-th].

[5] V. Chandrasekaran, R. Longo, G. Penington, and E. Witten, "An algebra of observables for de Sitter space," *JHEP* **02** (2023) 082, arXiv:2206.10780 [hep-th].

[6] T. Faulkner and A. J. Speranza, "Gravitational algebras and the generalized second law," *JHEP* **11** (2024) 099, arXiv:2405.00847 [hep-th].

[7] J. De Vuyst, S. Eccles, P. A. Höhn, and J. Kirklin, "Crossed products and quantum reference frames: on the observer-dependence of gravitational entropy," *JHEP* **07** (2025) 063, arXiv:2412.15502 [hep-th].

[8] J. De Vuyst, S. Eccles, P. A. Höhn, and J. Kirklin, "Gravitational entropy is observer-dependent," *JHEP* **07** (2025) 146, arXiv:2405.00114 [hep-th].

[9] M. P. Heller, A. Papalini, and T. Schuhmann, "Krylov spread complexity as holographic complexity beyond JT gravity," *Phys. Rev. Lett.* **135** (2025) 151602, arXiv:2412.17785 [hep-th].

[10] J. J. Bisognano and E. H. Wichmann, "On the duality condition for a Hermitian scalar field," *J. Math. Phys.* **17** (1976) 303.

[11] A. Connes and C. Rovelli, "Von Neumann algebra automorphisms and time-thermodynamics relation in generally covariant quantum theories," *Class. Quantum Grav.* **11** (1994) 2899–2917.

[12] N. Vardian, "Modular Krylov Complexity as a Boundary Probe of Area Operator and Entanglement Islands," arXiv:2602.02675 [hep-th] (February 2026).

[13] A. Avdoshkin and A. Dymarsky, "Euclidean operator growth and quantum chaos," *Phys. Rev. Research* **2** (2020) 043234, arXiv:1911.09672 [cond-mat.stat-mech].

[14] H. A. Camargo, V. Jahnke, K.-Y. Kim, and M. Nishida, "Krylov Complexity in Free and Interacting Scalar Field Theories with Bounded Power Spectrum," *JHEP* **05** (2023) 226, arXiv:2212.14702 [hep-th].

[15] Sreeram PG, J. Bharathi Kannan, R. Modak, and S. Aravinda, "Dependence of Krylov complexity on the initial operator and state," *Phys. Rev. E* **112** (2025) L032203, arXiv:2503.03400 [quant-ph].

[16] V. I. Kolobov, K. Golubkov, J. R. Muñoz de Nova, and J. Steinhauer, "Observation of stationary spontaneous Hawking radiation and the time evolution of an analogue black hole," *Nat. Phys.* **17** (2021) 362–367, arXiv:1910.09363 [gr-qc].

[17] J. R. Muñoz de Nova, K. Golubkov, V. I. Kolobov, and J. Steinhauer, "Observation of thermal Hawking radiation at the Hawking temperature in an analogue black hole," *Nature* **569** (2019) 688–691, arXiv:1809.00913 [gr-qc].

[18] D. D. Solnyshkov, I. Paquelier, A. Balmisse, and G. Malpuech, "Polariton-condensate analog black-hole merger" (working title), arXiv:2603.01664 (2026).

[19] A. Chandran and U. R. Fischer, "UV-finite volume-law entanglement negativity" (working title), arXiv:2604.02075 (2026).

[20] T. Torres, S. Patrick, A. Coutant, M. Richartz, E. W. Tedford, and S. Weinfurtner, "Rotational superradiant scattering in a vortex flow," *Nat. Phys.* **13** (2017) 833–836.

[21] [ECI author(s)], "Entanglement–Complexity–Information: a phenomenological architecture for quantum gravity phenomenology (v6.0.53)," Zenodo record `10.5281/zenodo.20030684` (2026).