The lemniscate period of $\mathbb{Q}(i)$ and full rationality of Damerell ladders for CM weight-5 newforms

Kevin Remondiere

Preprint, May 6, 2026

Abstract

For the CM newform $f = 4.5.b.a$ of weight 5, level 4, with complex multiplication by $K = \mathbb{Q}(i)$, we show computationally (PARI/GP, 80-digit precision) that all four normalized critical L-values $\alpha_m = L(f,m) \cdot \pi^{4-m}/\Omega_K^4$ lie in $\mathbb{Q}$:

$$(\alpha_1, \alpha_2, \alpha_3, \alpha_4) = \left(\tfrac{1}{10}, \tfrac{1}{12}, \tfrac{1}{24}, \tfrac{1}{60}\right).$$

Here $\Omega_K = \Gamma(1/4)^2/(2\sqrt{2\pi})$ is the Chowla–Selberg lemniscate period of $\mathbb{Q}(i)$. In contrast, for the CM weight-5 newforms $27.5.b.a$ and $12.5.c.a$ with complex multiplication by $\mathbb{Q}(\omega)$ ($\omega = e^{2\pi i/3}$), the odd-index values α_1, α_3 lie in $\mathbb{Q}(\sqrt{3}) \setminus \mathbb{Q}$, while only $\alpha_2, \alpha_4 \in \mathbb{Q}$.

The Ω-independent ratio $R(f) = \pi \cdot L(f,1)/L(f,2)$, which cancels the period, equals $6/5 \in \mathbb{Q}$ for $f = 4.5.b.a$ but equals $3\sqrt{3}$ or $3\sqrt{3}/2$ (both irrational) for the $\mathbb{Q}(\omega)$-CM forms. We identify the structural reason: the Chowla–Selberg period $\Omega_{\mathbb{Q}(i)} = \Gamma(1/4)^2/(2\sqrt{2\pi})$ is free of $\sqrt{3}$, while $\Omega_{\mathbb{Q}(\omega)} = \Gamma(1/3)^3/(2\pi\sqrt{3})$ carries an explicit $1/\sqrt{3}$ factor that propagates to odd-indexed critical values via the unit-group parity of the Hecke Größencharacter. We propose a conjecture extending this dichotomy to all class-number-1 imaginary quadratic fields. Additional PARI 80-digit computations for the CM newforms $7.5.b.a$ ($K = \mathbb{Q}(\sqrt{-7})$) and $11.5.b.a$ ($K = \mathbb{Q}(\sqrt{-11})$) find $R(f) = \frac{21}{32}\sqrt{7}$ and $R(f) = \frac{11}{15}\sqrt{11}$, respectively, corroborating Conjecture 3.3(c) for $d \in \{7, 11\}$.

1 Introduction

The algebraicity of special values of L-functions of CM modular forms at critical integers is a classical theme, going back to Damerell [2, 3] and Shimura [4]. For a CM newform f of weight k with complex multiplication by an imaginary quadratic field K, Damerell proved that the normalized values

$$\alpha_m(f) = \frac{L(f,m) \cdot \pi^{k-1-m}}{\Omega_K^{k-1}}, \qquad 1 \le m \le k-1, \tag{1}$$

are algebraic, where Ω_K is the Chowla–Selberg period of K [1]. The Shimura refinement [4] places α_m in the ray class field of K.

The natural question of *when* $\alpha_m \in \mathbb{Q}$ for all critical values simultaneously does not appear to be addressed in the existing literature. For weight $k = 5$ there are four critical integers $m \in \{1, 2, 3, 4\}$. We call $(\alpha_1, \alpha_2, \alpha_3, \alpha_4)$ the *Damerell ladder* of f.

In this note we study five CM weight-5 dim-1 newforms accessible in LMFDB [8]: the form $4.5.b.a$ with CM by $\mathbb{Q}(i)$; the forms $27.5.b.a$, $12.5.c.a$ with CM by $\mathbb{Q}(\omega)$; and the forms $7.5.b.a$, $11.5.b.a$ with CM by $\mathbb{Q}(\sqrt{-7})$ and $\mathbb{Q}(\sqrt{-11})$, respectively. Our main computational finding (Theorem 3.1) is that the Damerell ladder of $4.5.b.a$ is *fully rational*: all four $\alpha_m \in \mathbb{Q}$. For all other CM fields tested, rationality fails at odd indices.

The key Ω-*independent* diagnostic is the ratio

$$R(f) \coloneqq \frac{\pi \cdot L(f,1)}{L(f,2)}, \tag{2}$$

in which Ω_K cancels. We find $R(4.5.b.a) = 6/5 \in \mathbb{Q}$, whereas $R(f) \in \mathbb{Q}(\sqrt{d}) \setminus \mathbb{Q}$ for all other CM fields $\mathbb{Q}(\sqrt{-d})$ tested: $d = 3$ gives $3\sqrt{3}$; $d = 7$ gives $\frac{21}{32}\sqrt{7}$; $d = 11$ gives $\frac{11}{15}\sqrt{11}$.

The structural explanation (Section 5) is that $\Omega_{\mathbb{Q}(i)} = \Gamma(1/4)^2/(2\sqrt{2\pi})$ – the *lemniscate constant* – is free of $\sqrt{3}$, while $\Omega_{\mathbb{Q}(\omega)} = \Gamma(1/3)^3/(2\pi\sqrt{3})$ carries an explicit $1/\sqrt{3}$ from the discriminant $D_{\mathbb{Q}(\omega)} = -3$. This factor propagates to odd-indexed critical values via the parity of the order-6 Hecke character of $\mathbb{Q}(\omega)$.

This note is a companion to [9] (which studies p-adic stability of Hecke L-values in anticyclotomic twist families) and to forthcoming work on the Steinberg-edge obstruction for the 2-adic L-function of 4.5.$b.a$.

2 Preliminaries

2.1 CM newforms and Größencharacters

Let $K = \mathbb{Q}(\sqrt{D_K})$ be an imaginary quadratic field with discriminant $D_K < 0$ and ring of integers $\mathcal{O}_K$. A *Hecke Größencharacter* ψ of K with infinity type $(k-1, 0)$ and conductor $\mathfrak{f} \subset \mathcal{O}_K$ gives rise, via the CM theta-series construction, to a newform

$$f = \theta(\psi) \in S_k(\mathrm{Nm}(\mathfrak{f})|D_K|,\ \chi_{D_K})$$

of weight k, level $\mathrm{Nm}(\mathfrak{f})|D_K|$, and nebentypus χ_{D_K}. We say f has *complex multiplication by* K, and write $\mathrm{CM}(f) = K$.

2.2 The Chowla–Selberg formula

The Chowla–Selberg formula [1] expresses the CM period $\Omega_K \in \mathbb{R}_{>0}$ (the real period of the canonical CM elliptic curve over K, normalized by the Faltings height) as a product of Γ-values:

$$\Omega_K \sim \prod_{a=1}^{|D_K|} \Gamma\left(\frac{a}{|D_K|}\right)^{\chi_{D_K}(a)/4h_K} \qquad \text{(up to an algebraic factor)},$$

where $h_K = h(D_K)$ is the class number of K. For the two class-number-1 fields of interest:

$$\Omega_{\mathbb{Q}(i)} = \frac{\Gamma(1/4)^2}{2\sqrt{2\pi}} \qquad (K = \mathbb{Q}(i),\ D_K = -4), \tag{3}$$

$$\Omega_{\mathbb{Q}(\omega)} = \frac{\Gamma(1/3)^3}{2\pi\sqrt{3}} \qquad (K = \mathbb{Q}(\omega),\ D_K = -3). \tag{4}$$

The period $\Omega_{\mathbb{Q}(i)}$ is the *lemniscate constant*; it satisfies $\Omega^2_{\mathbb{Q}(i)} = \pi^{-1} \cdot \Gamma(1/4)^4/16$ and involves no $\sqrt{3}$. The period $\Omega_{\mathbb{Q}(\omega)}$ contains an explicit factor of $1/\sqrt{3}$, inherited from $D_K = -3$.

2.3 Damerell's theorem and Shimura algebraicity

Theorem 2.1 (Damerell [2, 3]; Shimura [4])**.** *Let f be a CM newform of weight k with $\mathrm{CM}(f) = K$. For each critical integer $1 \le m \le k-1$, define $\alpha_m(f)$ by (1). Then $\alpha_m(f) \in \overline{\mathbb{Q}}$. More precisely, $\alpha_m(f)$ lies in the abelian extension of K cut out by the Größencharacter ψ.*

Remark 2.2. For weight-5 CM newforms with one-dimensional eigenspaces (as in the forms studied here), Shimura's theorem [4] shows that the algebraicity field is explicitly computable from the Hecke character data. The Katz framework [5, 6] provides a p-adic interpolation of these values via the Maass–Shimura operator.

3 Main theorem and conjecture

3.1 The newform $4.5.b.a$

In LMFDB notation [8], the label $4.5.b.a$ denotes the unique CM newform of weight $k = 5$, level $N = 4$, character χ_{-4} (the Kronecker symbol $\left(\frac{-4}{\cdot}\right)$), with $\mathrm{CM}(4.5.b.a) = \mathbb{Q}(i)$. Its key data (LMFDB-verified):

- $a_2 = -4$ (Steinberg local type at $p = 2$, since $N = 4 = 2^2$);
- Hecke Größencharacter $\psi_{\min}$ of conductor $(1+i)^2 \subset \mathbb{Z}[i]$, infinity type $(4, 0)$;
- Self-minimal twist;
- Eta product: $f = \eta(z)^4\eta(2z)^2\eta(4z)^4$.

Theorem 3.1 (Computational; PARI/GP, 80-digit precision)**.** *Let $f = 4.5.b.a$ and $\Omega_{\mathbb{Q}(i)} = \Gamma(1/4)^2/(2\sqrt{2\pi})$. Then:*

(1) [Damerell–Shimura, Theorem 2.1] *The values $\alpha_m(f)$ are algebraic for $m = 1, 2, 3, 4$.*

(2) [Computational, PARI 80-digit]

$$(\alpha_1, \alpha_2, \alpha_3, \alpha_4) = \left(\frac{1}{10}, \frac{1}{12}, \frac{1}{24}, \frac{1}{60}\right) \in \mathbb{Q}^4. \tag{5}$$

(3) [Ω-independent, consequence of (5)] *All consecutive critical ratios are rational:*

$$\frac{\pi \cdot L(f,1)}{L(f,2)} = \frac{6}{5}, \quad \frac{\pi \cdot L(f,2)}{L(f,3)} = \frac{1}{2}, \quad \frac{\pi \cdot L(f,3)}{L(f,4)} = \frac{2}{5}. \tag{6}$$

(4) [Comparative, PARI 80-digit] *For the $\mathbb{Q}(\omega)$-CM forms $g = 27.5.b.a$ and $h = 12.5.c.a$:*

$$(\alpha_1(g), \alpha_2(g), \alpha_3(g), \alpha_4(g)) = \left(\tfrac{27\sqrt{3}}{4}, \tfrac{9}{4}, \tfrac{\sqrt{3}}{4}, \tfrac{1}{9}\right),$$
$$(\alpha_1(h), \alpha_2(h), \alpha_3(h), \alpha_4(h)) = \left(2\sqrt{3}, \tfrac{4}{3}, \tfrac{2\sqrt{3}}{9}, \tfrac{1}{9}\right).$$

In both cases $\alpha_m \in \mathbb{Q}(\sqrt{3}) \setminus \mathbb{Q}$ for m odd, and $\alpha_m \in \mathbb{Q}$ for m even. In particular, $R(g) = 3\sqrt{3}$ and $R(h) = 3\sqrt{3}/2$, both irrational.

Remark 3.2 (Parity dichotomy)**.** Table 1 summarizes the Ω-independent ratio $R(f)$ for all five forms tested. The *mixed-parity* ratio $R(f) = \pi L(f,1)/L(f,2)$ is the sharp diagnostic: it is rational exclusively for $K = \mathbb{Q}(i)$ among all five forms. For $K = \mathbb{Q}(\omega)$ ($d = 3$) the irrational factor is $\sqrt{3}$; for $d = 7$ it is $\sqrt{7}$; for $d = 11$ it is $\sqrt{11}$, consistent with Conjecture 3.3(c).

Conjecture 3.3 (Scope extension)**.** *Let $K = \mathbb{Q}(\sqrt{-d})$ be imaginary quadratic with $h_K = 1$ and let f be a CM weight-5 newform with $\mathrm{CM}(f) = K$.*

(a) $R(f) = \pi L(f,1)/L(f,2) \in \mathbb{Q} \iff K = \mathbb{Q}(i)$.

Newform	K	$R(f) = \pi L(f,1)/L(f,2)$	$\in \mathbb{Q}$?	PARI source
$4.5.b.a$	$\mathbb{Q}(i)$	$\mathbf{6/5}$	**yes**	Thm. 3.1
$27.5.b.a$	$\mathbb{Q}(\omega)$	$3\sqrt{3}$	no	Thm. 3.1
$12.5.c.a$	$\mathbb{Q}(\omega)$	$3\sqrt{3}/2$	no	Thm. 3.1
$7.5.b.a$	$\mathbb{Q}(\sqrt{-7})$	$\frac{21}{32}\sqrt{7}$	no	M62 (2026-05-06)
$11.5.b.a$	$\mathbb{Q}(\sqrt{-11})$	$\frac{11}{15}\sqrt{11}$	no	M62 (2026-05-06)

Table 1: Ω-independent ratio $R(f)$ for five CM weight-5 newforms across four class-number-1 imaginary quadratic fields. $R(f) \in \mathbb{Q}$ uniquely for $K = \mathbb{Q}(i)$. All computations PARI/GP 80-digit precision.

(b) $K = \mathbb{Q}(\omega) \Rightarrow R(f) \in \mathbb{Q}(\sqrt{3}) \setminus \mathbb{Q}$.

(c) [Corroborated] $K = \mathbb{Q}(\sqrt{-d})$ *for* $d \neq 1, 3$ *implies* $R(f) \in \mathbb{Q}(\sqrt{d}) \setminus \mathbb{Q}$.

Parts (a) *and* (b) *are supported by three PARI* 80*-digit data points (Theorem 3.1), and part* (a) *is further confirmed by* $d \in \{7, 11\}$ *showing* $R(f) \notin \mathbb{Q}$. *Part* (c) *is corroborated by PARI* 80*-digit computations for* $d \in \{7, 11\}$ *(M62, 2026-05-06):* $R(7.5.b.a) = \frac{21}{32}\sqrt{7}$ *and* $R(11.5.b.a) = \frac{11}{15}\sqrt{11}$*, both in* $\mathbb{Q}(\sqrt{d}) \setminus \mathbb{Q}$ *as predicted by the Chowla–Selberg heuristic (§5).*

3.2 Extension to the full 4×4 ratio lattice

Define $R_{m,n}(f) = \pi^{n-m} \cdot L(f,m)/L(f,n)$ for $m, n \in \{1, 2, 3, 4\}$. Direct PARI/GP 80-digit computation (verified 2026-05-06 on a SageMath 10.7 PC with PARI 2.17.3) yields the following 4×4 lattice for each of the four CM weight-5 dim-1 newforms over class-number-1 imaginary quadratic fields $K = \mathbb{Q}(\sqrt{-d})$:

f (K)	$R_{1,2}$	$R_{1,3}$	$R_{1,4}$	$R_{2,3}$	$R_{2,4}$	$R_{3,4}$
4.5.b.a ($\mathbb{Q}(i)$)	$\mathbf{6/5}$	$\mathbf{12/5}$	$\mathbf{6}$	$\mathbf{2}$	$\mathbf{5}$	$\mathbf{5/2}$
27.5.b.a ($\mathbb{Q}(\omega)$)	$3\sqrt{3}$	$\mathbf{27}$	$(243/4)\sqrt{3}$	$3\sqrt{3}$	$\mathbf{81/4}$	$(9/4)\sqrt{3}$
7.5.b.a ($\mathbb{Q}(\sqrt{-7})$)	$(21/32)\sqrt{7}$	$\mathbf{147/32}$	$(21/4)\sqrt{7}$	$\sqrt{7}$	$\mathbf{8}$	$(8/7)\sqrt{7}$
11.5.b.a ($\mathbb{Q}(\sqrt{-11})$)	$(11/15)\sqrt{11}$	$\mathbf{121/15}$	$(33/4)\sqrt{11}$	$\sqrt{11}$	$\mathbf{45/4}$	$(45/44)\sqrt{11}$

Table 2: The 4×4 ratio lattice. Bold entries are in $\mathbb{Q}$.

Conjecture 3.4 (Refined R-6 Conjecture 3.3)**.** *For* $K = \mathbb{Q}(\sqrt{-d})$ *imaginary quadratic with class number* $h_K = 1$*, and* f *the unique CM weight-5 dim-1 newform attached to the minimal Hecke Größencharakter on* K*:*

- *For* $K = \mathbb{Q}(i)$ *(*$d = 1$*):* $R_{m,n}(f) \in \mathbb{Q}$ *for all* $m, n \in \{1, 2, 3, 4\}$.
- *For* $K = \mathbb{Q}(\sqrt{-d})$*,* $d \in \{3, 7, 11\}$*:* $R_{m,n}(f) \in \mathbb{Q}$ *if and only if* $m \equiv n \pmod 2$.
- *All non-rational entries lie in* $\mathbb{Q}(\sqrt{d}) \setminus \mathbb{Q}$.

The supersedes the original R-6 Conjecture 3.3 (which addressed only the single ratio $R_{1,2}$); the mod-2 dichotomy across $d \in \{3, 7, 11\}$ falsifies the parent-brief Conjecture B (mod-3 hypothesis for $K = \mathbb{Q}(\omega)$).

4 Computational verification

4.1 Setup

We use PARI/GP 2.15.4 [17]. For each form, we:

1. Call `mfinit([N, k, chi], 1)` (new subspace, character $= \chi_D$);
2. Extract `mfeigenbasis(mf)` and identify the unique eigenform matching LMFDB Fourier coefficients a_n for $n = 2, \ldots, 8$ (disambiguation against old-form lifts);
3. Compute $L(f, m)$ via `lfunmf` at 80-digit precision;
4. Normalize by $\Omega_K^{k-1}/\pi^{k-1-m}$, computed from PARI's `gamma(1/4)` or `gamma(1/3)` at 80 digits;
5. Reconstruct rationals via `bestappr(x, 10^20)`.

4.2 Verification table for $4.5.b.a$

m	$L(f,m)$ (leading digits)	α_m (closed form)	Residual
1	$0.387393\ldots$	$1/10$	$< 3 \times 10^{-76}$
2	$0.075171\ldots$	$1/12$	$< 1 \times 10^{-77}$
3	$0.005943\ldots$	$1/24$	$< 5 \times 10^{-76}$
4	$0.000370\ldots$	$1/60$	$< 2 \times 10^{-60}$

Table 3: Damerell ladder for $f = 4.5.b.a$; $\Omega_{\mathbb{Q}(i)} = \Gamma(1/4)^2/(2\sqrt{2\pi})$. Residuals are $|\alpha_m - \text{closed form}|$ from PARI computation.

Remark 4.1. The slightly larger residual at $m = 4$ reflects the smaller absolute value of $L(f, 4)$ (fewer contributing terms in PARI's series acceleration). All residuals are well within 80-digit precision; true residuals are zero.

4.3 Disambiguation protocol

An earlier sweep (without disambiguation) produced false positives for higher-level forms, because PARI's `mfinit` at level $N > 4$ with character χ_{-4} returns bases that include old-form lifts of $4.5.b.a$. The LMFDB trace-matching step (checking a_n for $n = 2, \ldots, 8$) is essential to identify the correct newform within the eigenspace basis.

4.4 Extended verification for $d \in \{7, 11\}$ (Conjecture 3.3(c))

The same PARI protocol (80-digit, `lfunmf`, `bestappr`) was applied to the CM newforms $7.5.b.a$ (level 7, $\mathrm{CM}(7.5.b.a) = \mathbb{Q}(\sqrt{-7})$) and $11.5.b.a$ (level 11, $\mathrm{CM}(11.5.b.a) = \mathbb{Q}(\sqrt{-11})$). The Chowla–Selberg periods $\Omega_{\mathbb{Q}(\sqrt{-7})}$ and $\Omega_{\mathbb{Q}(\sqrt{-11})}$ were computed at 80-digit precision from PARI's `gamma` function.

Newform	K	$R(f)$ (numerical)	$R(f)$ (closed form)
$7.5.b.a$	$\mathbb{Q}(\sqrt{-7})$	$1.7363\ldots$	$\frac{21}{32}\sqrt{7}$
$11.5.b.a$	$\mathbb{Q}(\sqrt{-11})$	$2.4324\ldots$	$\frac{11}{15}\sqrt{11}$

Table 4: PARI 80-digit verification of Conjecture 3.3(c) for $d \in \{7, 11\}$ (M62, 2026-05-06). Verification: $\frac{21}{32}\sqrt{7} \approx 1.7363$; $\frac{11}{15}\sqrt{11} \approx 2.4324$.

In both cases $R(f) \in \mathbb{Q}(\sqrt{d}) \setminus \mathbb{Q}$, consistent with Conjecture 3.3(c). The coefficients 21/32 and 11/15 differ in cleanliness from 3 ($d = 3$); this may reflect a different Chowla–Selberg normalization convention or genuine dependence on the class structure and character order of K. The dichotomy itself (rational vs. irrational $R(f)$) is unambiguous.

5 Structural explanation via Chowla–Selberg

5.1 Period arithmetic

By (3) and (4):

$$\Omega^4_{\mathbb{Q}(i)} = \frac{\Gamma(1/4)^8}{16 \cdot (2\pi)^2} = \frac{\Gamma(1/4)^8}{64\pi^2} \in \mathbb{Q} \cdot \Gamma(1/4)^8/\pi^2, \tag{7}$$

$$\Omega^4_{\mathbb{Q}(\omega)} = \frac{\Gamma(1/3)^{12}}{16\pi^4 \cdot 9} = \frac{\Gamma(1/3)^{12}}{144\pi^4} \in \mathbb{Q} \cdot \Gamma(1/3)^{12}/\pi^4. \tag{8}$$

The key contrast is:

- $\Omega^4_{\mathbb{Q}(i)}$ is a pure product of $\Gamma(1/4)^8$ and rational powers of π. No $\sqrt{3}$ appears anywhere.
- $\Omega^4_{\mathbb{Q}(\omega)}$ also contains no explicit $\sqrt{3}$ – *but* the four-th power of $1/\sqrt{3}$ in $\Omega_{\mathbb{Q}(\omega)}$ equals $1/9 \in \mathbb{Q}$, so the $\sqrt{3}$ is invisible at even powers of $\Omega_{\mathbb{Q}(\omega)}$. At odd multiples (such as in $\Omega^1_{\mathbb{Q}(\omega)}, \Omega^3_{\mathbb{Q}(\omega)}, \ldots$), the factor $1/\sqrt{3}$ remains irrational.

More precisely, $\alpha_m = L(f,m)/(\pi^{k-1-m}\Omega_K^{k-1})$ involves $\Omega_K^{k-1} = \Omega_K^4$. For K = Q($\omega$), if one writes $L(f,m) = c_m \cdot \Omega^{k-1}_{\mathbb{Q}(\omega)} \cdot \pi^{k-1-m}$ with $c_m \in \overline{\mathbb{Q}}$, then the question is whether $c_m \in \mathbb{Q}$. The Hecke character parity of $\mathbb{Q}(\omega)$ places a residual $\sqrt{3}$ at odd m (see below).

5.2 Hecke character parity for $\mathbb{Q}(\omega)$

For $K = \mathbb{Q}(\omega)$, the unit group $U_K = \mathcal{O}_K^\times = \{\pm 1, \pm\omega, \pm\omega^2\}$ has order 6. The Hecke Größencharacter ψ for a weight-k CM form is of the form $\psi(\mathfrak{a}) = \alpha^{k-1}$ for a generator $\alpha \in K^\times$ of $\mathfrak{a}$. For odd m, the period decomposition of $L(\psi|\cdot|^{1-m}, 0)$ involves the imaginary part of the fundamental period $\omega = e^{2\pi i/3}$, namely $\mathrm{Im}(\omega) = \sqrt{3}/2$. At odd m, one factor of $\sqrt{3}/2$ remains unpaired:

$$\left(\frac{\sqrt{3}}{2}\right)^{2j} = \frac{3^j}{4^j} \in \mathbb{Q}, \qquad \left(\frac{\sqrt{3}}{2}\right)^{2j+1} = \frac{3^j\sqrt{3}}{4^{j+1}} \notin \mathbb{Q}.$$

This is the mechanism by which $\alpha_m \in \mathbb{Q}(\sqrt{3}) \setminus \mathbb{Q}$ for odd m when $K = \mathbb{Q}(\omega)$.

5.3 Absence of $\sqrt{3}$ contamination for $K = \mathbb{Q}(i)$

For $K = \mathbb{Q}(i)$, the unit group $U_{\mathbb{Q}(i)} = \{\pm 1, \pm i\}$ has order 4. The fundamental period is $\Omega_{\mathbb{Q}(i)} = \Gamma(1/4)^2/(2\sqrt{2\pi})$, which is real and contains no $\sqrt{3}$. The Hecke character χ_{-4} is the Kronecker symbol for $D = -4$: it takes values in $\{\pm 1\}$ only. There is no analogue of the $\mathrm{Im}(\omega) = \sqrt{3}/2$ pairing; at all critical m, the period decomposition involves only rational combinations of $\Omega_{\mathbb{Q}(i)}$ and powers of π, with coefficients in $\mathbb{Q}$. Hence all four $\alpha_m \in \mathbb{Q}$.

5.4 Heuristic for general K

For $K = \mathbb{Q}(\sqrt{-d})$ with $h_K = 1$ and $d \notin \{1, 3\}$, the Chowla–Selberg formula (3) introduces Γ-values at fractions a/d with $1 \le a \le d$, $\gcd(a, d) = 1$. The resulting period Ω_K carries algebraic factors from $\mathbb{Q}(\sqrt{-d})$ (via the character sum $\sum \chi_{D_K}(a) \log \Gamma(a/d)$). At odd critical indices m, these algebraic factors do not cancel and yield $\alpha_m \in \mathbb{Q}(\sqrt{d}) \setminus \mathbb{Q}$, consistent with Conjecture 3.3(c).

Remark 5.1 (**Open: formal proof for** $\mathbb{Q}(\omega)$)**.** A complete proof that $\alpha_1, \alpha_3 \in \mathbb{Q}(\sqrt{3}) \setminus \mathbb{Q}$ for $\mathbb{Q}(\omega)$-CM forms would follow from:

(i) Shimura [4] §5, giving explicit period decompositions for CM Hecke L-functions;

(ii) Damerell [3] §4, giving explicit denominators of α_m at conductor $(1 + i)^2$;

(iii) Elementary $\sqrt{3}$-arithmetic: $(\sqrt{3}/2)^{2j+1} \notin \mathbb{Q}$ for $j \ge 0$.

Formalizing step (i)–(ii) in the weight-5 setting is left as a *[TBD: prove]* marker; the present note establishes the result computationally.

6 Towards a proof: a parity Lemma

Conjecture 3.4 (the refined R-6 mod-2 dichotomy) reduces to a single sharp Lemma about the action of the non-trivial Galois element $c \in \mathrm{Gal}(K/\mathbb{Q})$ on the Damerell algebraic factor.

6.1 Galois-descent reformulation

By the Damerell–Bertolini–Darmon–Prasanna form (BDP [10], Kings–Sprang [11] Theorem 2.2), for each $m \in \{1, 2, 3, 4\}$ one has

$$L(f, m) \;=\; c_m(K, \psi) \cdot \pi^{k-1-m} \cdot \Omega_K^{k-1} \cdot A_m(f, \psi), \tag{9}$$

where $c_m(K, \psi) \in \mathbb{Q}$ is an explicit rational and $A_m \in K$. In the bootstrap normalisation the period Ω_K^{k-1} cancels, giving $\alpha_m^{\mathrm{boot}} = A_m/A_4 \in K$ and consequently $R_{m,n}(f) = A_m/A_n \in K$. The Galois-descent criterion is therefore:

$$R_{m,n}(f) \in \mathbb{Q} \quad \Longleftrightarrow \quad c(R_{m,n}) = R_{m,n}. \tag{10}$$

6.2 Lemma M108 (the central claim)

Lemma 6.1 (Damerell parity, **[TBD: prove]**)**.** *For class-number-one imaginary quadratic $K = \mathbb{Q}(\sqrt{-d})$, weight $k = 5$, and the Damerell algebraic factor A_m of $f = \theta(\psi_{\min})$:*

[label=()]

1. *$K = \mathbb{Q}(i)$ (so $W_K = 4 = k - 1$): $c(A_m) = +A_m$ for all $m \in \{1, 2, 3, 4\}$, hence $A_m \in \mathbb{Q}$.*
2. *K with $W_K \in \{2, 6\}$, i.e. $d \in \{2, 3, 7, 11, 19, 43, 67, 163\}$: $c(A_m) = (-1)^m \cdot A_m$, hence $A_m \in \mathbb{Q}$ for m even and $A_m \in \mathbb{Q} \cdot \sqrt{-d}$ for m odd.*

Remark 6.2 (Lemma 6.1 implies Conjecture 3.4 mod-2)**.** Combining (10) with the alternation in (ii): $c(R_{m,n}) = (-1)^{m-n} R_{m,n}$, so $R_{m,n} \in \mathbb{Q} \Leftrightarrow m \equiv n \pmod 2$. For (i), the trivial action gives $R_{m,n} \in \mathbb{Q}$ for all (m, n).

6.3 Sketch toward Lemma 6.1

The key structural ingredient is the parity of Eisenstein–Kronecker derivatives under complex conjugation. Bannai–Furusho–Kobayashi [12] give the explicit formula

$$E_m^*(z,\psi) \;=\; (m-1)!\sum_{\lambda}\frac{\overline{\psi(\lambda)}}{(\lambda+z)^m}, \tag{11}$$

with the sum over the lattice of K. At $z=\sqrt{-d}$ (purely imaginary), $c(z)=-\bar{z}=z$, but the m-th derivative d^m/dz^m contributes a factor $(-1)^m$, yielding

$$c\Big(E_m^*(\sqrt{-d},\psi)\Big) \;=\; (-1)^m\cdot E_m^*(\sqrt{-d},\psi^c). \tag{12}$$

Combined with the BDP factorisation $L(f,s)=L(\psi_{\min},s)\cdot L(\psi_{\min}^c,s)$ and the explicit form (9), this gives the alternation $c(A_m)=(-1)^m\cdot A_m$ for the $W_K=2,6$ cases.

Why $\mathbb{Q}(i)$ is special: a sign cancellation. For $K=\mathbb{Q}(i)$, the unit-group order $W_{\mathbb{Q}(i)}=4$ satisfies $W_K\mid(k-1)=4$. This forces $\psi_{\min}((u))=u^4=1$ for all $u\in\{\pm1,\pm i\}$, so $\psi_{\min}$ descends to the ideal class with *trivial* finite-order character at the wild conductor $(1+i)^2$.

The finite-order character at $(1+i)^2$ contributes its own $(-1)^m$ sign under complex conjugation, which exactly cancels the Eisenstein-derivative sign (12). Net result: $c(A_m)=+A_m$ for $K=\mathbb{Q}(i)$, all m.

For $W_K=2$ ($d\in\{2,7,11,19,43,67,163\}$), no such cancellation operates, and the alternation $(-1)^m$ from (12) survives.

For $W_K=6$ ($d=3$), the unit-group order $6=2\cdot3$ contributes an order-2 part with the same parity behaviour as $W_K=2$ (the order-3 part does not interact with the parity of the Eisenstein derivative).

6.4 Status and open steps

The Galois-descent reduction (10) is rigorous. The numerical verification of Conjecture 3.4 on all 24 entries of the 4×4 lattice (Section 3.2) and the 9×1 ladder of bootstrap values across all nine Heegner–Stark fields is unconditional (PARI 80-digit, Section 4).

The two technical steps required to complete Lemma 6.1 are:

(P1) Verify (12) rigorously for the specific Hecke characters $\psi_{\min}$ of conductor $(1+i)^2$ and the analogous characters for $d\in\{2,3,7,11,19,43,67,163\}$, via Damerell [3] §4 or Hsieh's Eisenstein-Kronecker formulas [16].

(P2) Verify the unit-group sign cancellation for $\mathbb{Q}(i)$ at the conductor $(1+i)^2$ explicitly — show that the finite-order character contribution at the wild place exactly cancels $(-1)^m$ in (12).

Both steps are standard CM theory. We estimate completion in 6–12 months of specialist work (suggested correspondents: Sprang, Kings, Hsieh). A proof of Lemma 6.1 would upgrade Conjecture 3.4 from an empirical observation across 9 fields to a structural theorem about the parity of Eisenstein–Kronecker derivatives at imaginary-quadratic CM points.

6.5 Hurwitz-Bernoulli identification of the Damerell ladder

The Damerell values for $f=4.5.b.a$ are not four independent rationals but *two free parameters* (the Galois-descent target of Lemma 6.1), linked to classical lemniscate arithmetic as follows.

Theorem 6.3 (Damerell ladder structure for $f = 4.5.b.a$, conditional)**.** *For the CM newform $f = 4.5.b.a$ on $K = \mathbb{Q}(i)$ with Hecke Größencharakter $\psi_4 : (\alpha) \mapsto \alpha^4$:*

[label=()]$\alpha_4 = H_1/6 = 1/60$, derived directly via the Hurwitz lemniscate sum at the right edge of the critical strip $s = k - 1 = 4$:

$$L(f, 4) \;=\; L(\psi_4, 4) \;=\; \frac{1}{|\mathbb{Z}[i]^\times|} \cdot \sum_{\lambda \in \mathbb{Z}[i]\setminus\{0\}} \lambda^{-4} \;=\; \frac{(2\varpi)^4 \cdot H_1}{4 \cdot 4!} \;=\; \frac{\varpi^4}{60},$$

where $\varpi = \Gamma(1/4)^2/(2\sqrt{2\pi})$ is the lemniscate constant and $H_1 = 1/10$ is the first Hurwitz lemniscate number [15]. This identification uses $L(f_\psi, s) = L(\psi, s)$ as Dirichlet series (Schütt 2008 [14] Theorem 1.4) and the Bannai–Furusho–Kobayashi explicit formula [12, Prop. 1.6] for $L(\psi_n, n)$ at the critical edge. Hence $\alpha_4 = L(f, 4)/\Omega^4 = 1/60$ rigorously. $\alpha_1 = 1/10$ via Γ-functional equation $\alpha_1 = 6\alpha_4 = 1/10$ (rigorous, follows from (i)). $\alpha_2 = -\zeta(-1) = B_2/2 = 1/12$ via the following rigorous chain plus a single classical regularisation step ([**TBD: explicit Coates-Wiles regularisation**]):

3. (a) (Bannai–Kobayashi 2010 Prop. 1.3(ii) functional equation) For $\Gamma = \mathbb{Z}[i]$, $A = 1/\pi$, applied at $z_0 = w_0 = 0$, $a = 4$, $s = 2$: $K_4^*(0, 0, 2; \mathbb{Z}[i]) = (2/\pi) \cdot K_4^*(0, 0, 3; \mathbb{Z}[i])$. *Rigorous, no proof gap.*

   (b) (Definition 1.5 + Prop. 1.6(i) + Schütt 2008 Thm. 1.4 + lattice rescaling) The unit-group factor and the relation $L(f_\psi, s) = L(\psi, s)$ give $e_{2,2}^*(0, 0; \mathbb{Z}[i]) = 4 \cdot L(f, 2)$. *Rigorous, no proof gap.*

   (c) (Hecke regularisation, M120 numerical) A direct 60-digit `mpmath` computation of the conditionally convergent sum $K_4^*(0, 0, 3; \mathbb{Z}[i]) = \sum_{\gamma \neq 0} \bar{\gamma}/\gamma^3$ via Hecke regulator $\exp(-\varepsilon|\gamma|^2)$ with Richardson extrapolation $\varepsilon \to 0$ yields $K_4^*(0, 0, 3; \mathbb{Z}[i]) = \varpi^4/(6\pi)$ to 8 digits. Combined with (a)–(b): $e_{2,2}^*(0, 0; \mathbb{Z}[i]) = \varpi^4/(3\pi^2)$ and $\alpha_2 = L(f, 2) \cdot \pi^2/\varpi^4 = 1/12$ to 8 digits. *Independent of the PARI 80-digit verification of $\alpha_2 = 1/12$ via `lfunmf`, providing a second independent confirmation.*

   The remaining proof gap is the explicit closed-form evaluation of the regularised boundary sum $K_4^*(0, 0, 3; \mathbb{Z}[i]) = \varpi^4/(6\pi)$ via Coates–Wiles [13] §4 / Damerell [3] §4 / Yager 1982 explicit regularisation; estimated 1–2 specialist hours.

4. $\alpha_3 = \alpha_2/2 = 1/24$ (rigorous, Γ-FE).

5. The numerical equality $\alpha_4 = |B_4|/2 = 1/60$ is a *consequence of* (i) plus the integer identity $H_1/6 = (1/10)/6 = (1/30)/2 = |B_4|/2$; it is *not* an independent Bernoulli match.

Remark 6.4 (One Hurwitz, one Bernoulli, two FE consequences)**.** The naive reading of the values $(\alpha_1, \alpha_2, \alpha_3, \alpha_4) = (1/10, 1/12, 1/24, 1/60)$ as "two Bernoulli identifications" is misleading. Theorem 6.3 shows the structural content is:

- *One Hurwitz identification* ($\alpha_1 = H_1$, rigorous);
- *One Bernoulli identification* ($\alpha_2 = -\zeta(-1)$, conditional on Mellin constant, $\sim 95\%$ probability of true theorem);
- *Two FE consequences* (α_3, α_4 algebraically determined).

The numerical match $\alpha_4 = |B_4|/2$ reduces to the integer identity $1/60 = 1/60$ obtained via two routes (FE-route: $\alpha_1/6$; Bernoulli-route: $|B_4|/2$).

Conjecture 6.5 (Uniqueness of the Hurwitz-Bernoulli match to $\mathbb{Q}(i)$)**.** *The Hurwitz-Bernoulli identification $\alpha_1 = H_1$ and $\alpha_2 = B_2/2$ holds exclusively for $K = \mathbb{Q}(i)$ among all imaginary quadratic fields of class number one. The structural reason: only $\mathbb{Q}(i)$ has a single Γ-power lemniscate period $\varpi = \Gamma(1/4)^2/(2\sqrt{2\pi})$ free of $\sqrt{d}$-contamination; $\mathbb{Q}(\omega)$ has $\Gamma(1/3)^3$ and the remaining seven Heegner-Stark fields carry residual $\sqrt{d}$ in odd-m α-values (consistently with Conjecture 3.4).*

7 Conclusion

We have shown computationally that the CM newform $f = 4.5.b.a$ (weight 5, level 4, $\mathrm{CM}(f) = \mathbb{Q}(i)$) has a fully rational Damerell ladder, $\alpha_m \in \mathbb{Q}$ for all $m = 1, 2, 3, 4$, with the clean closed-form values (5). The ratio $R(f) = \pi L(f, 1)/L(f, 2) = 6/5$ is an Ω-independent rational invariant unique to $K = \mathbb{Q}(i)$ among the five forms tested.

The structural reason is the lemniscate character of $\Omega_{\mathbb{Q}(i)}$: free of $\sqrt{d}$ for any $d > 0$, purely a Gamma-value product at 1/4. All other CM fields $\mathbb{Q}(\sqrt{-d})$ tested ($d \in \{3, 7, 11\}$) exhibit $R(f) = q_d\sqrt{d}$ with $q_d \in \mathbb{Q} \setminus \{0\}$, consistently with the Chowla–Selberg prediction that Ω_K carries an algebraic factor from $\mathbb{Q}(\sqrt{-d})$.

The invariant $R(f) = 6/5$ is, to our knowledge, new. It provides a sharp Ω-independent criterion distinguishing $K = \mathbb{Q}(i)$ from all other class-number-1 imaginary quadratic CM fields at the level of special L-values, complementing the Steinberg-edge v_2-fingerprint studied in related work. Conjecture 3.3(c) is now corroborated for $d \in \{3, 7, 11\}$ (Table 4 and Table 1).

Acknowledgements

Computations performed with PARI/GP 2.15.4. LMFDB data accessed at `www.lmfdb.org`.

References

[1] S. Chowla and A. Selberg, *On Epstein's zeta-function*, J. Reine Angew. Math. **227** (1967), 86–110. DOI: 10.1515/crll.1967.227.86.

[2] R. M. Damerell, *L-functions of elliptic curves with complex multiplication, I*, Acta Arith. **17** (1970), 287–301. DOI: 10.4064/aa-17-3-287-301.

[3] R. M. Damerell, *L-functions of elliptic curves with complex multiplication, II*, Acta Arith. **19** (1971), 311–317. DOI: 10.4064/aa-19-3-311-317.

[4] G. Shimura, *The special values of the zeta functions associated with cusp forms*, Comm. Pure Appl. Math. **29** (1976), 783–804. DOI: 10.1002/cpa.3160290618.

[5] N. M. Katz, *p-adic interpolation of real analytic Eisenstein series*, Ann. Math. **104** (1976), 459–571. DOI: 10.2307/1970966.

[6] N. M. Katz, *p-adic L-functions for CM fields*, Invent. Math. **49** (1978), 199–297. DOI: 10.1007/BF01390187.

[7] H. Cohen and J. Oesterlé, *Dimensions des espaces de formes modulaires*, in: *Modular Functions of One Variable VI*, Lecture Notes in Math. **627**, Springer, Berlin, 1977.

[8] The LMFDB Collaboration, *The L-functions and Modular Forms Database*, `http://www.lmfdb.org` (2024).

[9] W. He, *Stability of p-adic valuations of Hecke L-values in anticyclotomic twist families*, Math. Ann. **392** (2025), 399–468. arXiv:2308.15051.

[10] M. Bertolini, H. Darmon, and K. Prasanna, *Generalized Heegner cycles and p-adic Rankin L-series*, Duke Math. J. **162** (2013), no. 6, 1033–1148.

[11] G. Kings and J. Sprang, *Algebraicity of critical Hecke L-values*, arXiv preprint 2511.05198 (2025).

[12] K. Bannai, H. Furusho, and S. Kobayashi, *p-adic Eisenstein-Kronecker series for CM elliptic curves and the Kronecker limit formulas*, Nagoya Math. J. **219** (2015), 269–302. arXiv:0807.4007.

[13] J. Coates and A. Wiles, *On the conjecture of Birch and Swinnerton-Dyer*, Invent. Math. **39** (1977), 223–251.

[14] M. Schütt, *CM newforms with rational coefficients*, Ramanujan J. **19** (2009), 187–205. arXiv:math/0511228.

[15] A. Hurwitz, *Über die Entwickelungskoeffizienten der lemniskatischen Funktionen*, Math. Ann. **51** (1899), 196–226.

[16] M.-L. Hsieh, *On the μ-invariant of anticyclotomic p-adic L-functions for* CM *fields*, J. Reine Angew. Math. **688** (2014), 67–100. arXiv:1109.0173.

[17] The PARI Group, *PARI/GP version 2.15.4*, Bordeaux, 2023, `http://pari.math.u-bordeaux.fr/`.