

`viet@vvvqmr.com`
`https://vvvqmr.com`
Independent Researcher, Vietnam

Supplemental Material: Have Optical Wigner's Friend Experiments Been Blind to a Geometric Degree of Freedom?

VietVunVut (Viet – Nguyen Xuan)

2026-05-27

S1 Literature Search and Algebraic Proof

S1.1 Full Proof of Equatorial Cancellation Theorem

Theorem. Let Friend F measure in z -basis and Superobserver W at Bloch angles (θ, ϕ) .

With $f_{\perp}(b, d) = 1 - |\langle b|d \rangle|^2$:

$$f_{\perp}(+1, H) - f_{\perp}(-1, H) = -\cos \theta \quad (\text{S1})$$

Vanishes iff $\theta = \pi/2$.

Step 1: W 's measurement basis

$$|b = +1\rangle = \cos(\theta/2)|H\rangle + e^{i\phi} \sin(\theta/2)|V\rangle, \quad (\text{S2})$$

$$|b = -1\rangle = \sin(\theta/2)|H\rangle - e^{i\phi} \cos(\theta/2)|V\rangle. \quad (\text{S3})$$

Step 2: Overlaps with F 's outcomes

$$|\langle b = +1|H\rangle|^2 = \cos^2(\theta/2), \quad |\langle b = +1|V\rangle|^2 = \sin^2(\theta/2), \quad (\text{S4})$$

$$|\langle b = -1|H\rangle|^2 = \sin^2(\theta/2), \quad |\langle b = -1|V\rangle|^2 = \cos^2(\theta/2). \quad (\text{S5})$$

ϕ drops out: $|e^{i\phi}|^2 = 1$. Overlaps depend *only* on θ .

Step 3: $f_{\perp}$ values

$$f_{\perp}(+1, H) = 1 - \cos^2(\theta/2) = \sin^2(\theta/2), \quad (\text{S6})$$

$$f_{\perp}(-1, H) = 1 - \sin^2(\theta/2) = \cos^2(\theta/2), \quad (\text{S7})$$

$$f_{\perp}(+1, V) = 1 - \sin^2(\theta/2) = \cos^2(\theta/2), \quad (\text{S8})$$

$$f_{\perp}(-1, V) = 1 - \cos^2(\theta/2) = \sin^2(\theta/2). \quad (\text{S9})$$

Step 4: Outcome-dependence parameter

$$f_{\perp}(+1, H) - f_{\perp}(-1, H) = \sin^2(\theta/2) - \cos^2(\theta/2) \quad (\text{S10})$$

$$= -(\cos^2(\theta/2) - \sin^2(\theta/2)) \quad (\text{S11})$$

$$= -\cos \theta. \quad (\text{S12})$$

Vanishes iff $\theta = \pi/2$. At this angle, all four $f_{\perp} = 1/2$, constant across all outcome pairs. $\square$

Generality

At $\theta = \pi/2$, $|\langle b|d \rangle|^2 = 1/2$ for all b, d . Any $g(|\langle b|d \rangle|^2)$ therefore takes the same value for all outcome pairs. The equatorial plane is a fixed point for the entire class — the cancellation is not specific to $f_{\perp}$.

S1.2 BSM Equivalence for Proietti et al. (2019)

Proietti et al. use Bell-state measurement (BSM) rather than projective measurement. BSM projects onto four Bell states $\{|\Phi^+\rangle, |\Phi^-\rangle, |\Psi^+\rangle, |\Psi^-\rangle\}$. For the singlet-state source, each Bell outcome occurs with equal probability $1/4$, and conditioned on any Bell outcome b , the Friend’s recorded outcome d is equally likely to be H or V . Consequently $|\langle b|d \rangle|^2 = 1/2$ for all b, d , identical to the equatorial condition $\theta = \pi/2$ derived in the main text.

S1.3 Literature Search Methodology

Search date: 2026-05-15 to 2026-05-24.

Databases: Google Scholar, arXiv (quant-ph), Web of Science, InspireHEP.

Search strings (Boolean combinations):

- **Group A (subject):** “Wigner’s friend” OR “extended Wigner” OR “Frauchiger-Renner” OR “Local Friendliness”
- **Group B (parameter):** “equatorial measurement” OR “Bloch sphere polar angle” OR “outcome dependence” OR “geometric constraint” OR “measurement basis angle”

Inclusion criteria:

1. Optical EWF implementation with Friend+Superobserver structure
2. Published in peer-reviewed venue or arXiv
3. Reports measurement settings from which polar angle θ can be determined

Results: ~ 200 titles screened $\rightarrow$ 47 full-text examined $\rightarrow$ 2 published optical EWF experiments identified. Both operate at equatorial geometry ($\theta = \pi/2$ or equivalent via BSM). No published experiment varies θ from $\pi/2$ for any purpose.

Additional sources examined: Supplemental Material of Bong et al. (2020); Methods and Supplementary Information of Proietti et al. (2019); LF derivations in Frauchiger-Renner (2018) and Wiseman-Cavalcanti-Rieffel (2023); Bell/LF review by Brunner et al. (2014); multipartite, sequential, and possibilistic LF extensions.

Limitation: We cannot rule out unpublished results or implementations outside our database scope. The survey establishes that within the indexed, peer-reviewed literature, θ has not been varied.

S2 Numerical Methods and Statistical Robustness

S2.1 Exact Numerical Integration Method

All numerical predictions use the density matrix for the singlet state with finite visibility under the SPDC noise model (where noise is confined to the $\{|HV\rangle, |VH\rangle\}$ subspace):

$$\rho_\mu = \mu|\Phi^-\rangle\langle\Phi^-| + \frac{1-\mu}{2}(|HV\rangle\langle HV| + |VH\rangle\langle VH|). \quad (\text{S13})$$

For the overlap-dependent model Eq. (2–3) of the main text:

$$P(a, b \mid x, y) = P_{\text{QM}}(a, b \mid x, y) \cdot [1 - \beta f_\perp(b, d)]/Z, \quad (\text{S14})$$

with full renormalization $Z = \sum_{a,b} P_{\text{QM}}(a, b \mid x, y) \cdot [1 - \beta f_\perp(b, d)]$.

All computations performed via `K9S12_proposal.py`, `statistical_significance.py`. The algebraic cancellation theorem is independently verified via symbolic computation (`universal_theorem_lf_check.py`).

S2.2 Derivation of the Sensitivity Formula and First-Order Overestimate Warning

S2.2.1 $f_\perp$ values at polar angle θ

The Superobserver measurement basis at (θ, ϕ) is:

$$|b = +1\rangle = \cos(\theta/2)|H\rangle + e^{i\phi} \sin(\theta/2)|V\rangle, \quad (\text{S15})$$

$$|b = -1\rangle = \sin(\theta/2)|H\rangle - e^{i\phi} \cos(\theta/2)|V\rangle. \quad (\text{S16})$$

The squared overlaps with the Friend’s z -basis outcomes (where ϕ drops out since $|e^{i\phi}|^2 = 1$) are:

$$|\langle b = +1|H\rangle|^2 = \cos^2(\theta/2), \quad |\langle b = +1|V\rangle|^2 = \sin^2(\theta/2), \quad (\text{S17})$$

$$|\langle b = -1|H\rangle|^2 = \sin^2(\theta/2), \quad |\langle b = -1|V\rangle|^2 = \cos^2(\theta/2). \quad (\text{S18})$$

Using the incompatibility function $f_\perp(b, d) = 1 - |\langle b|d\rangle|^2$, we obtain:

$$f_\perp(+1, H) = \sin^2(\theta/2), \quad f_\perp(+1, V) = \cos^2(\theta/2), \quad (\text{S19})$$

$$f_\perp(-1, H) = \cos^2(\theta/2), \quad f_\perp(-1, V) = \sin^2(\theta/2). \quad (\text{S20})$$

The outcome-dependence difference parameter is:

$$|f_{\perp}(+1, H) - f_{\perp}(-1, H)|/2 = |\cos \theta|/2. \quad (\text{S21})$$

At the tilted angle $\theta = 31^\circ$, this yields:

$$|\cos 31^\circ|/2 = 0.8572/2 = 0.4286. \quad (\text{S22})$$

S2.2.2 QM correlator for mixed setting ($x = 1, y = 2$)

For the SPDC noise model (where noise is confined to the $\{|HV\rangle, |VH\rangle\}$ subspace), both the pure singlet and the noise term yield identical z -vs-tilted correlators ($-\cos \theta$), so the mixed-setting correlator is independent of visibility μ . At the reference angle $\theta = 31^\circ$:

$$\langle A_1 B_2 \rangle_{\text{QM}} = -\cos(31^\circ) = -0.8572. \quad (\text{S23})$$

S2.2.3 K9_E deformed correlator

For the multiplicative model, the K9_E probability is:

$$P_{\text{K9E}}(a, b|x, y) \propto P_{\text{QM}}(a, b|x, y) \cdot [1 - \beta f_{\perp}(b, d)], \quad (\text{S24})$$

with the full normalization $Z = \sum_{a,b} P_{\text{QM}} \cdot [1 - \beta f_{\perp}(b, d)]$ ensuring probability conservation. Because the normalization factor Z depends on both β and θ , the shift in the correlator is coupled across all outcome pairs. Numerical results at $\theta = 31^\circ, \mu = 0.95$ yield:

β	$\langle A_1 B_2 \rangle_{\text{QM}}$	$\langle A_1 B_2 \rangle_{\text{K9E}}$	δ
0.10	-0.8572	-0.8687	-0.0115
0.30	-0.8572	-0.8927	-0.0355
0.50	-0.8572	-0.9180	-0.0609

The deformation shifts the correlator to more negative values (enhancing anti-correlation) because the renormalization shifts probability weight toward outcome pairs with larger $f_{\perp}$ values, amplifying the geometric asymmetry.

S2.2.4 First-order expansion and overestimate warning

Expanding the renormalized probability to first order in β (before full renormalization by Z):

$$\delta \langle A_1 B_2 \rangle \approx -\beta \cdot |\cos \theta| \cdot \langle A_1 B_2 \rangle_{\text{QM}}^2 + O(\beta^2). \quad (\text{S25})$$

At $\theta = 31^\circ$, this unrenormalized expansion yields:

$$\delta \approx -\beta \cdot 0.8572 \cdot 0.7347 = -0.6298\beta. \quad (\text{S26})$$

Warning: This unrenormalized first-order expansion overestimates $|\delta|$ by approximately a factor of 5.5. The ratio of the numerical value to the leading-order expansion is nearly constant across all β :

β	δ (LO)	δ (numerical)	ratio
0.01	-0.0063	-0.0011	0.181
0.07	-0.0441	-0.0080	0.182
0.10	-0.0630	-0.0115	0.183
0.30	-0.1889	-0.0355	0.188
0.50	-0.3149	-0.0609	0.193

The discrepancy arises because the normalization factor $Z = \sum P_{\text{QM}}[1 - \beta f_{\perp}]$ is also $O(\beta)$, leading to a cancellation between the numerator and denominator corrections. Therefore, the unrenormalized formula captures the correct qualitative structural dependence (vanishing iff $\theta = \pi/2$) but severely overestimates its magnitude. All manuscript values use exact numerical computation.

S2.3 Complete Correlator Table ($\theta = 31^\circ$, $\mu = 0.95$)

Table S1: All nine $\langle A_x B_y \rangle$ correlators with uncertainties ($N = 91,000$).

(x, y)	$\langle AB \rangle_{\text{QM}}$	σ
(1, 1)	-1.0000	0.0000
(1, 2)	-0.8572	0.0017
(1, 3)	-0.8572	0.0017
(2, 1)	-0.8572	0.0017
(2, 2)	-0.5045	0.0029
(2, 3)	-0.8933	0.0015
(3, 1)	-0.8572	0.0017
(3, 2)	-0.8933	0.0015
(3, 3)	-0.8829	0.0016

S2.4 Grid Search Optimization

A grid search over $(\theta, \phi_2, \phi_3, \beta_{\text{Bob}})$ maximizes:

$$\text{FOM}(\theta) = \min \left(n_{\sigma}^{\text{LF}}(\theta), n_{\sigma}^{\text{signal}}(\theta, \beta) \right), \quad (\text{S27})$$

yielding $\theta = 31^\circ$, $\phi_2 = 112^\circ$, $\phi_3 = 217^\circ$, $\beta_{\text{Bob}} = 20^\circ$. FOM remains above 5σ for $\theta \in [20^\circ, 45^\circ]$ at $\beta = 0.30$ and for $\theta \in [35^\circ, 46^\circ]$ at $\beta = 0.07$, $N = 91,000$.

S2.5 Monte Carlo Validation

10,000 runs: Gen LF 1 $\geq 5\sigma$ in 99.97%. $\beta = 0.07$: >99% power (single); $\beta = 0.05$: $\sim 90\%$ (combined). Conservative Bayesian analysis (20% inflation): $\beta_{\text{min}} \approx 0.046$ (combined).

S2.6 Angular Stability

LF significance varies $<3\%$ across $\Delta\theta = \pm 5^\circ$. $\delta\langle AB \rangle$ (exact numerical θ -dependence): $<1.5\%$ variation at $\pm 1^\circ$ (Bong’s reported precision).

S2.7 ϕ -Scramble Control Protocol

Rotate HWP to $N_\phi \geq 10$ random azimuthal angles $\phi \in [0^\circ, 360^\circ)$, fit $\delta\langle AB \rangle(\phi) = A + B \cos(2\phi) + C \sin(2\phi)$. Test: $A \neq 0$ with $B, C \approx 0$ indicates geometric θ -dependent signal; B or $C \neq 0$ indicates birefringence/alignment artifacts.

S3 Interpretations and Physical Context

S3.1 Relation to Contextuality

Eq. (2) of the main text is not a contextual hidden-variable model. There is no unobserved λ ; the Friend outcome d is an observed macroscopic record. Dependence is on registration geometry ($|\langle b|d \rangle|^2$) rather than on which observables are co-measured.

Table S2: Comparison of dependence frameworks.

Framework	Hidden able?	vari-	Dependence on	Status of d
KS Contextuality	Yes (λ)		Measurement context	Unobserved
Standard contextuality	No		Co-measured observables	N/A
Retrocausal models	Varies		Future choices	Varies
Eq. (2) registration	No		$ \langle b d \rangle ^2$ geometry	Observed record

S3.2 Physical Picture

When a Friend measures in the z -basis, they produce a macroscopic record with a definite orientation on the Bloch sphere. A Superobserver measuring at (θ, ϕ) probes this record at a relative angle $|\langle b|d \rangle|^2$. Standard QM assumes that once the Friend’s outcome is registered, subsequent measurements factorize. Eq. (2) parametrizes a possible residual dependence on this geometric relationship.

S3.3 Quantum Interpretations

Copenhagen. Silent on registration-layer structure — the experiment tests a structural degree of freedom that no standard interpretation addresses.

Many-Worlds (Everett). $\beta \neq 0$ would provide the first quantitative probe of inter-branch structural correlations; a null result constrains such effects to $\beta < 0.04$.

Relational QM (Rovelli). Tests whether relational outcomes leave geometry-dependent measurable traces — a signature current RQM formulations neither predict nor preclude.

Objective Collapse. CSL and Diosi-Penrose models modify dynamics, not registration geometry. The overlap-dependent class is structurally orthogonal.

QBism. Parametrizes whether agent experiences exhibit structural coupling across agents at the registration layer. A null result constrains agent-agent coupling at $\sim 10^{-2}$.

S3.4 General Probabilistic Theories (GPT) Context

Within the GPT framework, the overlap-only deformation defines a one-parameter family of neighboring GPTs deviating from the quantum point along the registration-geometry direction. This is a structural degree of freedom orthogonal to standard nonlocality.

S3.5 Weak Measurement and Postselection

In weak measurement, the weak value $A_w = \langle \psi_f | A | \psi_i \rangle / \langle \psi_f | \psi_i \rangle$ depends explicitly on the pre-postselection overlap. The basis-alignment dependence in Eq. (2) shares the structural feature of geometric coupling between measurement records at different observational levels. The analogy is structural, not dynamical.

S3.6 Scale Comparison

Table S3: Phenomenological parameter scales.

Parameter class	Typical constraint	Reference
Photon-sector SME	$< 10^{-23}$	Colladay & Kostelecký (1997)
CSL collapse rate λ	$< 10^{-16} \text{ s}^{-1}$	Various
Weak-measurement anomalies	$\sim 10^{-2}$	Aharonov et al. (1988)
Overlap-dependent β (this work)	$\sim 10^{-2}$ (proposed)	—

S3.7 Multi-Observer Extension (Preliminary)

For $N > 2$ observers, the geometric cancellation structure generalizes. Preliminary analysis suggests $\sim 11\times$ amplification at $\beta = 0.3$ for three observers, conditional on bridge theorems connecting registration-layer structure to QM observables for $N > 2$. Rigorous derivation is left for future work.