

When Does a Physical Interaction Become a Valid Registered Measurement?

A VVV-QMRF Registration-Layer Extension of Quantum Mechanics with a Testable Prediction for Extended Wigner's Friend Scenarios

Working Paper v2.0 Author: Viet Nguyen Xuan (VietVunVut) **Repository:** <https://github.com/AIhugART/VVV-QMRF-VietVunVut-Quantum-Measurement-Registration-Framework> **Status:** Working paper submitted for community feedback. All formal claims are Class D (proposed) or Class C (conjecture) unless stated otherwise. Critique is explicitly invited.

Abstract

The quantum measurement problem has remained unresolved for nearly a century. Standard quantum mechanics specifies, via postulate P3, that a measurement of observable A on state $|\psi\rangle$ yields eigenvalue a_k with probability $|\langle a_k | \psi \rangle|^2$. What P3 does not specify is when a physical interaction constitutes a valid registered measurement event. This silence generates the von Neumann chain problem and leaves the Heisenberg cut formally undefined.

This paper proposes VVV-QMRF (VietVunVut Quantum Measurement Registration Framework): a registration-layer extension of quantum mechanics grounded in Buddhist Pramāṇa epistemology (Dignāga–Dharmakīrti tradition). The framework adds a K-side registration space K , separate from the physical Hilbert space H , and proposes a six-condition test for valid registered measurement. Applied to Extended Wigner's Friend (EWF) scenarios, this yields a specific conjecture: K-side registration spaces of two observers are incommensurable ($K_F \not\sqsubseteq K_W$) when a joint registration space K_{joint} is structurally required but cannot satisfy all six conditions simultaneously.

This conjecture is not trivially true for all frameworks. It separates ordinary Bell-type nonclassicality from LF-level Wigner-friend inconsistency, as illustrated by data from Bong et al. (2020) at $\mu = 0.80, 0.81$ where Bell non-LF violation is reported while Local Friendliness inequalities remain unviolated. Existing data from Proietti et al. (2019) and Bong et al. (2020) are compatible with the proposed structural reading. The framework does not replace Standard Quantum Mechanics, revise the Born rule, or invoke consciousness. All formal claims are classified by evidence level. A purpose-designed experiment with a predefined operational criterion is required for confirmation.

1. The Registration Layer Gap

1.1 What Quantum Mechanics Specifies

Standard quantum mechanics is built on four postulates:

- **P1 (State):** A physical system is represented by a density operator $\rho \in D(H)$.
- **P2 (Observables):** Physical quantities are represented by self-adjoint operators A on H .
- **P3 (Measurement):** Measurement of A on state $|\psi\rangle$ yields eigenvalue a_k with probability $|\langle a_k | \psi \rangle|^2$, after which the state updates to the corresponding eigenstate.
- **P4 (Dynamics):** Between measurements, the state evolves unitarily via the Schrödinger equation.

P3 specifies what outcome appears and with what probability. P3 does not specify when a physical interaction counts as a measurement, what structural conditions distinguish a measurement from a non-measurement

interaction, or what stops the measurement chain.

This is the registration layer gap.

1.2 The Von Neumann Chain Problem

Von Neumann (1932) observed that if apparatus A1 measures system S, A1 becomes entangled with S. If apparatus A2 then measures A1, A2 becomes entangled with A1+S. No postulate of standard QM specifies where this chain terminates. Every proposed stopping point — macroscopic size, decoherence, consciousness — is external to the formalism.

Decoherence (Zurek 2003) explains why superpositions become effectively classical at the macroscopic scale. It does not explain when a physical interaction constitutes a registration event in the sense of a definite recorded outcome. The von Neumann chain problem is a registration-layer problem, not a decoherence problem.

1.3 The Heisenberg Cut

Copenhagen quantum mechanics (Bohr 1935; Heisenberg 1958) assigns measurement authority to “the classical apparatus” without formally defining what makes an apparatus classical, why it has registration authority, or what structural conditions produce a definite outcome. The Heisenberg cut is a boundary without a formal location.

VVV-QMRF addresses both problems by introducing a registration layer K, separate from the physical layer ρ , and proposing formal conditions for valid registration events within K.

2. The K-side Registration Space

2.1 Layer Separation

VVV-QMRF introduces a strict separation between two layers:

- **Physical layer (p-side):** Physical states, observables, and dynamics as described by Standard QM. This layer is not modified. $\rho \in D(H)$, H is the standard Hilbert space.
- **Registration layer (K-side):** The space of registration events, recording conditions, and validity states. $K \neq H$. K-side structure does not alter ρ -side dynamics.

This separation is the core architectural commitment of VVV-QMRF.

2.2 Minimal K-state

A registration event is represented by a minimal K-state tuple:

$k = \langle M, o, \text{cert}, t, v \rangle$

where:

M = measurement act identifier
 o = registered outcome
 cert = certification status (self-certified or not)
 t = registration time
 v = validity status $\in \{0, 1\}$

The K-side registration space K is the collection of such tuples produced by a registering system R over time.

Provisional and final validity (Class D proposed):

Because E7 permits a later registration act to revise the validity of an earlier act, every validity assignment in K-side is provisional until the registration process closes. Formally:

$V_prov(M) = 1$ iff M satisfies Conditions 1-5 and no known later act has contradicted M within its K-side up to the current registration time.

$V_prov(M)$ may be revised to 0 if a later act M' with valid cross-registration authority contradicts M per E7.

$V_final(M) = 1$ iff $V_prov(M) = 1$ and no future registration act within K_R will contradict M (a limit property).

For the purposes of $requires_K_joint$ and $\square_K$, provisional validity is the operative notion unless stated otherwise.

2.3 Source: Buddhist Pramāṇa Epistemology

The registration-layer architecture is derived structurally from Buddhist Pramāṇa epistemology (Dignāga, 5th century; Dharmakīrti, 7th century), as systematized by Prasad (2023). Pramāṇa theory addresses the formal conditions under which a cognition constitutes a valid knowledge event. The Buddhist source relation is not used as empirical evidence for quantum mechanics. It functions as a structural source for the registration-layer postulates E1, E6, and E7; the move from Pramāṇa theory to K-side registration is a project-level formal extraction, not an established claim in Buddhist studies. The structural extraction used here concerns:

- **Svasaṃvedana:** Self-certifying cognition — a cognition that certifies its own occurrence without requiring a second-order cognition to validate it. Source for E1.
- **Arthakriyā:** Validity as functional success — a cognition is valid if it enables successful engagement with its object. Source for validity conditions in E7.
- **Anumāna/Pratyakṣa:** Inference and perception as distinct registration modes. Source for E11, E14.

This is a structural extraction, not a claim that Buddhist epistemology proves quantum mechanics or that quantum mechanics validates Buddhist doctrine.

3. The Valid Registered Measurement Test

This section states the central contribution of the paper: a six-condition test for when a physical interaction constitutes a valid registered measurement.

3.1 The Six Conditions

Interaction X is a valid registered measurement for registering system R if and only if:

Condition 1 (Physical):

X occurs at the physical ρ -side.

[Standard QM governs this layer. VVV-QMRF does not modify it.]

Condition 2 (Admission):

X is admitted into K-side as measurement act M_X for system R.
[The interaction crosses the layer boundary into K.]

Condition 3 (Process membership):

$M_X \sqsubseteq R$, where $R = \{M_{R1}, M_{R2}, \dots\}$ is the registering process of system R ordered by registration time. [E6]

Condition 4 (Self-certification):

$\sigma_R(M_X) = 1$: M_X self-certifies its occurrence within K_R , determined intrinsically, not by any external $M' \neq M_X$. [E1]

Condition 5 (Default validity):

$V(M_X) = 1$ by default upon admission. [E7]

Condition 6 (Non-invalidation):

No later registration act M' in K_R contradicts M_X such that $V(M_X)$ is revised to 0. [E7]

All six conditions must hold for X to count as a valid registered measurement.

3.2 The K-side Stopping Proposition

Proposition (K-side stopping, Class D proposed):

If Conditions 1–6 hold for measurement act M_X of registering system R, then no further K-side registration act is required to certify M_X . The von Neumann registration regress terminates at M_X on the K-side without modifying p-side dynamics.

Why this addresses the Heisenberg cut: Condition 4 (self-certification) provides the formal property that Copenhagen assigns to “classical apparatus” without definition. An apparatus has registration authority because $\sigma_R(M) = 1$ is determined intrinsically within K_R , requiring no external certifier. This is the registration-layer account of why the chain stops.

Claim class: D (proposed). Formal proof that Conditions 1–6 are jointly sufficient for chain termination is an open item.

3.3 Observer-Indexed Self-Certification

For multi-observer scenarios, the self-certification function carries an observer index:

$$\sigma_R(M) \sqsubseteq \{0, 1\}$$

$\sigma_R(M) = 1$ iff M occurred as a K-side registration event of registering system R, determined intrinsically within K_R , not by any $M' \neq M$ and not by any $R' \neq R$.

Independence: $\sigma_F(M_F)$ is determined within K_F independently of K_W , and vice versa.

The original $\sigma(M)$ is the single-observer special case: $\sigma(M) = \sigma_R(M)$ where R is unique. All properties of the single-observer case are preserved. [E01 Section 11.2, Class D]

4. Extended Wigner's Friend and K-side Incommensurability

4.1 The Extended Wigner's Friend Setup

In the Extended Wigner's Friend scenario (Frauchiger and Renner 2018; Brukner 2018), two observer pairs interact with a shared quantum system:

- **Friend F** measures quantum system S inside a sealed laboratory. From F's perspective, a definite outcome o_F is recorded: $\sigma_F(M_F) = 1$.
- **Wigner W** models the entire laboratory ($F + S$) as a unitarily evolving quantum state. From W's perspective, $F+S$ is in superposition before W's measurement. W then performs an interference measurement M_W on F's laboratory.

Standard QM describes both perspectives as valid without providing a formal account of their structural incompatibility. This is where the registration layer offers new structure.

4.2 Applying the Six Conditions to Each Observer

Friend F:

$X_F \rightarrow M_F \sqsubseteq R_F$ [Condition 3, E6]
 $\sigma_F(M_F) = 1$ [Condition 4, E1]
 $V_F(M_F) = 1$ [Condition 5, E7]
 No M' contradicts M_F [Condition 6, E7]
 $\rightarrow M_F$ is a valid registered measurement within K_F .

Wigner W:

$X_W \rightarrow M_W \sqsubseteq R_W$ [Condition 3, E6]
 $\sigma_W(M_W) = 1$ [Condition 4, E1]
 $V_W(M_W) = 1$ [Condition 5, E7]
 No M' contradicts M_W [Condition 6, E7]
 $\rightarrow M_W$ is a valid registered measurement within K_W .

Both observers satisfy the six conditions independently within their own K-sides.

4.3 The requires_K_joint Predicate

The question is whether a joint registration space K_{joint} can contain both K_F and K_W as jointly valid entries.

Definition (admissible joint K-side registration space, Class D proposed):

For two K-side registration structures A and B , an admissible $K_{\text{joint}}(A, B)$ is a proposed K-side registration space that can host both structures under a single validity architecture without changing the validity rules that made either structure valid in its own K-side. Formally:

$\text{AdmJoint}(K_{\text{joint}}; A, B) = 1$
 iff there exist embeddings $i_A: A \rightarrow K_{\text{joint}}$ and $i_B: B \rightarrow K_{\text{joint}}$

such that:

- (i) i_A and i_B preserve the registered act, outcome, certification, registration time/order, and validity status of A and B;
the embedding respects the internal time-order of each structure
(if $e1 <_A e2$ then $i_A(e1) <_{joint} i_A(e2)$), and the combined order in K_joint is the transitive closure of the two embedded orders plus any cross-structure temporal relations implied by the shared laboratory history;
- (ii) self-certification remains intrinsic to each embedded act;
- (iii) the Conditions 1-6 validity test remains satisfied for each embedded registration structure;
- (iv) no later registration-state update required by K_joint invalidates either embedded structure while both are still claimed as jointly valid;
- (v) K_joint does not import an external certifier M' as the source of the original self-certification for either side.

K_joint is therefore not a physical Hilbert space, not an observer's private K-space, and not a mere collection of records. It is a registration-layer candidate for joint validity preservation.

Definition (requires_K_joint predicate, Class D proposed):

For two K-side registration structures A and B, $requires_K_joint(A, B)=1$ means that the framework is not merely comparing two private records; it is asking whether both records can support one shared registration-validity claim. Formally:

$requires_K_joint(A, B) = 1$

iff A and B are each valid or provisionally valid within their own K-side

AND A and B are brought under a shared validity demand D_joint

AND D_joint requires both A and B to be assessed as parts of the same registration target, history, counterfactual claim, or validity claim

AND the truth or validity of D_joint cannot be evaluated while leaving A and B in fully independent K-side spaces

AND preserving D_joint requires a candidate K_joint in which A and B are embedded as jointly valid entries satisfying Conditions 1-6.

$requires_K_joint(A, B) = 0$

iff no shared validity demand D_joint is imposed, or D_joint can be evaluated without embedding A and B into one candidate K_joint .

Formally, D_joint is a predicate over pairs of K-side registration structures and a comparison architecture:

$D_joint(A, B, Arch) \in \{0, 1\}$

It evaluates to 1 when Arch demands that A and B support one shared registration-validity claim. The comparison architecture Arch is specified by the experimental design (EWF setup, LF inequality, direct comparison protocol, etc.). D_joint is not imposed by any single observer; it is a structural feature of the comparison architecture.

For the EWF case:

$\text{requires_K_joint}(F, W, M_F, M_W) = 1$
 iff $k_F = \langle M_F, o_F, \text{cert}_F, t_F, V_F \rangle$ and
 $k_W = \langle M_W, o_W, \text{cert}_W, t_W, V_W \rangle$
 are both claimed as valid entries for one laboratory registration
 history, one cross-observer comparison, or one LF-style
 observer-independent fact constraint.

Here D_{joint} is the demand that the two K-side structures be jointly valid under one comparison architecture. It is not a demand that both physical interactions be globally identical, and it is not a new physical postulate.

Operational sufficient conditions for $\text{requires_K_joint} = 1$:

- **Condition A (Wigner interference):** W performs an interference measurement on the laboratory containing F+S. M_W registers a superposition description of F+S; M_F registers a definite outcome o_F for S within the same laboratory registration history. Both claims concern the same registration history but place incompatible K-side validity requirements on that history. $\rightarrow \text{requires_K_joint} = 1$.
- **Condition B (Direct comparison):** F and W directly compare registration records and a logical contradiction is detectable. $\rightarrow \text{requires_K_joint} = 1$.
- **Condition B2 (LF observer-independent fact constraint):** an LF inequality or no-go argument requires F-side and W-side outcome claims to be assigned simultaneous cross-observer validity. $\rightarrow \text{requires_K_joint} = 1$.

Operational sufficient conditions for $\text{requires_K_joint} = 0$:

- **Condition C (No interference, no comparison):** W does not perform interference measurement on F's laboratory. K_F and K_W remain causally and inferentially isolated. $\rightarrow \text{requires_K_joint} = 0$.
- **Condition D (Separable state):** the shared quantum state $|\psi\rangle$ is separable and M_F, M_W act on non-overlapping subsystems. $\rightarrow \text{requires_K_joint} = 0$.
- **Condition E (Independent bookkeeping only):** A and B are stored in the same database, report, or external archive, but no shared validity demand is imposed on their contents. $\rightarrow \text{requires_K_joint} = 0$.

Claim class: D (proposed). The D_{joint} definition gives the necessary-and-sufficient structural core for when a joint K-side check is required. Conditions A-E remain operational sufficient cases and require experiment-specific instantiation.

4.4 Formal Definition of K-side Incommensurability

The symbol $\square_K$ denotes a registration-layer incommensurability relation, not a physical contradiction and not an event-level null value. It is defined relative to a proposed joint K-side space.

Definition (K-side incommensurability relation, Class D proposed):

Let A and B be K-side registration structures, where each may be an individual registration act (M_A, M_B) or an observer-indexed registration space (K_A, K_B). Let V_A, V_B be their respective validity conditions, including self-certification, registration order, and non-invalidation constraints. Then:

$A \square_K B$
 iff $\text{requires_K_joint}(A, B) = 1$

AND there exists no admissible K_joint such that

- (i) A and B both embed into K_joint ,
- (ii) their respective self-certifications are preserved,
- (iii) their respective validity conditions remain satisfied, and
- (iv) no required registration-state update in K_joint invalidates either A or B while both are still claimed as jointly valid.

For observer-indexed registration spaces, this yields:

$K_A \sqsubseteq K_B$
iff $requires_K_joint(K_A, K_B) = 1$
AND no admissible K_joint preserves both K_A and K_B
as jointly valid registration spaces under Conditions 1-6.

For individual registration acts inside a required joint comparison, use the E7 registered contradiction relation, not the space-level incommensurability symbol. The contradiction relation depends on a comparison context defined as follows:

Definition (K-side comparison context, Class D proposed):

A K-side comparison context C_K is a minimal shared frame in which two registration acts can be evaluated for compatibility. C_K requires:

- (a) both acts are admitted into the same comparison domain;
- (b) both acts are indexed to the same registration target or history;
- (c) the comparison does not presuppose that both acts are already jointly valid.

C_K is strictly weaker than $AdmJoint$: it enables the comparison without requiring that joint validity be preserved. $AdmJoint$ requires C_K but adds preservation of Conditions 1-6 and validity for both sides.

Definition (cross-registration authority, Class D proposed):

In a comparison context C_K containing acts $M_earlier$ and M_later :

M_later has valid cross-registration authority w.r.t. $M_earlier$ iff:

- (a) M_later is a valid registered measurement within its own K-side ($\sigma(M_later) = 1$, $V(M_later) = 1$ per Conditions 1-6);
- (b) M_later 's registration content directly concerns the same registration target or history as $M_earlier$;
- (c) M_later was produced by a measurement act that, within the comparison architecture, is structurally required to register the state of the same laboratory or composite system that $M_earlier$ registered as a subsystem;
- (d) the comparison architecture does not arbitrarily privilege one observer's K-side over the other – it only identifies which act is later in registration time and whether its content is structurally incompatible with the earlier act's content.

Definition (act-level registered contradiction, Class D proposed; grounded in E7):

$M_A \sqsubseteq M_B$
iff M_A and M_B share a K-side comparison context C_K ,
concern the same registration target, history, or validity claim,
cannot both satisfy the relevant validity conditions there,

and the later act has valid cross-registration authority with respect to the earlier act.

Act-level contradiction can support a conclusion that no admissible K_joint preserves the relevant K-side spaces, but it is not identical to the space-level relation $K_A \sqsubseteq K_B$.

Boundary clauses:

- $\square_K$ does not assert that either physical event fails to occur on the ρ -side.
- $\square_K$ does not mean that either observer's registered outcome is false within its own K-side.
- $\square_K$ is not equivalent to an event-level registration-null value. If an event fails to enter a valid K-registration domain, that should be written separately as $Null_K(e)$ or $R_K(e) = \square$, not conflated with $A \sqsubseteq K B$.
- $\square_K$ is a registration-layer relation proposed by VVV-QMRF; it is not a new postulate of Standard Quantum Mechanics.
- $\square_K$ is defined only for K-side structures that are each valid or provisionally valid within their own K-side. If either structure fails admission or is already invalidated within its own K-side, $\square_K$ does not apply; the situation should be described using the relevant single-K-side invalidation language of E7.

4.5 K_joint Failure and K-side Incommensurability

Bridge lemma (D_joint to E7 contradiction, Class D proposed; application status: Class C):

In an EWF/LF configuration, D_joint necessarily instantiates the E7 registered contradiction $M_W \sqsubseteq M_F$ when all of the following hold:

```
Bridge_EWF( $D\_joint$ ;  $M\_F$ ,  $M\_W$ ) = 1
iff  $D\_joint$  requires F-side and W-side registrations to be evaluated
    as jointly valid parts of one laboratory registration history
AND  $M\_F$  registers a definite friend-side outcome  $o\_F$ 
AND  $M\_W$  registers the same laboratory as a coherent superposition
    in which no definite  $o\_F$  is preserved as a W-side valid claim
AND the LF/no-go comparison requires both claims to support one
    cross-observer validity constraint
AND no reinterpretation inside the same  $K\_joint$  can preserve both
    registered contents without changing the validity claim of at
    least one side.
```

```
Bridge_EWF( $D\_joint$ ;  $M\_F$ ,  $M\_W$ ) = 1  $\rightarrow$   $M\_W \sqsubseteq M\_F$ .
```

This bridge is not triggered by mere difference of perspective. It is triggered only when the comparison architecture requires both registrations to support the same cross-observer validity claim.

Relativization defense and response (Class D proposed):

A possible objection: could K_joint host meta-level facts such as “within K_F , M_F registered $|h\rangle$ ” and “within K_W , M_W registered $|\Psi+\rangle$ ” without contradiction? Under this reading, both meta-descriptions are jointly true, and the original registration contents are never directly compared.

This relativization defense is rejected here because D_joint demands joint validity of the original registration claims, not meta-descriptions of them. If K_joint only hosts relativized meta-descriptions indexed

to different K-spaces, it does not satisfy D_{joint} : the LF/no-go constraint requires F's outcome and W's outcome to be assigned simultaneous cross-observer validity as facts about the same laboratory history, not as facts-about-facts whose validity is relativized to different K-side perspectives. Relativizing the contents abandons D_{joint} rather than satisfying it. The bridge is therefore not defeated by second-order redescription.

Conditional lemma (K_{joint} failure, Class D proposed; proof audit status: conditional):

If $requires_K_{joint} = 1$ via D_{joint} , $Bridge_EWF(D_{joint}; M_F, M_W) = 1$, and the later act has valid cross-registration authority with respect to the earlier act, then no admissible K_{joint} exists such that: - (i) $\sigma_F(M_F) = 1$ holds within K_{joint} [Condition 4 for F] - (ii) $\sigma_W(M_W) = 1$ holds within K_{joint} [Condition 4 for W] - (iii) $V(M_F) = 1$ and $V(M_W) = 1$ simultaneously remain preserved [Condition 5+6 for both] - (iv) no required registration-state update invalidates either embedded structure while both are still claimed as jointly valid [AdmJoint condition (iv)]

Proof audit:

Step 1 [E6]: F and W are distinct registering processes R_F and R_W with distinct K-side time sequences.

Step 2 [E1, observer-indexed]: $\sigma_F(M_F) = 1$ is intrinsic to K_F .
 $\sigma_W(M_W) = 1$ is intrinsic to K_W . Both hold independently.

Step 3 [$D_{joint} + Bridge_EWF$]: D_{joint} requires both M_F and M_W to support one cross-observer validity constraint. M_F registers o_F as definite. M_W registers F+S as coherent superposition with no definite o_F preserved as a W-side valid claim. Under $Bridge_EWF$, the two contents instantiate $M_W \sqsubseteq M_F$ inside the required comparison context.

Step 4 [E7 + AdmJoint condition (iv)]: E7 says that if $M' \sqsubseteq M$ and M' has valid cross-registration authority in the comparison context, then $V(M)$ may be revised to 0. Therefore, if $Bridge_EWF$ makes $M_W \sqsubseteq M_F$ a necessary registration-state update inside K_{joint} , then K_{joint} cannot also satisfy AdmJoint condition (iv), which requires that no embedded side be invalidated while both are still claimed as jointly valid.

Step 5 [Conditional conclusion]: Under those assumptions, no admissible K_{joint} preserving both K_F and K_W as jointly valid exists.

K-side incommensurability:

$K_F \sqsubseteq_K K_W$
iff $requires_K_{joint} = 1$
AND $Bridge_EWF(D_{joint}; M_F, M_W) = 1$
AND no admissible K_{joint} preserves both K_F and K_W as jointly valid under Conditions 1-6 and AdmJoint condition (iv).

Claim class of Step 4: C/D boundary. $Bridge_EWF$ is a proposed Class D bridge lemma, but its application to concrete EWF/LF experiments remains Class C until the paper supplies a full semantic proof that no admissible reinterpretation can preserve both claims inside the same K_{joint} , and until operational data criteria are supplied.

4.6 The Falsifiable Prediction

VVV-QMRF conjectures:

In Extended Wigner’s Friend experiments, configurations satisfying $\text{requires_K_joint} = 1$ are the natural candidates for LF-level violation, provided the quantum state and measurement settings are strong enough to expose the failure of a single joint registration space. Configurations satisfying $\text{requires_K_joint} = 0$ are predicted not to generate LF-level violation by the K-side registration mechanism alone.

$\text{requires_K_joint} = 1$ is therefore proposed as a necessary registration-layer condition for LF-level contradiction to be exposed, but it is not by itself sufficient for an observed LF inequality violation. Empirical violation also depends on entanglement strength, measurement settings, and experimental noise.

This is not a theorem of Standard Quantum Mechanics and is not yet a necessary-and-sufficient characterization of all LF violations.

Operational data criterion (ODC_K, Class C proposed):

K_joint is not directly observed by a detector. It is operationally tested through the existence or failure of a joint registration model for the observed EWF/LF probability data.

```
ODC_K(Data, Cfg) = K_joint_exists
  iff there exists a joint registration model J_K such that:
    (i) J_K assigns jointly valid entries to the relevant F-side and
        W-side registrations selected by Cfg;
    (ii) J_K preserves Conditions 1-3, 5, and 6 for each embedded
        registration; Condition 4 (self-certification) is treated as
        a structural postulate satisfied by construction when the
        registering system satisfies the architectural definition
        of a K-side observer (Section 3.3);
    (iii) J_K preserves AdmJoint condition (iv): no required
        registration-state update invalidates either embedded side
        while both are still claimed as jointly valid;
    (iv) J_K reproduces the observed probability distribution within
        the predeclared statistical tolerance  $\tau$ ;
    (v) J_K does not reclassify a Bridge_EWF configuration as a merely
        independent bookkeeping case.
```

```
ODC_K(Data, Cfg) = K_joint_fails
  iff Cfg satisfies  $\text{requires\_K\_joint} = 1$  via  $\text{D\_joint}$  and  $\text{Bridge\_EWF}$ ,
  AND no J_K satisfying (i)-(v) fits the observed data within  $\tau$ .
```

Here Cfg is the experimental configuration, including measurement settings, observer roles, and the prior classification of requires_K_joint . The tolerance τ must be fixed before evaluating the data; otherwise the criterion becomes post hoc.

Interpretation: $\text{ODC_K} = \text{K_joint_fails}$ does not mean that Standard QM fails. It means the data do not support one registration-layer joint-validity model preserving both K-side claims under the VVV-QMRF constraints.

τ sensitivity: The choice of statistical tolerance τ affects the sensitivity of ODC_K . A τ that is too large

risks false-negative $K_{\text{joint_fails}}$ conclusions (accepting poor models); a τ that is too small risks false-positive $K_{\text{joint_fails}}$ conclusions (rejecting models due to experimental noise). Calibration of τ requires specification before data collection and should be guided by the experiment’s statistical power and expected effect size. This paper does not fix τ ; it is a parameter to be set by the experimental design.

Claim class: C (conjecture). Pending experimental operationalization and verification.

5. Experimental Contact

5.1 What Distinguishes VVV-QMRF from Existing Frameworks

The key asymmetry is this: $\text{requires_K_joint} = 1$ is a necessary but not sufficient condition for LF violation. The quantum state must also be sufficiently entangled for K_{joint} failure to produce an observable inequality violation. This generates a two-regime structure:

- **Regime 1:** $\text{requires_K_joint} = 1$ AND state weakly entangled $\rightarrow$ Condition A structurally active, K_{joint} structurally required, but empirical LF violation may be absent.
- **Regime 2:** $\text{requires_K_joint} = 1$ AND state strongly entangled $\rightarrow$ K_{joint} failure is empirically visible as LF violation.

A decoherence-only framework does not distinguish these two regimes at the registration-layer level, because it contains no predicate equivalent to requires_K_joint for separating Bell non-LF violation from LF-level joint-validity failure.

This is not trivially true for all frameworks. It generates an observable expectation: Bell non-LF violation can appear in Regime 1 while LF inequalities remain unviolated, and LF violation appears in Regime 2 when entanglement is sufficient.

5.2 Proietti et al. (2019)

Reference: Proietti, M. et al. Experimental test of local observer independence. *Science Advances* 5(9), eaaw9832 (2019). DOI: 10.1126/sciadv.aaw9832.

Setup: Six-photon optical experiment with four observers (Alice’s Friend, Bob’s Friend, Alice, Bob). Two measurement choices per external observer:

- A0 / B0: read Friend’s memory record (no interference).
- A1 / B1: Wigner-type joint measurement via nonclassical interference on a 50/50 beam splitter.

Applying requires_K_joint :

Term	Configuration	W interference?	Condition	requires_K_joint	Role in S
$\square A1B1$	$\square$ Both Wigner-type	Both sides	A	1	Added
$\square A1B0$	$\square$ Alice Wigner, Bob record	Alice side	A	1	Added
$\square A0B1$	$\square$ Alice record, Bob Wigner	Bob side	A	1	Added
$\square A0B0$	$\square$ Both record readout	None	C	0	Subtracted

Experimental values (alternative observable protocol):

$$S = \langle A_1 B_1 \rangle + \langle A_1 B_0 \rangle + \langle A_0 B_1 \rangle - \langle A_0 B_0 \rangle$$

$$S = 0.571 + 0.577 + 0.573 - 0.662$$

$$S = 2.407 \quad (\text{reported: } 2.407 \pm 0.073, \text{ violation} > 5\sigma)$$

VVV-QMRF reading: The three terms with $\text{requires_K_joint} = 1$ contribute positively to the violated expression. The one term with $\text{requires_K_joint} = 0$ is subtracted. The allocation of Wigner-type and record-readout settings within the aggregate Bell-Wigner expression is compatible with the VVV-QMRF reading. However, the violation belongs to the aggregate S value, not to any single correlator term. This is a term-role interpretation of an aggregate inequality, not a per-configuration experimental confirmation.

5.3 Bong et al. (2020)

Reference: Bong, K.W. et al. A strong no-go theorem on the Wigner’s friend paradox. Nature Physics 16, 1199–1205 (2020). DOI: 10.1038/s41567-020-0990-x. arXiv: 1907.05607.

Setup: Extended Wigner’s Friend scenario with four observers (Charlie, Debbie, Alice, Bob). Three measurement settings per external observer. Friends implemented as photon paths within beam-displacer interferometers. State family:

$$\rho_{\mu} = \mu |\Phi_{HHHH}\Phi\rangle + (1-\mu)/2 (|HVHHHV\rangle + |VHHHHV\rangle)$$

where $\mu \in [0,1]$ controls singlet fraction.

Applying requires_K_joint to settings:

Setting pair	Alice side	Bob side	W interference?	Condition	requires_K_joint
$x=1, y=1$	Ask Charlie	Ask Debbie	None	C	0
$x=1, y=2/3$	Ask Charlie	Superobserve Bob side		A	1
$x=2/3, y=1$	Superobserver	Ask Debbie	Alice side	A	1
$x=2/3, y=2/3$	Superobserver	Superobserve Both sides		A	1

μ -threshold behavior and VVV-QMRF reading:

μ regime	Reported result	VVV-QMRF reading
Low μ	No inequalities violated	$\text{requires_K_joint} = 1$ structurally active in $x=2/3$ settings; LF violation absent because state too weakly entangled
$\mu = 0.80, 0.81$	Bell non-LF violated; LF inequalities NOT violated	Regime 1: Condition A active, K_joint required, but entanglement insufficient for LF-level failure
$\mu \approx 0.87$	First LF violation (Semi-Brukner)	Transition to Regime 2: K_joint failure becomes empirically visible
High μ	All categories violated including Genuine LF	Regime 2: $x=2/3, y=2/3$ Condition A strongly active, K_joint failure fully exposed

The asymmetric prediction: At $\mu = 0.80, 0.81$, Bell non-LF violation appears while LF inequalities remain unviolated. VVV-QMRF accounts for this via the two-regime structure: `requires_K_joint = 1` in superobserver settings (Condition A), but the quantum state at this μ value is not sufficiently entangled for `K_joint` failure to produce LF-level violation. A framework without an explicit analogue of `requires_K_joint` does not provide this particular registration-layer distinction between Bell-type violation and LF-type joint-validity failure at different μ thresholds.

Applying `requires_K_joint` and `Bridge_EWF` to individual LF facet terms:

Bong et al. report the LF polytope for $N=3, O=2$ has 932 facets grouped into 9 inequivalent classes. Five classes are relevant to the VVV-QMRF mapping:

Facet class	Settings	LF facet?	Bell facet?	<code>requires_K_joint</code> per term	<code>Bridge_EWF</code> per term	VVV-QMRF reading
Brukner	(1i, 1j)	Yes	Yes	Mixed: (1,1) $\rightarrow$ 0; (1,j),(i,1),(i,j) $\rightarrow$ 1	(1,1) $\rightarrow$ 0; others $\rightarrow$ 1	Violation driven by su- perobserver terms with re- <code>quires_K_joint=1</code> Violation $\rightarrow$ <code>K_joint</code> fails for all contributing terms
Semi- Brukner	(1i, 23)	Yes	Yes	All terms $\rightarrow$ 1 (no (1,1) term)	All terms $\rightarrow$ 1 ($\geq$ 1 su- perobserver per term)	Violation is Bell-level; does NOT constitute LF joint- validity failure
Bell non-LF	(23, 23)	No	Yes	All terms $\rightarrow$ 1	All terms $\rightarrow$ 1	Mixed; LF violation requires superobserver- contributed terms
I3322	(123, 123)	Yes	Yes	Mixed: (1,1) $\rightarrow$ 0; others $\rightarrow$ 1	(1,1) $\rightarrow$ 0; others $\rightarrow$ 1	Strongest test: violation $\rightarrow$ pure LF failure without Bell- analogue support
Genuine LF	(123, 123)	Yes	No	Mixed: (1,1) $\rightarrow$ 0; others $\rightarrow$ 1	(1,1) $\rightarrow$ 0; others $\rightarrow$ 1	

Per-setting-pair `Bridge_EWF` classification:

Bridge_EWF = 1 for $(x \in \{2,3\}, y=1)$: Bob reads Debbie $\rightarrow$ definite outcome.

Alice superobserver $\rightarrow$ registers superposition of Debbie's lab $\rightarrow M_{\text{Alice}} \sqsubset M_{\text{Debbie}}$.

Bridge_EWF = 1 for $(x=1, y \in \{2,3\})$: symmetric.

Bridge_EWF = 1 for $(x \in \{2,3\}, y \in \{2,3\})$: both register superpositions of both labs.

Bridge_EWF = 0 for $(x=1, y=1)$: both ask friends; no cross-observer validity conflict.

ODC_K prediction per facet class:

Facet class	$\mu=0.80\text{-}0.81$ prediction	High- μ prediction	Mechanism
Bell non-LF	K_joint_fails (violated)	K_joint_fails (violated)	Bell-level; superobserver correlators alone suffice
Brukner	K_joint_exists (NOT violated)	K_joint_fails (violated)	Friend-read terms at low μ preserve at least one J_K
Semi-Brukner	K_joint_exists (NOT violated)	K_joint_fails (violated)	Single-superobserver- side insufficient at low μ
Genuine LF	K_joint_exists (NOT violated)	K_joint_fails (violated)	All-setting joint model preservable at low μ ; fails at high μ

This is the per-facet operationalization of the two-regime structure: at $\mu=0.80\text{-}0.81$, Bell non-LF alone is violated because only superobserver-superobserver correlators are strong enough. The mixed-setting LF facets (Brukner, Semi-Brukner, Genuine LF) are not yet violated because the friend-read correlators dilute the joint K-side failure when entanglement is insufficient. As μ increases, the superobserver-contributed terms dominate and the full LF facets become violated.

Result: Setting-level, μ -threshold-level, and now per-facet-term-level classification is structurally compatible with the published Bong et al. data. The prediction hierarchy (Bell non-LF violates first, then Semi-Brukner, then Genuine LF) matches the observed μ -threshold ordering. Not yet a facet-level confirmation because the full ODC_K model-fit test (stage 3) has not been performed against the published correlator data.

5.4 Operational Use of ODC_K

For existing EWF/LF datasets, ODC_K should be applied in three stages:

1. **Configuration classification:** classify each measurement setting or LF facet term as `requires_K_joint = 1` or 0 using `D_joint` and the operational conditions in Section 4.3.
2. **Bridge classification:** for settings with `requires_K_joint = 1`, test whether `Bridge_EWF(D_joint; M_F, M_W) = 1` is satisfied by the registration contents required by that setting.
3. **Model-fit test:** attempt to construct a joint registration model J_K satisfying Conditions 1-6 and AdmJoint condition (iv), then compare its predicted probabilities with the observed data under a pre-declared tolerance τ .

A dataset supports the VVV-QMRF conjecture only if the settings classified as `requires_K_joint = 1` and `Bridge_EWF = 1` are precisely the settings where ODC_K returns `K_joint_fails`, while set-

tings classified as $\text{requires_K_joint} = 0$ remain compatible with K_joint_exists or require no joint K-side test.

This criterion is stronger than the current aggregate compatibility checks in Sections 5.2-5.3. Those checks show structural compatibility, not confirmation.

5.5 Purpose-Designed Experiment Specification (Class C proposed)

The existing-data checks in Sections 5.2–5.3 are compatibility checks, not confirmation. A purpose-designed experiment must vary requires_K_joint directly as an independent variable.

Core design principle: The same physical platform (photonic EWF) can switch between $\text{requires_K_joint} = 0$ and $\text{requires_K_joint} = 1$ by changing only the external observers’ measurement settings, while keeping the source, friends, and state identical.

Platform: Four-party photonic EWF setup with two friends (Charlie, Debbie, encoded as photon paths) and two superobservers (Alice, Bob), following the Bong et al. (2020) architecture:

- Friend-read setting ($x=1$ or $y=1$): external observer reads friend’s memory record directly. No interference on friend’s laboratory. $\rightarrow \text{requires_K_joint} = 0$ (Condition C or E).
- Superobserver setting ($x \in \{2,3\}$ or $y \in \{2,3\}$): external observer performs interference measurement on friend’s laboratory. $\rightarrow \text{requires_K_joint} = 1$ (Condition A) + $\text{Bridge_EWF} = 1$.

Independent variables:

Variable	How varied	Values
requires_K_joint	Measurement setting pairs (x,y)	0: ($x=1, y=1$); 1: ($x \in \{2,3\}, y=1$), ($x=1, y \in \{2,3\}$), ($x \in \{2,3\}, y \in \{2,3\}$)
Entanglement strength	State parameter $\mu \in [0,1]$	Scan from $\mu \approx 0.5$ to $\mu \approx 1.0$ in fine steps ($\Delta\mu \leq 0.05$)

Dependent variable: $\text{ODC_K}(\text{Data}, \text{Cfg})$ as defined in Section 4.6. For each (μ , setting-pair) combination:

1. Collect coincident photon counts.
2. Attempt to fit a joint registration model $\mathcal{J_K}$ satisfying Conditions 1-6 and AdmJoint condition (iv).
3. Compare model fit to data under predeclared tolerance τ .
4. Return K_joint_exists or K_joint_fails .

Predeclared tolerance τ : Must be fixed before data collection. Recommended: χ^2 goodness-of-fit test at $\alpha = 0.05$, or Bayesian model comparison with predeclared prior and Bayes factor threshold.

Predicted outcome (two-regime structure):

μ range	($x=1, y=1$) re- quires_K_joint=0	($x \in \{2,3\}, y \in \{2,3\}$) re- quires_K_joint=1	(mixed) re- quires_K_joint=1
Low μ (< 0.80)	K_joint_exists	K_joint_exists	K_joint_exists

μ range	$(x=1,y=1)$ re- quires_K_joint=0	$(x \in \{2,3\}, y \in \{2,3\})$ re- quires_K_joint=1	(mixed) re- quires_K_joint=1
Mid μ (0.80–0.87)	K_joint_exists	K_joint_fails (Bell non-LF)	K_joint_exists (LF not yet violated)
High μ (> 0.87)	K_joint_exists	K_joint_fails	K_joint_fails (LF violated)

The critical test is the mid- μ regime: Bell non-LF violation appears while LF inequalities remain unviolated. This is the Regime 1 $\rightarrow$ Regime 2 transition that a decoherence-only framework does not predict as a registration-layer distinction.

Falsification (any one suffices):

1. A setting with `requires_K_joint = 1` and `Bridge_EWF = 1` yields `ODC_K = K_joint_exists` at any μ where the corresponding LF inequality is independently violated.
2. A setting with `requires_K_joint = 0` yields `ODC_K = K_joint_fails`.
3. The predicted μ -hierarchy (Bell non-LF first, then Semi-Brukner, then Genuine LF) is inverted or absent.

Claim class: C (conjectural experiment specification). The specification is derived from the Class D formal chain but has not been peer-reviewed as an experimental protocol. Implementation requires an experimental collaboration with access to a photonic EWF platform.

5.6 Overall Condition 3 Status

Both Proietti et al. (2019) and Bong et al. (2020) are compatible with the VVV-QMRF conjecture. This is an existing-data compatibility check, not confirmation. The experiment specification in Section 5.5 defines what a confirmation-level test requires.

6. Positioning Against Existing Interpretations

Interpretation	Response to EWF paradox	VVV-QMRF difference
Copenhagen	WF is ill-posed; classical apparatus required	E1/E7 formalize why apparatus has registration authority: $\sigma_R(M) = 1$ intrinsically
Many-Worlds	All outcomes occur; no paradox	Registration events are singular per K-side, not globally branching
QBism	Facts are agent-relative; no paradox	Agrees on agent-relativity; adds formal K-side structure via six conditions
Relational QM	Facts are relation-relative; no paradox	Closest existing framework; VVV-QMRF adds formal registration machinery that Relational QM does not provide

Interpretation	Response to EWF paradox	VVV-QMRF difference
Objective Collapse	Physical collapse restores absolute facts	VVV-QMRF does not require physical collapse mechanism on ρ -side

Relation to Relational Quantum Mechanics: Rovelli (1996) argues that quantum states are relative to observers. VVV-QMRF adds the formal registration-layer structure that explains why they are relative: $\sigma_R(M)$ operates intrinsically within K_R , independently of $K_{\{R'\}}$. $K_F \sqsubseteq K_W$ is the registration-layer account of Rovelli’s observer-relativity. This is a genuine extension, not a restatement: VVV-QMRF supplies the formal conditions (E1, E6, E7) and the requires_K predicate that Relational QM does not.

7. Scope, Limitations, and Open Items

7.1 What This Paper Does Not Claim

- Does not replace Standard Quantum Mechanics or revise P1–P4.
- Does not revise the Born rule or unitary evolution.
- Does not claim Buddhist epistemology proves quantum mechanics.
- Does not claim K-side incommensurability is experimentally confirmed.
- Does not claim that consciousness plays any role in registration.
- Does not claim the EWF paradox is fully resolved on the ρ -side.
- Does not claim requires_K is a necessary-and-sufficient condition for all LF violations; it is proposed only as a necessary registration-layer condition for the K-side mechanism described here.

7.2 Formal Phase Summary and Open Items

The $\sqsubseteq_K$ formal definition chain is now complete at the proposed level (Class D/C). Each dependent layer has a registry-level definition and a working paper location:

Formal layer	Symbol	Class	Defined in
K-side incommensurability	$\sqsubseteq_K$	D	Section 4.4
Admissible joint K-space	$K_{\text{joint}} / \text{AdmJoint}$	D	Section 4.3
Joint-validity demand	$D_{\text{joint}} / \text{requires}_K$	D	Section 4.3
E7 registered contradiction	$M' \sqsubseteq M$	D	E7 framework §3b; Section 4.4
Bridge lemma	$\text{Bridge}_{\text{EWF}}$	D/C	Section 4.5
Operational data criterion	ODC_K	C	Section 4.6
LF facet-term classification	—	C	Section 5.3

Deferred proof items (not required for current claim class):

Deferred item	Reason deferred
Full semantic proof for <code>Bridge_EWF</code>	Requires formal semantics of “no admissible reinterpretation”; paper currently uses operational sufficient conditions
Full formal proof for $\square_K$ as a mathematical relation	Requires axiomatized K-space (topology, order); paper currently uses registration-layer structural definition
<code>AdmJoint</code> necessary-and-sufficient conditions	Currently sufficient conditions A-E; full characterization requires completed K-space axiomatization
Equivalence of $\sigma(M)$ and $R^\wedge_svasa$ formalisms Axiomatize K as a full mathematical structure	Separate research track (E01 §11.5) Long-term architectural task; not needed for current Class C/D claims → Addressed 2026-05-19: K1-K8 core axioms + T1-T4 bridge theorems formalized in <code>documents/research_documents/meta_architecture/1</code> (VVV-QMRF §K-AXIOM v1.5). Layer 1 (K1-K8) frozen; Layer 2 (T1-T4) pending Level 4 freeze. Concrete model (minimal EWF, 2 observers) completed: K1-K8 consistency verified, Level 4 derivation chain verified, T2 proof attempt done with 2 remaining gaps (relativization defense G1, Level 4 $\square$ freeze G3). Full Phase 1–6 RCA audit complete; all blocking issues resolved.

Next-step operational items (required for confirmation):

Next item	Current status	Priority
Purpose-designed VVV-QMRF experiment	Section 5.5	Class C experiment specification completed; requires experimental collaboration for implementation
ODC_K stage 3 model-fit test on published correlator data	Section 5.3 — classification done; fit not performed	High — would upgrade from structural compatibility to quantitative test
Apply <code>requires_K_joint</code> to Proietti raw count-level data	Section 5.2 — aggregate level only	Medium
Explicit separability verification for Bong ρ_μ regimes	Section 5.3 — not done	Medium

7.3 Falsification Condition (Restated)

This paper’s central conjecture is falsified if:

An Extended Wigner’s Friend experiment produces results consistent with a single `K_joint` registration model satisfying Conditions 1–6 for both observers simultaneously, for a configuration independently classified as `requires_K_joint` = 1 via `D_joint` and `Bridge_EWF`.

Operationally, this requires a predefined data criterion connecting observed probabilities to the existence of such a joint registration model. If K_joint exists empirically, E7 Axiom 2 or the Step 4 conditional bridge requires revision, and the K_joint failure proposition does not hold.

7.4 Architectural Constraints of the K-side/p-side Separation

The core architectural commitment of VVV-QMRF is $K \neq H$: the registration layer is structurally distinct from the physical layer (Section 2.1). This separation is what enables the framework to address the registration layer gap. It also imposes two architectural constraints that are not fixable by refinement — they are inherent consequences of the layer separation, not oversights.

Constraint 1 — Self-certification is not a p-side observable. Condition 4 requires $\sigma_R(M) = 1$ to be determined intrinsically within K_R . This cannot be a p-side observable because any physical measurement of σ would itself require a second-order registration act, re-entering the von Neumann regress that E1 is designed to terminate. Making σ directly testable from p-side data would collapse the K-side back into the p-side and remove the framework’s reason for existing. The operational compromise is stated in ODC_K condition (ii): self-certification is treated as a structural postulate satisfied by construction when the registering system meets the architectural definition of a K-side observer. What is tested is not whether $\sigma = 1$, but whether — given that assumption — the full joint registration model can reproduce the observed data.

Constraint 2 — D_joint is a framework-level classification, not a physical observable. The predicate $D_joint(A, B, Arch)$ classifies whether a comparison architecture demands joint validity. Experimental architectures are designed and described in the language of Standard QM plus assumptions (O, L, F, AOE, NSD); they do not carry intrinsic VVV-QMRF labels. The bridge from physical setup to K-side predicate is therefore interpretive. This does not make D_joint arbitrary: operational sufficient conditions A–E (Section 4.3) connect observable features of the setup (interference vs. readout, comparison vs. no comparison, separable vs. entangled state) to D_joint values in a reproducible way. But the mapping is a VVV-QMRF classification of the experimental architecture, not a measurement of a physical quantity.

These constraints are typical of the measurement problem, not unique to VVV-QMRF. Every interpretation or extension of quantum mechanics that addresses measurement must introduce a primitive concept that is not directly observable within the formalism:

Framework	Central concept	Why not directly observable
Copenhagen	Classical apparatus / Heisenberg cut	No formal definition of what makes an apparatus classical
Many-Worlds (Everett)	Branching of worlds	No physical observable distinguishes one branch from another
QBism	Agent’s degree of belief	Subjective probability is not a physical quantity
Relational QM (Rovelli)	Facts relative to observer	Relata are only defined within the relation
Objective Collapse (GRW)	Collapse mechanism	Collapse events are postulated, not yet detected
VVV-QMRF	Self-certification / D_joint	K-side properties are not p-side observables ($K \neq H$)

This is not a methodological coincidence. If the measurement problem could be resolved using only concepts

that are directly observable within the standard physical formalism, it would not have remained open for nearly a century. The presence of an unobservable primitive in every listed framework reflects the depth of the problem, not a defect unique to any one proposal.

The presence of architectural constraints does not distinguish VVV-QMRF from existing frameworks. What distinguishes it is that the constraints are explicitly named, traced to a single architectural commitment, and given operational bridges (ODC_K, conditions A–E) that define what can and cannot be tested under the current formalism.

References

1. von Neumann, J. (1932). *Mathematical Foundations of Quantum Mechanics*. Princeton University Press.
2. Bohr, N. (1935). Can Quantum-Mechanical Description of Physical Reality Be Considered Complete? *Physical Review* 48, 696–702.
3. Heisenberg, W. (1958). *Physics and Philosophy*. Harper & Row.
4. Wigner, E.P. (1961). Remarks on the mind-body question. In: *The Scientist Speculates*, Good (ed.), pp. 284–302.
5. Rovelli, C. (1996). Relational quantum mechanics. *International Journal of Theoretical Physics* 35, 1637–1678.
6. Prasad, H.S. (2023). Buddhist Pramāṇa-Epistemology. *Studia Humana* 12(1-2), 21–52. DOI: 10.2478/sh-2023-0004.
7. Zurek, W.H. (2003). Decoherence, einselection, and the quantum origins of the classical. *Reviews of Modern Physics* 75, 715–775.
8. Frauchiger, D. and Renner, R. (2018). Quantum theory cannot consistently describe the use of itself. *Nature Communications* 9, 3711.
9. Brukner, C. (2018). A no-go theorem for observer-independent facts. *Entropy* 20(5), 350.
10. Proietti, M. et al. (2019). Experimental test of local observer independence. *Science Advances* 5(9), eaaw9832. DOI: 10.1126/sciadv.aaw9832.
11. Bong, K.W. et al. (2020). A strong no-go theorem on the Wigner’s friend paradox. *Nature Physics* 16, 1199–1205. DOI: 10.1038/s41567-020-0990-x.
12. Jordan, A.N. and Siddiqi, I.A. (2024). *Quantum Measurement Theory and Practice*. Cambridge University Press. DOI: 10.1017/9781009103909.

Working Paper v2.0 — VVV-QMRF Posted to Zenodo for community feedback All formal claims are Class D (proposed) or Class C (conjecture) Falsification condition stated in Section 4.6 and Section 7.3 Experiment specification in Section 5.5; requires experimental collaboration for implementation