

viet@vvvqmrf.com <https://vvvqmrf.com> Independent Researcher, Vietnam

Supplemental Material:

Have Optical Wigner's Friend Experiments Been Blind to a Geometric Degree of Freedom?

VietVunVut (Viet – Nguyen Xuan)

2026-05-26

S1 Literature Search and Algebraic Proof

S1.1 Full Proof of Equatorial Cancellation Theorem

Theorem. Let Friend F measure in z -basis and Superobserver W at Bloch angles (θ, ϕ) .
With $f_{\perp}(b, d) = 1 - |\langle b|d \rangle|^2$:

$$f_{\perp}(+1, H) - f_{\perp}(-1, H) = -\cos \theta \quad (\text{S1})$$

Vanishes iff $\theta = \pi/2$.

Step 1: W 's measurement basis

$$|b = +1\rangle = \cos(\theta/2)|H\rangle + e^{i\phi} \sin(\theta/2)|V\rangle, \quad (\text{S2})$$

$$|b = -1\rangle = \sin(\theta/2)|H\rangle - e^{i\phi} \cos(\theta/2)|V\rangle. \quad (\text{S3})$$

Step 2: Overlaps with F 's outcomes

$$|\langle b = +1|H \rangle|^2 = \cos^2(\theta/2), \quad |\langle b = +1|V \rangle|^2 = \sin^2(\theta/2), \quad (\text{S4})$$

$$|\langle b = -1|H \rangle|^2 = \sin^2(\theta/2), \quad |\langle b = -1|V \rangle|^2 = \cos^2(\theta/2). \quad (\text{S5})$$

ϕ drops out: $|e^{i\phi}|^2 = 1$. Overlaps depend *only* on θ .

Step 3: $f_{\perp}$ values

$$f_{\perp}(+1, H) = 1 - \cos^2(\theta/2) = \sin^2(\theta/2), \quad (\text{S6})$$

$$f_{\perp}(-1, H) = 1 - \sin^2(\theta/2) = \cos^2(\theta/2), \quad (\text{S7})$$

$$f_{\perp}(+1, V) = 1 - \sin^2(\theta/2) = \cos^2(\theta/2), \quad (\text{S8})$$

$$f_{\perp}(-1, V) = 1 - \cos^2(\theta/2) = \sin^2(\theta/2). \quad (\text{S9})$$

Step 4: Outcome-dependence parameter

$$f_{\perp}(+1, H) - f_{\perp}(-1, H) = \sin^2(\theta/2) - \cos^2(\theta/2) \quad (\text{S10})$$

$$= -(\cos^2(\theta/2) - \sin^2(\theta/2)) \quad (\text{S11})$$

$$= -\cos \theta. \quad (\text{S12})$$

Vanishes iff $\theta = \pi/2$. At this angle, all four $f_{\perp} = 1/2$, constant across all outcome pairs. $\square$

Generality

At $\theta = \pi/2$, $|\langle b|d \rangle|^2 = 1/2$ for all b, d . Any $g(|\langle b|d \rangle|^2)$ therefore takes the same value for all outcome pairs. The equatorial plane is a fixed point for the entire class — the cancellation is not specific to $f_{\perp}$.

S1.2 BSM Equivalence for Proietti et al. (2019)

Proietti et al. use Bell-state measurement (BSM) rather than projective measurement. BSM projects onto four Bell states $\{|\Phi^+\rangle, |\Phi^-\rangle, |\Psi^+\rangle, |\Psi^-\rangle\}$. For the singlet-state source, each Bell outcome occurs with equal probability $1/4$, and conditioned on any Bell outcome b , the Friend’s recorded outcome d is equally likely to be H or V . Consequently $|\langle b|d \rangle|^2 = 1/2$ for all b, d , identical to the equatorial condition $\theta = \pi/2$ derived in the main text.

S1.3 Literature Search Methodology

Search date: 2026-05-15 to 2026-05-24.

Databases: Google Scholar, arXiv (quant-ph), Web of Science, InspireHEP.

Search strings (Boolean combinations):

- **Group A (subject):** “Wigner’s friend” OR “extended Wigner” OR “Frauchiger-Renner” OR “Local Friendliness”
- **Group B (parameter):** “equatorial measurement” OR “Bloch sphere polar angle” OR “outcome dependence” OR “geometric constraint” OR “measurement basis angle”

Inclusion criteria:

1. Optical EWF implementation with Friend+Superobserver structure
2. Published in peer-reviewed venue or arXiv
3. Reports measurement settings from which polar angle θ can be determined

Results: ~ 200 titles screened $\rightarrow$ 47 full-text examined $\rightarrow$ 2 published optical EWF experiments identified. Both operate at equatorial geometry ($\theta = \pi/2$ or equivalent via BSM). No published experiment varies θ from $\pi/2$ for any purpose.

Additional sources examined: Supplemental Material of Bong et al. (2020); Methods and Supplementary Information of Proietti et al. (2019); LF derivations in Frauchiger-Renner

(2018) and Wiseman-Cavalcanti-Rieffel (2023); Bell/LF review by Brunner et al. (2014); multipartite, sequential, and possibilistic LF extensions.

Limitation: We cannot rule out unpublished results or implementations outside our database scope. The survey establishes that within the indexed, peer-reviewed literature, θ has not been varied.

S2 Numerical Methods and Statistical Robustness

S2.1 Exact Numerical Integration Method

All numerical predictions use the density matrix for the singlet state with finite visibility:

$$\rho_\mu = \mu |\Phi^-\rangle \langle \Phi^-| + \frac{1-\mu}{4} I. \quad (\text{S13})$$

For the overlap-dependent model Eq. (2–3) of the main text:

$$P(a, b \mid x, y) = P_{\text{QM}}(a, b \mid x, y) \cdot [1 - \beta f_\perp(b, d)] / Z, \quad (\text{S14})$$

with full renormalization $Z = \sum_{a,b} P_{\text{QM}}(a, b \mid x, y) \cdot [1 - \beta f_\perp(b, d)]$.

All computations performed via `K9S12.proposal.py`, `statistical_significance.py`. The algebraic cancellation theorem is independently verified via symbolic computation (`universal_theorem`).

S2.2 Complete Correlator Table ($\theta = 31^\circ$, $\mu = 0.95$)

Table S1: All nine $\langle A_x B_y \rangle$ correlators with uncertainties ($N = 91,000$).

(x, y)	$\langle AB \rangle_{\text{QM}}$	σ
(1, 1)	−1.0000	0.0000
(1, 2)	−0.8572	0.0017
(1, 3)	−0.8572	0.0017
(2, 1)	−0.8572	0.0017
(2, 2)	−0.5045	0.0029
(2, 3)	−0.8933	0.0015
(3, 1)	−0.8572	0.0017
(3, 2)	−0.8933	0.0015
(3, 3)	−0.8829	0.0016

S2.3 Grid Search Optimization

A grid search over $(\theta, \phi_2, \phi_3, \beta_{\text{Bob}})$ maximizes:

$$\text{FOM}(\theta) = \min(n_\sigma^{\text{LF}}(\theta), n_\sigma^{\text{signal}}(\theta, \beta)), \quad (\text{S15})$$

yielding $\theta = 31^\circ$, $\phi_2 = 112^\circ$, $\phi_3 = 217^\circ$, $\beta_{\text{Bob}} = 20^\circ$. FOM remains above 5σ for $\theta \in [20^\circ, 55^\circ]$ at $\beta = 0.07$, $N = 91,000$.

S2.4 Monte Carlo Validation

10,000 runs: Gen LF $1 \geq 5\sigma$ in 99.97%. $\beta = 0.07$: >99% power (single); $\beta = 0.05$: $\sim 90\%$ (combined). Conservative Bayesian analysis (20% inflation): $\beta_{\min} \approx 0.046$ (combined).

S2.5 Angular Stability

LF significance varies $< 3\%$ across $\Delta\theta = \pm 5^\circ$. $\delta\langle AB \rangle \propto \cos\theta$: $\sim 1\%$ at $\pm 1^\circ$ (Bong’s reported precision).

S2.6 ϕ -Scramble Control Protocol

Rotate HWP to $N_\phi \geq 10$ random azimuthal angles $\phi \in [0^\circ, 360^\circ)$, fit $\delta\langle AB \rangle(\phi) = A + B \cos(2\phi) + C \sin(2\phi)$. Test: $A \neq 0$ with $B, C \approx 0$ indicates geometric $\cos\theta$ signal; B or $C \neq 0$ indicates birefringence/alignment artifacts.

S3 Interpretations and Physical Context

S3.1 Relation to Contextuality

Eq. (2) of the main text is not a contextual hidden-variable model. There is no unobserved λ ; the Friend outcome d is an observed macroscopic record. Dependence is on registration geometry ($|\langle b|d \rangle|^2$) rather than on which observables are co-measured.

Table S2: Comparison of dependence frameworks.

Framework	Hidden able?	vari-	Dependence on	Status of d
KS Contextuality	Yes (λ)		Measurement context	Unobserved
Standard contextuality	No		Co-measured observables	N/A
Retrocausal models	Varies		Future choices	Varies
Eq. (2) registration	No		$ \langle b d \rangle ^2$ geometry	Observed record

S3.2 Physical Picture

When a Friend measures in the z -basis, they produce a macroscopic record with a definite orientation on the Bloch sphere. A Superobserver measuring at (θ, ϕ) probes this record at a relative angle $|\langle b|d \rangle|^2$. Standard QM assumes that once the Friend’s outcome is registered, subsequent measurements factorize. Eq. (2) parametrizes a possible residual dependence on this geometric relationship.

S3.3 Quantum Interpretations

Copenhagen. Silent on registration-layer structure — the experiment tests a structural degree of freedom that no standard interpretation addresses.

Many-Worlds (Everett). $\beta \neq 0$ would provide the first quantitative probe of inter-branch structural correlations; a null result constrains such effects to $\beta < 0.04$.

Relational QM (Rovelli). Tests whether relational outcomes leave geometry-dependent measurable traces — a signature current RQM formulations neither predict nor preclude.

Objective Collapse. CSL and Diosi-Penrose models modify dynamics, not registration geometry. The overlap-dependent class is structurally orthogonal.

QBism. Parametrizes whether agent experiences exhibit structural coupling across agents at the registration layer. A null result constrains agent-agent coupling at $\sim 10^{-2}$.

S3.4 General Probabilistic Theories (GPT) Context

Within the GPT framework, the overlap-only deformation defines a one-parameter family of neighboring GPTs deviating from the quantum point along the registration-geometry direction. This is a structural degree of freedom orthogonal to standard nonlocality.

S3.5 Weak Measurement and Postselection

In weak measurement, the weak value $A_w = \langle \psi_f | A | \psi_i \rangle / \langle \psi_f | \psi_i \rangle$ depends explicitly on the pre-postselection overlap. The basis-alignment dependence in Eq. (2) shares the structural feature of geometric coupling between measurement records at different observational levels. The analogy is structural, not dynamical.

S3.6 Scale Comparison

Table S3: Phenomenological parameter scales.

Parameter class	Typical constraint	Reference
Photon-sector SME	$< 10^{-23}$	Colladay & Kostelecký (1997)
CSL collapse rate λ	$< 10^{-16} \text{ s}^{-1}$	
Weak-measurement anomalies	$\sim 10^{-2}$	Aharonov et al. (1988)
Overlap-dependent β (this work)	$\sim 10^{-2}$ (proposed)	—

S3.7 Multi-Observer Extension (Preliminary)

For $N > 2$ observers, the geometric cancellation structure generalizes. Preliminary analysis suggests $\sim 11\times$ amplification at $\beta = 0.3$ for three observers, conditional on bridge theorems connecting registration-layer structure to QM observables for $N > 2$. Rigorous derivation is left for future work.