

(Cytoscapeを使った) データの可視化

ライフサイエンス統合データベースセンター

岡本 忍

ネットワーク表現の利用

論文の情報を可視化する

Induction of Pluripotent Stem Cells from Mouse Embryonic and Adult Fibroblast Cultures by Defined Factors

Kazutoshi Takahashi¹ and Shinya Yamanaka^{1,2,*}¹Department of Stem Cell Biology, Institute for Frontier Medical Sciences, Kyoto University, Kyoto 606-8507, Japan²CREST, Japan Science and Technology Agency, Kawaguchi 332-0012, Japan

*Contact: yamanaka@frontier.kyoto-u.ac.jp

DOI 10.1016/j.cell.2006.07.024

SUMMARY

Differentiated cells can be reprogrammed to an embryonic-like state by transfer of nuclear contents into oocytes or by fusion with embryonic stem (ES) cells. Little is known about factors that induce this reprogramming. Here, we demonstrate induction of pluripotent stem cells from mouse embryonic or adult fibroblasts by introducing four factors, Oct3/4, Sox2, c-Myc, and Klf4, under ES cell culture conditions. Unexpectedly, Nanog was dispensable. These cells, which we designated iPS (induced pluripotent stem) cells, exhibit the morphology and growth properties of ES cells and express ES cell marker genes. Subcutaneous transplantation of iPS cells into nude mice resulted in tumors containing a variety of tissues from all three germ layers. Following injection into blastocysts, iPS cells contributed to mouse embryonic development. These data demonstrate that pluripotent stem cells can be directly generated from fibroblast cultures by the addition of only a few defined factors.

INTRODUCTION

Embryonic stem (ES) cells, which are derived from the inner cell mass of mammalian blastocysts, have the ability to grow indefinitely while maintaining pluripotency and the ability to differentiate into cells of all three germ layers (Evans and Kaufman, 1981; Martin, 1981). Human ES cells might be used to treat a host of diseases, such as Parkinson's disease, spinal cord injury, and diabetes (Thomson et al., 1998). However, there are ethical difficulties regarding the use of human embryos, as well as the problem of tissue rejection following transplantation in patients. One way to circumvent these issues is the generation of pluripotent cells directly from the patients' own cells.

Somatic cells can be reprogrammed by transferring their nuclear contents into oocytes (Wilmut et al., 1997)

or by fusion with ES cells (Cowan et al., 2005; Tada et al., 2001), indicating that unfertilized eggs and ES cells contain factors that can confer totipotency or pluripotency to somatic cells. We hypothesized that the factors that play important roles in the maintenance of ES cell identity also play pivotal roles in the induction of pluripotency in somatic cells.

Several transcription factors, including Oct3/4 (Nichols et al., 1998; Niwa et al., 2000), Sox2 (Avilion et al., 2003), and Nanog (Chambers et al., 2003; Mitsui et al., 2003), function in the maintenance of pluripotency in both early embryos and ES cells. Several genes that are frequently upregulated in tumors, such as Stat3 (Matsuda et al., 1999; Niwa et al., 1998), E-Ras (Takahashi et al., 2003), c-myc (Cartwright et al., 2005), Klf4 (Li et al., 2005), and β -catenin (Kelman et al., 2002; Sato et al., 2004), have been shown to contribute to the long-term maintenance of the ES cell phenotype and the rapid proliferation of ES cells in culture. In addition, we have identified several other genes that are specifically expressed in ES cells (Maruyama et al., 2005; Mitsui et al., 2003).

In this study, we examined whether these factors could induce pluripotency in somatic cells. By combining four selected factors, we were able to generate pluripotent cells, which we call induced pluripotent stem (iPS) cells, directly from mouse embryonic or adult fibroblast cultures.

RESULTS

We selected 24 genes as candidates for factors that induce pluripotency in somatic cells, based on our hypothesis that such factors also play pivotal roles in the maintenance of ES cell identity (see Table S1 in the Supplemental Data available with this article online). For β -catenin, c-Myc, and Stat3, we used active forms, S33Y- β -catenin (Sadot et al., 2002), T58A-c-Myc (Chang et al., 2000), and Stat3-C (Bromberg et al., 1999), respectively. Because of the reported negative effect of Grb2 on pluripotency (Burdon et al., 1999; Cheng et al., 1998), we included its dominant-negative mutant Grb2 Δ SH2 (Miyamoto et al., 2004) as 1 of the 24 candidates.

Oct3/4

Sox2

c-Myc

Klf4


Nanog

Stat3

E-Ras

 β -catenin



Grb2



シアノバクテリア, 藍藻

Synechocystis sp. PCC 6803

- 光合成によって酸素を生み出す
- 光合成研究のモデル生物
- 1996年に日本で世界で光合成生物としてはじめてゲノムが決定された



シアノバクテリアの研究論文と遺伝子のネットワーク図

