

2 **Population Genetics Based Phylogenetics Under**
3 **Stabilizing Selection for an Optimal Amino Acid**
4 **Sequence: A Nested Modeling Approach**

5 JEREMY M. BEAULIEU^{1,2,3}, BRIAN C. O'MEARA^{2,3}, RUSSELL ZARETZKI⁴,
6 CEDRIC LANDERER^{2,3}, JUANJUAN CHAI^{2,5}, AND MICHAEL
7 A. GILCHRIST^{2,3,*}

8 ¹Department of Biological Sciences, University of Arkansas, Fayetteville, AR 72701

9 ²Department of Ecology & Evolutionary Biology, University of Tennessee, Knoxville, TN
10 37996-1610

11 ³National Institute for Mathematical and Biological Synthesis, Knoxville, TN 37996-3410

12 ⁴Department of Business Analytics & Statistics, Knoxville, TN 37996-0532

13 ⁵Current address: 50 Main St, Suite 1039, White Plains, NY 10606

14 *Corresponding author. E-mail: mikeg@utk.edu

ABSTRACT

16 We present a novel phylogenetic approach based of substitution rates generated from a
17 nested population genetics model. Unlike many simpler codon models, which assume a
18 single substitution matrix for all sites, our model more realistically represents the evolution
19 of protein-coding DNA under the assumption of consistent, stabilizing selection for a gene
20 specific, optimal amino acid sequence. By modeling the cost-benefit function of an amino
21 acid sequence, our model naturally links the strength of stabilizing selection to protein
22 synthesis levels, which, in turn, can be estimated. Our new set of models, which we
23 collectively call SelAC (Selection on Amino acids and Codons), fit phylogenetic data much
24 better than popular models, suggesting strong potential for more accurate inference of
25 phylogenetic trees and branch lengths from existing data. SelAC also demonstrates that a
26 large amount of biologically meaningful information is accessible when using a nested set of
27 mechanistic models. For example, SelAC prediction of gene specific protein synthesis rates
28 correlates well with both empirical ($r = 0.34 - 0.48$) and other theoretical predictions
29 ($r = 0.59 - 0.64$) for multiple species. SelAC also provides estimates of which amino acid is
30 optimal for a given site. Finally, because SelAC is a nested approach based on clearly
31 stated biological assumptions, it can be simplified or expanded as needed, such as including
32 shifts in the optimal amino acid sequence within or across lineages.

Phylogenetic analyses plays a critical role in most aspects of biology, particularly in the fields of ecology, evolution, paleontology, medicine, and conservation. While the scale and impact of phylogenetic studies has increased substantially over the past two decades, the realism of the mathematical models on which these analyses are based has changed relatively little by comparison. The most popular models of DNA substitution used by molecular phylogenetics are simple nucleotide models that are indifferent to the type of sequences to which they are applied. For example, when evaluating protein-coding sequences these models are inherently agnostic with regards to the different amino acid substitutions and their impact on gene function (e.g. F81, F84, HYK85, TN93, and GTR, see Yang (2014) for an overview) and, as a result, cannot describe the behavior of natural selection at the amino acid or protein level.

To address this critical shortcoming, Goldman and Yang (1994) and Muse and Gaut (1994) independently introduced models which assumed that differences in the physicochemical properties between amino acids, or physicochemical distances for short, could affect substitution rates. These physicochemical approaches as originally described have rarely been used for empirical data; instead these models have served as the basis for an array of simpler and, in turn, more popular models that, starting with Yang and Nielsen (1998); Nielsen and Yang (1998), typically assume an equal fixation probability for *all* non-synonymous mutations. Thus, these simpler models initially employed a single term ω to model the differences in fixation probability between nonsynonymous and synonymous changes at all sites. To improve their realism, more complex forms have been developed that allow ω to vary between sites or branches (as cited in Anisimova 2012) and include selection on different synonyms for the same amino acid (e.g. Yang and Nielsen 2008)

In Goldman and Yang (1994); Yang and Nielsen (1998); Nielsen and Yang (1998) and later studies based on their work, ω is suggested to indicate whether a given site within a protein sequence is under consistent ‘stabilizing ($\omega < 1$) or ‘diversifying’ ($\omega > 1$) selection.

However, the model's actual behavior is inconsistent with how these terms are typically defined and understood (e.g. see Pellmyr 2002). Because synonymous substitutions have a higher substitution rate than any possible non-synonymous substitutions when $\omega < 1$, the model behaves as if the resident amino acid i at a given site is favored by natural selection. Even when ω is allowed to vary between sites, the symmetrical nature of the model means that for any given site the strength of selection for the resident amino acid i over its 19 alternatives is equally strong regardless of their physicochemical properties. Paradoxically, selection for amino acid i persists *until* a substitution for another amino acid, j , occurs. As soon as amino acid j fixes, but not before, selection now favors amino acid j equally over all other amino acids, including amino acid i . This is now the opposite scenario from when i was the resident. Similarly, when $\omega > 1$, synonymous substitutions have a lower substitution rate than any possible non-synonymous substitutions from the resident amino acid. Again due to the model's symmetry, the selection *against* the resident amino acid i is equally strong relative to alternative amino acids. The selection against the resident amino acid i persists until a substitution occurs at which point selection now *favors* amino acid i , as well as the 18 other amino acids, to the same degree i was previously disfavored.

Thus, the simplest and most consistent interpretation of ω is that it represents the rate at which the *selective environment itself* changes, and this change in selection perfectly coincides with the fixation of a new amino acid. This, in turn, implies that the rate of shifts in the optimal (or pessimal) amino acid is on the time scale as the rate of substitution. Contrary to popular belief, ω does not describe whether a site is evolving under a constant regime of stabilizing or diversifying selection, but instead how a very particular selective environment changes over time. Given this behavior, ω based models are likely to only reasonably approximate a subset of scenarios such as perfectly symmetrical over-/under-dominance or positive/negative frequency dependent selection (Hughes and Nei 1988; Nowak 2006).

Here we propose an approach where selection explicitly favors minimizing the cost-benefit function η of a protein whose relative performance is determined by the order and physicochemical properties of its amino acids. Our approach, which we call SelAC (Selection on Amino acids and Codons), is developed in the same vein as previous phylogenetic applications of the Wright-Fisher process (e.g. Muse and Gaut 1994; Halpern and Bruno 1998; Yang and Nielsen 2008; Rodrigue et al. 2005; Koshi and Goldstein 1997; Koshi et al. 1999; Dimmic et al. 2000; Thorne et al. 2012; Lartillot and Philippe 2004; Rodrigue and Lartillot 2014). Similar to Lartillot’s work (Lartillot and Philippe 2004; Rodrigue and Lartillot 2014), we assume there is a finite set of rate matrices describing the substitution process and that each position within a protein is assigned to a particular rate matrix category. Unlike Lartillot’s work, we assume *a priori* there are 20 different families of rate matrices, one family for when a given amino acid is favored at a site. The key parameters underlying these matrices are shared across genes except for gene expression. As a result, SelAC allows us to quantitatively evaluate the support for a particular amino acid being favored at a particular position within the protein sequence.

MATERIALS & METHODS

Overview

We model the substitution process as a classic Wright-Fisher process which includes the forces of mutation, selection, and drift (Fisher 1930; Kimura 1962; Wright 1969; Iwasa 1988; Berg and Lässig 2003; Sella and Hirsh 2005; McCandlish and Stoltzfus 2014). For simplicity, we ignore linkage effects and, as a result of this and other assumptions, sequences evolve in a site independent manner.

Because SelAC requires twenty families of 61×61 matrices, the number of parameters needed to implement SelAC would, without further assumptions, be extremely large (i.e. on the order of 74,420 parameters). To reduce the number of parameters needed, while still maintaining a high degree of biological realism, we construct our gene and amino acid specific substitution matrices using a submodel nested within our substitution model, similar to approaches in Gilchrist (2007); Shah and Gilchrist (2011); Gilchrist et al. (2015).

One advantage of a nested modeling framework is that it requires only a handful of genome-wide parameters such as nucleotide specific mutation rates (scaled by effective population size N_e), amino acid side chain physicochemical weighting parameters, and a shape parameter describing the distribution of site sensitivities. In addition to these genome-wide parameters, SelAC requires a gene g specific expression parameter ψ_g which describes the average rate at which the protein's functionality is produced by the organism or a gene's 'average functionality production rate' for short (for notational simplicity, we will ignore the gene specific indicator $_g$, unless explicitly needed). Currently, ψ is fixed across the phylogeny, though relaxing this assumption is a goal of future work. The gene specific parameter ψ is multiplied by additional model terms to make a composite term ψ' which scales the strength and efficacy of selection for the optimal amino acid sequence relative to drift (see Implementation below). In terms of the functionality of the protein encoded, we assume that for any given gene there exists an optimal amino acid sequence \vec{a}^* and that, by definition, a complete, error free peptide consisting of \vec{a}^* provides one unit of the gene's functionality. We also assume that natural selection favors genotypes that are able to synthesize their proteome more efficiently than their competitors and that each savings of an high energy phosphate bond per unit time leads to a constant proportional gain in fitness A_0 . SelAC also requires the specification (as part of parameter optimization) of an optimal amino acid a^* at each position within a coding sequence. This requirement of one a^* per site makes our \vec{a}^* the largest category of parameters SelAC estimates. Despite

the need to specify a^* for each site, because we use a submodel to derive our substitution matrices, SelAC estimates a relatively small number of the parameters when compared to more general approaches where the fitness of each amino acid is allowed to vary freely of any physicochemical properties (Halpern and Bruno 1998; Lartillot and Philippe 2004; Rodrigue and Lartillot 2014).

As with other phylogenetic methods, SelAC generates estimates of branch lengths and nucleotide specific mutation rates. In addition, the method can also be used to make quantitative inferences on the optimal amino acid sequence of a given protein as well as the realized average synthesis rate of each protein used in the analysis. The mechanistic basis of SelAC also means it can be easily extended to include more biological realism and test more explicit hypotheses about sequence evolution.

Mutation Rate Matrix μ

We begin with a 4x4 nucleotide mutation matrix μ that describes mutation rates between different bases and, in turn, different codons. For our purposes, we rely on the general unrestricted model (UNREST from Yang 1994) because it imposes no constraints on the instantaneous rate of change between any pair of nucleotides. More constrained models, such as the Jukes-Cantor (JC), Hasegawa-Kishino-Yano (HKY), or the general time-reversible model (GTR), could also be used.

The 12 parameter UNREST model defines the relative rates of change between a pair of nucleotides. Thus, we arbitrarily set the G→T mutation rate to 1, resulting in 11 free mutation rate parameters in the 4x4 mutation nucleotide mutation matrix. The nucleotide mutation matrix is also scaled by a diagonal matrix π whose entries, $\pi_{i,i}$, correspond to the equilibrium frequencies of each base. These equilibrium nucleotide frequencies are determined by analytically solving $\pi \times \mathbf{Q} = 0$. We use this \mathbf{Q} to populate a 61×61 codon mutation matrix μ , whose entries $\mu_{i,j}$ $i \neq j$ describes the mutation rate from

codon i to j and $\mu_{i,i} = -\sum_j \mu_{i,j}$. We generate this matrix using a “weak mutation” assumption, such that evolution is mutation limited, codon substitutions only occur one nucleotide at a time. As a result, the rate of change between any pair of codons that differ by more than one nucleotide is zero.

While the overall model does not assume equilibrium, we still need to scale our mutation matrices μ by a scaling factor S . As traditionally done, we rescale our time units such that at equilibrium, one unit of branch length represents one expected mutation per site (which equals the substitution rate under neutrality, but would not with selection). More explicitly, $S = -(\sum_{i \in \text{codons}} \mu_{i,i} \pi_{i,i})$ where the final mutation rate matrix is the original mutation rate matrix multiplied by $1/S$.

Protein Synthesis Cost-Benefit Function η

SelAC links fitness to the product of the cost-benefit function of a gene η and the organism’s average target synthesis rate of the functionality provided by gene ψ . This is because the average flux energy an organism spends to meet its target functionality provided by the gene is, by definition, $\eta \times \psi$. Compensatory changes that allow an organism to maintain functionality even with loss of one or both copies of a gene are widespread (reviewed in ¹); here we assume that for finer scale problems than entire loss (for example, a 10% loss of functionality) the compensation is more production of the protein. In order to link genotype to our cost-benefit function $\eta = \mathbf{C}/\mathbf{B}$, we begin by defining our benefit function \mathbf{B} .

¹From Cruft: There is evidence of compensation for protein function. Metabolism with gene expression models (ME-models) link those factors to successfully make predictions about response to perturbations in a cell <https://www.nature.com/articles/ncomms1928>, <https://www.sciencedirect.com/science/article/pii/S0958166914002316>. For example, an ME-model for *E. coli* successfully predicted gene expression levels in vivo <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0045635>.

MIKE:
Brian, please
provide
references or
cut. BRIAN:
Cedric had
suggestions,
or we can
pull from
footnote

Benefit: Our benefit function \mathbf{B} measures the functionality of the amino acid sequence \vec{a}_i encoded by a set of codons \vec{c}_i , i.e. $a(\vec{c}_i) = \vec{a}_i$ relative to that of an optimal sequence \vec{a}^* . By definition, $\mathbf{B}(\vec{a}^*|\vec{a}^*) = 1$ and $\mathbf{B}(\vec{a}_i|\vec{a}^*) < 1$ for all other sequences. We assume all amino acids within the sequence contribute to protein function and that this contribution declines as an inverse function of physicochemical distance between each amino acid and the optimal one. Formally, we assume that

$$\mathbf{B}(\vec{a}|\vec{a}^*) = \left(\frac{1}{n} \sum_{p=1}^n (1 + G_p d(a_p, a_p^*)) \right)^{-1} \quad (1)$$

where n is the length of the protein, $d(a_p, a_p^*)$ is a weighted physicochemical distance between the amino acid encoded at a given position p and a_p^* is the optimal amino acid for that position. For simplicity, we assume all nonsense mutations are lethal by defining the physicochemical distance between a stop codon and a sense codon as ∞ . The term G_p describes the sensitivity of the protein's function to physicochemical deviation from the optimum at site position p . There are many possible measures for physicochemical distance; we use Grantham (1974) distances by default, though others may be chosen. We assume that $G_p \sim \text{Gamma}(\text{shape} = \alpha_G, \text{rate} = \alpha_G)$ in order to ensure $\mathbb{E}(G_p) = 1$. Given the definition of the Gamma distribution, the variance in G_p is equal to $\text{shape}/\text{rate}^2 = 1/\alpha_G$. Further, at the limit of $\alpha_G \rightarrow \infty$, the model becomes equivalent to assuming uniform site sensitivity where $G_p = 1$ for all positions p . Finally, we note that $\mathbf{B}(\vec{a}_i|\vec{a}^*)$ is inversely proportional to the average physicochemical deviation of an amino acid sequence \vec{a}_i from the optimal sequence \vec{a}^* weighted by each site's sensitivity to this deviation. $\mathbf{B}(\vec{a}_i|\vec{a}^*)$ can be generalized to include second and higher order terms of the distance measure d .

Cost: Protein synthesis involves both direct and indirect assembly costs. Direct costs consist of the high energy phosphate bonds $\sim P$ of ATP or GTP's used to assemble the ribosome on the mRNA, charge tRNA's for elongation, move the ribosome forward along the transcript, and terminate protein synthesis. As a result, direct protein assembly costs are the same for all proteins of the same length. Indirect costs of protein assembly are potentially numerous and could include the cost of amino acid synthesis as well the cost and efficiency with which the protein assembly infrastructure such as ribosomes, aminoacyl-tRNA synthetases, tRNAs, and mRNAs are used. When these indirect costs are combined with sequence specific benefits, the probability of a mutant allele fixing is no longer independent of the rest of the sequence (Gilchrist et al. 2015) and, as a result, model fitting becomes substantially more complex. Thus for simplicity, in this study we ignore indirect costs of protein assembly that vary between genotypes and define,

$$\mathbf{C}(\vec{c}_i) = \text{Energetic cost of protein synthesis.} \quad (2)$$

$$= A_1 + A_2 n \quad (3)$$

193 where, A_1 and A_2 represent the direct cost, in high energy phosphate bonds, of ribosome
194 initiation and peptide elongation, respectively, where $A_1 = A_2 = 4 \sim P$.

195 *Defining Physicochemical Distances*

Assuming that functionality declines with an amino acid a_i 's physicochemical distance from the optimum amino acid a^* at each site provides a biologically defensible way of mapping genotype to protein function that requires relatively few free parameters. In addition, SelAC naturally lends itself to model selection since we can compare the quality of SelAC fits using different mixtures of physicochemical properties. Following Grantham (1974), we focus on using composition c , polarity p , and molecular volume v of each amino

acid's side chain residue to define our distance function, but the model and its implementation can flexibly handle a variety of properties. We use the Euclidian distance between residue properties where each property c , p , and v has its own weighting term, α_c , α_p , α_v , respectively, which we refer to as 'Grantham weights'. Because physicochemical distance is ultimately weighted by a gene's specific average protein synthesis rate ψ , another parameter we estimate, there is a problem with parameter identifiability. The scale of gene expression is affected by how we measure physicochemical distances which, in turn, is determined by our choice of Grantham weights. As a result, by default we set $\alpha_v = 3.990 \times 10^{-4}$, the value originally estimated by Grantham, and recognize that our estimates of α_c and α_p and ψ are scaled relative to this choice for α_v . More specifically,

$$d(a_i, a^*) = \left(\alpha_c [c(a_i) - c(a^*)]^2 + \alpha_p [p(a_i) - p(a^*)]^2 + \alpha_v [v(a_i) - v(a^*)]^2 \right)^{1/2}.$$

Linking Protein Synthesis to Allele Substitution

Next we link the protein synthesis cost-benefit function η of an allele with its fixation probability. First, we assume that each protein encoded within a genome provides some beneficial function and that the organism needs that functionality to be produced at a target average rate ψ . Again, by definition, the optimal amino acid sequence for a given gene, \vec{a}^* , produces one unit of functionality, i.e. $\mathbf{B}(\vec{a}^*) = 1$. Second, we assume that the actual average rate a protein is synthesized ϕ is regulated by the organism to ensure that functionality is produced at rate ψ . As a result, it follows that $\phi = \psi / \mathbf{B}(\vec{a} | \vec{a}^*)$ and the cost of a suboptimal amino acid increases the more it decreases the protein's functionality, \mathbf{B} . In other words, the average production rate of a protein \vec{a} with relative functionality $\mathbf{B}(\vec{a}) < 1$ must be $1/\mathbf{B}(\vec{a} | \vec{a}^*)$ times higher than the production rate needed if the optimal amino acid

sequence \vec{a}^* was encoded since $\mathbf{B}(\vec{a}^*|\vec{a}^*) = 1$. For example, a cell with an allele \vec{a} where $\mathbf{B}(\vec{a}|\vec{a}^*) = 9/10$ would have to produce the protein at rate $\phi = 10/9 \times \psi = 1.11\psi$. Similarly, a cell with an allele \vec{a} where $\mathbf{B}(\vec{a}|\vec{a}^*) = 1/2$ will have to produce the protein at $\phi = 2\psi$. In contrast, a cell with the optimal allele \vec{a}^* would have to produce the protein at rate $\phi = \psi$.

Third, we assume that every additional high energy phosphate bond, $\sim P$, spent per unit time to meet the organism's target function synthesis rate ψ leads to a slight and proportional decrease in fitness W . This assumption, in turn, implies

$$W_i(\vec{c}) \propto \exp[-A_0 \eta(\vec{c}_i)\psi]. \quad (4)$$

where A_0 , again, describes the proportional decline in fitness with every $\sim P$ wasted per unit time. Because A_0 shares the same time units as ψ and ϕ and only occurs in SelAC in conjunction with ψ , we do not need to explicitly identify our time units. Instead, we recognize that our estimates of ψ share an unknown scaling term.

Correspondingly, the ratio of fitness between two genotypes is,

$$W_i/W_j = \exp[-A_0 \eta(\vec{c}_i)\psi] / \exp[-A_0 \eta(\vec{c}_j)\psi] \quad (5)$$

$$= \exp[-A_0 (\eta(\vec{c}_i) - \eta(\vec{c}_j)) \psi] \quad (6)$$

$$(7)$$

Given our formulations of \mathbf{C} and \mathbf{B} , the fitness effects between sites are multiplicative and, therefore, the substitution of an amino acid at one site can be modeled independently of the amino acids at the other sites within the coding sequence. As a result, the fitness ratio

for two genotypes differing at a multiple site simplifies to

$$W_i/W_j = \exp \left[- \left(\frac{A_0 (A_1 + A_2 n_g)}{n_g} \right) \sum_{p \in \mathbb{P}} [d(a_{i,p}, a_p^*) - d(a_{j,p}, a_p^*)] G_p \psi \right]$$

where \mathbb{P} represents the codon positions in which \vec{c}_i and \vec{c}_j differ. Fourth, we make a weak mutation assumption, such that alleles can differ at only one position at any given time, i.e. $|\mathbb{P}| = 1$, and that the population is evolving according to a Wright-Fisher process. As a result, the probability a new mutant, j , introduced via mutation into a resident population i with effective size N_e will go to fixation is,

$$\begin{aligned} u_{i,j} &= \frac{1 - (W_i/W_j)^b}{1 - (W_i/W_j)^{2N_e}} \\ &= \frac{1 - \exp \left\{ -\frac{A_0}{n_g} (A_1 + A_2 n_g) [d(a_i, a^*) - d(a_j, a^*)] G_p \psi b \right\}}{1 - \exp \left\{ -\frac{A_0}{n_g} (A_1 + A_2 n_g) [d(a_i, a^*) - d(a_j, a^*)] G_p \psi 2N_e \right\}} \end{aligned}$$

where $b = 1$ for a diploid population and 2 for a haploid population (Kimura 1962; Wright 1969; Iwasa 1988; Berg and Lässig 2003; Sella and Hirsh 2005). Finally, assuming a constant mutation rate between alleles i and j , $\mu_{i,j}$, the substitution rate from allele i to j can be modeled as,

$$q_{i,j} = \frac{2}{b} \mu_{i,j} N_e u_{i,j}.$$

215 where, given the substitution model's weak mutation assumption, $N_e \mu \ll 1$. In the end,
 216 each optimal amino acid has a separate 64 x 64 substitution rate matrix \mathbf{Q}_a , which
 217 incorporates selection for the amino acid (and the fixation rate matrix this creates) as well
 218 as the common mutation parameters across optimal amino acids. This results in the

creation of 20 \mathbf{Q} matrices, one for each amino acid and each with 3,721 entries which are based on a relatively small number of model parameters (one to 11 mutation rates, two free Grantham weights, the cost of protein assembly, A_1 and A_2 , the gene specific target functionality synthesis rate ψ , and optimal amino acid at each position p , a_p^*). These model parameters can either be specified *a priori* and/or estimated from the data.

Given our assumption of independent evolution among sites, it follows that the probability of the whole data set is the product of the probabilities of observing the data at each individual site. Thus, the likelihood \mathcal{L} of amino acid a being optimal at a given site position p is calculated as

$$\mathcal{L}(\mathbf{Q}_a|\mathbf{D}_p, \mathbf{T}) \propto \mathbf{P}(\mathbf{D}_p|\mathbf{Q}_a, \mathbf{T}) \quad (8)$$

In this case, the data, \mathbf{D}_p , are the observed codon states at position p for the tips of the phylogenetic tree with topology \mathbf{T} . For our purposes we take \mathbf{T} as given but it could be estimated as well. The pruning algorithm of Felsenstein (1981) is used to calculate $\mathcal{L}(\mathbf{Q}_a|\mathbf{D}_p, \mathbf{T})$. The log of the likelihood is maximized by estimating the genome scale parameters which consist of 11 mutation parameters which are implicitly scaled by $2N_e/b$, and two Grantham distance parameters, α_c and α_p , and the sensitivity distribution parameter α_G . Because A_0 and ψ_g always co-occur and are scaled by N_e , for each gene g we estimate a composite term $\psi'_g = \psi_g A_0 b N_e$ and the optimal amino acid for each position a_p^* of the protein. When estimating α_G , the likelihood then becomes the average likelihood which we calculate using the generalized Laguerre quadrature with $k = 4$ points (Felsenstein 2001).

Finally, we note that because we infer the ancestral state of the system, our approach does not rely on any assumptions of model stationarity. Nevertheless, as our branch lengths grow the probability of observing a particular amino acid a at a given site

approaches a stationary value proportional to $W(a)^{2N_e-b}$ and any effects of mutation bias (Sella and Hirsh 2005).

Implementation

All methods described above are implemented in the new R package, **selac** available through GitHub (<https://github.com/bomeara/selac>) [it will be uploaded to CRAN once peer review has completed]. Our package requires as input a set of fasta files that each contain an alignment of coding sequence for a set of taxa, and the phylogeny depicting the hypothesized relationships among them. In addition to the SelAC models, we implemented the GY94 codon model of Goldman and Yang (1994), the FMutSel mutation-selection model of Yang and Nielsen (2008), and the standard general time-reversible nucleotide model that allows for Γ distributed rates across sites. These likelihood-based models represent a sample of the types of popular models often fit to codon data.

For the SelAC models, the starting guess for the optimal amino acid at a site comes from ‘majority’ rule, where the initial optimum is the most frequently observed amino acid at a given site (ties resolved randomly). Our optimization routine utilizes a four stage hill climbing approach. More specifically, within each stage a block of parameters are optimized while the remaining parameters are held constant. The first stage optimizes the block of branch length parameters. The second stage optimizes the block of gene specific composite parameters $\psi'_g = A_0\psi_g N_e b$. The third stage optimizes SelAC’s parameters shared across the genome α_c and α_p , and the sensitivity distribution parameter α_G . The fourth stage estimates the optimal amino acid at each site a^* . This entire four stage cycle is repeated six more times, using the estimates from the previous cycle as the initial conditions for the new one. The search is terminated when the improvement in the log-likelihood between cycles is less than 10^{-8} at which point we consider the ML solution found and the search is terminated. For optimization of a given set of parameters, we rely

MIKE: I thought we updated this to be based on the change in LLik function.
JMB: Is this clear now?
MAC: Yes

on a bounded subplex routine (Rowan 1990) in the package **NLOpt** (Johnson 2012) to maximize the log-likelihood function. To help the optimization navigate through local peaks, we perform a set of independent analyses with different sets of naive starting points with respect to the gene specific composite ψ' parameters, α_c , and α_p . Confidence in the parameter estimates can be generated by an ‘adaptive search’ procedure that we implemented to provide an estimate of the parameter space that is some pre-defined likelihood distance (e.g., 2 lnL units) from the maximum likelihood estimate (MLE), which follows Beaulieu and OMeara (2016) and Edwards (1984).

We note that our current implementation of SelAC is painfully slow, and is best suited for data sets with relatively few number of taxa (i.e. < 10). This limitation is largely due to the size and quantity of matrices we create and manipulate to calculate the log-likelihood of an individual site. Ongoing work will address the need for speed, with the eventual goal of implementing SelAC in popular phylogenetic inference toolkits, such as RevBayes (Hhna et al. 2016), PAML (Yang 2007) and RAxML (Stamatakis 2006).

Simulations

We evaluated the performance of our codon model by simulating datasets and estimating the bias of the inferred model parameters from these data. Our ‘known’ parameters under a given generating model were based on fitting SelAC to the 106 gene data set and phylogeny of Rokas et al. (2003). The tree used in these analyses is outdated with respect to the current hypothesis of relationships within *Saccharomyces*, but we rely on it simply as a training set that is separate from our empirical analyses (see section below). Bias in the model parameters were assessed under two generating models: one where we assumed a model of SelAC assuming uniform sensitivity across sites (i.e. $G_p = 1$ for all sites, i.e. $\alpha_G = \infty$), and one where we used the Gamma distribution joint shape and rate parameter α_G estimated from the empirical data. Under each of these two scenarios, we

used parameter estimates from the corresponding empirical analysis and simulated 50 five-gene data sets. For the gene specific composite parameter ψ'_g the ‘known’ values used for the simulation were five evenly spaced points along the rank order of the estimates across the 106 genes. The MLE estimate for a given replicate were taken as the fit with the highest log-likelihood after running five independent analyses with different sets of naive starting points with respect to the composite ψ'_g parameter, α_c , and α_p . All analyses were carried out in our **selac** R package.

Analysis of yeast genomes & tests of model adequacy

We focus our empirical analyses on the large yeast data set and phylogeny of Salichos and Rokas (2013). The yeast genome is an ideal system to examine our phylogenetic estimates of gene expression and its connection to real world measurements of these data within individual taxa. The complete data set of Salichos and Rokas (2013) contain 1070 orthologs, where we selected 100 at random for our analyses. We also focus our analyses on *Saccharomyces sensu stricto* and their sister taxon *Candida glabrata*, and we used the phylogeny depicted in Fig. 1 of Salichos and Rokas (2013) for our fixed tree. We fit the two SelAC models described above (i.e., SelAC and SelAC+ Γ), as well as two codon models, GY94 and FMutSel, and a standard GTR + Γ nucleotide model. The FMutSel model assumes that the amino acid frequencies are determined by functional requirements of the protein while the other models make no assumptions about amino acid frequencies. In all cases, we assumed that the model was partitioned by gene, but with branch lengths linked across genes.

For SelAC, we compared our estimates of $\phi' = \psi'/\mathbf{B}$, which represents the average protein synthesis rate of a gene, to estimates of gene expression from empirical data. Specifically, we obtained gene expression data for five of the six species used - four species were measured during log-growth phase, whereas the other was measured at the beginning

of the stationary phase (*S. kudriavzevii*) from the Gene Expression Omnibus (GEO). Gene expression in this context corresponds to mRNA abundances which were measured using either microarrays (*C. glabrata*, *S. castellii*, and *S. kudriavzevii*) or RNA-Seq (*S. paradoxus*, *S. mikatae*, and *S. cerevisiae*).

For further comparison, we also predicted the average protein synthesis rate for each gene ϕ by analyzing gene and genome-wide patterns of synonymous codon usage using ROC-SEMPPR (Gilchrist et al. 2015) for each individual genome. While, like SelAC, ROC-SEMPPR uses codon level information, it does not rely on any inter-specific comparisons and, unlike SelAC, uses only the intra- and inter-genic frequencies of synonymous codon usage as its data. Nevertheless, ROC-SEMPPR predictions of gene expression ϕ correlates strongly (Pearson $r = 0.53 - 0.74$) with a wide range of laboratory measurements of gene expression (Gilchrist et al. 2015).

While one of our main objectives was to determine the improvement of fit that SelAC has with respect to other standard phylogenetic models, we also evaluated the adequacy of SelAC. Model fit, measured with assessments such as the Akaike Information Criterion (AIC), can tell which model is least bad as an approximation for the data, but it does not reveal whether a model is actually doing a good job of representing the data. An adequate model does the latter, one measure of which is that data generated under the model resemble real data (Goldman 1993). For example, Beaulieu et al. (2013) assessed whether parsimony scores and the size of monomorphic clades of empirical data were within the distributions of simulated data under a new model and the best standard model; if the empirical summaries were outside the range for each, it would have suggested that neither model was adequately modeling this part of the biology.

In order to test adequacy for a given gene we first remove a particular taxon from the data set and the phylogeny. A marginal reconstruction of the likeliest sequence across all remaining nodes is conducted under the model, including the node where the pruned

taxon attached to the tree. The marginal probabilities of each site are used to sample and assemble the starting coding sequence. This sequence is then evolved along the branch, periodically being sampled and its current functionality assessed. We repeat this process 100 times and compare the distribution of trajectories against the observed functionality calculated for the gene. For comparison, we also conducted the same test, by simulating the sequence under the standard GTR + Γ nucleotide model, which is often used on these data but does not account for the fact that the sequences are protein coding, and under FMutSel, which includes selection on codons but in a fundamentally different way as our model.

The appropriate estimator of bias for AIC

As part of the model set described above, we also included a reduced form of each of the two SelAC models, SelAC and SelAC+ Γ . Specifically, rather than optimizing the amino acid at any given site, we assume the the most frequently observed amino acid at each site is the optimal amino acid a^* . We refer to these ‘majority rule’ models as SelAC_M and SelAC_M + Γ and the majority rule parameterization accelerates model fitting.

Since these majority rule models assume that the optimal amino acids are known prior to fitting of our model, it is tempting to reduce the count of estimated parameters in the model by the number of parameters estimated using majority rule. Despite having become standard behavior in the field of phylogenetics, this reduction is statistically inappropriate unless one uses an additional dataset for this purpose, something we have not seen. Thus, although using majority rule does not necessarily provide the most likely parameter estimate, it still uses the data to generate the estimate and, thus, represents a parameter estimated from the data. Because the difference in the number of parameters K when counting or not counting the number of nucleotide sites drops out when comparing nucleotide models with AIC, this statistical issue does not apply to nucleotide models. It

364 does, however, matter for AICc, where K and the sample size n combine in the penalty
365 term. This also matters in our case, where the number of estimated parameters for the
366 majority rule estimation differs based on whether one is looking at codons or single
367 nucleotides.

368 In phylogenetics two variants of AICc are used. In comparative methods
369 (e.g. Butler and King 2004; O’Meara et al. 2006; Beaulieu et al. 2013) the number of data
370 points, n , is taken as the number of taxa. More taxa allow the fitting of more complex
371 models, given more data. However, in DNA evolution, which is effectively the same as a
372 discrete character model used in comparative methods, the n is taken as the number of
373 sites. Obviously, both cannot be correct. This uncertainty was highlighted by Posada and
374 Buckley (2004): they chose to use number of sites, but mentioned in their discussion that
375 sample size also depends on the number of taxa. Sullivan and Joyce (2005) also mention
376 that the number of sites is often taken as sample size, but whether that is appropriate in
377 phylogenetics is not entirely clear. One approach incorporating both number of taxa and
378 sites is in calculating AICc is the program SURFACE implemented by Ingram and Mahler
379 (2013), which uses multiple characters and taxa. While its default is to use AIC to
380 compare models, if one chooses to use AICc, the number of samples is taken as the product
381 of number of sites and number of taxa.

382 Recently, Jhwueng et al. (2014) performed an analysis that investigated what
383 variant of AIC and AICc worked best as an estimator, but the results were inconclusive.
384 Here, we have adopted and extended the simulation approach of Jhwueng et al. (2014) in
385 order to examine a large set of different penalty functions and how well they approximate
386 the remaining portion of the Kullback-Liebler (KL) divergence between two models after
387 accounting for the deviance (i.e., $-2\mathcal{L}$) (see Appendix 1 for more details).

388 RESULTS

By linking transition rates $q_{i,j}$ to gene expression ψ , our approach allows use of the same model for genes under varying degrees of stabilizing selection. Specifically, we assume the strength of stabilizing selection for the optimal sequence, \vec{a}^* , is proportional to the average protein synthesis rate ϕ , which we can estimate for each gene. In regards to model fit, our results clearly indicated that linking the strength of stabilizing selection for the optimal sequence to gene expression substantially improves our model fit. Further, including the shape parameter α_G for the random effects term $G \sim \text{Gamma}(\text{shape} = \alpha_G, \text{rate} = \alpha_G)$ to allow for heterogeneity in this selection between sites within a gene improves the ΔAICc of SelAC+ Γ over the simpler SelAC models by over 22,000 AIC units. Using either ΔAICc or AIC_w as our measure of model support, the SelAC models fit extraordinarily better than GTR + Γ , GY94, or FMutSel (Table 1). This is in spite of the need for estimating the optimal amino acid at each position in each protein, which accounts for 49,881 additional model parameters. Even when compared to the next most parameter rich codon model in our model set, FMutSel, SelAC+ Γ model shows over 160,000 AIC unit improvement over FMutSel.

The analysis building upon Jhwueng et al. (2014) suggests that using the number of taxa times the number of sites as the sample size performs best as a small sample size correction for estimating Kullback-Liebler distance in phylogenetic models. This also has intuitive appeal: in models that have at least some parameters shared across sites and some parameters shared across taxa, increasing the number of sites and/or taxa should be adding more samples for the parameters to estimate. This is consistent in considering how likelihood is calculated for phylogenetic models: the likelihood for a given site is the sum of the probabilities of each observed state at each tip, which is then multiplied across sites. It is arguable that the conventional approach in comparative methods is calculating AICc in the same way. That is, if only one column of data (or "site") is examined, as remains remarkably common in comparative methods, when we refer to sample size, it is technically

the number of taxa multiplied by number of sites, even though it is referred to simply as the number of taxa.

With respect to estimates of ϕ within SelAC, they were strongly correlated with both our empirical measurements (Pearson $r = 0.34 - 0.48$) and theoretical predictions (Pearson $r = 0.59 - 0.64$) of gene expression (Figure 1 and Figures S1-S2, respectively). In other words, using only codon sequences, our model can predict which genes have high or low expression levels. The estimate of the α_G parameter, which describes the site-specific variation in sensitivity of the protein's functionality, indicated a moderate level of variation in gene expression among sites. Our estimate of $\alpha_G = 1.36$, produced a distribution of sensitivity terms G ranged from 0.342-7.32, but with more than 90% of the weight for a given site-likelihood being contributed by the 0.342 and 1.50 rate categories. In simulation, however, of all the parameters in the model, only α_G showed a consistent bias, in that the MLE were generally lower than their actual values (see Supporting Materials). Other parameters in the model, such as the Grantham weights, provide an indication as to the physicochemical distance between amino acids. Our estimates of these weights only strongly deviate from Grantham's 1974 original estimates in regards to composition weight, α_c , which is the ratio of noncarbon elements in the end groups to the number of side chains. Our estimate of the composition weighting factor of $\alpha_c=0.459$ is 1/4th the value estimate by Grantham which suggests that the substitution process is less sensitive to this physicochemical property when shared ancestry and variation in stabilizing selection are taken into account.

It is important to note that the nonsynonymous/synonymous mutation ratio, or ω , which we estimated for each gene under the FMutSel model strongly correlated with our estimates of $\phi' = \psi'/\mathbf{B}$ where \mathbf{B} depends on the sequence of each taxa. In fact, ω showed similar, though slightly reduced correlations, with the same empirical estimates of gene expression described above (Figure 2) This would give the impression that the same

conclusions could have been gleaned using a much simpler model, both in terms of the number of parameters and the assumptions made. However, as we discussed earlier, not only is this model greatly restricted in terms of its biological feasibility, SelAC clearly performs better in terms of its fit to the data and biological realism.

For example, when we simulated the sequence for *S. cerevisiae*, starting from the ancestral sequence under both GTR + Γ and FMutSel, the functionality of the simulated sequence moves away from the observed sequence, whereas SelAC remains near the functionality of the observed sequence (Figure 3b). This is somewhat unsurprising, given that both GTR + Γ and FMutSel are agnostic to the functionality of the gene, but it does highlight the improvement in biological realism in amino acid sequence evolution that SelAC provides. We do note that the adequacy of the SelAC model does vary among individual taxa, and does not always match the observed functionality. For instance, our simulations of *S. castellii* gene function is consistently higher than estimated from the data (Figure 3c). We suspect this is an indication that assuming a single set of optimal amino acid across all taxa is too simplistic. However, we cannot rule out violations of SelAC's other model assumptions such as: a single set of Grantham weights, a single α_G , or reductions in protein functionality \mathbf{B} being solely a function of physicochemical distances d between sites.

Finally, we note that our simulation analysis suggested that the best measure of dataset size for estimating KL distance uses a scaled value of the product of number of sites and number of characters. The model comparison approach described above included this assumption. For more details on the simulation approach, see Appendix 1.

DISCUSSION

Biologically, we know protein-coding DNA sequences largely evolve through the introduction of new mutations that either become fixed or lost due to selection and/or

drift. Selection on protein coding regions can take many forms, but can generally be related to the cost of producing the protein and the functional benefit (or potential harm) caused by the protein product. The gene specific cost of protein synthesis can be affected by the amino acids used, the direct and indirect costs of peptide assembly by the ribosome, the use of chaperones to aid in folding, and even the expected lifespan of the protein. Importantly, these costs can be computed to varying degrees of realism (e.g. Wagner 2005; Lynch and Marinov 2015). We have previously presented models of protein synthesis costs that, alternatively, take into account the cost of ribosome pausing (Shah and Gilchrist 2011) or premature termination errors (Gilchrist and Wagner 2006; Gilchrist 2007; Gilchrist et al. 2009).

Protein benefit or ‘function’ can be affected by the amino acids at each site and their interactions. As a result, amino acid substitutions can affect the functionality at key catalytic sites or, more broadly, the probability of a particular protein fold and, in turn, the expected functionality of the protein. Linking amino acid sequence to protein function is a daunting task; thus for simplicity, we assume that for any given desired biological function to be carried out by a protein, that (a) the biological importance of this protein function is invariant across the tree, (b) single optimal amino acid sequence that carries out this function best, and (c) the functionality of alternative amino acid sequences declines with their physicochemical distance from the optimum on a site by site basis.

We readily acknowledge that sequence space may have more than one local optimum, that physicochemical distance from the optimal primary amino acid sequence is likely a poor model of protein function, and that the biological importance of a function can vary over time. Nevertheless, we believe our cost-benefit approach to be a substantial advance of the more simplistic ω models, is complementary to the work of others in the field (e.g. Thorne et al. 2012; Rodrigue and Lartillot 2014), and, in turn, lays the foundation for more realistic work in the future.

For instance, by assuming there is an optimal amino acid for each site, SelAC naturally leads to a non-symmetrical and, thus, more cogent model of protein sequence evolution. Because the strength of selection depends on an additive function of amino acid physicochemical properties, an amino acid more similar to the optimum has a higher probability of replacing a more dissimilar amino acid than the converse situation. Further, SelAC does not assume the system is always at the optimum or pessimum point of the fitness landscape, as occurs when $\omega < 1$ or > 1 , respectively.

Importantly, the cost-benefit approach underlying SelAC allows us to link the strength of selection on a protein sequence to its gene's expression level. Despite its well recognized importance in determining the rate of protein evolution (e.g. Drummond et al. 2005, 2006), phylogenetic models have ignored the fact that expression levels vary between genes. In order to link gene expression and the strength of stabilizing selection on protein sequences, we simply assume that the strength of selection on a gene is proportional to the average protein synthesis rate of the gene.

One possible mechanism that generates a linear relationship between the strength of selection and gene expression is the assumption of compensatory gene expression. That is, the assumption that any reduction in protein function is compensated for by an increase in the protein's production rate and, in turn, abundance. For example, a mutation which reduces the functionality of the protein to 90% of the optimal protein, would require $1/0.9 = 1.11$ of these suboptimal proteins to be produced relative to the optimal protein in order to maintain the same amount of that protein's functionality in the cell. Because the energetic cost of an 11% increase in a protein's synthesis rate is proportional to its target synthesis rate, our assumptions naturally link changes in protein functionality and changes in gene expression and its associated costs. Under what circumstances cells actually respond in this manner, remains to be determined. The fact that our method allows us to explain 13-23% of the variation in gene expression measured using RNA-Seq, suggests that

this assumption is a reasonable starting point. More importantly, by linking the strength of stabilizing selection for an optimal amino acid sequence to gene expression, we can effectively weight the phylogenetic information encoded in high and low expression genes which tend to evolve at different rates.

Because SelAC infers the optimal amino acid for each site, it is substantially more parameter rich than more commonly used models such as GTR+ Γ , GY94, and FMutSel. Despite this increase in number of model parameters, SelAC drastically outperforms these models with AICc values on the order of 10,000s to 100,000s. We predict that SelAC's performance could be improved even further if we use a hierarchical approach where the optimal amino acid is not estimated on a per site basis, but rather as a vector of probability an amino acid is optimal at the gene level.

SelAC makes inferences about the tree, but also about population genetic parameters, and we can validate the assumptions indirectly by comparing our inferences to other empirical data, such as we do with protein synthesis data. More generally, SelAC's assumptions lead to mechanistic and, thus, testable hypothesis about the relationship between mutation, protein function, gene expression, and rates of evolution. More importantly, alternative hypotheses could be used in place of ours and, in turn, phylogenetic and other types of data could be used to evaluate the support of these alternative models. Our hope is that by moving away from the more phenomenological models we can better connect population genetics, molecular biology, and phylogenetics allowing each area inform the others more effectively.

A central goal in evolutionary biology is to quantify the nature, strength, and, ultimately, shifts in the forces of natural selection relative to genetic drift and mutation. As data set size and complexity increase, so does the amount of potential information on these forces and their dynamics. As a result, there is a need for more complex and realistic models (Goldman et al. 1996; Thorne et al. 1996; Goldman et al. 1998; Halpern and Bruno

MAG: Seems like some merging of the above text moved from the intro and the following text that previously opened the intro could be done. I will take a stab at later today.

1998; Lartillot and Philippe 2004) to accomplish this goal. Although extremely popular due to their elegance and computational efficiency, the utility of ω based models in helping us reach this goal is substantially more limited than commonly recognized. Because these ω models use a single substitution matrix, they are only applicable for situations in which the substitution process and shifts in the selective environment are intrinsic to the sequence, such as with positive or negative frequency dependent selection; these models do not describe stabilizing or diversifying selection as commonly envisioned (Endler 1986; Pelmyr 2002).

Starting with Halpern and Bruno (1998), a number of researchers have developed methods for linking site-specific selection on protein sequence and phylogenetics (e.g. Koshi et al. 1999; Dimmic et al. 2000; Koshi and Goldstein 2000; Robinson et al. 2003; Lartillot and Philippe 2004; Thorne et al. 2012; Rodrigue and Lartillot 2014) Halpern and Bruno (1998) calculated a vector of 19 expected amino acid frequencies for each amino acid site, making it the most general and most parameter rich of these methods. This generality, however, comes at the cost of being purely descriptive; there is no explicit biological mechanism proposed to explain the site specific amino acid frequencies estimated. By grouping together amino sites with similar evolutionary behaviors, Lartillot and colleagues retained the descriptive nature of Halpern and Bruno (1998) work while greatly reduced the number of model parameters needed (Lartillot and Philippe 2004; Rodrigue and Lartillot 2014). SelAC follows in this tradition of using multiple substitution matrices, but includes some key advances.

First, by nesting a model of a sequence's cost-benefit function \mathbf{C}/\mathbf{B} within a broader model, SelAC allows us to formulate and test a hierarchical, mechanistic models of stabilizing selection. More precisely, our nested approach allows us to relax the assumption that physicochemical deviations from the optimal sequence \vec{a}^* are equally disruptive at all sites within a protein. We found strong support for SelAC's hypothesis that the strength of

570 stabilizing selection against physicochemical deviations from \bar{a}^* varies between sites
 571 ($\Delta\text{AICc} = 20,983$). Second, because our substitution matrices are built on a formal
 572 description of a sequence's cost-benefit function \mathbf{C}/\mathbf{B} , we are able to efficiently
 573 parameterize 20 different matrices using a relatively small number of genome-wide
 574 parameters – e.g. our physicochemical weighting and G distribution shape parameters, and
 575 one gene specific gene expression parameter ψ . While the \mathbf{C}/\mathbf{B} function on which SelAC
 576 currently rests is very simple, nevertheless, it leads to a dramatic increase in our ability to
 577 explain the sequence data we analyzed. Importantly, because SelAC uses a formal
 578 description of a sequence's \mathbf{C}/\mathbf{B} , replacing our assumptions with more sophisticated ones
 579 in the future is relatively straightforward. Conceptually, our work lies in between that of
 580 Lartillot's and Thorne's, where the latter is utilizing even more detailed models of protein
 581 structure as a means of linking amino acid substitutions and stabilizing selection. Third,
 582 our use of nested models also allows us to make biologically meaningful and testable
 583 predictions. By linking a gene's expression level to the strength of purifying selection it
 584 experiences, we are able to provide coarse estimates of gene expression. This also suggests
 585 that ω is best explained as a proxy for gene expression, rather than the nature of selection
 586 on a sequence.

587 One simplifying assumption we make is that the organism can and does compensate
 588 for any reduction in protein function by simply increasing the protein's production rate.
 589 While this production compensation assumption will clearly not hold in many situations, it
 590 does allow us to connect protein function and energetic costs in a simple and biologically
 591 plausible manner. Of course, researchers could employ and test other assumptions within
 592 our framework, namely, by utilizing more detailed, gene specific knowledge about the
 593 relationship between protein function and organism fitness. For example, suppose a protein
 594 for a glucose transporter is far less efficient than usual. One possible response and the one
 595 envisioned here is that the protein is thus produced at a higher rate to compensate. This

would leave the overall ability to transport glucose unchanged. An alternative is that the cell is just less able to transport glucose across membranes. In biology, it is likely that a mixture of such effects exists. However, the production compensation mechanism is likely to have the same costs across proteins, making it a useful first approximation, while the same expression but reduced functionality will have gene specific effects more difficult to model generally (e.g., how does the cost of having glucose transport slow by half compare to the cost of underproducing an anthocyanin for flower color or fewer taste receptor proteins?). Moreover, there is evidence that cells do compensate for lower protein function by increasing gene expression. Nevertheless, by assuming that fitness declines with extraneous energy flux, SelAC explicitly links the variation in the strength of stabilizing selection for the optimal protein sequence among genes, to the variation among genes in their target expression levels ψ .

MIKE: Cut
unless we
provide
citations.

Furthermore, by linking expression and selection, SelAC provides a natural framework for combining information from protein coding genes with very different rates of evolution; from low expression genes providing information on shallow branches to high expression genes providing information on deep branches. This is in contrast to a more traditional approach of concatenating gene sequences together, which is equivalent to assuming the same average protein synthesis rate ψ for all of the genes, or more recent approaches where different models are fitted to different genes. Our results indicate that including a gene specific ψ value vastly improves SelAC fits (Table 1). Perhaps more convincingly, we find that the target expression level ψ and realized average protein synthesis rate ϕ are reasonably well correlated with laboratory measurements and theoretical predictions of gene expression (Pearson $r = 0.34 - 0.64$; Figures 1, S1, and S2). The idea that quantitative information on gene expression is embedded within intra-genomic patterns of synonymous codon usage is well accepted; our work shows that this information can also be extracted from comparative data at the amino acid level.

Of course, given the general nature of SelAC and the complexity of biological systems, other biological forces besides selection for reducing energy flux likely contribute to intergenic variation in the magnitude of stabilizing selection. Similarly, other physicochemical properties besides composition, volume, and charge likely contribute to site specific patterns of amino acid substitution. Thus, a larger and more informative set of physicochemical weights might improve our model fit and reduce the noise in our estimates of ϕ . Even if other physicochemical properties are considered, the idea of a consistent, genome wide physicochemical weighting of these terms seems highly unlikely. Since the importance of an amino acid's physicochemical properties likely changes with its position in a folded protein, one way to incorporate such effects is to test whether the data supports multiple sets of physicochemical weights for either subsets of genes or regions within genes, rather than a single set.

Both of these points highlight the advantage of the detailed, mechanistic modeling approach underlying SelAC. Because there is a clear link between protein expression, synthesis cost, and functionality, SelAC can be extended by increasing the realism of the mapping between these terms and the coding sequences being analyzed. For example, SelAC currently assumes the optimal amino acid for any site is fixed along all branches. This assumption can be relaxed by allowing the optimal amino acid to change during the course of evolution along a branch. From a computational standpoint, the additive nature of selection between sites is desirable because it allows us to analyze sites within a gene largely independently of each other. From a biological standpoint, this additivity between sites ignores any non-linear interactions between sites, such as epistasis, or between alleles, such as dominance. Thus, our work can be considered a first step to modeling these more complex scenarios.

For example, our current implementation ignores any selection on synonymous codon usage bias (CUB) (c.f. Yang and Nielsen 2008; Pouyet et al. 2016). Including such

selection is tricky because introducing the site-specific cost effects of CUB, which is consistent with the hypothesis that codon usage affects the efficiency of protein assembly or **C**, into a model where amino acids affect protein function or **B**, results in a cost-benefit ratio **C/B** with epistatic interactions between all sites. These epistatic effects can likely be ignored under certain conditions or reasonably approximated based on an expectation of codon specific costs (e.g. Kubatko et al. 2016). Nevertheless, it is difficult to see how one could identify such conditions without modeling the way in which codon and amino acid usage affects **C/B**.

This work also points out the potential importance of further investigation into model choice in phylogenetics. For likelihood models, use of AICc has become standard. However, how one determines the appropriate number of parameters estimated in a model is more complicated than generally recognized. Common sense suggests that dataset size is increased by adding taxa and/or sites. In other words, a dataset of 1000 taxa and 100 sites must have more information on substitution models than a dataset of 4 taxa and 100 sites. Our simple analyses agree that the number of observations in a dataset (number of sites \times number of taxa) should be taken as the sample size for AICc, but this conclusion likely only applies when there is sufficient independence between taxa. For instance, one could imagine a phylogeny where one taxon is sister to a polytomy of 99 taxa that have zero length terminal branches. Absent measurement error or other intraspecific variation, one would have 100 species but only two unique trait values, and the only information about the process of evolution comes from what happens on the path connecting the lone taxon to the polytomy. Although this is a rather extreme example, it seems prudent for researchers to use a simulation based approach similar to the one we take here to determine the appropriate means for calculating the effective number of data points in their data.

There are still significant shortcomings in the approach outlined here. Most worrisome are biological oversimplifications in SelAC. For example, at its heart, SelAC

assumes that suboptimal proteins can be compensated for, at a cost, simply by producing more of them. However, this is likely only true for proteins reasonably close to the optimal sequence. Different enough proteins will fail to function entirely: the active site will not sufficiently match its substrates, a protein will not properly pass through a membrane, and so forth. Yet, in our model, even random sequences still permit survival, just requiring more protein production. Other oversimplifications include the assumption of no selection on codon usage, no change of optimal amino acids through time, and no change of the effect of physiochemical properties on fitness through time. However, because we take a mechanistic approach, all of these assumptions can be relaxed through further extension of our model.

There are also deficiencies in our implementation. Though reasonable to use for a given topology with a modest number of species, it is currently too slow for practical use for tree search. Our work serves as a proof of concept, or of utility for targeted questions where a more realistic model may be of use (placement of particular taxa, for example). Future work will encode SelAC models into a variety of mature, popular tree-search programs. SelAC also represents a challenging optimization problem: the nested models reduce parameter complexity vastly, but there are still numerous parameters to optimize, including the discrete parameter of the optimal amino acid at each site. A different implementation, more parameter-rich, would optimize values of three (or more) physiochemical properties per site. This would have the practical advantage of continuous parameter optimization rather than discrete, and biologically would be more realistic (as it is the properties that selection "sees", not the identity of the amino acid itself).

In spite of these difficulties, SelAC represents an important step in uniting phylogenetic and population genetic models. While Koshi et al. (1999); Dimmic et al. (2000); Koshi and Goldstein (2000); Robinson et al. (2003); Lartillot and Philippe (2004); Thorne et al. (2012); Rodrigue and Lartillot (2014) are all models of constant, stabilizing selection, SelAC can be generalized further to include diversifying selection. Specifically, by

700 letting SelAC's Grantham weighting term G , which we now assume is ≥ 0 , to take on
701 negative values, SelAC will behave as if there is a pessimal, rather than optimal, amino
702 acid for the given site. In this diversifying selection scenario, amino acids with
703 physicochemical qualities more dissimilar to the pessimal amino acid are increasingly
704 favored, potentially resulting in multiple fitness peaks.

705 This ability to extend our model and, in turn, sharpen our thinking about the
706 nature of natural selection on amino acid sequences illustrates the value of moving from
707 descriptive to more mechanistic models in general and phylogenetics in particular. How
708 frequently diversifying selection of this nature occurs is an open, but addressable, question.
709 Regardless of the frequency at which diversifying selection occurs, it leads to the question,
710 "How often does the optimal/pessimal amino sequence change along any given branch?"
711 Due to its mechanistic nature, SelAC can also be extended to include changes in the
712 optimal/pessimal sequence over a phylogeny using a hidden markov modelling approach.
713 Extending SelAC in these ways, will allow researchers to explicitly model shifts in selection
714 on protein sequences and, in turn, quantify their frequency and magnitude.

715 In summary, SelAC allows biologically relevant population genetic parameters to be
716 estimated from phylogenetic information, while also dramatically improving fit and
717 accuracy of phylogenetic models. By explicitly modeling the optimal/pessimal sequence of
718 a gene, SelAC can be extended to include shifts in the optimal/pessimal sequence over
719 evolutionary time. Extending this model in this way will allow researchers to describe not
720 only the dynamic shifts in natural selection, but evaluate how well a given dataset supports
721 such a model. Moreover, it demonstrates that there remains substantially more information
722 in the coding sequences used for phylogenetic analysis than other methods can access.
723 Given the enormous amount of efforts expended to generate sequence datasets, it makes
724 sense for researchers to continue developing more realistic models of sequence evolution in
725 order to extract the biological information embedded in these datasets. The cost-benefit

726 model we develop here is just one of many possible paths of mechanistic model
727 development.

728

ACKNOWLEDGEMENTS

729 This work was supported in part by NSF Awards MCB-1120370 (MAG and RZ) and
730 DEB-1355033 (BCO, MAG, and RZ) with additional support from The University of
731 Tennessee Knoxville and University of Arkansas (JMB). JJC and JMB received support as
732 Postdoctoral Fellows and CL received support as a Graduate Student Fellow at the
733 National Institute for Mathematical and Biological Synthesis, an Institute sponsored by the
734 National Science Foundation through NSF Award DBI-1300426, with additional support
735 from UTK. The authors would like to thank Premal Shah, Todd Oakley, and two
736 anonymous reviewers for their helpful criticisms and suggestions for this work.

REFERENCES

- 739 Anisimova, M. 2012. Parametric models of codon evolution. Pages 12–33 *in* Codon
 740 Evolution: Mechanisms and Models (G. M. Cannarozzi and A. Schneider, eds.). Oxford
 741 University Press, Oxford, UK.
- 742 Beaulieu, J. M., B. C. O’Meara, and M. J. Donoghue. 2013. Identifying Hidden Rate
 743 Changes in the Evolution of a Binary Morphological Character: The Evolution of Plant
 744 Habit in Campanulid Angiosperms. *Systematic Biology* 62:725–737.
- 745 Beaulieu, J. M. and B. C. O’Meara. 2016. Detecting Hidden Diversification Shifts in Models
 746 of Trait-Dependent Speciation and Extinction. *Systematic Biology* 65:583–601.
- 747 Berg, J. and M. Lässig. 2003. Stochastic Evolution and Transcription Factor Binding Sites.
 748 *Biophysics* 48:S36–S44.
- 749 Butler, M. A. and A. A. King. 2004. Phylogenetic comparative analysis: a modeling
 750 approach for adaptive evolution. *American Naturalist* 164:683–695.
- 751 Dimmic, M. W., D. P. Mindell, and R. A. Goldstein. 2000. Modeling evolution at the
 752 protein level using an adjustable amino acid fitness model. *Pacific Symposium on*
 753 *Biocomputing* 5:18–29.
- 754 Drummond, D. A., J. D. Bloom, C. Adami, C. O. Wilke, and F. H. Arnold. 2005. Why
 755 highly expressed proteins evolve slowly. *Proceedings of the National Academy of Sciences*
 756 *of the United States of America* 102:14338–14343.
- 757 Drummond, D. A., A. Raval, and C. O. Wilke. 2006. A single determinant dominates the
 758 rate of yeast protein evolution. *Molecular Biology and Evolution* 23:327–337.

759 Edwards, A. 1984. Likelihood. Cambridge science classics Cambridge University Press.

760 Endler, J. A. 1986. Natural Selection in the Wild Pages 16–17. No. 21 in Monographs in
761 Population Biology Princeton University Press, Princeton, NJ reference for definition of
762 diversifying selection.

763 Felsenstein, J. 1981. Evolutionary trees from DNA-sequences - a maximum-likelihood
764 approach. Journal of Molecular Evolution 17:368–376.

765 Felsenstein, J. 2001. Taking Variation of Evolutionary Rates Between Sites into Account in
766 Inferring Phylogenies. Journal of Molecular Evolution 53:447–455.

767 Fisher, S., Ronald A. 1930. The Genetical Theory of Natural Selection. Oxford University
768 Press, Oxford.

769 Gilchrist, M., P. Shah, and R. Zaretzki. 2009. Measuring and detecting molecular
770 adaptation in codon usage against nonsense errors during protein translation. Genetics
771 183:1493–1505.

772 Gilchrist, M. A. 2007. Combining Models of Protein Translation and Population Genetics
773 to Predict Protein Production Rates from Codon Usage Patterns. Molecular Biology and
774 Evolution 24:2362–2373.

775 Gilchrist, M. A., W.-C. Chen, P. Shah, C. L. Landerer, and R. Zaretzki. 2015. Estimating
776 Gene Expression and Codon-Specific Translational Efficiencies, Mutation Biases, and
777 Selection Coefficients from Genomic Data Alone. Genome Biology and Evolution
778 7:1559–1579.

779 Gilchrist, M. A. and A. Wagner. 2006. A model of protein translation including codon bias,
780 nonsense errors, and ribosome recycling. Journal of Theoretical Biology 239:417–434.

781 Goldman, N. 1993. Statistical tests of models of DNA substitution. *Journal of molecular*
782 *evolution* 36:182–198.

783 Goldman, N., J. L. Thorne, and D. T. Jones. 1996. Using Evolutionary Trees in Protein
784 Secondary Structure Prediction and Other Comparative Sequence Analyses. *Journal of*
785 *Molecular Biology* 263:196 – 208.

786 Goldman, N., J. L. Thorne, and D. T. Jones. 1998. Assessing the Impact of Secondary
787 Structure and Solvent Accessibility on Protein Evolution. *Genetics* 149:445–458.

788 Goldman, N. and Z. H. Yang. 1994. Codon-based model of nucleotide substitution for
789 protein-coding DNA-sequences. *Molecular Biology and Evolution* 11:725–736.

790 Grantham, R. 1974. Amino acid difference formula to help explain protein evolution.
791 *Science* 185:862–864.

792 Halpern, A. L. and W. J. Bruno. 1998. Evolutionary distances for protein-coding sequences:
793 Modeling site-specific residue frequencies. *Molecular Biology And Evolution* 15:910–917.

794 Hughes, A. L. and M. Nei. 1988. Pattern of nucleotide substitution at major
795 histocompatibility complex class-i loci reveals overdominant selection. *Nature*
796 335:167–170.

797 Hhna, S., M. J. Landis, T. A. Heath, B. Boussau, N. Lartillot, B. R. Moore, J. P.
798 Huelsenbeck, and F. Ronquist. 2016. RevBayes: Bayesian Phylogenetic Inference Using
799 Graphical Models and an Interactive Model-Specification Language. *Systematic Biology*
800 65:726.

801 Ingram, T. and D. L. Mahler. 2013. SURFACE: detecting convergent evolution from data
802 by fitting Ornstein-Uhlenbeck models with stepwise Akaike Information Criterion.
803 *Methods in ecology and evolution* 4:416–425.

804 Iwasa, Y. 1988. Free fitness that always increases in evolution. *Journal of Theoretical*
805 *Biology* 135:265–281.

806 Jhvueng, D.-C., H. Snehalata, B. C. O’Meara, and L. Liu. 2014. Investigating the
807 performance of AIC in selecting phylogenetic models. *Statistical applications in genetics*
808 *and molecubr biology* 13:459–475.

809 Johnson, S. G. 2012. The NLOpt nonlinear-optimization package. Version 2.4.2 – Released
810 20 May 2014.

811 Kimura, M. 1962. on the probability of fixation of mutant genes in a population. *Genetics*
812 47:713–719.

813 Koshi, J. M. and R. A. Goldstein. 1997. Mutation matrices and physical-chemical
814 properties: Correlations and implications. *Proteins-Structure Function And Genetics*
815 27:336–344.

816 Koshi, J. M. and R. A. Goldstein. 2000. Analyzing site heterogeneity during protein
817 evolution. Pages 191–202 *in* *Biocomputing 2001*. World Scientific.

818 Koshi, J. M., D. P. Mindell, and R. A. Goldstein. 1999. Using physical-chemistry-based
819 substitution models in phylogenetic analyses of HIV-1 subtypes. *Molecular biology and*
820 *evolution* 16:173–179.

821 Kubatko, L., P. Shah, R. Herbei, and M. A. Gilchrist. 2016. A codon model of nucleotide
822 substitution with selection on synonymous codon usage. *Molecular Phylogenetics and*
823 *Evolution* 94:290 – 297.

824 Lartillot, N. and H. Philippe. 2004. A Bayesian mixture model for across-site
825 heterogeneities in the amino-acid replacement process. *Molecular Biology And Evolution*
826 21:1095–1109.

827 Lynch, M. and G. K. Marinov. 2015. The bioenergetic costs of a gene. *Proceedings Of The*
828 *National Academy Of Sciences Of The United States Of America* 112:15690–15695.

829 Mayrose, I., N. Friedman, and T. Pupko. 2005. A Gamma mixture model better accounts
830 for among site rate heterogeneity. *Bioinformatics* 21:ii151–ii158.

831 McCandlish, D. M. and A. Stoltzfus. 2014. Modeling evolution using the probability of
832 fixation: History and implications. *The Quarterly Review of Biology* 89:225–252.

833 Muse, S. V. and B. S. Gaut. 1994. A likelihood approach for comparing synonymous and
834 nonsynonymous nucleotide substitution rates, with application to the chloroplast
835 genome. *Molecular Biology and Evolution* 11:715–724.

836 Nielsen, R. and Z. H. Yang. 1998. Likelihood models for detecting positively selected amino
837 acid sites and applications to the HIV-1 envelope gene. *Genetics* 148:929–936.

838 Nowak, M. A. 2006. *Evolutionary Dynamics: Exploring the Equations of Life*. Belknap of
839 Harvard University Press, Cambridge, MA.

840 O’Meara, B. C., C. Ane, M. J. Sanderson, and W. P.C. 2006. Testing for different rates of
841 continuous trait evolution using likelihood. *Evolution* 60:922–933.

842 Pellmyr, O. 2002. Microevolution. Pages 731–732 *in* *Encyclopedia of Evolution* (M. Pagel,
843 ed.). Oxford University Press, Oxford, UK.

844 Pellmyr, O. 2002. Microevolution. Pages 731–732 *in* *Encyclopedia of Evolution* (M. Pagel,
845 ed.) vol. 2. Oxford University Press, Oxford, UK.

846 Posada, D. and T. R. Buckley. 2004. Model Selection and Model Averaging in
847 Phylogenetics: Advantages of Akaike Information Criterion and Bayesian Approaches
848 over Likelihood Ratio Tests. *Systematic Biology* 53:793–808.

- Pouyet, F., M. Bailly-Bechet, D. Mouchiroud, and L. Guguen. 2016. SENCA: A Multilayered Codon Model to Study the Origins and Dynamics of Codon Usage. *Genome Biology and Evolution* 8:2427–2441.
- Robinson, D. M., D. T. Jones, H. Kishino, N. Goldman, and J. L. Thorne. 2003. Protein evolution with dependence among codons due to tertiary structure. *Molecular Biology And Evolution* 20:1692–1704.
- Rodrigue, N. and N. Lartillot. 2014. Site-heterogeneous mutation-selection models within the PhyloBayes-MPI package. *Bioinformatics* 30:1020–1021.
- Rodrigue, N., N. Lartillot, D. Bryant, and H. Philippe. 2005. Site interdependence attributed to tertiary structure in amino acid sequence evolution. *Gene* 347:207–217.
- Rokas, A., B. L. Williams, N. King, and S. B. Carroll. 2003. Genome-scale approaches to resolving incongruence in molecular phylogenies. *Nature* 425:798–804.
- Rowan, T. 1990. Functional Stability Analysis of Numerical Algorithms. Ph.D. thesis University of Texas, Austin.
- Salichos, L. and A. Rokas. 2013. Inferring ancient divergences requires genes with strong phylogenetic signals. *Nature* 497:327–331.
- Sella, G. and A. E. Hirsh. 2005. The application of statistical physics to evolutionary biology. *Proceedings of the National Academy of Sciences of the United States of America* 102:9541–9546.
- Shah, P. and M. A. Gilchrist. 2011. Explaining complex codon usage patterns with selection for translational efficiency, mutation bias, and genetic drift. *Proceedings of the National Academy of Sciences of the United States of America* 108:10231–10236.

871 Stamatakis, A. 2006. RAxML-VI-HPC: maximum likelihood-based phylogenetic analyses
872 with thousands of taxa and mixed models. *Bioinformatics* 22:2688–2690.

873 Sullivan, J. and P. Joyce. 2005. Model Selection in Phylogenetics. *Annual Review of*
874 *Ecology, Evolution, and Systematics* 36:445–466.

875 Thorne, J. L., N. Goldman, and D. T. Jones. 1996. Combining protein evolution and
876 secondary structure. *Molecular Biology and Evolution* 13:666–673.

877 Thorne, J. L., N. Lartillot, N. Rodrigue, and S. C. Choi. 2012. Codon models as a vehicle
878 for reconciling population genetics with inter-specific sequence data. *Codon Evolution:*
879 *Mechanisms And Models* Pages 97–110 D2 10.1093/acprof:osobl/9780199601165.001.0001
880 ER.

881 Wagner, A. 2005. Energy constraints on the evolution of gene expression. *Molecular*
882 *Biology and Evolution* 22:1365–1374.

883 Wright, S. 1969. *Evolution and the genetics of populations. Vol. 2. The theory of gene*
884 *frequencies. vol. 2.* University of Chicago Press.

885 Yang, Z. 2014. *Molecular Evolution: A Statistical Approach.* Oxford University Press, New
886 York.

887 Yang, Z. H. 1994. Maximum-likelihood phylogenetic estimation from DNA-sequences with
888 variable rates over sites - approximate methods. *Journal Of Molecular Evolution*
889 39:306–314.

890 Yang, Z. H. 2007. PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular*
891 *Biology And Evolution* 24:1586–1591.

892 Yang, Z. H. and R. Nielsen. 1998. Synonymous and nonsynonymous rate variation in
893 nuclear genes of mammals. *Journal Of Molecular Evolution* 46:409–418.

894 Yang, Z. H. and R. Nielsen. 2008. Mutation-selection models of codon substitution and
895 their use to estimate selective strengths on codon usage. *Molecular Biology and*
896 *Evolution* 25:568–579.

TABLE

Model	logLik	Parameters Estimated	AIC	AICc	Δ AICc	Model Weight
GTR+ Γ	-655,166.4	610	1,311,553	1,311,554	284,240	<0.001
GY94	-612,670.4	111	1,225,563	1,225,563	198,249	<0.001
FMutSel	-597,140.7	178	1,194,637	1,194,638	167,324	<0.001
SelAC _{<i>M</i>}	-478,302.4	50,004	1,056,613	1,076,674	49,360	<0.001
SelAC	-464,114.8	50,004	1,028,238	1,048,299	20,985	<0.001
SelAC _{<i>M</i>} + Γ	-465,106.9	50,005	1,030,189	1,050,286	22,972	<0.001
SelAC+ Γ	-453,620.8	50,005	1,007,252	1,027,314	0	>0.999

Table 1: Comparison of model fits using AIC, AICc, and AIC_w. Note the subscripts *M* indicate model fits where the most common or ‘majority rule’ amino acid was fixed as the optimal amino acid a^* for each site. As discussed in text, despite the fact that a^* for each site was not fitted by our algorithm, its value was determined by examining the data and, as a result, represent an additional parameter estimated from the data and are accounted for in our table. Also, the sample size used in the calculation of AICc is assumed to be equal to the size of the matrix (number of taxa x number of sites).

FIGURES




Figure 1: Comparisons between estimates of average protein translation rate $\hat{\phi}_{\text{SelAC}}$ obtained from SelAC+ Γ and direct measurements of expression for individual yeast taxa across the 100 selected genes from Salichos and Rokas (2013). Estimates of $\hat{\phi}_{\text{SelAC}}$ were generated by dividing the composite term ψ' by $\mathbf{B}(\vec{a}_i|\vec{a}^*)$. Gene expression was measured using either RNA-Seq (a)-(c) or microarray (d). The equations in the upper left hand corner of each panel represent the regression fit and the Pearson correlation coefficient r .




Figure 2: Comparisons between ω_{FMutSel} , which is the nonsynonymous/synonymous mutation ratio in FMutSel, SelAC+ Γ estimates of protein functionality production rates $\hat{\psi}_{\text{SelAC}}$ (a), RNA-Seq based measurements of mRNA abundance $\phi_{\text{RNA-seq}}$ (b), and ROC-SEMPER's estimates of protein translation rates ϕ_{ROC} , which are based solely on *S. cerevisiae*'s patterns of codon usage bias (c), for *S. cerevisiae* across the 100 selected genes from Salichos and Rokas (2013). As in Figure 1, the equations in the upper right hand corner of each panel provide the regression fit and correlation coefficient.




Figure 3: (a) Maximum likelihood estimates of branch lengths under SelAC+ Γ for 100 selected genes from Salichos and Rokas (2013). Tests of model adequacy for *S. cerevisiae* (b) and *S. castellii* (c) indicated that, when these taxa are removed from the tree, and their sequences are simulated, the parameters of SelAC+ Γ exhibit functionality $\mathbf{B}(\vec{a}_{\text{obs}}|\vec{a}^*)$ that is far closer to the observed (dashed black line) than data sets produced from parameters of either FMutSel or GTR + Γ .

SUPPORTING MATERIALS

Supporting Materials for *Population Genetics Based Phylogenetics Under Stabilizing Selection for an Optimal Amino Acid Sequence: A Nested Modeling Approach* by Beaulieu *et al.* (In Review).

Comparisons of SelAC gene expression estimates with empirical measurements

In our model, the parameter ϕ measures the realized average protein synthesis rate of a gene. We compared our estimates of ϕ to two separate measures of gene expression, one empirical (Figure S1), and one model-based prediction that does not account for shared ancestry, for individual yeast taxa across the same set of genes. Our estimates of ϕ are positively correlated with both measures, which are also reasonably well correlated with each other (Figure 1 - S2) On the whole, these comparisons indicate not only a high degree of consistency among all three measures, but also, importantly, that estimates of ϕ obtained from SelAC provide real biological insight into the expression level of a gene.




Figure S1: Comparisons between estimates of ϕ obtained from SelAC+ Γ and the predicted gene expression from the ROC SEMPER model (Gilchrist et al. (2015)) for individual yeast taxa across the 100 selected genes from Salichos and Rokas (2013). As with figures in the main text, estimates of ϕ were obtained by solving for ψ based on estimates of ψ' , and then dividing by $\mathbf{B}(\vec{a}_i|\vec{a}^*)$. The equations in the upper left hand corner of each panel represent the regression fit and correlation coefficient.




Figure S2: Comparisons of predicted gene expression from the ROC SEMPER model (Gilchrist et al. (2015)) and direct measurements of expression from RNA-Seq or microarray data for individual yeast taxa across the 100 selected genes from Salichos and Rokas (2013). The equations in the upper left hand corner of each panel represent the regression fit and correlation coefficient.

Simulations

Overall, the simulation results indicate that the SelAC model can reasonably recover the known values of the generating model (Figure S3 - S6). This includes not only the

parameters in SelAC, but also the optimal amino acids for a given sequence as well as the estimates of the branch lengths. There are a few observations to note. First, the ability to accurately recover the true optimal amino acid sequence will largely depend on the magnitude of the realized average protein synthesis rate of the gene ϕ . This is, of course, intuitive, given that ϕ sets the strength of stabilizing selection towards an optimal amino acid at a site. However, the inclusion of α_G into SelAC, appears to generally increase values of ϕ and generally improves the ability to recover the optimal amino acids even for the gene with the lowest baseline ϕ . Second, we found a strong downward bias in estimates of α_G , which actually translates to greater variation among the rate categories. The choice of a gamma distribution to represent site-specific variation in sensitivity was based on mathematical convenience and convention, rather than on biological reality. Nevertheless, we suspect that this bias is in large part due to the difficulty in determining the baseline ψ for a given gene and the value of α_G that globally satisfies the site-specific variation in sensitivity across all genes, as indicated by the slight upward bias in estimates of ψ . A reviewer pointed out that it may also be difficult for SelAC to account for changing amino-acid, which we agree may also play a role. It has been suggested, in studies of the behavior of the gamma distribution in applications of nucleotide substitution model, that increasing the number of rate categories can often improve accuracy of the shape parameter (Mayrose et al. (2005)). Future work will address this issue.




Figure S3: Summary of a 5-gene simulation for a SelAC model where we assume $\alpha_G = \infty$, and thus, no site-specific sensitivity in the generating model. The ‘known’ parameters were based on fitting the same SelAC to the 106 gene data set and phylogeny of Rokas et al. (2003), with gene choice being based on five evenly spaced points along the rank order of the gene specific composite parameter ψ'_g . The points and associated uncertainty in the estimates of the gene-specific average protein synthesis rate, or ψ (calculated from ψ')(a), nucleotide mutation rates under the UNREST model (b), proportion of correct optimal amino acids for a given gene (c), and estimates of the individual edge lengths are based the mean and 2.5% and 97.5% quantiles across all 50 simulated datasets (d). Gene index on the x-axis refers to the arbitrary number assigned to the simulated gene.




Figure S4: The distribution of estimates of the Grantham weights, α_c and α_p , in a SelAC model, where we assume $\alpha_G = \infty$, and thus no site-specific sensitivity in the generating model. The dashed line represents the value used in the generating model.




Figure S5: Same figure as in Figure S3, except the generating model includes site-specific sensitivity in the generating model (i.e., α_G).




Figure S6: Same figure as in Figure S4, except the generating model includes site-specific sensitivity in the generating model (i.e., α_G). Unlike, Grantham weights, which showed no systematic bias, there is a downward bias in estimates of α_G .