

Rutgers University: Complex Variables and Advanced Calculus Written Qualifying Exam

January 2015: Problem 2 Solution

Exercise. Use contour integration to evaluate the integral

$$\int_0^{2\pi} \frac{d\theta}{a + \sin \theta}, \quad \text{for } a > 1.$$

Solution.

$$\begin{aligned} z &= e^{i\theta} & dz &= ie^{i\theta} d\theta \implies d\theta = \frac{dz}{ie^{i\theta}} = \frac{dz}{iz} & C : |z| &= 1 & e^{i\theta} & & 0 \leq \theta \leq 2\pi \\ \sin \theta &= \frac{z - z^{-1}}{2i} \end{aligned}$$

$$\begin{aligned} \int_0^{2\pi} \frac{d\theta}{a + \sin \theta} &= \int_C \frac{1}{a + \frac{z - z^{-1}}{2i}} \left(\frac{1}{iz} \right) dz \\ &= \int_C \frac{1}{iaz + \frac{z^2 - 1}{2}} dz \\ &= 2 \int_C \frac{1}{z^2 + 2aiz - 1} dz \\ &= 2 \int_C \frac{-2ai \pm \sqrt{-4a^2 - 4(-1)}}{2} dz \\ &= -ai \pm i\sqrt{a^2 - 1} \end{aligned}$$

$$\implies f(z) = \frac{1}{z^2 + 2aiz - 1} \text{ has simple poles at } z = (-a + \sqrt{a^2 - 1})i \text{ and } z = (-a - \sqrt{a^2 - 1})i$$

But $|(-a - \sqrt{a^2 - 1})i| = a + \sqrt{a^2 - 1} > 1$ since $a > 1$.

So only $(-a + \sqrt{a^2 - 1})i$ lies inside contour $|z| = 1$ and $|(-a + \sqrt{a^2 - 1})i| = -a + \sqrt{a^2 - 1}$

$$\begin{aligned} \int_C f(z) dz &= 2\pi i \operatorname{Res}(f, (-a + \sqrt{a^2 - 1})i) \\ &= 2\pi i \cdot \frac{1}{(-a + \sqrt{a^2 - 1} + a + \sqrt{a^2 - 1})i} \\ &= \frac{2\pi}{2\sqrt{a^2 - 1}} \\ &= \frac{\pi}{\sqrt{a^2 - 1}} \\ \int_0^{2\pi} \frac{d\theta}{a + \sin \theta} &= 2 \int_C f(z) dz \\ &= 2 \left(\frac{\pi}{\sqrt{a^2 - 1}} \right) \\ &= \boxed{\frac{2\pi}{\sqrt{a^2 - 1}}} \end{aligned}$$