

Rutgers University: Algebra Written Qualifying Exam

August 2018: Problem 2 Solution

Exercise. Let p and q be distinct primes. Let $\bar{q} \in \mathbb{Z}/p\mathbb{Z}$ denote the class q modulo p and let k denote the order of \bar{q} as an element of $(\mathbb{Z}/p\mathbb{Z})^*$. Prove that no group of order pq^ℓ with $1 \leq \ell \leq k$ is simple.

Solution.

Let $|G| = pq^\ell$ such that $1 \leq \ell \leq k$.

Since k is the order of \bar{q} in \mathbb{Z}_p^* ,

$$\begin{aligned} q^k &\equiv 1 \pmod{p}, \text{ and} \\ q^j &\not\equiv 1 \pmod{p} \text{ for } 1 \leq j < k \end{aligned}$$

$$\begin{aligned} n_p &\equiv 1 \pmod{p} \text{ and } n_p \mid q^\ell \\ \implies n_p &= 1 \text{ or } n_p = q^k \text{ and } \ell = k \end{aligned}$$

If $n_p = 1$, there is a unique Sylow p -subgroup
 \implies the Sylow p -subgroup is normal in G
 $\implies G$ is not simple.

If $n_p = q^k$, then $\ell = k$ and $|G| = pq^k$

Since the Sylow p -subgroups have prime order, they are cyclic

If P and Q are two Sylow p -subgroups

$$P \cap Q = \{e\} \text{ or } P.$$

This is because $P \cap Q$ is a subgroup of both P and Q and its order must divide p .

If $n_p = q^k$ then there are q^k Sylow p -subgroups that pairwise only intersect with the identity element.

So, the q^k Sylow p -subgroups contain a total of $(q^k - 1)q^k = q^{k+1} - q^k$ elements of order p .

The remaining q^k elements must be elements of the Sylow q -subgroup:

$$\begin{aligned} \implies n_q &= 1 \\ \implies \text{The Sylow } q\text{-subgroup is normal} \\ \implies G &\text{ is not simple} \end{aligned}$$