

OUTLIER REMOVAL

- 1. PERCENTILE
- 2. IQR
- 3. STD DEV

EG : HEIGHT
describe(): max and min
min = 3 feet
max = 7.5 feet

HEIGHT
0.025 feet (lower exceptions)
100 feet (higher exceptions)

PERCENTILE
(DOMAIN KNOWLEDGE)

25% = 5
50% (MEDIAN)
75%

Inter Quartile
Range

2% = 2 feet
2.5% = 2.10 feet
3% = 2.7 feet
4% = 2.98 feet
4.8% = 3 feet

lower_limit = 4.8% = 3 feet

95% = 6 feet
97% = 7 feet
97.8% = 7.55 feet

upper_limit = 97.8% = 7.55 feet

ll = 3 (df.height.quantile(0.048))
ul = 7.55(df.height.quantile(0.978))

df[(df.height >= ll)&(df.height <= ul)]

df.height.quantile(0.048)

IQR
Q1 = 25%
Q2 = 50
Q3 = 75%

IQR = Q3 - Q1
Q1 (Q3)=
df.height.quantile(0.25)0.75

IQR = Q3- Q1

LL = Q1 - 1.5 * IQR
UL = Q3 + 1.5 * IQR

df[(df.height >= ll)&(df.height <= ul)]

STD DEV

MEAN = 66
STD = 2

DP 1= 68 = MEAN + (X) STD
68 = 66 + (X) 2
X = 1
68 = 68

MEAN = df.height.mean()
STD = df.height.std()

LL = MEAN -(3 * STD)
UL = MEAN + (3 * STD)

df[(df.height >= ll)&(df.height <= ul)]

