Therapeutic strategies to deplete macrophages in atherosclerotic plaques

Inge De Meyer, Wim Martinet & Guido R. Y. De Meyer

Division of Physiopharmacology, University of Antwerp, Antwerp, Belgium

Correspondence

Professor Dr Guido R.Y. De Meyer, Division of Physiopharmacology, University of Antwerp, Universiteitsplein 1, B-2610 Antwerp, Belgium.

Tel.: +32 3265 2737 Fax: +32 3265 2567

E-mail: guido.demeyer@ua.ac.be

Keywords

adhesion molecules, atherosclerosis, cell death, macrophages, reverse cholesterol transport

Received

5 October 2011

Accepted

31 January 2012

Accepted Article Published Online

6 February 2012

Macrophages can be found in all stages of atherosclerosis and are major contributors of atherosclerotic plaque development, progression and destabilization. Continuous recruitment of monocytes drives this chronic inflammatory disease, which can be intervened by several strategies: reducing the inflammatory stimulus by lowering circulating lipids and promoting cholesterol efflux from plaque, direct and indirect targeting of adhesion molecules and chemokines involved in monocyte adhesion and transmigration and inducing macrophage death in atherosclerotic plaques in combination with anti-inflammatory drugs. This review discusses the outlined strategies to deplete macrophages from atherosclerotic plaques to promote plaque stabilization.

Macrophages in atherosclerosis

Atherosclerosis is a chronic inflammatory disease of the arterial wall and is the foremost cause of death in developed countries. This progressive disease is characterized by a pathological lipid deposition within the vessel wall of medium-sized and large arteries. Over the years, studies have implicated an important role of macrophages in plaque development, progression and destabilization [1–3]. Initially, monocytes selectively bind on adhesion molecules, such as vascular adhesion molecule 1 (VCAM-1), intercellular adhesion molecule 1 (ICAM-1), and selectins on endothelial cells. These adhesion molecules are induced by mediators associated with risk factors (hypertension, diabetes, hyperlipidaemia, smoking, obesity and inflammation) [4]. Once bound, transmigration of inflammatory cells into the vessel wall is mediated by the endothelial release of chemokines, such as monocyte attractant protein 1 (MCP-1), which binds to its receptor on monocytes [5-7]. Infiltrated monocytes mature into macrophages by macrophage colony-stimulating factor and express scavenger receptors that allow them to engulf modified lipoproteins [7, 8]. Accumulation of lipid-laden macrophages and to a lesser extent T-lymphocytes beneath the endothelium constitutes a fatty streak [2, 9].

The evolution of a fatty streak to an atherosclerotic plaque involves activation of macrophages by T lymphocytes through interferon-y (IFN-y) and CD40 [2, 7]. Once macrophages are activated, they produce cytokines, chemokines and growth factors, promoting smooth muscle cell (SMC) migration and proliferation in the intimal layer, even though they also induce focal necrosis [10]. The collagen-synthesizing SMCs form a fibrous cap that surrounds the lipid and necrotic core. The fibrous cap prevents exposition of the thrombogenic lipid core to arterial blood and provides mechanical strength and stability to the plaque [11]. Accumulation of activated inflammatory cells, such as macrophages and T lymphocytes, in the shoulder region of the plaque amplifies the inflammatory process by the release of pro-inflammatory cytokines. Macrophages also release matrix metalloproteinases (MMPs) [12, 13] that degrade the extracellular matrix, and excessive amounts of nitric oxide (NO) [14] that induce SMC apoptosis. All these events weaken the fibrous cap and render the plaque rupture-prone [15]. Another mechanism that contributes to plaque destabilization is apoptosis, which occurs more frequently in regions with a high density of macrophages and foam cells, and is also strongly related to macrophage infiltration [16]. Vulnerable plaques typically display a large lipid core, a thin fibrous cap, a reduced number of SMCs and large numbers of macrophages and T cells [17, 18].



Clinical manifestations of atherosclerosis are determined by cellular plague composition and risk of thrombus formation. Stenotic fibrous lesions without thrombus risk restrict blood flow and provoke symptoms such as stable angina. Acute coronary syndromes, such as unstable angina and acute myocardial infarction, are caused by plague rupture or endothelial erosion with mural thrombus formation [19]. Histopathological comparison between symptomatic and asymptomatic endarterectomy specimens reveals that the number of inflammatory cells is significantly increased in symptomatic plaques vs. asymptomatic ones [20–23]. Removal of macrophages from the plaque could attenuate the inflammatory response and subsequent plaque destabilization. This review focuses on current and possible new therapeutic targets and drugs that provoke macrophage elimination in different stages of atherosclerosis, either by reducing lipids in the circulation or from the plaque, decreasing monocyte recruitment, inhibiting macrophage activation or inducing macrophage death in atherosclerotic plagues.

Systemic LDL lowering

Statins or 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase inhibitors block the conversion of acetoacetyl-CoA and acetyl-CoA to mevalonate, a key intermediate in cholesterol synthesis in liver and other tissues (Figure 1 (1)). The resulting decrease in hepatic cholesterol content leads to a compensatory increase in hepatic low density lipoprotein (LDL) receptors, which clear LDL cholesterol from the circulation [24]. Accordingly, statins reduce the incidence of acute coronary syndromes in primary and secondary prevention in patients with high cholesterol concentrations as well as those at increased risk for cardiovascular events, independent of their cholesterol concentrations, suggesting that they also exhibit beneficial effects beyond lipid lowering (pleiotropic effects) [25-30]. A further reduction of major vascular events is safely produced by more intensive statin therapy (40-80 mg statin compared with 20-40 mg statin daily) [31]. Despite the efficacy of statins, a subpopulation of patients does not benefit from current lipid lowering strategies [32]. Therefore, overall morbidity and mortality from atherosclerotic vascular disease remain substantial and development of new strategies is still essential to reduce disease burden.

Other cholesterol reducing agents, such as ezetimibe, have not yet been proven to reduce the risk of cardiovascular events [33–36]. Ezetimibe inhibits intestinal cholesterol absorption via inhibition of the Niemann-Pick C1-Like 1 (NPC1L1) sterol transporter leading to a compensatory upregulation of hepatic LDL receptors and clearance of circulating LDL (Figure 1 (2)) [37].

Experimental studies

Preclinical studies demonstrate that the lipid lowering properties of ezetimibe as monotherapy or on top of statin

therapy inhibit atherogenesis or reduce plaque burden [38–42].

Clinical studies

Clinical trials confirm the LDL cholesterol lowering effects of ezetimibe added to statin treatment, but whether ezetimibe enhances plaque regression and reduces clinical events has been controversial [34, 35, 43, 44]. The ongoing IMPROVE-IT trial attempts to detect benefits of ezetimibe as a second LDL cholesterol lowering drug added to aggressive statin therapy [45]. However, a recent study with statin-naïve patients shows no change in plaque volume between patients treated with simvastatin or simvastatin plus ezetimibe despite greater LDL cholesterol lowering in the latter group [46]. Moreover, adding ezetimibe to patients previously on statin therapy reduces LDL cholesterol but paradoxically induces plaque progression [46].

Interestingly, proprotein convertase subtilisin kexin 9 (PCSK9) regulates circulating LDL concentrations by inhibiting hepatic LDL receptor-mediated LDL uptake. In humans and mice, loss of function mutations in the gene encoding PCSK9 is associated with lifelong low LDL concentrations [47, 48]. Clinical trials with anti-PCSK9 antibody are ongoing [49].

Increased cholesterol efflux from macrophages

Recent studies focus on pathways to reverse cholesterol transport from atherosclerotic plaques to the liver for excretion into bile and faeces, whereby high density lipoprotein (HDL) and its major apolipoprotein, apolipoprotein A-I (apoA-I) play key roles in these processes. HDL is formed as discoidal pre-β₁-HDL by addition of phospholipid to lipid poor apoA-I, which is synthesized in liver and intestines. These particles take up free cholesterol from cells through adenosine triphosphate-binding cassette transporter-A1 (ABCA1). The free cholesterol is converted to cholesteryl ester (CE) by lecithin cholesterol acyltransferase (LCAT) to form spherical α -HDL particles, which can take up more free cholesterol through ABCG1 and ABCG4 that in turn is converted to CE by LCAT. The CEs are returned to the liver either directly via scavenger receptor B1 or via LDL receptors after transfer to apolipoprotein B lipoproteins in exchange for triglycerides by the cholesteryl ester transfer protein (CETP) (Figure 1). Studies have shown a strong inverse correlation between plasma HDL concentrations and atherosclerotic cardiovascular risk in humans [50, 51]. Several strategies have been explored to increase cholesterol efflux in order to stabilize atherosclerotic plagues including (i) raising HDL concentrations, (ii) inhibiting CETP, (iii) promoting expression of cholesterol transporters and (iv) other mechanisms (e.g. probucol).




Figure 1

Schematic overview of pharmacological targets involved in cholesterol uptake (solid line) and reverse cholesterol transport (bold line). Cholesterol is synthesized in the liver via HMG-CoA reductase, a pharmacological target of statins (1). Ezetimibe inhibits cholesterol uptake in the intestines by blocking sterol transporter NPC1L1 (2). Chylomicrons transport dietary triglycerides and cholesterol from the gastrointestinal tract to the peripheral tissues, whereby chylomicron remnants are taken up by the liver. Lipid-poor pre- β_1 -HDL (origin indicated with dashed line) promotes efflux of free cholesterol of macrophages via ABCA1. The expression of this cholesterol transporter is up-regulated by PPAR ligands thiozolidinediones and LXR agonists T0901317 and ATI-829 (3). In α -HDL particles, cholesterol is converted to cholesteryl esters by LCAT and further cholesterol efflux from macrophages is promoted by ABCG1 and ABCG4. Cholesterol is returned to the liver by SR-B1 or transferred to apoB lipoproteins VLDL and LDL via CETP. HDL concentrations are effectively raised by CETP inhibitors, torcetrapib and anacetrapib, and niacin (4). Finally, hepatic cholesterol is oxidized to bile acids or secreted in the bile unaltered where it is secreted in the faeces unless it is reabsorbed by the intestine. Probucol promotes cholesterol efflux by enhancing faecal excretion of HDL-derived cholesterol (5). ABCA1, adenosine triphosphate-binding cassette transporter-A1; apoA-I, apolipoprotein A-I; BA, bile acids; CE, cholesteryl ester; CETP, cholesteryl ester transfer protein; FC, free cholesterol; HDL, high density lipoprotein; HMG-CoA, 3-hydroxy-3-methylglutaryl coenzyme A; LCAT, lecithin cholesterolacyltransferase; LDL, low density lipoprotein; LXR, liver X receptors; NPC1L1, Niemann-Pick C1-Like 1; PPAR, peroxisome proliferator-activated receptor; SR-B1, scavenger receptor B1; TG, triglycerides; VLDL, very low density lipoprotein

HDL infusion/raising

Experimental studies Animal studies support the concept of raising HDL by administering recombinant HDL and apoA-I Milano or overexpressing apoA-I for inhibition of atherosclerosis and show significant reductions in lesion and macrophage area by mobilizing tissue cholesterol and by reducing plaque lipids [52–55].

Clinical studies Short term infusion of reconstituted HDL containing recombinant apoA-I Milano in humans induces a substantial regression of coronary atherosclerosis after only 5 weeks of treatment [56]. Despite the positive outcome, the small sample size and dissimilarities in plaque burden at baseline are limitations of this study. In another investigation, infusion of reconstituted HDL

containing plasma-derived apoA-I significantly improves plaque characteristics as demonstrated by intravascular ultrasound, but there is no significant difference in lesion size [57]. This study was hampered by a small sample size and presence of liver abnormalities in the high dose group. In the search for HDL-raising drugs, fibrates, such as gemfibrozil, demonstrate improved cardiovascular outcomes in human clinical trials [58, 59], whereas subsequent trials with bezafibrate and fenofibrate have disappointing results with regard to amelioration in cardiovascular mortality outcomes [60-62]. The ARBITER-6-HALTS trial shows that HDL-raising by niacin as add-on therapy in patients receiving long term statin therapy is superior to ezetimibe with respect to changes in carotid intimamedia thickness and the incidence of major adverse cardiovascular events [36]. These beneficial effects of niacin along with statin therapy on atherosclerosis development are further investigated in the AIM-HIGH study [63], but this study was prematurely discontinued in 2011 due to lack of benefits in reducing cardiovascular events of combined treatment and due to unexpectedly increased ischemic stroke rates.

CETP inhibition

CETP tends to influence negatively the reverse cholesterol transport pathway by lowering HDL and consequently scavenging of cholesterol from tissues and macrophages declines (Figure 1).

Clinical studies CETP inhibitors are capable of producing higher HDL concentrations than niacin, but clinical studies with the CETP inhibitor torcetrapib dramatically failed due to production of dysfunctional HDL and off-target toxicity [64]. Clinical trials with other CETP inhibitors, anacetrapib and dalcetrapib, are ongoing and these drugs are so far well tolerated (Figure 1 (4)) [65,66]. Nevertheless, increased HDL concentrations as such are not a sensitive indicator of removal of lesional cholesterol to the liver, but rather the distribution of functional HDL particles. Recently, it has been shown that sera of patients with similar HDL concentrations have different cholesterol efflux capacities in macrophages because of differences in concentration of pre- β_1 -HDL [67].

Up-regulation of cholesterol transporters

Besides the composition and concentration of extracellular cholesterol acceptors, the cholesterol efflux capacity is also dependent on the level of expression of cholesterol transporters ABCA1 and ABCG1. Expression of cholesterol transporters can be modulated by liver X receptors (LXR), which are ligand-activated transcription factors that regulate cholesterol transport and lipid metabolism in liver, intestines and macrophages in response to oxysterols (Figure 1 (3)). Furthermore, synthetic ligands of peroxisome proliferator-activated receptor (PPAR), fibrates (PPARα) and thiozolidinediones (PPARγ), are clinically used for treat-

ment of dyslipidaemia and type II diabetes, respectively. In addition, these drugs are thought to have anti-atherogenic functions due to cholesterol efflux.

Experimental studies Bone marrow transplantations of ABCA1-/- and ABCG1-/- mice in hypercholesterolaemic mice demonstrate a cooperative role of ABCA1 and ABCG1 in macrophage reverse cholesterol transport since deficiency of both transporters in macrophages increases foam cell formation and accelerates atherosclerosis to a greater extent than bone marrow from either single knockout alone [68]. The natural LXR agonist 24(S), 25-epoxycholesterol induces selective upregulation of ABCA1, ABCG1 and apoE in macrophages thereby affecting the macrophage cholesterol transport [69]. Synthetic LXR agonists, such as T0901317, increase ABCA1 and ABCG1 expression, inhibit plaque progression, induce regression of established lesions and alter plaque composition favouring plaque stability, i.e. reducing macrophage content and increasing collagen content [70]. Moreover, studies in macrophage-selective LXR-deficient hypercholesterolaemic mice confirm that macrophage LXR expression is necessary for the atheroprotective actions of an LXR agonist [70]. The major drawback of these drugs is the effect on the liver resulting in severe hypertriglyceridaemia and hepatic steatosis. The tissue selective LXR agonist ATI-829 has overcome these side effects and is a potent inducer of ABCA1 expression in macrophages, supporting the possibility of using LXR agonists for the treatment of atherosclerosis [71]. PPAR activation augments cholesterol efflux from macrophages by controlling the transcriptional activity of genes regulating LXR, which induces ABCA1 expression, and raises HDL concentrations by stimulating hepatic apoA-I expression (Figure 1 (3)) [72, 73]. Bone marrow transplantation of PPAR- γ^{-1} into atherosclerotic mice results in a significant increase in atherosclerosis. PPAR-y agonists in atherosclerotic mice attenuate atherosclerosis [74] and PPAR- α activation by fenofibrate in hypercholesterolaemic rabbits is associated with improvements in lipoprotein levels by decreasing triglyceride concentrations, plaque regression and changes in plaque composition towards a stable plaque phenotype [75].

Clinical studies In a small human study, patients treated with the PPAR- γ ligand pioglitazone show a significant enhanced cholesterol efflux capacity [76]. Although cholesterol efflux from macrophages represents only a small fraction of reverse transport cholesterol, it is probably the component most relevant to atheroprotection [77].

Other mechanisms

Probucol is a potent anti-oxidant drug that has been used in the past for the treatment and prevention of cardiovascular events, but was withdrawn from the European and American market due to its effects in reducing HDL concentrations and possible QT interval prolongation. Nevertheless, recent studies show that probucol promotes reverse cholesterol transport [78, 79].

Experimental studies In animal models probucol reduces atherosclerosis despite reducing HDL cholesterol concentrations [80, 81]. The anti-atherogenic effects of probucol are attributed to the enhancement of macrophage reverse cholesterol transport and faecal excretion of HDL-derived cholesterol. Probucol inhibits hepatic ABCA1, thereby reducing HDL concentrations but promoting reverse cholesterol transport without the involvement of scavenger receptor B1 (Figure 1 (5)) [78]. The mechanism of hepatic ABCA1 inhibition remains to be investigated but it is possible that probucol inhibits ABCA1 translocation to the plasma membrane, as shown in cultured macrophages [82].

Clinical studies A recent study in Japan shows a reduction in cardiovascular risk with long term probucol treatment in patients with familial hypercholesterolaemia [79]. In a very small study of 10 hypercholesterolaemic patients it was shown that probucol increases CETP activity, suggesting that reduced HDL concentrations are a direct consequence of CETP stimulation [83].

Decreased monocyte recruitment

Decreased monocyte recruitment at the level of adhesion molecules

Adhesion molecules (selectins and integrins) are locally up-regulated on the endothelium and leucocytes by cytokines due to risk factors like oxLDL, smoking, hypertension and diabetes, and are modulated by haemodynamic stress including arterial pressure and shear stress. E- and P-selectins mediate the initial monocyte tethering and rolling along endothelium, whereas vascular adhesion molecules VCAM-1 and ICAM-1 are involved in leucocyte adhesion by binding to the activated integrins on leucocytes, very late antigen-4 (VLA-4) and lymphocyte function associated antigen-1 (LFA-1), respectively.

Experimental studies Studies with atherosclerotic knockout mice have shown that altering the expression of E-selectin, P-selectin, VCAM-1 and ICAM-1 reduces lesion size as well as the number of macrophages in these lesions without affecting lipoprotein profiles and numbers of circulating monocytes (Table 1) [84–89]. Moreover, blocking antibodies towards leucocyte integrin VLA-4 or towards their endothelial counterparts VCAM-1 and ICAM-1 is effective in reducing macrophage recruitment to atherosclerotic plaques in mice [90–92]. Evidence for VCAM-1 expression on the endothelium preceding atherosclerotic plaque formation and most pronounced inhibition of mononuclear cell accumulation and reduction in lesions by VCAM-1 blockade further imply a dominant role of VCAM-1 in monocyte recruitment as well as in lesion initiation, progression and expansion.

Clinical studies Despite the promising results in animal studies, clinical anti-VCAM-1 therapy directed at lymphocyte trafficking with AGI-1067 in patients with acute coronary syndrome shows no significant cardiovascular benefits [93]. Clinical trials with VCAM-1 inhibitors, CAM741 and HUN-7293, are ongoing to assess the possible beneficial effects.

The anti-inflammatory effects of statins are in part due to inhibition of formation of mevalonate, a precursor of isoprenoids that prenylate small signal transducing guanosine triphosphatases Ras, Rho, Rab and Rae. Consequently, activation of nuclear factor-κB, expression of adhesion molecules and vascular inflammation is inhibited [94, 95]. Other anti-atherogenic effects of statins include direct binding of these drugs to the pro-inflammatory mediator LFA-1 on leucocytes, thereby blocking ICAM-1 for leucocyte adhesion [96]. Thus, statins may also inhibit leucocyte recruitment in atherosclerotic plaques by an HMG-CoA independent pathway.

Decreased monocyte recruitment at the level of junctional proteins

Transmigration of leucocytes across endothelial cells, also known as diapedesis, is determined by junctional proteins, such as junctional adhesional molecules (JAM) and connexins.

Experimental studies Enhanced expression of JAM-A is already present in the early stages of atherosclerosis in the endothelium of apoE-deficient mice on a high cholesterol diet, and the soluble form of JAM-A significantly reduces mononuclear cell recruitment to ex vivo perfused atherosclerotic carotid arteries [97]. Genetic deletion of JAM-A in apoE-deficient mice corroborates that JAM-A is implicated in leucocyte recruitment since JAM-A deficiency is associated with a decrease in macrophage content without altering the SMC content and endothelial cell recovery after wire injury of carotid arteries [98]. Moreover, JAM-A deficiency diminishes the deposited delivery of the chemokine regulated upon activation, normal T cell expressed and secreted (RANTES) on activated endothelium and subsequent monocyte recruitment and plaque formation [98]. Connexins are proteins forming gap junctions that play a role in direct intercellular communication whereby connexin 43 is upregulated in atherosclerotic lesions. Reducing connexin 43 by a genetic or pharmacological approach using statins in hypercholesterolaemic mice results in a decreased neointimal formation with concurrent reduced number of atheroma-associated inflammatory cells and increased thickness of the fibrous cap covering the lesions [99]. This suggests that gap junction communication between endothelial cells and leucocytes



 Table 1

 Molecular targets affecting monocyte recruitment and macrophage accumulation in atherosclerotic plaques

Molecular target	Mechanism	Model	Effect	Reference
P-selectin	Tethering and rolling	P-selectin ^{-/-} on C57BL/6 background on western diet or apoE ^{-/-} double knock out	Reduces macrophage number, reduces lesion size at initial stages of atherosclerosis and less development to advanced lesions	85–87
		P-selectin ^{-/-} on C57BL/6 background	Reduces neointimal formation without infiltration of inflammatory cells	84, 89
VCAM-1	Adhesion	VCAM-1- ^{-/-} on LDLR- ^{-/-} or apoE- ^{-/-} background	Reduces lesion size and less adherence of mononuclear cells at initial stages of atherosclerosis	88, 169
		Monoclonal antibody against VCAM-1	Reduces monocyte infiltration and neointimal formation	91
ICAM-1	Adhesion	ICAM-1-/- on C57BL/6 background on western diet	Slightly reduces lesion size at initial stages of atherosclerosis	85
		Monoclonal antibody against ICAM-1 in apoE-/- mice	Reduces macrophage homing by 65%	90
VLA-4	Adhesion	Monoclonal antibody against VLA-4 in apoE ^{-/-} mice	Reduces macrophage recruitment by 48% and reduces neointimal formation by 72%	92
		Monoclonal antibody against VLA-4 in apoE-/- mice	Reduces macrophage homing by 75%	90
JAM-A	Transmigration	Ex vivo perfused carotid arteries by soluble JAM-A in apoE ^{-/-} mice on western diet	Inhibits mononuclear cell recruitment upon pretreatment	97
		JAM-A- ^{-/-} on apoE- ^{-/-} mice background on western diet	Decreases macrophage content and neointimal formation after wire injury	98
Connexin 43	Transmigration	Cx43 $^{\pm}$ on LDLR-/- mice background on western diet	Decreases development of atherosclerosis with concurrent decrease in macrophage number	99
CCL2(MCP-1)/ CCR2	Chemotaxis	CCL2- ^{f-} on LDLR- ^{f-} mice background on western diet CCR2- ^{f-} on apoE- ^{f-} mice background	Reduces macrophage recruitment by 54% Reduces development of atherosclerosis from fatty streak to more advanced lesions	5 101
CCL5(RANTES)/ CCR5	Chemotaxis	Receptor antagonist Met-RANTES in LDLR ^{-/-} mice on western diet	Reduces macrophage content by 43%	105
		Mutant [⁴⁴ AANA ⁴⁷]-RANTES in LDLR ^{-/-} mice on western diet	Limits plaque formation and reduces macrophage infiltration	106
CX3CL1(fractal- kine)/CX3CR1	Chemotaxis	CX3CL1- ⁴ -on apoE- ⁴ - or LDLR- ⁴ - background	Reduces accumulation of macrophages and dramatically reduces lesion area in brachiocephalic artery in apoE ^{-/-} mice as compared with aortic root and LDLR ^{-/-} mice	104
		CX3CR1- ^{f-} and apoE- ^{f-} double knockout mice on western diet	Decreases atherosclerotic lesion formation and macrophage infiltration by 40%	103
		CX3CR1 ^{-/-} and apoE ^{-/-} double knockout mice on chow diet	Decreases atherosclerotic lesion formation and macrophage infiltration by 50%	102
MIF	Arrest and migration	Monoclonal antibody against MIF in apoE- ⁷⁻ mice on western diet	Reduces neointimal macrophage infiltration after wire injury and inhibits transformation of macrophages into foam cells	115
		Monoclonal antibody against MIF in apoE ^{-/-} mice on chow diet	Strongly reduces intimal macrophage content and inflammatory mediators	116

promotes leucocyte extravasation and may represent a therapeutic target for atherosclerosis treatment (Table 1).

Decreased monocyte recruitment at the level of chemokines

Chemokines are chemotactic proteins that are secreted by SMCs and endothelial cells in response to proinflammatory stimuli. These proteins interact with cell surface receptors on leucocytes and, to a lesser extent, with glycosylaminoglycans of the extracellular matrix and endothelial surfaces.

Experimental studies Several chemokines and their receptors have been identified to take part actively in atherogenesis, including CCL2(MCP-1)/CCR2, CCL5(RANTES)/

CCR5 and CX3CL1 (fractalkine)/CX3CR1 (Table 1) [100]. Genetic deletion of these ligands or their receptors drastically reduces monocyte recruitment and development of diet-induced atherosclerosis [5,101–106]. Simultaneously targeting three chemokine/chemokine receptor pairs almost abolishes atherosclerosis [107]. *In vivo* antagonism of RANTES, either by the receptor antagonist Met-RANTES or by mutant [44AANA47]-RANTES that impairs essential chemokine binding and oligomerization by glycosylaminoglycans, reduces the progression of established atherosclerosis in LDLR7/- mice [105, 106]. Both treatments result in reduction of lesion size and a more stable plaque phenotype via direct inhibition of leucocyte infiltration and subsequent reduction of MMP-9. Furthermore, the increased SMC content is associated with higher amounts of intersti-

tial collagen within lesions. Given the complexity of the C-C chemokine system, blocking one chemokine/chemokine receptor pair might affect other pairs. For example, oligomerization of RANTES on glycosylaminoglycans is required for its *in vivo* function, but also for CCR1-mediated leucocyte adhesion and activation [108]. Although CCL5/RANTES binds to CCR5, which is expressed at high levels on T cells, Met-RANTES binds to CCR5 and CCR1, the latter being strongly expressed on macrophages [105].

Clinical studies In patients with increased cardiovascular risk a phase II clinical trial with MLN1202, a human monoclonal antibody towards CCR2, provides evidence that CCR2 antagonism reduces C-reactive protein concentrations, a biomarker of inflammation associated with coronary artery disease. However, MLN1202 administration results in an increase in serum MCP-1 concentrations and a decrease in circulating monocytes due to a decreased clearance by CCR2 and decreased emigration of inflammatory monocytes from the bone marrow by CCR2, respectively [109, 110]. Consequently, reduced plaque progression by CCR2 inhibition is partly attributed to direct inhibition of monocyte recruitment and partly to decreased blood monocyte count, which might increase infection risk.

Decreased monocyte recruitment at the level of the macrophage migration inhibitory factor

Macrophage migration inhibitory factor (MIF) is a cytokine with potent inflammatory functions and is highly upregulated in atherosclerotic plaques, especially at shoulder regions where most macrophages and foam cells reside.

Experimental studies MIF affects monocyte adhesion, migration and activation by upregulating adhesion molecules VCAM-1 and ICAM-1 [111], influencing chemotaxis via binding with its receptors CXCR2, CXCR4 and CD74 [112], and regulating innate immune responses via Toll-like receptor (TLR) 4 [113]. This evidence supporting the role of MIF in atherosclerosis development has encouraged the exploration of MIF inhibition as a potential therapeutic strategy. MIF-deficient mice or treatment with MIF-antibody foster changes in plaque biology towards stabilized atherosclerotic plaques by reducing macrophage and foam cell number and by increasing SMC and collagen content [114–116].

Inhibited monocyte activation by platelets

Activated platelets release chemokines, such as RANTES and platelet factor 4, and interact with monocytes through ligation of various adhesion molecules, including P-selectin, CD40L and triggering receptor expressed on myeloid cells 1 (TREM-1) ligand [117–121]. Platelet-

monocyte interaction induces activation of monocytes resulting in an increase in monocyte adhesive capacity by upregulation of β -integrins and secretion of proinflammatory cytokines, chemokines and tissue factor [122–124]. Furthermore, platelet-monocyte complexes tether and roll on endothelial cells with a very high avidity thereby enhancing endothelial activation and further monocyte recruitment [125, 126]. Accordingly, inhibition of platelet activation might reduce the number of plaque macrophages.

Experimental studies

Administration of the antiplatelet drug aspirin to hyper-cholesterolaemic mice reduces the development and progression of atherosclerotic plaques with an increase in SMCs and a decrease in macrophages [127–129]. The ADP P2Y₁₂ receptor antagonist, clopidrogrel, inhibits platelet activation and aggregation, and significantly reduces plaque size and induces a stable plaque phenotype in atherosclerotic mice [130]. Moreover, clopidogrel decreases the expression of P-selectin and CD40L on platelets and, therefore, reduces platelet-leucocyte interaction and platelet-dependent leucocyte activation [131–133].

Inhibited macrophage activation by T lymphocytes

Inhibited macrophage activation at the level of interferon-γ

The most important signal for macrophage activation is IFN- γ , a cytokine with predominantly pro-atherogenic properties, which is highly expressed in T lymphocytes of atherosclerotic plaques. IFN- γ mediates differentiation of monocytes into macrophages, pro-inflammatory cytokine release from macrophages and T cells, recruitment of leucocytes through up-regulation of chemokines and adhesion molecules, and foam cell formation through up-regulation of scavenger receptors.

Experimental studies Exogenously administered IFN-γ increases the number of T cells and MHC II-positive cells, including macrophages and dendritic cells, in the significantly expanded lesions [134]. In contrast, genetic ablation of IFN-γ or its receptor in atherosclerotic mice reduces lesions together with cellularity but increases extracellular collagen content [135]. Inhibition of IFN-γ function by gene transfer of a soluble mutant IFN-γ receptor not only prevents plaque progression but also stabilizes advanced plaques by interfering with plaque composition and inflammation [136]. The phenotypic shift towards stabilization involves decreased expression of cytokines, chemokines, adhesion molecules, MMPs and CD40. Moreover, statins and agonists of nuclear receptors, PPAR and LXR, partly exert their anti-inflammatory effects by limiting production of pro-inflammatory cytokines,



such as IFN- γ . These studies provide evidence that IFN- γ is a potential therapeutic target involved in early and late development of atherosclerosis as well as in plaque destabilization, but systemically inhibiting IFN- γ limits its clinical use due to impaired immune surveillance and function.

Inhibited macrophage activation at the level of the CD40 ligand

Another signal for macrophage activation is the CD40 ligand (CD40L) expressed on the majority of immune cells, including lymphocytes, monocytes, dendritic cells, neutrophils and mast cells, which binds CD40 on the macrophage cell surface, but it is also present on endothelial cells and SMCs. Activation of CD40-CD40L signalling in atherosclerosis mediates various functions including expression of the pro-inflammatory cytokines IL-6 and TNF- α , the pro-atherogenic chemokines CCL2, CCL5 and IL-8, MMPs, procoagulants, and the adhesion molecules VCAM-1, ICAM-1 and E-selectin. These downstream effects are associated with monocyte recruitment, inflammation, expansion of the lipid core, degradation of the fibrous cap and risk for thrombosis. Accordingly, interruption of CD40-CD40L signalling might reduce atherosclerosis and its complications.

Experimental studies Atherosclerotic plagues of CD40L^{-/-} apoE^{-/-} mice show a decrease in plague area, lipids, macrophages and T lymphocytes and an increase in collagen content [137]. Administration of anti-CD40L antibody in apoE^{-/-} and LDLR^{-/-} mice changes plague phenotype into a lipid- and macrophage-poor collagen-rich stable plague. Most profound effects are present in advanced lesions in the delayed treatment group indicating the important role of the CD40-CD40L pathway in late atherosclerotic plaques [138, 139]. Furthermore, deficiency of CD40tumour necrosis factor receptor-associated factor 6 (CD40-TRAF6) signalling in atherosclerotic mice abolishes atherosclerosis, impairs recruitment of monocytes to the arterial wall and polarizes macrophages towards an antiinflammatory phenotype [140]. However, systemic modulation of the immune system compromises host defence. Therefore, cell-type specific targeting of the CD40-CD40L pathway is under extensive investigation to overcome this adverse effect.

Selective induction of macrophage death in atherosclerotic plaques

Pharmacological depletion of macrophages via selective induction of cell death (apoptosis, autophagy) may be a promising alternative strategy to alter plaque composition in order to increase plaque stability [141]. The main challenges of this approach are (i) to initiate macrophage death in a selective way, without inducing inflammation and affecting other cell types in the plaque, in particular SMCs and endothelial cells and (ii) local application of the

drug to avoid systemic loss of macrophages. Moreover, the drug-induced type of cell death might determine possible complications. Apoptosis is a type of cell death characterized by caspase activation leading to chromatin condensation, nucleosomal DNA degradation, budding of apoptotic bodies and exposure of signalling molecules targeting for its phagocytosis. Engulfment of apoptotic bodies by phagocytic cells has been considered immunologically silent, but inefficient removal of apoptotic cells may lead to induction of postapoptotic necrosis and inflammatory responses [142]. Therefore, the complications of macrophage apoptosis depend on efficiency of apoptotic cell phagocytosis, which is defective in advanced atherosclerotic lesions further promoting plague instability [143]. Autophagy features sequestration of cytoplasmic material within double-membrane vacuoles, called autophagosomes, that mature into degrading autolysosomes after fusion with lysosomes. During autophagy, cells progressively digest the cytoplasmic content so that activation of (post-)inflammatory responses is believed to be limited. Moreover, autophagy in macrophage foam cells may enhance cholesterol efflux by delivering lipid droplets to the lysosomes where free cholesterol is generated for ABCA1-dependent-efflux [144, 145]. Therefore, induction of macrophage autophagy may have several beneficial effects towards plaque stabilization [146].

Experimental studies

By using drug-containing liposomes, cytotoxic drugs such as clodronate can be intracellularly delivered into macrophages via phagocytosis, and these liposomes are not cytotoxic to nonphagocytic cells [147]. The intracellularly metabolized clodronate inhibits mitochondrial oxygen consumption leading to apoptosis (Figure 2 (A)). This macrophage 'suicide' approach has been successful in reducing neointimal formation by reducing macrophages in vascular injury models in mice, rats and rabbits [148, 149]. However, the systemic administration of clodronatecontaining liposomes also reduces blood monocytes, which increases the risk of immunosuppression and infection. Interestingly, this adverse effect is not observed when recombinant tumour necrosis factor-related apoptosisinducing ligand (TRAIL) is systemically used in diabetic apoE^{-/-} mice. TRAIL induces apoptosis in infiltrating macrophages in atherosclerotic plaques, but not in circulating macrophages, and markedly attenuates atherosclerosis development (Figure 2 (C)) [150]. The stabilizing effects of TRAIL are also attributed to preservation of endothelial cell coverage and the enrichment of SMCs in the fibrous cap within atherosclerotic plagues. Therefore, TRAIL might represent a promising therapeutic in atherosclerosis treatment. Moreover, serum concentrations of soluble TRAIL are significantly lower in patients with cardiovascular disease, further supporting a protective role of TRAIL in atherosclerosis [151].




Figure 2

Schematic overview of targets to induce selective macrophage death. Clodronate-containing liposomes are ingested by phagocytes and the released clodronate induces apoptosis (A). Interfering with phosphatidylinositol cell signalling through inhibition of IMPase by lithium chloride leads to decreased free inositol. The subsequent decrease in IP₃ concentrations triggers apoptotic cell death (B). Systemic administration of recombinant TRAIL induces apoptosis via its death receptor in infiltrating monocytes in atherosclerotic plaques but not in circulating monocytes (C). Treatment of atherosclerotic plaques with an NO donor clears macrophages via apoptosis. Induction of macrophage apoptosis is based on inhibition of *de novo* protein synthesis and induction of ER stress (CHOP expression). ER stress inducers, thapsigargin and tunicamycin, also initiate selective macrophage apoptosis (D). Protein translation inhibitors, cycloheximide and anisomycin, selectively clear macrophages from atherosclerotic plaques via apoptosis (E). mTOR inhibitor, everolimus, induces autophagic cell death in macrophages via activation of Atg genes that attribute to autophagosome and ultimately autolysosome formation (F). Stimulation of the endosomal receptor TLR7 with imiquimod initiates autophagy exclusively in macrophages (G). TLR7 stimulation by imiquimod, as well as mTOR inhibition by everolimus, leads to the release of pro-inflammatory cytokines that can be abolished by adding anti-inflammatory drugs, such as glucocorticoids. CHOP, CCAAT/-enhancer-binding protein homologous protein; ER, endoplasmic reticulum; IMPase, inositol monophosphatase; IP₃, inositol triphosphate; mTOR, mammalian target of rapamycin; NO, nitric oxide; TLR, Toll-like receptor; TRAIL, tumour necrosis factor-related apoptosis-inducing ligand

Selective targeting of macrophages in atherosclerotic plaques is also achieved in molsidomine-treated hypercholesterolaemic rabbits. The NO donor molsidomine preferentially kills activated macrophages, e.g. inducible NO synthase (iNOS)-expressing macrophages in atherosclerotic plaques, without influencing circulating monocytes/ macrophages. Indeed, systemic treatment molsidomine during lipid lowering decreases the ratio between macrophages and SMCs favouring plaque stability [152]. In an in vitro follow-up study, it was shown that other NO donors, spermineNONOate or S-nitroso-Nacetylpenicillamine, induce endoplasmic reticulum (ER) stress, as confirmed by upregulation of CCAAT/-enhancerbinding protein homologous protein (CHOP) expression and inhibition of de novo protein synthesis, leading to macrophage apoptosis, which is likely to be independent of iNOS expression (Figure 2 (D)) [153]. Since CHOP expression was increased in molsdomine-treated atherosclerotic plagues and ER stress inducers thapsigargin and tunicamycin also initiated selective macrophage death, these data confirm the finding that induction of ER stress during treatment with an NO donor initiates selective macrophage apoptosis (Figure 2 (D)). Although ER stress is present in both macrophages and SMCs upon NO donor treatment, only macrophages undergo apoptosis. This finding might be explained by the initial presence of ER stress in intimal macrophages that is balanced by compensatory cell survival pathways. Activation of an additional mediator and/or pathway, such as ER stress by NO donors, provokes macrophage apoptosis.

On the other hand, local administration of the protein synthesis inhibitors, cycloheximide or anisomycin, to atherosclerotic plaques via osmotic minipumps selectively induces apoptosis in macrophages without inducing ER stress and without affecting SMCs and endothelial cells (Figure 2 (E)) [154,155]. The selectivity is most likely due to the high metabolic activity of macrophages, which are therefore more sensitive to protein synthesis inhibitors than SMCs, and to

modulation of SMCs towards a quiescent contractile phenotype upon protein translation inhibition, which renders these cells less sensitive to cell death. However, protein synthesis inhibition does not always induce apoptosis. Stentbased delivery of everolimus in atherosclerotic plagues of cholesterol-fed rabbits leads to a marked reduction of macrophages via autophagic cell death without altering the SMC content [156]. Everolimus-induced autophagy is mediated by inhibition of mammalian target of rapamycin (mTOR), an important serine/threonine protein kinase that is involved in cell survival (Figure 2 (F)). Besides mTOR inhibitors, a novel pathway for pharmacological modulation of autophagy has been described [157]. Lithium, an inhibitor of inositol monophosphatase, induces autophagy in nonneuronal and precursor neuronal cells by depleting inositol and consequent inositol triphosphate (IP3) concentrations. Recent experiments in our laboratory show that local treatment of atherosclerotic plaques with lithium triggers selective macrophage apoptosis, and not autophagy, by depletion of IP₃ in macrophages only (Figure 2 (B)) [158]. Another example of selective induction of macrophage autophagy is based on TLR7 expression, since it is expressed on macrophages but not on SMCs and endothelial cells (Figure 2 (G)). Stimulation of TLR7 by imidazoguinoline compound imiguimod induces autophagic cell death in macrophages without affecting other cells, at least in vitro (De Meyer, unpublished data, 2011). In vivo, however, imiquimodinduced macrophage autophagy in atherosclerotic plagues stimulates plague progression by cytokine release and enhances infiltration of inflammatory cells by increased VCAM-1 expression. These adverse pro-inflammatory effects could be abolished by adding dexamethasone. Recent in vitro results show that everolimus-induced autophagy in macrophages may also stimulate cytokine secretion [159]. These findings corroborate that the consequences of induction of selective macrophage autophagy in atherosclerotic plaques have to be taken into account, and that additional anti-inflammatory drugs are recommended to prevent inflammation and infiltration of new monocytes.

Targeting plaque macrophages

After identifying therapeutic targets to eliminate macrophages from atherosclerotic plaques, the use of drugs has been stymied by adverse effects, including impaired host defence, due to unselective drug availability. Therefore, local delivery systems to target plaque macrophages are being developed to overcome this therapeutic drawback.

Experimental studies

As previously mentioned, liposomes are effective drug carriers targeting macrophages via phagocytosis. Interestingly, target specificity can be augmented by adding ligands, such as peptides, antibodies and lectins, on the

surface of liposomes. Cyclopentanone prostaglandins (CP-PGs)-containing liposomes coated with anti-VCAM-1 antibodies specifically target VCAM-1, which is highly expressed on endothelial and foam cells at atherosclerotic sites, and deliver the anti-inflammatory CP-PGs [160]. Administration of these endothelium-directed CP-PGscontaining liposomes reverses atherosclerotic lesions via anti-inflammatory, antiproliferative, anticholesterologenic and cytoprotective effects [160]. Besides active targeting of lesional macrophages, liposomal ligands compete with the natural ligands making them viable antiatherosclerotic agents. For example, Kelly et al. suggested that hexarelin containing-liposomes directed against scavenger receptor CD36 on macrophages in atherosclerosis [161] can deliver therapeutic agents to the lesions and block LDL uptake at the same time [162].

Besides liposomes, nanoparticles are also phagocytozed by macrophages and these polymer-based nanostructures can be designed to allow controlled drug or gene delivery [163, 164]. Furthermore, they can be functionalized by ligands for active targeting towards atherosclerosis sites upon systemic administration [164, 165].

Clinical studies

Drug-eluting stents have proved to be a useful strategy for local drug delivery and prevention of in-stent restenosis [166, 167], but are compromised by complications, such as late stent thrombosis. The impaired healing of the arterial wall that is responsible for late stent thrombosis is predominantly attributed to delayed restoration of endothelial coverage and function in combination with persistent inflammation around stent struts. These adverse effects have led to the development of vascular restoration therapy by bioresorbable drug-eluting scaffolds. By degradation of the polymers and release of everolimus, the vessel lumen is significantly enlarged after 2 years and the vessel wall is characterized by infiltration of functional SMCs, absence of remodelling, restoration of endothelial cell structure and function, and reduction in plaque/media [168]. It has been hypothesized that selective induction of macrophage autophagy by everolimus may contribute to the reduction in plague size [156, 168].

Conclusion

Macrophages play a key role in the initiation and progression of atherosclerosis as well as in plaque destabilization. Continuous infiltration of monocytes into atherosclerotic plaques and macrophage survival drives disease progression and, therefore, therapies that affect overall macrophage accumulation would likely improve plaque stabilization. Despite massive use of effective drugs such as statins, the risk for cardiovascular events remains substantial. The plaque stabilizing effects of statins are attributed

to their lipid lowering properties, and partly to their anti-inflammatory effects. Therefore, additional antiinflammatory therapeutic interventions that result in reduced plague macrophage content are of great interest. New LDL lowering, HDL raising drugs, compounds that target adhesion molecules or chemokines or drugs that directly induce macrophage death tackle vascular inflammation by reducing the inflammatory stimulus in atherosclerotic plagues, promoting cholesterol efflux, inhibiting monocyte recruitment and reducing the number of plaque macrophages, respectively, and may offer new perspectives. Even though results with these new drugs are promising in animal studies, clinical trials are so far disappointing due to side effects or the lack of additional benefits. Targeting plague macrophages has been the focus of recent research, not only for cell-specific delivery of drugs, but also for the detection and imaging of vulnerable plagues. Recent progress in macrophage-targeted nanostructures and bioresorbable drug-eluting scaffolds may facilitate therapeutic atherosclerosis treatment with minimal adverse effects.

Competing Interests

There are no competing interests to declare.

This work was supported by the Agency for Innovation by Science and Technology in Flanders (IWT), the Fund for Scientific Research (FWO)-Flanders and the University of Antwerp (TOP-BOF). Illustrations are adapted from Servier Medical Art. The authors are indebted to Hans De Loof for fruitful discussions.

REFERENCES

- 1 Lendon CL, Davies MJ, Born GV, Richardson PD. Atherosclerotic plaque caps are locally weakened when macrophages density is increased. Atherosclerosis 1991; 87: 87–90.
- **2** Libby P. Inflammation in atherosclerosis. Nature 2002; 420: 868–74.
- 3 Libby P, Ridker PM, Maseri A. Inflammation and atherosclerosis. Circulation 2002; 105: 1135–43.
- 4 Nakashima Y, Raines EW, Plump AS, Breslow JL, Ross R. Upregulation of VCAM-1 and ICAM-1 at atherosclerosis-prone sites on the endothelium in the ApoE-deficient mouse. Arterioscler Thromb Vasc Biol 1998; 18: 842–51.
- **5** Gu L, Okada Y, Clinton SK, Gerard C, Sukhova GK, Libby P, Rollins BJ. Absence of monocyte chemoattractant protein-1 reduces atherosclerosis in low density lipoprotein receptor-deficient mice. Mol Cell 1998; 2: 275–81.
- **6** Inoue S, Egashira K, Ni W, Kitamoto S, Usui M, Otani K, Ishibashi M, Hiasa K, Nishida K, Takeshita A. Anti-monocyte

- chemoattractant protein-1 gene therapy limits progression and destabilization of established atherosclerosis in apolipoprotein E-knockout mice. Circulation 2002; 106: 2700–6.
- **7** Boyle JJ. Macrophage activation in atherosclerosis: pathogenesis and pharmacology of plaque rupture. Curr Vasc Pharmacol 2005; 3: 63–8.
- **8** Qiao JH, Tripathi J, Mishra NK, Cai Y, Tripathi S, Wang XP, Imes S, Fishbein MC, Clinton SK, Libby P, Lusis AJ, Rajavashisth TB. Role of macrophage colony-stimulating factor in atherosclerosis: studies of osteopetrotic mice. Am J Pathol 1997; 150: 1687–99.
- **9** Stary HC, Chandler AB, Glagov S, Guyton JR, Insull W Jr, Rosenfeld ME, Schaffer SA, Schwartz CJ, Wagner WD, Wissler RW. A definition of initial, fatty streak, and intermediate lesions of atherosclerosis. A report from the committee on vascular lesions of the council on arteriosclerosis, american heart association. Circulation 1994; 89: 2462–78.
- 10 Takahashi K, Takeya M, Sakashita N. Multifunctional roles of macrophages in the development and progression of atherosclerosis in humans and experimental animals. Med Electron Microsc 2002: 35: 179–203.
- **11** Libby P, Theroux P. Pathophysiology of coronary artery disease. Circulation 2005; 111: 3481–8.
- **12** Newby AC, George SJ, Ismail Y, Johnson JL, Sala-Newby GB, Thomas AC. Vulnerable atherosclerotic plaque metalloproteinases and foam cell phenotypes. Thromb Haemost 2009; 101: 1006–11.
- **13** Back M, Ketelhuth DF, Agewall S. Matrix metalloproteinases in atherothrombosis. Prog Cardiovasc Dis 2010; 52: 410–28.
- 14 Boyle JJ, Weissberg PL, Bennett MR. Human macrophage-induced vascular smooth muscle cell apoptosis requires NO enhancement of Fas/Fas-L interactions. Arterioscler Thromb Vasc Biol 2002; 22: 1624–30.
- **15** Hansson GK. Inflammation, atherosclerosis, and coronary artery disease. N Engl J Med 2005; 352: 1685–95.
- **16** Kockx MM, De Meyer GRY, Muhring J, Jacob W, Bult H, Herman AG. Apoptosis and related proteins in different stages of human atherosclerotic plaques. Circulation 1998; 97: 2307–15.
- 17 van der Wal AC, Becker AE, van der Loos CM, Das PK. Site of intimal rupture or erosion of thrombosed coronary atherosclerotic plaques is characterized by an inflammatory process irrespective of the dominant plaque morphology. Circulation 1994; 89: 36–44.
- **18** Libby P, Geng YJ, Aikawa M, Schoenbeck U, Mach F, Clinton SK, Sukhova GK, Lee RT. Macrophages and atherosclerotic plaque stability. Curr Opin Lipidol 1996; 7: 330–5.
- 19 Davies MJ, Thomas AC. Plaque fissuring—the cause of acute myocardial infarction, sudden ischaemic death, and crescendo angina. Br Heart J 1985; 53: 363–73.
- **20** Moreno PR, Falk E, Palacios IF, Newell JB, Fuster V, Fallon JT. Macrophage infiltration in acute coronary syndromes. Implications for plaque rupture. Circulation 1994; 90: 775–8.



- 21 Burke AP, Farb A, Malcom GT, Liang YH, Smialek J, Virmani R. Coronary risk factors and plaque morphology in men with coronary disease who died suddenly. N Engl J Med 1997; 336: 1276–82.
- 22 Jander S, Sitzer M, Schumann R, Schroeter M, Siebler M, Steinmetz H, Stoll G. Inflammation in high-grade carotid stenosis: a possible role for macrophages and T cells in plaque destabilization. Stroke 1998; 29: 1625–30.
- 23 Ward MR, Pasterkamp G, Yeung AC, Borst C. Arterial remodeling. Mechanisms and clinical implications. Circulation 2000; 102: 1186–91.
- 24 Vaughan CJ, Gotto AM Jr, Basson CT. The evolving role of statins in the management of atherosclerosis. J Am Coll Cardiol 2000; 35: 1–10.
- 25 Takemoto M, Liao JK. Pleiotropic effects of 3-hydroxy-3-methylglutaryl coenzyme a reductase inhibitors. Arterioscler Thromb Vasc Biol 2001; 21: 1712–9.
- **26** Amarenco P, Labreuche J, Lavallee P, Touboul PJ. Statins in stroke prevention and carotid atherosclerosis: systematic review and up-to-date meta-analysis. Stroke 2004; 35: 2902–9.
- 27 Gotto AM Jr. Review of primary and secondary prevention trials with lovastatin, pravastatin, and simvastatin. Am J Cardiol 2005; 96: 34F–8F.
- 28 Beckman JA, Creager MA. The nonlipid effects of statins on endothelial function. Trends Cardiovasc Med 2006; 16: 156–62.
- 29 Nassief A, Marsh JD. Statin therapy for stroke prevention. Stroke 2008; 39: 1042–8.
- **30** Liu PY, Liu YW, Lin LJ, Chen JH, Liao JK. Evidence for statin pleiotropy in humans: differential effects of statins and ezetimibe on rho-associated coiled-coil containing protein kinase activity, endothelial function, and inflammation. Circulation 2009; 119: 131–8.
- 31 Baigent C, Blackwell L, Emberson J, Holland LE, Reith C, Bhala N, Peto R, Barnes EH, Keech A, Simes J, Collins R. Efficacy and safety of more intensive lowering of LDL cholesterol: a meta-analysis of data from 170,000 participants in 26 randomised trials. Lancet 2010; 376: 1670–81.
- **32** Superko HR, King S. Lipid management to reduce cardiovascular risk: a new strategy is required. Circulation 2008; 117: 560–8.
- 33 Kastelein JJ, Sager PT, de GE, Veltri E. Comparison of ezetimibe plus simvastatin versus simvastatin monotherapy on atherosclerosis progression in familial hypercholesterolemia. Design and rationale of the Ezetimibe and Simvastatin in Hypercholesterolemia Enhances Atherosclerosis Regression (ENHANCE) trial. Am Heart J 2005: 149: 234–9.
- 34 Rossebo AB, Pedersen TR, Boman K, Brudi P, Chambers JB, Egstrup K, Gerdts E, Gohlke-Barwolf C, Holme I, Kesaniemi YA, Malbecq W, Nienaber CA, Ray S, Skjaerpe T, Wachtell K, Willenheimer R. Intensive lipid lowering with simvastatin and ezetimibe in aortic stenosis. N Engl J Med 2008; 359: 1343–56.

- **35** Taylor AJ, Villines TC, Stanek EJ, Devine PJ, Griffen L, Miller M, Weissman NJ, Turco M. Extended-release niacin or ezetimibe and carotid intima-media thickness. N Engl J Med 2009; 361: 2113–22.
- **36** Villines TC, Stanek EJ, Devine PJ, Turco M, Miller M, Weissman NJ, Griffen L, Taylor AJ. The ARBITER 6-HALTS Trial (Arterial biology for the investigation of the treatment effects of reducing cholesterol 6-HDL and LDL Treatment Strategies in Atherosclerosis): final results and the impact of medication adherence, dose, and treatment duration. J Am Coll Cardiol 2010; 55: 2721–6.
- 37 Garcia-Calvo M, Lisnock J, Bull HG, Hawes BE, Burnett DA, Braun MP, Crona JH, Davis HR Jr, Dean DC, Detmers PA, Graziano MP, Hughes M, Macintyre DE, Ogawa A, O'neill KA, Iyer SP, Shevell DE, Smith MM, Tang YS, Makarewicz AM, Ujjainwalla F, Altmann SW, Chapman KT, Thornberry NA. The target of ezetimibe is Niemann-Pick C1-Like 1 (NPC1L1). Proc Natl Acad Sci U S A 2005; 102: 8132–7.
- **38** Van Heek M, France CF, Compton DS, McLeod RL, Yumibe NP, Alton KB, Sybertz EJ, Davis HR Jr. *In vivo* metabolism-based discovery of a potent cholesterol absorption inhibitor, SCH58235, in the rat and rhesus monkey through the identification of the active metabolites of SCH48461. J Pharmacol Exp Ther 1997; 283: 157–63.
- 39 Van Heek M, Compton DS, Davis HR. The cholesterol absorption inhibitor, ezetimibe, decreases diet-induced hypercholesterolemia in monkeys. Eur J Pharmacol 2001; 415: 79–84.
- 40 Davis HR Jr, Compton DS, Hoos L, Tetzloff G. Ezetimibe, a potent cholesterol absorption inhibitor, inhibits the development of atherosclerosis in ApoE knockout mice. Arterioscler Thromb Vasc Biol 2001; 21: 2032–8.
- **41** Davis HR Jr, Pula KK, Alton KB, Burrier RE, Watkins RW. The synergistic hypocholesterolemic activity of the potent cholesterol absorption inhibitor, ezetimibe, in combination with 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors in dogs. Metabolism 2001; 50: 1234–41.
- **42** Patel R, Janoudi A, Vedre A, Aziz K, Tamhane U, Rubinstein J, Abela OG, Berger K, Abela GS. Plaque rupture and thrombosis are reduced by lowering cholesterol levels and crystallization with ezetimibe and are correlated with fluorodeoxyglucose positron emission tomography. Arterioscler Thromb Vasc Biol 2011; 31: 2007–14.
- **43** Kastelein JJ, Akdim F, Stroes ES, Zwinderman AH, Bots ML, Stalenhoef AF, Visseren FL, Sijbrands EJ, Trip MD, Stein EA, Gaudet D, Duivenvoorden R, Veltri EP, Marais AD, de GE. Simvastatin with or without ezetimibe in familial hypercholesterolemia. N Engl J Med 2008; 358: 1431–43.
- **44** Bays HE, Davidson MH, Massaad R, Flaim D, Lowe RS, Tershakovec AM, Jones-Burton C. Safety and efficacy of ezetimibe added on to rosuvastatin 5 or 10 mg versus up-titration of rosuvastatin in patients with hypercholesterolemia (the ACTE Study). Am J Cardiol 2011; 108: 523–30.
- **45** Cannon CP, Giugliano RP, Blazing MA, Harrington RA, Peterson JL, Sisk CM, Strony J, Musliner TA, McCabe CH,

- Veltri E, Braunwald E, Califf RM. Rationale and design of IMPROVE-IT (IMProved Reduction of Outcomes: vytorin Efficacy International Trial): comparison of ezetimbe/simvastatin versus simvastatin monotherapy on cardiovascular outcomes in patients with acute coronary syndromes. Am Heart J 2008; 156: 826–32.
- **46** West AM, Anderson JD, Meyer CH, Epstein FH, Wang H, Hagspiel KD, Berr SS, Harthun NL, Dimaria JM, Hunter JR, Christopher JM, Chew JD, Winberry GB, Kramer CM. The effect of ezetimibe on peripheral arterial atherosclerosis depends upon statin use at baseline. Atherosclerosis 2011; 218: 156–62.
- 47 Cohen J, Pertsemlidis A, Kotowski IK, Graham R, Garcia CK, Hobbs HH. Low LDL cholesterol in individuals of African descent resulting from frequent nonsense mutations in PCSK9. Nat Genet 2005; 37: 161–5.
- **48** Rashid S, Curtis DE, Garuti R, Anderson NN, Bashmakov Y, Ho YK, Hammer RE, Moon YA, Horton JD. Decreased plasma cholesterol and hypersensitivity to statins in mice lacking Pcsk9. Proc Natl Acad Sci U S A 2005; 102: 5374–9.
- **49** Swergold G, Biedermann S, Renard R, Nadler D, Wu R, Lisbon EA, Gutierrez MJ, Mellis S. REGN727/SAR236553, a fully human proprotein convertase subtilisin kexin 9 (PCSK9) monoclonal antibody: effects on safety and lipid and lipoprotein profiles when administered subcutaneously. J Am Coll Cardiol 2011; 57: E2023.
- 50 Gordon DJ, Knoke J, Probstfield JL, Superko R, Tyroler HA. High-density lipoprotein cholesterol and coronary heart disease in hypercholesterolemic men: the Lipid Research Clinics Coronary Primary Prevention Trial. Circulation 1986; 74: 1217–25.
- 51 Boden WE. High-density lipoprotein cholesterol as an independent risk factor in cardiovascular disease: assessing the data from Framingham to the Veterans Affairs High-Density Lipoprotein Intervention Trial. Am J Cardiol 2000: 86: 19L-22L.
- 52 Ameli S, Hultgardh-Nilsson A, Cercek B, Shah PK, Forrester JS, Ageland H, Nilsson J. Recombinant apolipoprotein A-I Milano reduces intimal thickening after balloon injury in hypercholesterolemic rabbits. Circulation 1994; 90: 1935–41.
- 53 Dimayuga P, Zhu J, Oguchi S, Chyu KY, Xu XO, Yano J, Shah PK, Nilsson J, Cercek B. Reconstituted HDL containing human apolipoprotein A-1 reduces VCAM-1 expression and neointima formation following periadventitial cuff-induced carotid injury in apoE null mice. Biochem Biophys Res Commun 1999; 264: 465–8.
- **54** Tangirala RK, Tsukamoto K, Chun SH, Usher D, Pure E, Rader DJ. Regression of atherosclerosis induced by liver-directed gene transfer of apolipoprotein A-I in mice. Circulation 1999: 100: 1816–22.
- **55** Shah PK, Kaul S, Nilsson J, Cercek B. Exploiting the vascular protective effects of high-density lipoprotein and its apolipoproteins: an idea whose time for testing is coming, part I. Circulation 2001; 104: 2376–83.
- **56** Nissen SE, Tsunoda T, Tuzcu EM, Schoenhagen P, Cooper CJ, Yasin M, Eaton GM, Lauer MA, Sheldon WS, Grines CL,

- Halpern S, Crowe T, Blankenship JC, Kerensky R. Effect of recombinant ApoA-I Milano on coronary atherosclerosis in patients with acute coronary syndromes: a randomized controlled trial. JAMA 2003; 290: 2292–300.
- 57 Tardif JC, Gregoire J, L'Allier PL, Ibrahim R, Lesperance J, Heinonen TM, Kouz S, Berry C, Basser R, Lavoie MA, Guertin MC, Rodes-Cabau J. Effects of reconstituted high-density lipoprotein infusions on coronary atherosclerosis: a randomized controlled trial. JAMA 2007; 297: 1675–82.
- **58** Frick MH, Elo O, Haapa K, Heinonen OP, Heinsalmi P, Helo P, Huttunen JK, Kaitaniemi P, Koskinen P, Manninen V. Helsinki Heart Study: primary-prevention trial with gemfibrozil in middle-aged men with dyslipidemia. Safety of treatment, changes in risk factors and incidence of coronary heart disease. N Engl J Med 1987; 317: 1237–45.
- **59** Rubins HB, Robins SJ, Collins D, Fye CL, Anderson JW, Elam MB, Faas FH, Linares E, Schaefer EJ, Schectman G, Wilt TJ, Wittes J. Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol. Veterans Affairs High-Density Lipoprotein Cholesterol Intervention Trial Study Group. N Engl J Med 1999; 341: 410–8.
- **60** The BIP study group. Secondary prevention by raising HDL cholesterol and reducing triglycerides in patients with coronary artery disease: the Bezafibrate Infarction Prevention (BIP) study. Circulation 2000; 102: 21–7.
- 61 Keech A, Simes RJ, Barter P, Best J, Scott R, Taskinen MR, Forder P, Pillai A, Davis T, Glasziou P, Drury P, Kesaniemi YA, Sullivan D, Hunt D, Colman P, d'Emden M, Whiting M, Ehnholm C, Laakso M. Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): randomised controlled trial. Lancet 2005; 366: 1849–61.
- **62** Ginsberg HN, Elam MB, Lovato LC, Crouse JR III, Leiter LA, Linz P, Friedewald WT, Buse JB, Gerstein HC, Probstfield J, Grimm RH, Ismail-Beigi F, Bigger JT, Goff DC Jr, Cushman WC, Simons-Morton DG, Byington RP. Effects of combination lipid therapy in type 2 diabetes mellitus. N Engl J Med 2010; 362: 1563–74.
- 63 AIM-HIGH investigators. The role of niacin in raising high-density lipoprotein cholesterol to reduce cardiovascular events in patients with atherosclerotic cardiovascular disease and optimally treated low-density lipoprotein cholesterol Rationale and study design. The Atherothrombosis Intervention in Metabolic syndrome with low HDL/high triglycerides: impact on Global Health outcomes (AIM-HIGH). Am Heart J 2011; 161: 471–7.
- **64** Nissen SE, Tardif JC, Nicholls SJ, Revkin JH, Shear CL, Duggan WT, Ruzyllo W, Bachinsky WB, Lasala GP, Tuzcu EM. Effect of torcetrapib on the progression of coronary atherosclerosis. N Engl J Med 2007; 356: 1304–16.
- 65 Schwartz GG, Olsson AG, Ballantyne CM, Barter PJ, Holme IM, Kallend D, Leiter LA, Leitersdorf E, McMurray JJ, Shah PK, Tardif JC, Chaitman BR, Duttlinger-Maddux R, Mathieson J. Rationale and design of the dal-OUTCOMES trial: efficacy and safety of dalcetrapib in patients with recent acute coronary syndrome. Am Heart J 2009; 158: 896–901.



- **66** Cannon CP, Shah S, Dansky HM, Davidson M, Brinton EA, Gotto AM, Stepanavage M, Liu SX, Gibbons P, Ashraf TB, Zafarino J, Mitchel Y, Barter P. Safety of anacetrapib in patients with or at high risk for coronary heart disease. N Engl J Med 2010; 363: 2406–15.
- 67 Llera-Moya M, Drazul-Schrader D, Asztalos BF, Cuchel M, Rader DJ, Rothblat GH. The ability to promote efflux via ABCA1 determines the capacity of serum specimens with similar high-density lipoprotein cholesterol to remove cholesterol from macrophages. Arterioscler Thromb Vasc Biol 2010; 30: 796–801.
- **68** Yvan-Charvet L, Ranalletta M, Wang N, Han S, Terasaka N, Li R, Welch C, Tall AR. Combined deficiency of ABCA1 and ABCG1 promotes foam cell accumulation and accelerates atherosclerosis in mice. J Clin Invest 2007; 117: 3900–8.
- 69 Beyea MM, Heslop CL, Sawyez CG, Edwards JY, Markle JG, Hegele RA, Huff MW. Selective up-regulation of LXR-regulated genes ABCA1, ABCG1 and APOE in macrophages through increased endogenous synthesis of 24(S),25-epoxycholesterol. J Biol Chem 2007; 282: 5207–16.
- **70** Levin N, Bischoff ED, Daige CL, Thomas D, Vu CT, Heyman RA, Tangirala RK, Schulman IG. Macrophage liver X receptor is required for antiatherogenic activity of LXR agonists. Arterioscler Thromb Vasc Biol 2005; 25: 135–42.
- 71 Peng D, Hiipakka RA, Dai Q, Guo J, Reardon CA, Getz GS, Liao S. Antiatherosclerotic effects of a novel synthetic tissue-selective steroidal liver X receptor agonist in low-density lipoprotein receptor-deficient mice. J Pharmacol Exp Ther 2008; 327: 332–42.
- 72 Chawla A, Boisvert WA, Lee CH, Laffitte BA, Barak Y, Joseph SB, Liao D, Nagy L, Edwards PA, Curtiss LK, Evans RM, Tontonoz P. A PPAR gamma-LXR-ABCA1 pathway in macrophages is involved in cholesterol efflux and atherogenesis. Mol Cell 2001; 7: 161–71.
- 73 Chinetti G, Lestavel S, Bocher V, Remaley AT, Neve B, Torra IP, Teissier E, Minnich A, Jaye M, Duverger N, Brewer HB, Fruchart JC, Clavey V, Staels B. PPAR-alpha and PPAR-gamma activators induce cholesterol removal from human macrophage foam cells through stimulation of the ABCA1 pathway. Nat Med 2001; 7: 53–8.
- 74 Li AC, Brown KK, Silvestre MJ, Willson TM, Palinski W, Glass CK. Peroxisome proliferator-activated receptor gamma ligands inhibit development of atherosclerosis in LDL receptor-deficient mice. J Clin Invest 2000; 106: 523–31.
- 75 Corti R, Osende J, Hutter R, Viles-Gonzalez JF, Zafar U, Valdivieso C, Mizsei G, Fallon JT, Fuster V, Badimon JJ. Fenofibrate induces plaque regression in hypercholesterolemic atherosclerotic rabbits: in vivo demonstration by high-resolution MRI. Atherosclerosis 2007; 190: 106–13.
- 76 Khera AV, Cuchel M, Llera-Moya M, Rodrigues A, Burke MF, Jafri K, French BC, Phillips JA, Mucksavage ML, Wilensky RL, Mohler ER, Rothblat GH, Rader DJ. Cholesterol efflux capacity, high-density lipoprotein function, and atherosclerosis. N Engl J Med 2011; 364: 127–35.
- **77** Lewis GF, Rader DJ. New insights into the regulation of HDL metabolism and reverse cholesterol transport. Circ Res 2005; 96: 1221–32.

- **78** Yamamoto S, Tanigawa H, Li X, Komaru Y, Billheimer JT, Rader DJ. Pharmacologic suppression of hepatic ATP-binding cassette transporter 1 activity in mice reduces high-density lipoprotein cholesterol levels but promotes reverse cholesterol transport. Circulation 2011; 124: 1382–90.
- 79 Yamashita S, Hbujo H, Arai H, Harada-Shiba M, Matsui S, Fukushima M, Saito Y, Kita T, Matsuzawa Y. Long-term probucol treatment prevents secondary cardiovascular events: a cohort study of patients with heterozygous familial hypercholesterolemia in Japan. J Atheroscler Thromb 2008; 15: 292–303.
- 80 Brasen JH, Koenig K, Bach H, Kontush A, Heinle H, Witting PK, Yla-Herttuala S, Stocker R, Beisiegel U. Comparison of the effects of alpha-tocopherol, ubiquinone-10 and probucol at therapeutic doses on atherosclerosis in WHHL rabbits. Atherosclerosis 2002; 163: 249–59.
- 81 Braun A, Zhang S, Miettinen HE, Ebrahim S, Holm TM, Vasile E, Post MJ, Yoerger DM, Picard MH, Krieger JL, Andrews NC, Simons M, Krieger M. Probucol prevents early coronary heart disease and death in the high-density lipoprotein receptor SR-Bl/apolipoprotein E double knockout mouse. Proc Natl Acad Sci U S A 2003; 100: 7283–8.
- **82** Favari E, Zanotti I, Zimetti F, Ronda N, Bernini F, Rothblat GH. Probucol inhibits ABCA1-mediated cellular lipid efflux. Arterioscler Thromb Vasc Biol 2004; 24: 2345–50.
- **83** Franceschini G, Chiesa G, Sirtori CR. Probucol increases cholesteryl ester transfer protein activity in hypercholesterolaemic patients. Eur J Clin Invest 1991; 21: 384–8.
- **84** Kumar A, Hoover JL, Simmons CA, Lindner V, Shebuski RJ. Remodeling and neointimal formation in the carotid artery of normal and P-selectin-deficient mice. Circulation 1997; 96: 4333–42.
- **85** Nageh MF, Sandberg ET, Marotti KR, Lin AH, Melchior EP, Bullard DC, Beaudet AL. Deficiency of inflammatory cell adhesion molecules protects against atherosclerosis in mice. Arterioscler Thromb Vasc Biol 1997; 17: 1517–20.
- **86** Collins RG, Velji R, Guevara NV, Hicks MJ, Chan L, Beaudet AL. P-Selectin or intercellular adhesion molecule (ICAM)-1 deficiency substantially protects against atherosclerosis in apolipoprotein E-deficient mice. J Exp Med 2000; 191: 189–94.
- **87** Dong ZM, Brown AA, Wagner DD. Prominent role of P-selectin in the development of advanced atherosclerosis in ApoE-deficient mice. Circulation 2000; 101: 2290–5.
- 88 Dansky HM, Barlow CB, Lominska C, Sikes JL, Kao C, Weinsaft J, Cybulsky MI, Smith JD. Adhesion of monocytes to arterial endothelium and initiation of atherosclerosis are critically dependent on vascular cell adhesion molecule-1 gene dosage. Arterioscler Thromb Vasc Biol 2001; 21: 1662–7
- **89** Manka D, Collins RG, Ley K, Beaudet AL, Sarembock IJ. Absence of P-selectin, but not intercellular adhesion

- molecule-1, attenuates neointimal growth after arterial injury in apolipoprotein e-deficient mice. Circulation 2001; 103: 1000–5.
- 90 Patel SS, Thiagarajan R, Willerson JT, Yeh ET. Inhibition of alpha4 integrin and ICAM-1 markedly attenuate macrophage homing to atherosclerotic plaques in ApoE-deficient mice. Circulation 1998; 97: 75–81.
- 91 Oguchi S, Dimayuga P, Zhu J, Chyu KY, Yano J, Shah PK, Nilsson J, Cercek B. Monoclonal antibody against vascular cell adhesion molecule-1 inhibits neointimal formation after periadventitial carotid artery injury in genetically hypercholesterolemic mice. Arterioscler Thromb Vasc Biol 2000; 20: 1729–36.
- 92 Barringhaus KG, Phillips JW, Thatte JS, Sanders JM, Czarnik AC, Bennett DK, Ley KF, Sarembock IJ. Alpha4beta1 integrin (VLA-4) blockade attenuates both early and late leukocyte recruitment and neointimal growth following carotid injury in apolipoprotein E (-/-) mice. J Vasc Res 2004; 41: 252–60.
- **93** Tardif JC, McMurray JJ, Klug E, Small R, Schumi J, Choi J, Cooper J, Scott R, Lewis EF, L'Allier PL, Pfeffer MA. Effects of succinobucol (AGI-1067) after an acute coronary syndrome: a randomised, double-blind, placebo-controlled trial. Lancet 2008; 371: 1761–8.
- **94** Ni W, Egashira K, Kataoka C, Kitamoto S, Koyanagi M, Inoue S, Takeshita A. Antiinflammatory and antiarteriosclerotic actions of HMG-CoA reductase inhibitors in a rat model of chronic inhibition of nitric oxide synthesis. Circ Res 2001; 89: 415–21.
- **95** Veillard NR, Braunersreuther V, Arnaud C, Burger F, Pelli G, Steffens S, Mach F. Simvastatin modulates chemokine and chemokine receptor expression by geranylgeranyl isoprenoid pathway in human endothelial cells and macrophages. Atherosclerosis 2006; 188: 51–8.
- **96** Weitz-Schmidt G, Welzenbach K, Brinkmann V, Kamata T, Kallen J, Bruns C, Cottens S, Takada Y, Hommel U. Statins selectively inhibit leukocyte function antigen-1 by binding to a novel regulatory integrin site. Nat Med 2001; 7: 687–92.
- 97 Ostermann G, Fraemohs L, Baltus T, Schober A, Lietz M, Zernecke A, Liehn EA, Weber C. Involvement of JAM-A in mononuclear cell recruitment on inflamed or atherosclerotic endothelium: inhibition by soluble JAM-A. Arterioscler Thromb Vasc Biol 2005; 25: 729–35.
- 98 Zernecke A, Liehn EA, Fraemohs L, von HP, Koenen RR, Corada M, Dejana E, Weber C. Importance of junctional adhesion molecule-A for neointimal lesion formation and infiltration in atherosclerosis-prone mice. Arterioscler Thromb Vasc Biol 2006; 26: e10–e13.
- **99** Kwak BR, Veillard N, Pelli G, Mulhaupt F, James RW, Chanson M, Mach F. Reduced connexin43 expression inhibits atherosclerotic lesion formation in low-density lipoprotein receptor-deficient mice. Circulation 2003; 107: 1033–9.
- **100** Gautier EL, Jakubzick C, Randolph GJ. Regulation of the migration and survival of monocyte subsets by chemokine

- receptors and its relevance to atherosclerosis. Arterioscler Thromb Vasc Biol 2009; 29: 1412–8.
- 101 Dawson TC, Kuziel WA, Osahar TA, Maeda N. Absence of CC chemokine receptor-2 reduces atherosclerosis in apolipoprotein E-deficient mice. Atherosclerosis 1999; 143: 205–11.
- 102 Combadiere C, Potteaux S, Gao JL, Esposito B, Casanova S, Lee EJ, Debre P, Tedgui A, Murphy PM, Mallat Z. Decreased atherosclerotic lesion formation in CX3CR1/apolipoprotein E double knockout mice. Circulation 2003; 107: 1009–16.
- **103** Lesnik P, Haskell CA, Charo IF. Decreased atherosclerosis in CX3CR1-/- mice reveals a role for fractalkine in atherogenesis. J Clin Invest 2003; 111: 333–40.
- 104 Teupser D, Pavlides S, Tan M, Gutierrez-Ramos JC, Kolbeck R, Breslow JL. Major reduction of atherosclerosis in fractalkine (CX3CL1)-deficient mice is at the brachiocephalic artery, not the aortic root. Proc Natl Acad Sci U S A 2004; 101: 17795–800.
- 105 Veillard NR, Kwak B, Pelli G, Mulhaupt F, James RW, Proudfoot AE, Mach F. Antagonism of RANTES receptors reduces atherosclerotic plaque formation in mice. Circ Res 2004; 94: 253–61.
- 106 Braunersreuther V, Steffens S, Arnaud C, Pelli G, Burger F, Proudfoot A, Mach F. A novel RANTES antagonist prevents progression of established atherosclerotic lesions in mice. Arterioscler Thromb Vasc Biol 2008; 28: 1090–6.
- 107 Combadiere C, Potteaux S, Rodero M, Simon T, Pezard A, Esposito B, Merval R, Proudfoot A, Tedgui A, Mallat Z. Combined inhibition of CCL2, CX3CR1, and CCR5 abrogates Ly6C(hi) and Ly6C(lo) monocytosis and almost abolishes atherosclerosis in hypercholesterolemic mice. Circulation 2008: 117: 1649–57.
- **108** Baltus T, Weber KS, Johnson Z, Proudfoot AE, Weber C. Oligomerization of RANTES is required for CCR1-mediated arrest but not CCR5-mediated transmigration of leukocytes on inflamed endothelium. Blood 2003; 102: 1985–8.
- 109 Serbina NV, Pamer EG. Monocyte emigration from bone marrow during bacterial infection requires signals mediated by chemokine receptor CCR2. Nat Immunol 2006; 7: 311–7.
- **110** Gilbert J, Lekstrom-Himes J, Donaldson D, Lee Y, Hu M, Xu J, Wyant T, Davidson M. Effect of CC chemokine receptor 2 CCR2 blockade on serum C-reactive protein in individuals at atherosclerotic risk and with a single nucleotide polymorphism of the monocyte chemoattractant protein-1 promoter region. Am J Cardiol 2011; 107: 906–11.
- **111** Amin MA, Haas CS, Zhu K, Mansfield PJ, Kim MJ, Lackowski NP, Koch AE. Migration inhibitory factor up-regulates vascular cell adhesion molecule-1 and intercellular adhesion molecule-1 via Src, PI3 kinase, and NFkappaB. Blood 2006; 107: 2252–61.
- 112 Kraemer S, Lue H, Zernecke A, Kapurniotu A, Andreetto E, Frank R, Lennartz B, Weber C, Bernhagen J. MIF-chemokine receptor interactions in atherogenesis are dependent on an N-loop-based 2-site binding mechanism. FASEB J 2011; 25: 894–906.



- **113** Roger T, David J, Glauser MP, Calandra T. MIF regulates innate immune responses through modulation of Toll-like receptor 4. Nature 2001; 414: 920–4.
- 114 Pan JH, Sukhova GK, Yang JT, Wang B, Xie T, Fu H, Zhang Y, Satoskar AR, David JR, Metz CN, Bucala R, Fang K, Simon DI, Chapman HA, Libby P, Shi GP. Macrophage migration inhibitory factor deficiency impairs atherosclerosis in low-density lipoprotein receptor-deficient mice. Circulation 2004; 109: 3149–53.
- 115 Schober A, Bernhagen J, Thiele M, Zeiffer U, Knarren S, Roller M, Bucala R, Weber C. Stabilization of atherosclerotic plaques by blockade of macrophage migration inhibitory factor after vascular injury in apolipoprotein E-deficient mice. Circulation 2004; 109: 380–5.
- 116 Burger-Kentischer A, Gobel H, Kleemann R, Zernecke A, Bucala R, Leng L, Finkelmeier D, Geiger G, Schaefer HE, Schober A, Weber C, Brunner H, Rutten H, Ihling C, Bernhagen J. Reduction of the aortic inflammatory response in spontaneous atherosclerosis by blockade of macrophage migration inhibitory factor (MIF). Atherosclerosis 2006; 184: 28–38.
- 117 Lievens D, Zernecke A, Seijkens T, Soehnlein O, Beckers L, Munnix IC, Wijnands E, Goossens P, van KR, Thevissen L, Boon L, Flavell RA, Noelle RJ, Gerdes N, Biessen EA, Daemen MJ, Heemskerk JW, Weber C, Lutgens E. Platelet CD40L mediates thrombotic and inflammatory processes in atherosclerosis. Blood 2010; 116: 4317–27.
- **118** Scheuerer B, Ernst M, Durrbaum-Landmann I, Fleischer J, Grage-Griebenow E, Brandt E, Flad HD, Petersen F. The CXC-chemokine platelet factor 4 promotes monocyte survival and induces monocyte differentiation into macrophages. Blood 2000; 95: 1158–66.
- **119** von HP, Weber KS, Huo Y, Proudfoot AE, Nelson PJ, Ley K, Weber C. RANTES deposition by platelets triggers monocyte arrest on inflamed and atherosclerotic endothelium. Circulation 2001; 103: 1772–7.
- **120** Huo Y, Schober A, Forlow SB, Smith DF, Hyman MC, Jung S, Littman DR, Weber C, Ley K. Circulating activated platelets exacerbate atherosclerosis in mice deficient in apolipoprotein E. Nat Med 2003; 9: 61–7.
- **121** Bouchon A, Dietrich J, Colonna M. Cutting edge: inflammatory responses can be triggered by TREM-1, a novel receptor expressed on neutrophils and monocytes. J Immunol 2000; 164: 4991–5.
- **122** da Costa Martins PA, van Gils JM, Mol A, Hordijk PL, Zwaginga JJ. Platelet binding to monocytes increases the adhesive properties of monocytes by up-regulating the expression and functionality of beta1 and beta2 integrins. J Leukoc Biol 2006; 79: 499–507.
- **123** Christersson C, Johnell M, Siegbahn A. Tissue factor and IL8 production by P-selectin-dependent platelet-monocyte aggregates in whole blood involves phosphorylation of Lyn and is inhibited by IL10. J Thromb Haemost 2008; 6: 986–94.
- **124** Rinder HM, Bonan JL, Rinder CS, Ault KA, Smith BR. Dynamics of leukocyte-platelet adhesion in whole blood. Blood 1991; 78: 1730–7.

- **125** van Gils JM, Zwaginga JJ, Hordijk PL. Molecular and functional interactions among monocytes, platelets, and endothelial cells and their relevance for cardiovascular diseases. J Leukoc Biol 2009: 85: 195–204.
- 126 da Costa MP, van den Berk N, Ulfman LH, Koenderman L, Hordijk PL, Zwaginga JJ. Platelet-monocyte complexes support monocyte adhesion to endothelium by enhancing secondary tethering and cluster formation. Arterioscler Thromb Vasc Biol 2004; 24: 193–9.
- **127** Cyrus T, Sung S, Zhao L, Funk CD, Tang S, Pratico D. Effect of low-dose aspirin on vascular inflammation, plaque stability, and atherogenesis in low-density lipoprotein receptor-deficient mice. Circulation 2002; 106: 1282–7.
- **128** Tous M, Ferre N, Vilella E, Riu F, Camps J, Joven J. Aspirin attenuates the initiation but not the progression of atherosclerosis in apolipoprotein E-deficient mice fed a high-fat, high-cholesterol diet. Basic Clin Pharmacol Toxicol 2004: 95: 15–9.
- **129** Cyrus T, Yao Y, Tung LX, Pratico D. Stabilization of advanced atherosclerosis in low-density lipoprotein receptor-deficient mice by aspirin. Atherosclerosis 2006; 184: 8–14.
- **130** Afek A, Kogan E, Maysel-Auslender S, Mor A, Regev E, Rubinstein A, Keren G, George J. Clopidogrel attenuates atheroma formation and induces a stable plaque phenotype in apolipoprotein E knockout mice. Microvasc Res 2009; 77: 364–9.
- **131** Cha JK, Jeong MH, Lee KM, Bae HR, Lim YJ, Park KW, Cheon SM. Changes in platelet P-selectin and in plasma C-reactive protein in acute atherosclerotic ischemic stroke treated with a loading dose of clopidogrel. J Thromb Thrombolysis 2002; 14: 145–50.
- **132** Hermann A, Rauch BH, Braun M, Schror K, Weber AA. Platelet CD40 ligand (CD40L)–subcellular localization, regulation of expression, and inhibition by clopidogrel. Platelets 2001; 12: 74–82.
- **133** Evangelista V, Manarini S, Dell'Elba G, Martelli N, Napoleone E, Di SA, Lorenzet PS. Clopidogrel inhibits platelet-leukocyte adhesion and platelet-dependent leukocyte activation. Thromb Haemost 2005; 94: 568–77.
- **134** Whitman SC, Ravisankar P, Elam H, Daugherty A. Exogenous interferon-gamma enhances atherosclerosis in apolipoprotein E-/- mice. Am J Pathol 2000; 157: 1819–24.
- **135** Gupta S, Pablo AM, Jiang X, Wang N, Tall AR, IF SC. N-gamma potentiates atherosclerosis in ApoE knock-out mice. J Clin Invest 1997; 99: 2752–61.
- **136** Koga M, Kai H, Yasukawa H, Yamamoto T, Kawai Y, Kato S, Kusaba K, Kai M, Egashira K, Kataoka Y, Imaizumi T. Inhibition of progression and stabilization of plaques by postnatal interferon-gamma function blocking in ApoE-knockout mice. Circ Res 2007; 101: 348–56.
- **137** Lutgens E, Gorelik L, Daemen MJ, de Muinck ED, Grewal IS, Koteliansky VE, Flavell RA. Requirement for CD154 in the progression of atherosclerosis. Nat Med 1999; 5: 1313–6.
- **138** Lutgens E, Cleutjens KB, Heeneman S, Koteliansky VE, Burkly LC, Daemen MJ. Both early and delayed anti-CD40L

- antibody treatment induces a stable plaque phenotype. Proc Natl Acad Sci U S A 2000; 97: 7464–9.
- **139** Schonbeck U, Sukhova GK, Shimizu K, Mach F, Libby P. Inhibition of CD40 signaling limits evolution of established atherosclerosis in mice. Proc Natl Acad Sci U S A 2000; 97: 7458–63.
- 140 Lutgens E, Lievens D, Beckers L, Wijnands E, Soehnlein O, Zernecke A, Seijkens T, Engel D, Cleutjens J, Keller AM, Naik SH, Boon L, Oufella HA, Mallat Z, Ahonen CL, Noelle RJ, de Winther MP, Daemen MJ, Biessen EA, Weber C. Deficient CD40-TRAF6 signaling in leukocytes prevents atherosclerosis by skewing the immune response toward an antiinflammatory profile. J Exp Med 2010; 207: 391–404.
- 141 Martinet W, Verheye S, De Meyer GRY. Selective depletion of macrophages in atherosclerotic plaques via macrophage-specific initiation of cell death. Trends Cardiovasc Med 2007: 17: 69–75.
- **142** Kockx MM, Herman AG. Apoptosis in atherosclerosis: beneficial or detrimental? Cardiovasc Res 2000; 45: 736–46.
- **143** Schrijvers DM, De Meyer GRY, Kockx MM, Herman AG, Martinet W. Phagocytosis of apoptotic cells by macrophages is impaired in atherosclerosis. Arterioscler Thromb Vasc Biol 2005; 25: 1256–61.
- **144** Ouimet M, Franklin V, Mak E, Liao X, Tabas I, Marcel YL. Autophagy regulates cholesterol efflux from macrophage foam cells via lysosomal acid lipase. Cell Metab 2011; 13: 655–67.
- **145** Ouimet M, Marcel YL. Regulation of lipid droplet cholesterol efflux from macrophage foam cells. Arterioscler Thromb Vasc Biol 2012; 32: 575–81.
- **146** Schrijvers DM, De Meyer GRY, Martinet W. Autophagy in atherosclerosis: a potential drug target for plaque stabilization. Arterioscler Thromb Vasc Biol 2011; 31: 2787–91.
- **147** Van Rooijen N, Sanders A, van den Berg TK. Apoptosis of macrophages induced by liposome-mediated intracellular delivery of clodronate and propamidine. J Immunol Methods 1996; 193: 93–9.
- 148 Danenberg HD, Fishbein I, Gao J, Monkkonen J, Reich R, Gati I, Moerman E, Golomb G. Macrophage depletion by clodronate-containing liposomes reduces neointimal formation after balloon injury in rats and rabbits. Circulation 2002; 106: 599–605.
- **149** Ayala-Lopez W, Xia W, Varghese B, Low PS. Imaging of atherosclerosis in apoliprotein e knockout mice: targeting of a folate-conjugated radiopharmaceutical to activated macrophages. J Nucl Med 2010; 51: 768–74.
- **150** Secchiero P, Candido R, Corallini F, Zacchigna S, Toffoli B, Rimondi E, Fabris B, Giacca M, Zauli G. Systemic tumor necrosis factor-related apoptosis-inducing ligand delivery shows antiatherosclerotic activity in apolipoprotein E-null diabetic mice. Circulation 2006; 114: 1522–30.
- **151** Michowitz Y, Goldstein E, Roth A, Afek A, Abashidze A, Ben GY, Keren G, George J. The involvement of tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) in atherosclerosis. J Am Coll Cardiol 2005; 45: 1018–24.

- **152** De Meyer GRY, Kockx MM, Knaapen MWM, Martinet W, De Cleen DM, Bult H, Herman AG. Nitric oxide donor molsidomine favors features of atherosclerotic plaque stability during cholesterol lowering in rabbits. J Cardiovasc Pharmacol 2003; 41: 970–8.
- **153** Martinet W, Croons V, Timmermans JP, Herman AG, De Meyer GRY. Nitric oxide selectively depletes macrophages in atherosclerotic plaques via induction of endoplasmic reticulum stress. Br J Pharmacol 2007; 152: 493–500.
- **154** Croons V, Martinet W, Herman AG, Timmermans JP, De Meyer GRY. Selective clearance of macrophages in atherosclerotic plaques by the protein synthesis inhibitor cycloheximide. J Pharmacol Exp Ther 2007; 320: 986–93.
- **155** Croons V, Martinet W, Herman AG, Timmermans JP, De Meyer GRY. The protein synthesis inhibitor anisomycin induces macrophage apoptosis in rabbit atherosclerotic plaques through p38 mitogen-activated protein kinase. J Pharmacol Exp Ther 2009; 329: 856–64.
- **156** Verheye S, Martinet W, Kockx MM, Knaapen MWM, Salu K, Timmermans JP, Ellis JT, Kilpatrick DL, De Meyer GRY. Selective clearance of macrophages in atherosclerotic plaques by autophagy. J Am Coll Cardiol 2007; 49: 706–15.
- **157** Sarkar S, Floto RA, Berger Z, Imarisio S, Cordenier A, Pasco M, Cook LJ, Rubinsztein DC. Lithium induces autophagy by inhibiting inositol monophosphatase. J Cell Biol 2005; 170: 1101–11.
- **158** De Meyer I, Martinet W, Van Hove CE, Schrijvers DM, Hoymans VY, Van Vaeck L, Fransen P, Bult H, De Meyer GRY. Inhibition of inositol monophosphatase by lithium chloride induces selective macrophage apoptosis in atherosclerotic plaques. Br J Pharmacol 2011; 162: 1410–23.
- 159 Martinet W, Verheye S, De Meyer I, Timmermans JP, Schrijvers DM, Van Brussel I, Bult H, De Meyer GRY. Everolimus triggers cytokine release by macrophages: rationale for stents eluting everolimus and a glucocorticoid. Arterioscler Thromb Vasc Biol 2012; doi:10.1161/ATVBAHA.112.245381.
- 160 Homem de Bittencourt PI Jr, Lagranha DJ, Maslinkiewicz A, Senna SM, Tavares AM, Baldissera LP, Janner DR, Peralta JS, Bock PM, Gutierrez LL, Scola G, Heck TG, Krause MS, Cruz LA, Abdalla DS, Lagranha CJ, Lima T, Curi R. LipoCardium: endothelium-directed cyclopentenone prostaglandin-based liposome formulation that completely reverses atherosclerotic lesions. Atherosclerosis 2007; 193: 245–58.
- 161 Demers A, McNicoll N, Febbraio M, Servant M, Marleau S, Silverstein R, Ong H. Identification of the growth hormone-releasing peptide binding site in CD36: a photoaffinity cross-linking study. Biochem J 2004; 382: 417–24.
- **162** Kelly C, Jefferies C, Cryan SA. Targeted liposomal drug delivery to monocytes and macrophages. J Drug Deliv 2011; 2011: 1–11.
- **163** Aouadi M, Tesz GJ, Nicoloro SM, Wang M, Chouinard M, Soto E, Ostroff GR, Czech MP. Orally delivered siRNA



- targeting macrophage Map4k4 suppresses systemic inflammation. Nature 2009; 458: 1180–4.
- 164 Chan JM, Zhang L, Tong R, Ghosh D, Gao W, Liao G, Yuet KP, Gray D, Rhee JW, Cheng J, Golomb G, Libby P, Langer R, Farokhzad OC. Spatiotemporal controlled delivery of nanoparticles to injured vasculature. Proc Natl Acad Sci U S A 2010: 107: 2213–8.
- **165** Winter PM, Neubauer AM, Caruthers SD, Harris TD, Robertson JD, Williams TA, Schmieder AH, Hu G, Allen JS, Lacy EK, Zhang H, Wickline SA, Lanza GM. Endothelial alpha(v)beta3 integrin-targeted fumagillin nanoparticles inhibit angiogenesis in atherosclerosis. Arterioscler Thromb Vasc Biol 2006; 26: 2103–9.
- **166** Rensing BJ, Vos J, Smits PC, Foley DP, van den Brand MJ, van der Giessen WJ, de Feijter PJ, Serruys PW. Coronary restenosis elimination with a sirolimus eluting stent: first European human experience with 6-month angiographic

- and intravascular ultrasonic follow-up. Eur Heart J 2001; 22: 2125–30.
- 167 Sousa JE, Costa MA, Abizaid A, Abizaid AS, Feres F, Pinto IM, Seixas AC, Staico R, Mattos LA, Sousa AG, Falotico R, Jaeger J, Popma JJ, Serruys PW. Lack of neointimal proliferation after implantation of sirolimus-coated stents in human coronary arteries: a quantitative coronary angiography and three-dimensional intravascular ultrasound study. Circulation 2001; 103: 192–5.
- **168** Onuma Y, Serruys PW. Bioresorbable scaffold: the advent of a new era in percutaneous coronary and peripheral revascularization? Circulation 2011; 123: 779–97.
- **169** Cybulsky MI, liyama K, Li H, Zhu S, Chen M, liyama M, Davis V, Gutierrez-Ramos JC, Connelly PW, Milstone DS. A major role for VCAM-1, but not ICAM-1, in early atherosclerosis. J Clin Invest 2001; 107: 1255–62.