

See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/51639234>

# Selective liver X receptor modulators (SLiMs): What use in human health?

ARTICLE *in* MOLECULAR AND CELLULAR ENDOCRINOLOGY · SEPTEMBER 2011

Impact Factor: 4.24 · DOI: 10.1016/j.mce.2011.08.036 · Source: PubMed

CITATIONS

37

DOWNLOADS

460

VIEWS

283

6 AUTHORS, INCLUDING:



**Emilie Viennois**

Georgia State University

37 PUBLICATIONS 249 CITATIONS

SEE PROFILE



**Julie Dufour**

12 PUBLICATIONS 138 CITATIONS

SEE PROFILE



**Silvère Baron**

Université Blaise Pascal - Clermont-Ferrand II

40 PUBLICATIONS 499 CITATIONS

SEE PROFILE

Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

## Molecular and Cellular Endocrinology

journal homepage: [www.elsevier.com/locate/mce](http://www.elsevier.com/locate/mce)

## Review

## Selective liver X receptor modulators (SLiMs): What use in human health?

Emilie Viennois<sup>a,b,c</sup>, Kevin Mouzat<sup>d</sup>, Julie Dufour<sup>a,b,c</sup>, Laurent Morel<sup>a,b,c</sup>, Jean-Marc Lobaccaro<sup>a,b,c,\*</sup>,  
Silvère Baron<sup>a,b,c</sup>

<sup>a</sup> Clermont Université, Université Blaise Pascal, Génétique Reproduction et Développement, BP 10448, F-63000 Clermont-Ferrand, France

<sup>b</sup> CNRS, UMR 6247, GReD, F-63177 Aubière, France

<sup>c</sup> Centre de Recherche en Nutrition Humaine d'Auvergne, F-63000 Clermont-Ferrand, France

<sup>d</sup> Laboratoire de Biochimie, Hôpital Universitaire Carêmeau, Place du Pr. Robert Debré, F-30029 Nîmes, France

## ARTICLE INFO

Article history:  
Available online xxx

Keywords:  
Liver X receptors  
Pharmacology  
SLiMs  
Human pathologies  
Cholesterol  
Nuclear receptors

## ABSTRACT

Liver X receptors (LXR) are members of the nuclear receptor family. As activated transcription factors, their putative association with human diseases makes them promising pharmacological targets because of the large potential to develop ligands. LXR are mainly considered as intracellular cholesterol “sensors” whose activation leads to decreased plasma cholesterol. They also modulate numerous physiological functions: fatty acid synthesis and metabolism, glucose homeostasis, steroidogenesis, immunity, and neurological homeostasis. LXR-deficiency in mouse results in several phenotypes mimicking pathological conditions in humans. This review will be focused on the various natural and synthetic LXR agonists and antagonists. Putative clinical targets including atherosclerosis, diabetes, Alzheimer's disease, skin disorders, and cancer will be covered.

© 2011 Elsevier Ireland Ltd. All rights reserved.

## Contents

1. Introduction	00
1.1. LXR structure	00
1.2. Mechanism of action	00
2. Activator ligands	00
2.1. Endogenous agonists: oxysterols and other sterols	00
2.2. Ligands from microbial and plants	00
2.3. Synthetic agonists	00
3. Antagonists and inhibitors ligands	00
3.1. Natural antagonists	00
3.2. Synthetic antagonists	00
4. From physiological function to pathology	00
4.1. LXR and cancer	00
4.1.1. LXR and hormone-dependant cancer	00
4.1.2. LXR and colon cancer	00
4.2. LXR and skin disorders	00
4.3. LXR and Alzheimer's disease	00
4.4. LXR and atherosclerosis	00
4.5. LXR and inflammation	00
4.6. LXR and diabetes	00
5. Conclusion	00
Acknowledgments	00
References	00

\* Corresponding author. Address: Génétique Reproduction et Développement, Unité Mixte de Recherche CNRS 6247 Clermont Université, 24 avenue des Landais, BP80026, 63171 Aubière Cedex, France. Tel.: +33 4 73 40 74 16; fax: +33 4 73 40 70 42.

E-mail address: [j-marc.lobaccaro@univ-bpclermont.fr](mailto:j-marc.lobaccaro@univ-bpclermont.fr) (J.-M. Lobaccaro).

## 1. Introduction

Nuclear receptors are molecular regulators of gene transcription and intracellular function. Some of them are ligand-activated proteins that control various biological events varying from cell differentiation and development to lipid metabolism and energy homeostasis. Among the nuclear receptors, liver X receptors (LXR)  $\alpha$  (Willy et al., 1995; NRIH3) and  $\beta$  (Shinar et al., 1994; Song et al., 1994; NRIH2) were discovered in the mid 90's. Although originally discovered as pivotal regulators of cholesterol homeostasis (Peet et al., 1998), the known roles of LXRs continue to grow with the study of many cell types and animal models. Thus far, there is evidence that LXR is involved in the following physiological functions: *de novo* synthesis of cholesterol (Wang et al., 2008a,b), excretion (reviewed in Repa and Mangelsdorf, 2000) and detoxification of bile acids (Barbier et al., 2009) or lipids (Volle et al., 2004), glucose homeostasis (Cha and Repa, 2007), immunity (reviewed in Zeller and Tontonoz, 2006), skin development and homeostasis (Demerjian et al., 2009) and neurological functions (Koldamova and Lefterov, 2007) and reviewed in (Leoni and Caccia, 2011). LXR $\alpha$  was initially described as highly expressed in a restricted subset of tissues known to play an important role in lipid metabolism such as liver, small intestine, kidney, spleen and adipose tissue whereas LXR $\beta$  was found to be expressed more ubiquitously ([www.nur-sa.org](http://www.nur-sa.org)). Since their identification as orphan receptors, oxidized derivatives of cholesterol, known as oxysterols (for a review see Schroepfer, 2000), deorphanized both LXRs (Janowski et al., 1996; Janowski et al., 1999). Because LXR deficiency has been associated with various human-like diseases in mice, these receptors are promising pharmacological targets due to the enormous potential to develop ligands. The present review will thus be focused on the various known LXR ligands and their potential use in human health.


### 1.1. LXR structure

As members of the nuclear receptor superfamily, LXRs are composed of four independent domains (for an extensive review about

LXR structure see (Viennois et al., 2011): (i) an amino-terminal domain (A/B domain) containing an activating function AF1, which permits the recruitment of ligand-independent co-activators; (ii) a DNA-Binding Domain (DBD) characterized by two zinc fingers; (iii) a hinge domain that permits the recruitment of co-repressors in absence of ligands; and (iv) a carboxy-terminal domain, containing a hydrophobic ligand-binding domain (LBD), which is required for dimerization, and a transactivation domain (AF-2), which recruits co-activators.

### 1.2. Mechanism of action

(Fig. 1) LXR $\alpha$  and LXR $\beta$  usually form obligatory heterodimers with retinoid X receptor, the receptor of 9-*cis* retinoid acid (Willy et al., 1995; Willy and Mangelsdorf, 1997). Both also bind to a non-canonical responsive element on the renin promoter to regulate renin transcription (Morello et al., 2005). In this conformation LXR $\alpha$  functions as a cAMP activated factor (Tamura et al., 2000) while LXR $\beta$  seems to be inversely affected by cAMP (Morello et al., 2005). When bound to RXR in absence of ligand, acetylated LXR is located on LXR responsive element (LXRE) in the promoter of the target genes. The consensus LXRE sequence consists of two direct repeats of hexameric nucleotides, AGGTCA, separated by four or one nucleotide(s) (DR4 or DR1). The heterodimer interacts with either a co-repressors, nuclear receptor co-repressor (N-CoR), or silencing mediator for retinoic acid and thyroid hormone receptors (SMRT) (Hu et al., 2003), which block transcription by recruitment of histone deacetylase through the interaction with proteins such as stress activated MAP kinase interacting protein 3A (Sin3A) (Lazar, 2003). When the ligand of one of the two partners binds to the heterodimer, it leads to conformational changes that induce the release of the co-repressors, the recruitment of specific co-activators such as activating signal cointegrator-2 (ASC-2) (Lee et al., 2008) on the AF-2 domain (Svensson et al., 2003), the interaction with histone acetyl transferase leading to the transcription of the target genes (reviewed in Viennois et al., 2011).



**Fig. 1.** Schematic view of LXR modulation of gene transcription. LXR $\alpha$  or LXR $\beta$  usually bind RXR, the receptor for 9-*cis* retinoic acid, on a DR4 or a DR1 element. In absence of ligand RXR/LXR heterodimer is silenced by corepressors. The binding of their respective ligands (9 *cis* RA and/or oxysterols) induces conformational changes that allow corepressor departure and coactivator recruitment. On the renin promoter, both LXR can bind as a monomer on a CNRE. Their transcriptional activity is modulated by cAMP. For more details, see text.

## 2. Activator ligands

Traditional pharmacology states that a ligand can be either classified as agonist (full or partial), antagonist or inverse agonist, though this last type has never been described in either of the LXR. Due to the various diseases LXR $\alpha$  and/or LXR $\beta$  are associated with (see above), these nuclear receptors are promising pharmacological targets. This explains why numerous drug companies have been developing a large variety of natural or synthetic agonists. However, LXR $\alpha$  and LXR $\beta$  could be expressed in the same cells. It is thus important to have molecules that could selectively modulate each isoform. The other important point is the necessity to target this modulation specifically in the affected organ and make the ligand bio-available. Hence, the development of such *selective Liver X modulators* (SLiMs) will be the challenge in the future. This will prevent deleterious side effects. From previous studies, it appears that activating or inhibiting characteristics of LXR-ligands are dependent of the amino acid sequence of each LXR isoform, despite a clear homology. The cellular context should also be taken into account: for example, the sequence of the response elements within the promoter or the intracellular enzymes that could modify ligand activity. In general, SLiMs may apply a similar philosophy as that of Selective Estrogen Receptor Modulators (SERMs) such as raloxifen and tamoxifen.


### 2.1. Endogenous agonists: oxysterols and other sterols

Oxidized or hydroxylated metabolites of cholesterol known as oxysterols (Schroepfer, 2000) have been described to be the *bona fide* ligands for LXRs (Janowski et al., 1996). Two sources of oxysterols should be considered on mammals: (i) endogenous, by enzymatic or chemical synthesis; (ii) exogenous, by nutritional supply. The first screenings of LXR ligand have been performed by double hybrid


experiments using the LXR $\alpha$  LBD associated with the DBD of GAL-4 (Janowski et al., 1996). Oxysterols showing the greatest induction factors were 22(R)-hydroxycholesterol, 20(S)-hydroxycholesterol, 24(S)-hydroxycholesterol, 7 $\alpha$ -hydroxycholesterol and 27-hydroxycholesterol (Fig. 2). Besides, 24-cetocholesterol, 25-hydroxycholesterol and 24(S),25-epoxycholesterol have also been shown to be good LXR $\alpha$  and LXR $\beta$  activators (Lehmann et al., 1997) (Fig. 2). Interestingly, the nature of oxysterols depends on the site of synthesis: 22(R)-hydroxycholesterol in steroidogenic tissues, 24(S)-hydroxycholesterol (or cerebrosterol) in brain and plasma, 24(S),25-epoxycholesterol in liver, and 27-hydroxycholesterol in plasma and macrophages. In 1999, Janowski et al. (1999) synthesized and tested a series of related compounds. Their work revealed that position-specific monooxidation of the sterol side chain was requisite for LXR high-affinity binding and activation. Enhanced binding and activation could also be achieved through the use of 24-oxo ligands that act as hydrogen bond acceptors in the side chain. In addition, introduction of an oxygen on the sterol B-ring results in a ligand with LXR $\alpha$  selectivity. The ultimate demonstration that oxysterols were the true LXR ligands *in vivo* came from Russell's group (Chen et al., 2007): forced expression of cholesterol sulfotransferase, an enzyme that metabolizes oxysterol ligands, led to inactivation of LXR signaling in cell lines and cholesterol-fed mice. Also, genetic elimination of three oxysterol biosynthetic enzymes attenuated the response of some, but not all, LXR target genes in mouse liver.

Other cholesterol-derived molecules, such as follicular fluid meiosis activating sterol (FF-MAS), may also activate LXR $\alpha$  (Ruan et al., 1998). Likewise, desmosterol, a cholesterol precursor produced from zymosterol, could also activate LXR (Yang et al., 2006) (Fig. 3A).


Bile acid pathway-derived molecules have also been proposed as putative LXR ligands. Indeed, even though bile acids are the nuclear receptor FXR ligands (Makishima et al., 1999), cholestenic



**Fig. 2.** Natural agonist ligands, the oxysterols. Cholesterol chemical structure is represented and carbon atoms are numbered. The name of each oxysterol depends on the hydroxyl group position.



**Fig. 3.** Natural agonists, other cholesterol deriving sterols. (A) FF-MAS and demosterol chemical structure, molecules derived from cholesterol biosynthesis pathway, upstream cholesterol. FF-MAS, Follicular Fluid Meiosis Activating Sterol. (B) LXR agonists derived from Bile acids biosynthesis pathway. CDCA, chenodeoxycholic acid. Note that CDCA does not activate LXR.



**Fig. 4.** LXR activators derived from plants (A) and fungi (B).

acid, chenodeoxycholic acid (CDCA) precursor, is able to induce transcriptional activity of LXR $\alpha$  and  $\beta$  on gene reporter assay (Song and Liao, 2000). Likewise, the same group described 6 $\alpha$ -hydroxylated bile acids (notably hyodeoxycholic acid) as selective activators of LXR $\alpha$  (Song et al., 2000) (Fig. 3B).

## 2.2. Ligands from microbial and plants

Over the last decade various plant sterols and stanols, which are structurally related to cholesterol, have been identified as LXR activators since they are the natural occurring equivalent of mammalian cholesterol (Fig. 4). Therefore, sitosterol and sitostanol, derived from the 4-desmethylsterol family, activated both LXR $\alpha$  and LXR $\beta$  and increased expression of ATP Binding Cassette A1, ABCA1 in Caco-2 cells (Plat et al., 2005). Gynosaponin TR1, isolated from the herbal medicine, *Gynostemma pentaphyllum*, an herbaceous vine of the *Cucurbitaceae* family, demonstrated greater selectivity toward activation of the LXR $\alpha$  isoform than LXR $\beta$  in HEK293 cells (Huang et al., 2005). Acanthoic acid from *rollinia*, an exotic tropical fruit, selectively activates LXR $\alpha$  in HEK-293 cells (Jayasuriya et al., 2005) and both isoforms in macrophages lead to cholesterol efflux (Traves et al., 2007). Podocarpic acid is a natural activator derived from plant resins. This molecule gets its name from Podocarpus from which it was extracted. It has been reported that podocarpic acid is able to bind to LXR $\alpha$  and  $\beta$ . In hamster and mice models, it increases plasma HDL with concomitant decrease of plasma LDL (Singh et al., 2005). More recently, honokiol, extracted from the bark of *Magnolia abovata*, a native Japanese Magniola, has been showed to be an activator of the LXR/RXR heterodimer and its ABCA1 target gene in peritoneal macrophages, excluding foam cells (Kotani et al., 2010). However, it is not known whether honokiol directly binds RXR and/or LXR $\alpha$ . Besides plants, some fungal derivatives may also activate LXR (Fig. 4), such as Paxillin from *Penicillium paxilli* (Bramlett et al., 2003), or ergostan-4,6,8,22-tetraen-3-one, a ergostane derviate from *Tolypocladium niveum*, a Norwegian soil fungus (Ondeyka et al., 2005).

## 2.3. Synthetic agonists

The discovery of the hypocholesterolemia potency of LXR at the end of the 90's (Peet et al., 1998) motivated many pharmacological companies to develop LXR ligands (for an exhaustive review see Li et al., 2010). Among them, a non steroidal compound, T0901317 (Fig. 5) is the most commonly used synthetic ligand in fundamental research. However, this compound is not completely selective for LXR since it is able to activate PXR or FXR at high concentrations (Houck et al., 2004; Mitro et al., 2007; Shenoy et al., 2004). T0901317 has also been found to be a very effective ROR $\alpha$  and ROR $\gamma$  inhibitor (Kumar et al., 2010). A screening of GlaxoSmithKline (GSK) compounds has permitted the identification of the GW3965 (Fig. 5) as a strong agonist of LXR $\alpha$  and LXR $\beta$ . In contrast with T0901317, GW3965 is more selective for LXR (Mitro et al., 2007). Unfortunately, human therapeutic use of these molecules is impossible because of their temporary hypertriglyceridemic effects (Joseph et al., 2002a,b). More recently, GSK has also developed an intestinal-specific LXR agonist closely related to GW3965, GW6340, which specifically activates LXR target genes (ABCA1, G5 and G8) in small intestine without modification of hepatic target genes expression or hepatic triglyceride content (Yasuda et al., 2010).

A promising synthetic ligand was also identified by Makishima's group: (22E)-ergost-22-ene-1 $\alpha$ ,2 $\beta$  diol (YT-32) (Fig. 5), derived from ergosterol, shows strong LXR agonist activity (Kaneko et al., 2003). In mice, the oral administration of YT-32 leads to an intestinal accumulation of the ABCA1, ABCG5 and ABCG8 encoding genes, and a reduction of plasma cholesterol without any modification of expression of these genes in liver. This ligand is the first tissue-specific LXR agonist. To explain this specificity, Kaneko et al. suggested that YT-32 may reduce the intestinal absorption of cholesterol, by inducing expression of ABCG5 and G8, which could excrete the compound from the apical membranes of mucosal cells. A second hypothesis was that YT-32 is secreted into bile by ABCG5 and G8 before it has a chance to induce expression of lipogenic




Fig. 5. Synthetic LXR agonists.



genes in liver. A third explanation could be a tissue-specific agonist activity depending on co-factor recruitment as observed for selective estrogen receptor modulators.

In the same way as YT-32, DMHCA (N,N-Dimethyl-3 $\beta$ -hydroxy-cholenamide) (Fig. 5) is an LXR agonist that is able to stimulate cholesterol transport through upregulation of LXR target genes, including ABCA1, in liver, small intestine, and peritoneal macrophages (Quinet et al., 2004). DMHCA's binding affinity to both LXR isoforms was calculated by scintillation proximity assay at 100–200 nM (Janowski et al., 1999). Compared with known nonsteroidal LXR agonists, DMHCA exhibits only limited activity to increase hepatic Sterol-response element binding protein 1c (srebp-1c) and Fatty acid synthase (fas) mRNA and does not alter circulating plasma triglycerides (Quinet et al., 2004). A recent study shows that DMHCA significantly reduces cholesterol absorption and uptake in the duodenum and jejunum of the small intestine and, in turn, leads to a reduction of plasma cholesterol. Interestingly, this study reveals that DMHCA also suppresses endogenous cholesterol biosynthesis (Pfeifer et al., 2011).

The first LXR agonist tested in human in a phase I clinical trial was a Wyeth's compound, the LXR-623 (Fig. 5): an orally bioavailable indazole highly selective for LXR (DiBlasio-Smith et al., 2008; Katz et al., 2009). After administration of LXR-623 to healthy volunteers, LXR target genes involved in the reverse cholesterol transport were upregulated in the macrophages without any induction of triglyceride synthesis. Unfortunately, adverse effects were observed on the central nervous system, ending the trial. Pharmaceutical companies have focused their efforts on the identification of molecules specific of each LXR isoform. Wyeth group developed a phenylsulfone-substituted quinoxaline, WYE-672 (Fig. 5) as an oral and tissue active LXR $\beta$  isoform selective agonist (Hu et al., 2010). Recently, a novel synthetic steroidal LXR agonist, ATI-111 compound, has been described to significantly decrease cholesterol and triglyceride levels in plasma in LDLR $^{-/-}$  mice. This observation is correlated to an increased expression of ABCA1, ABCG1 and ABCG5/G8 in intestine and macrophages without modification of acetyl-coenzyme A carboxylase (ACC) and FAS expressions in liver and intestine (Peng et al., 2011).

### 3. Antagonists and inhibitors ligands

Given the hypocholesterolemia potency of LXR, pharmaceutical companies have focused their effort on the discovery of LXR activators rather than LXR antagonists. Thus, very few natural or synthetic antagonist compounds have been identified or developed. However, use of LXR-isoform antagonists could help in further deciphering the physiological roles of LXR $\alpha$  and LXR $\beta$ .

#### 3.1. Natural antagonists

One of the most potent endogenous LXR $\alpha$  antagonists to date is 5 $\alpha$ ,6 $\alpha$ -epoxycholesterol. This oxysterol, present in high quantities in food, has been detected in atherosclerotic plaques. This compound induces the recruitment of cofactor peptides to both LXR $\alpha$  and LXR $\beta$ . Radiolabeled ligand displacement assay demonstrated that 5 $\alpha$ ,6 $\alpha$ -epoxycholesterol binds to LXR $\alpha$ . This molecule is able to antagonize LXR-mediated gene expression in different cell culture systems and notably in keratinocytes (Berrodin et al., 2010). Initially, it was only the sulfated form of this compound, 5 $\alpha$ ,6 $\alpha$ -epoxycholesterol-3-sulfate which was described to antagonize LXR transactivation on a LXR reporter gene assay (Song et al., 2001). It has been demonstrated to disturb the recruitment of certain co-activators and to bind directly to LXR-LBD. In the same study the authors have described another sulfated oxysterol, 7-

cetocholesterol-3-sulfate, as a LXR antagonist (Song et al., 2001) (Fig. 6).

Polyunsaturated fatty acids (PUFAs) are known to be PPAR $\alpha$  and PPAR $\delta$  agonists. Several PUFAs, especially arachidonic acid (Fig. 6), have a LXR-LBD-dependent antagonist effect on LXR $\alpha$  (Ou et al., 2001) and LXR $\beta$  (Yoshikawa et al., 2002). The physiological role of such antagonist activity has been found associated with the regulation of hepatic fatty acid elongase 5 (Qin et al., 2009). Indeed, PUFAs may feedback inhibit their own synthesis through repression of LXR $\alpha$  and blocking of *srebp1c* transcription, a master gene involved in fatty acid synthesis.

Herath et al. (2005) revealed, using a scintillation binding assay, that a new polyisoprenylated benzophenone named guttiferone I extract from *Garcinia humilis*, a small fruiting tree growing in Bolivia, is able to selectively bind LXR $\alpha$  (Herath et al., 2005) (Fig. 6). However, this compound is unable to provoke the recruitment of co-activators.

Riccardin is a compound deriving from the liverwort *Blasia pusilla*. Riccardin C is a natural antagonist of LXR $\beta$  and riccardin F is a natural antagonist of both LXR isoforms. Both compete with the endogenous or synthetic LXR ligand for LXR activation (Tamehiro et al., 2005) (Fig. 6).

A recent study highlights the LXR $\alpha$  antagonizing effect of naringenin (Fig. 6), a flavanone found in grapefruit, oranges and tomatoes (Goldwasser et al., 2010). Naringenin activates the ligand-binding domain of both PPAR $\alpha$  and PPAR $\delta$  while inhibiting LXR $\alpha$  in GAL4-fusion reporters. It inhibits LXR $\alpha$  association with the co-activator TRAP-222 in presence of T0901317. It also leads to an inhibition of LXR $\alpha$  target genes such as *fas*, *abca1*, *abcg1*, resulting in an increase of fatty acid oxidation in primary rat hepatocytes and a decrease of bile acid synthesis.

Whole leaf extract of *Parthenocissus tricuspidata* Virginia creeper from Japan, and *Euscaphis japonica*, a small Japanese tree, suppressed the transcriptional activity of LXR $\alpha$  as well as the expression of LXR $\alpha$  target genes (Kim et al., 2006). Furthermore, both extracts exerted repressive effects on adipocyte differentiation and on lipid metabolism *in vitro* without identification of the active compound.

#### 3.2. Synthetic antagonists


The 22(S)-hydroxycholesterol is able to interfere with the binding of other oxysterols (Janowski et al., 1999) to LXR-LBD and to inhibit the lipogenic activity of T0901317 (Lehmann et al., 1997) (Fig. 6).

Motoshima et al. (Motoshima et al., 2009) focused on the creation of LXR antagonists using a multi-template approach based on thalidomide. They found that several thalidomide-related phthalimide derivatives, including PP2P (Fig. 6), possess LXR-antagonistic activity (Noguchi-Yachide et al., 2007). This is based on the agonist property of glucose on LXR. This group develops other phenethyl-phenyl phthalimide derivatives (Motoshima et al., 2009). Likewise, GSK group identified GSK2033, a tertiary sulfonamide, as the first potent cell-active LXR antagonist described to date (Zuercher et al., 2010) (Fig. 6).

Finally, fibrates have been identified as PPAR $\alpha$  ligand leading to inhibition of circulating triglycerides. Esterified-fibrates such as fenofibrate (Fig. 6) block LXR $\alpha$  activity by occupation of the ligand-binding pocket without induction of co-factor recruitment (Thomas et al., 2003).

### 4. From physiological function to pathology

LXR functions have been schematically associated with several various physiological fields from reproduction to energy



**Fig. 6.** Natural (A) or synthetic (B) LXR antagonists. (A) The R1 group is OH for 5α,6α-epoxycholesterol and H<sub>2</sub>SO<sub>4</sub> for the 5α,6α-epoxycholesterol-3-sulfate, the compound initially described as LXR antagonist, R2 is OCH<sub>3</sub> for Riccardin F and OH for Riccardin C.


homeostasis, neuron physiology, immunity, skin development and apoptosis. Implication of LXR on these processes has been well documented (Cao et al., 2003; Cummins et al., 2006; El-Hajjaji et al., 2011; Joseph et al., 2004; Laffitte et al., 2003; Repa and Mangelsdorf, 2000; Schultz et al., 2000; Viennois et al., 2011; Volle and Lobaccaro, 2007). LXR<sup>−/−</sup> mouse analysis also pointed out that the lack of one or both isoform could be associated to the development of symptoms linked to human disease (Fig. 7). Hence, LXR are promising pharmacological targets.

#### 4.1. LXR and cancer

##### 4.1.1. LXR and hormone-dependant cancer

Breast and prostate cancers are the leading causes of cancer in women and men, respectively. They share many characteristics (Coffey, 2001). Incidence rates for both prostate and breast cancer are less than 10% compared to those observed in Occidental countries. As people migrate from Asia to the United States, the low rates observed in Asia begin to rise with time and subsequent generations toward those observed commonly in the general American





**Fig. 7.** Schematic representation of the diseases potentially targets of LXR ligands. Black circle, cancer.

population. Furthermore, as Asian countries adopt Western-style diets, their incidencies of prostate and breast cancer rise. In all of these situations, there is a close correlation between responses in the prostate and breast cancer rates. Moreover, the emergence of resistance is common, especially for locally advanced tumors and metastatic tumors (Rau et al., 2005).

**4.1.1.1. LXR $\alpha$  and prostate cancer.** As previously stated, prostate cancer is the most frequently diagnosed cancer and the second leading cause of cancer death in men (Gronberg, 2003). Among the various genetic and environmental risk factors, epidemiological analyses revealed a clear positive association between hypercholesterolemia and the development of prostate cancer (Bravi et al., 2006; Magura et al., 2008). Indeed, epidemiological studies have shown that Chinese populations, with initially low risk of prostate cancer, have an increased risk after migration to the United States. This environmental effect was mainly attributed to the deleterious impact of lipid consumption (Watanabe et al., 2000). Consistent with this hypothesis, a clinical study revealed that hypercholesterolemia increases prostate cancer risk while a statin treatment, inhibiting *de novo* cholesterol biosynthesis, is associated with lower risk of metastatic prostate cancer (Platz et al., 2006; Shannon et al., 2005). Even though LXR are key regulators of cholesterol homeostasis, their roles in prostate pathophysiology remain little known. Fukushima's group first reports the control of proliferation by LXR on LNCaP human prostate carcinoma cell line (Fukuchi

et al., 2004a,b). In their experiments, T0901317 decreases the number of cells in S-phase and induces the expression of ABCA1 (Fukuchi et al., 2004a,b). *In vitro* and *in vivo* analyses also revealed that modulation of LXR activity also triggers apoptosis in prostate cancer cells. This effect involves both increased cholesterol efflux by ABCA1 and the disruption of lipid-raft signaling activity (Pommier et al., 2010). Androgens also play a crucial role on prostate cancer development. Androgen ablation is also a common therapy for this type of cancer. Recently, androgens have been demonstrated to crosstalk with LXR in prostate cancer cell line LNCaP (Krycer and Brown, 2011). Hence LXR may also modulate prostate cancer development by decreasing the androgenic pathway.

**4.1.1.2. LXR and breast cancer.** Several studies have described LXR activity on modulation of breast cancer cell line proliferation and apoptosis. In several estrogen dependent breast cancer cell lines, Vedin et al. demonstrated that T0901317 treatment decreases the number of proliferating cells independently of their lipogenic activity (Vedin et al., 2009). In estrogen independent cell lines, this effect is abolished suggesting a crucial role of estrogen on this LXR effect. *In vivo*, LXR activation increases estrogen catabolism leading to estrogen deprivation. In this way, LXR inhibits estrogen-dependent breast cancer growth (Gong et al., 2007). Among the other female cancers, studies enlighten the anti-proliferative effects of LXR-ligand on ovarian cancer models (Rough et al., 2010).

#### 4.1.2. LXR and colon cancer

Several studies have described a correlation between colon cancer and LXR modulation. Uno et al. pointed that LXR ligand activation decreases the expression of several endogenous  $\beta$ -catenin target genes such as myc, bmp4 and mmp7 in colon cancer cells. This correlates with a reduction of cell proliferation without modification of apoptosis (Uno et al., 2009). In a recent review, Chuu (2011) hypothesized that LXR could be a biomarker or a therapeutic target in colon cancers. Phytosterol-rich diets were reported to decrease incidence of several cancer and  $\beta$ -sitosterol were shown to suppress proliferation of HT-29 colon cancer cells. Since  $\beta$ -sitosterol is an efficient LXR agonist, phytosterols may possibly suppress tumor growth partially through activation of LXR signaling (Chuu, 2011). A case-control study that associates soy (containing phytosterols) consumption and colon cancer onset supports this hypothesis (Spector et al., 2003).

#### 4.2. LXR and skin disorders

Both LXR $\alpha$  and LXR $\beta$  are expressed in human keratinocytes and fetal rat epidermis and LXR $\beta$  is predominantly expressed in mouse epidermis (Hanley et al., 2000; Komuves et al., 2002). Studies on LXR $^{-/-}$  mice pointed out that epidermal differentiation is regulated by LXR $\beta$  and that oxysterols can induce differentiation and inhibit proliferation through LXR $\beta$  activation (Komuves et al., 2002). Cholesterol homeostasis is essential for cutaneous barrier function. A study demonstrated that an increase in cholesterol stimulates both ABCA1 (Jiang et al., 2006) and ABCG1 (Jiang et al., 2010) expression, whereas inhibition of cholesterol synthesis by statins decreases ABCA1 and ABCG1 expression in cultured human keratinocytes. After acute permeability disruption, ABCA1 expression decreases keratinocyte cholesterol efflux to make more cholesterol available for regeneration of the barrier (Jiang et al., 2006). Topical treatment of murine skin with LXR activators also increases ABCG1 expression in murine epidermis (Jiang et al., 2010). In case of skin-aging resulting from photo-aging and normal chronological skin aging, an *in vivo* study showed that LXR agonists reverse aging associated syndrome in a photo-aging animal model. Moreover, LXR $^{-/-}$  mice could represent a great model of chronologically aged human skin (Chang et al., 2008). Furthermore, LXR

ligands reverse clinical symptoms of atopic dermatitis, an inflammatory, chronically relapsing and non-contagious skin disorder (Chang et al., 2008; Hatano et al., 2010).

Interestingly, Sticozzi et al. made a connection between cigarette smoke and LXR activity on the skin (Sticozzi et al., 2010). They showed an increase of ABCA1 expression on human keratinocytes, HaCaT cell line, after cigarette smoke exposure. This is the first study describing the negative potential effect of cigarette smoke on cholesterol skin content (Sticozzi et al., 2010). Finally, locally administered LXR agonists may be potentially useful in the treatment of skin conditions such as dermatitis or aging.

#### 4.3. LXR and Alzheimer's disease

The link between cholesterol and Alzheimer's disease (AD), an age-related dementia disorder, has been well established by two groups after they treated their patients with statins (Jick et al., 2000; Wolozin et al., 2000). Koldamova et al. evaluated the effect of the synthetic LXR ligand T0901317 on ABCA1 expression in the brain of APP transgenic mouse model of AD. T0901317 significantly enhanced ABCA1 expression and downregulated the production of A $\beta$  peptide, whose deposit leads to neuronal degeneration, suggesting that ABCA1 plays a protective role in AD progression (Koldamova et al., 2005). Likewise, the loss of *lrx $\alpha$*  or *lrx $\beta$*  in APP23 transgenic mice increases A $\beta$  deposit (Zelcer et al., 2007) while activation of LXR by T0901317 causes a significant reduction of memory deficits observed in APP23 model fed a high fat diet (Fitz et al., 2010). Recently, it has been demonstrated that beneficial effects, like memory recovery, of LXR agonists in an APP/presenilin1 (whose mutations are the most common cause of familial early-onset Alzheimer's disease) model of amyloidogenesis, is mediated by ABCA1 (Donkin et al., 2010). Zelcer et al. (2007) showed that LXR activation decreases the inflammatory response likely involved in AD. Likewise, seladin-1/DHCR24, a selective AD marker, is a LXR target gene (Wang et al., 2008a,b). At last, T0901317 significantly decreases amyloid pathology caused by a high-cholesterol diet and improves cognitive performance (Fitz et al., 2010).

Altogether, these data suggest that LXR may be promising pharmacological targets for the treatment of AD and other neurological disorders involving the disruption of cholesterol homeostasis. It is worthwhile to note that two single-nucleotide polymorphisms in LXR $\beta$  have been associated with AD (Adighibe et al., 2006; Infante et al., 2010), further suggesting that LXR is involved in this neurodegenerative disease in human. Leoni and Caccia (Leoni and Caccia, 2011) showed that 24(S)-hydroxycholesterol, a LXR-ligand (see above) is reduced in plasma and in cerebrospinal fluid of patients with neurodegenerative disease, while 27-hydroxycholesterol is increased. This study highlights the potential use of LXR endogenous ligand not only as a pharmacological compound to treat symptoms, but also as a biomarker of certain diseases.

#### 4.4. LXR and atherosclerosis

The LXR signaling pathway clearly displays anti-atherogenic properties, both by reducing cholesterol levels and by anti-inflammatory properties. Studies in various models of atherosclerosis such as LDLR $-/-$  or APOE $-/-$  mice have now clearly established that treatment with LXR agonist leads to a regression of atherosclerosis *in vivo*. Joseph et al. initially demonstrated that GW3965 treatment inhibited atherosclerotic lesions in APOE $-/-$  or in LDLR $-/-$  mice fed a high-cholesterol diet (Joseph et al., 2002a,b). LXR is also involved in the stimulation of macrophages transcription of ABC transporters resulting in release of cholesterol from the cells. This prevents the transformation of macrophages into foam cells in response to lipid loading. Moschetta's group (Lo Sasso et al., 2010) demonstrated that genetic intestinal-specific LXR activation

led to decreased intestinal cholesterol absorption, improved lipoprotein profile, and increased reverse cholesterol transport *in vivo* in the absence of hepatic steatosis thus protecting LDLR $-/-$  mice from atherosclerosis. Lo Sasso et al. identified the intestines as a key player in the LXR-driven protective environment against cardiovascular disease (Lo Sasso et al., 2010). The athero-protective role of LXR may also be mediated by the inhibition of platelet function and thrombus formation (Spyridon et al., 2011): LXR $\beta$  is present in human platelets and its ligand activation inhibits platelet aggregation, suggesting the anti-thrombotic role LXR $\beta$ .

Using a pharmacological approach with GW6340, Yasuda et al. (2010) showed that intestinal-specific LXR activation promotes macrophage reverse cholesterol transport. ABCA1, ABCG5 and ABCG8 were significantly upregulated in the small intestine, but not in the liver. Two independent mechanisms could account for these observations: increased intestinal HDL production and promotion of intestinal excretion of HDL-derived cholesterol (Yasuda et al., 2010). These data demonstrate that activation of LXR in intestine and macrophages efficiently prevents atherosclerosis. Likewise, and in contrast with T0901317, LXR-623 treatment is not associated with increased hepatic lipogenesis. In non-human primates with normal lipid levels, LXR-623 significantly reduces total and LDL-cholesterol in a time and dose-dependent manner. LXR-623 is also associated with increased expression of the target genes ABCA1/G1 in peripheral blood cells (Quinet et al., 2009). These observations suggest that LXR-623 is a promising pharmacological ligand in atherosclerosis. However, as discussed above, the first clinical trial with this synthetic LXR ligand showed adverse effects (Katz et al., 2009). In the rabbit model of atherosclerosis, LXR-623 and simvastatin, which inhibits endogenous synthesis of cholesterol, reduce the progression of atherosclerosis and induce plaque regression (Giannarelli et al., 2011). A novel LXR therapeutic approach could involve combination therapy for synergistic anti-atherotic effects.

Stein and Matter recently compiled data concerning the protective role of SIRT1 in atherosclerosis (Stein and Matter, 2011). SIRT1 directly deacetylates LXR $\alpha$ , regulating its transcriptional activity. In SIRT1 $-/-$  macrophages, cholesterol efflux is lower than in SIRT1 $+/+$  macrophages (Li et al., 2007). This study highlights the potential use of activators or inhibitors of LXR co-factors as a new way to modulate LXR.

Human genetics also illuminate LXR as putative pharmacological targets. Analysis of LXR gene sequences in patients with coronary heart disease identified three mutations in the LBD of LXR $\alpha$  (Dave et al., 2009). These mutations create conformational changes that may prevent activation of LXR $\alpha$  by its natural ligands.

#### 4.5. LXR and inflammation

Besides the anti-inflammatory impact of LXR in Alzheimer's disease, dermatitis or atherosclerosis, LXR has also a suppressive role in numerous inflammatory diseases such as, stroke, lung inflammation or rheumatoid arthritis.

Stroke or cerebrovascular accident causes irreversible neuronal injury. It can be due to ischemia (caused by thrombosis or arterial embolism) or hemorrhages. Recently, anti-inflammatory agents have shown to restraint inflammation during a stroke, enhancing neurogenesis and tissue repair. Previously in this review, we discussed about the anti-inflammatory role of LXR and their functional amelioration in Alzheimer's disease. Interestingly, Sironi et al. demonstrated that LXR activation with a single dose of GW3965 blocks ischemia-induced brain damage two hours after the induction of ischemia in rat model (Sironi et al., 2008). In another study, it has been shown that GW3965 or T0901317 improves stroke outcome in rats after ischemic occlusion. This

neuroprotection is correlated with a decrease of proinflammatory factors in brain, like *cox-2*, *iNos* or *NF-kB* (Morales et al., 2008).

Using a pre-clinical rat model of endotoxin-induced lung inflammation it has also been shown that LXR activation leads to a suppression of lung inflammatory response (Birrell et al., 2007). In a mouse model of carrageenan-induced pleurisy, LXR activation induces a significant reduction of numerous parameters of inflammation such as cytokine, *iNOS* formation in lung, lipid peroxidation and apoptosis (Crisafulli et al., 2010).

GW3965 prophylaxis significantly reduced arthritis incidence and attenuated the clinical and histological severity in collagen-induced arthritis mice. Moreover, GW3965 prophylaxis also significantly attenuated inflammatory mediator production in joint sections and serum pro-inflammatory cytokine levels in a dose-dependent manner (Park et al., 2010).

It can be noted that oxysterols may act as EBI2 (*Epstein-Barr virus-induced gene 2*, an orphan G-protein-coupled receptor) ligands and have an important role on adaptative immune response. By this way, oxysterols act as chemoattractant for immune cells, directing cell migration (Hannedouche et al., 2011; Liu et al., 2011).

Altogether, LXR agonists play various anti-inflammatory roles on different tissues like brain, lung immune cells. One way by which LXR exert its anti-inflammatory effect is by SUMOylation. Especially, LXR ligands inhibited STAT1-mediated inflammatory responses in IFN $\gamma$ -stimulated brain astrocytes. Ligand-dependent SUMOylations of LXRs are required for inhibitory action on STAT1, as STAT1 complexed with SUMOylated LXR become unable to bind to promoter regions of target genes, resulting in transcriptional failure (Lee et al., 2009 #136).

#### 4.6. LXR and diabetes

Diabetes mellitus type 2 is a metabolic disorder whose main risk factor is obesity. It is characterized by high blood glucose in the context of insulin resistance (Misra and Khurana, 2008). Treatment of diabetic rodents or mouse models of diet-induced obesity with LXR agonists improves glucose tolerance (Cao et al., 2003; Lafitte et al., 2003). However, *in vitro* experiments showed that the lipogenic effect of LXR agonists in the skeletal muscle of patients with type 2 diabetes is increased compared to healthy control patients (Kase et al., 2005). This study shows that LXR activation may promote triglyceride accumulation in the presence of high glucose concentration in skeletal muscle cells probably via induction lipogenic enzymes expression (Kase et al., 2005). This may prove to be problematic effect of LXR activation in muscle. Genetic linkage analyses suggest role of LXR in obesity (Dahlman et al., 2006) or pancreatic  $\beta$ -cell dysfunction (Ketterer et al., 2011).

## 5. Conclusion

This review enlightened the potential applications of LXR ligands as therapeutic agents in various pathologies. Thus far, few studies have addressed the effect of such agonists in human because of their adverse effects on lipogenesis. The first clinical trial in human was the use of LXR-623 against atherosclerosis in healthy patients. This compound does not activate the lipogenic enzymes in liver and acts specifically in macrophages. Unfortunately, this trial was rapidly aborted because of the adverse side effects of this compound on the central nervous system. Nonetheless, the way has been opened for the use of LXR agonists in human. The goal is to be able to activate LXR specifically in the target tissue. It is precisely the concept of develop new SLiMs that could activate or inhibit LXR $\alpha$  or LXR $\beta$  depending on the tissue. Future studies will be necessary to determine whether manipulations of these

pathways have utility in the treatment of the human diseases described above.

## Acknowledgments

We thank David H. Jiang, University of Virginia School of Medicine, Charlottesville, VA, for critically reading the manuscript.

This study was supported by Association de Recherche sur les Tumeurs Prostatiques, Ligue Allier contre le Cancer, Association de Recherches contre le Cancer, FRM and Fondation BNP-Paribas Research Grants, Contrat Nouveau Chercheur Auvergne (to S.B.). E. Viennois is funded by Région Auvergne and Fond Européen de Développement Régional (FEDER); J. Dufour is funded by MNERT.

## References

- Adighibe, O., Arepalli, S., Duckworth, J., Hardy, J., Wavrant-De Vrieze, F., 2006. Genetic variability at the LXR gene (NR1H2) may contribute to the risk of Alzheimer's disease. *Neurobiol. Aging* 27, 1431–1434.
- Barbier, O., Trottier, J., Kaeding, J., Caron, P., Verreault, M., 2009. Lipid-activated transcription factors control bile acid glucuronidation. *Mol. Cell. Biochem.* 326, 3–8.
- Berrodin, T.J., Shen, Q., Quinet, E.M., Yudit, M.R., Freedman, L.P., Nagpal, S., 2010. Identification of Salpha, 6alpha-epoxycholesterol as a novel modulator of liver X receptor activity. *Mol. Pharmacol.* 78, 1046–1058.
- Birrell, M.A., Catley, M.C., Hardaker, E., Wong, S., Willson, T.M., McCluskie, K., Leonard, T., Farrow, S.N., Collins, J.L., Haj-Yahia, S., Belvisi, M.G., 2007. Novel role for the liver X nuclear receptor in the suppression of lung inflammatory responses. *J. Biol. Chem.* 282, 31882–31890.
- Bramlett, K.S., Houck, K.A., Borchert, K.M., Dowless, M.S., Kulanthai, P., Zhang, Y., Beyer, T.P., Schmidt, R., Thomas, J.S., Michael, L.F., Barr, R., Montrose, C., Eacho, P.I., Cao, G., Burris, T.P., 2003. A natural product ligand of the oxysterol receptor, liver X receptor. *J. Pharmacol. Exp. Ther.* 307, 291–296.
- Bravi, F., Scotti, L., Bosetti, C., Talamini, R., Negri, E., Montella, M., Franceschi, S., La Vecchia, C., 2006. Self-reported history of hypercholesterolaemia and gallstones and the risk of prostate cancer. *Ann. Oncol.* 17, 1014–1017.
- Cao, G., Liang, Y., Broderick, C.L., Oldham, B.A., Beyer, T.P., Schmidt, R.J., Zhang, Y., Staybrook, K.R., Suen, C., Otto, K.A., Miller, A.R., Dai, J., Foxworthy, P., Gao, H., Ryan, T.P., Jiang, X.C., Burris, T.P., Eacho, P.I., Etgen, G.J., 2003. Antidiabetic action of a liver X receptor agonist mediated by inhibition of hepatic gluconeogenesis. *J. Biol. Chem.* 278, 1131–1136.
- Cha, J.Y., Repa, J.J., 2007. The liver X receptor (LXR) and hepatic lipogenesis. The carbohydrate-response element-binding protein is a target gene of LXR. *J. Biol. Chem.* 282, 743–751.
- Chang, K.C., Shen, Q., Oh, I.G., Jelinsky, S.A., Jenkins, S.F., Wang, W., Wang, Y., LaCava, M., Yudit, M.R., Thompson, C.C., Freedman, L.P., Chung, J.H., Nagpal, S., 2008. Liver X receptor is a therapeutic target for photoaging and chronological skin aging. *Mol. Endocrinol.* 22, 2407–2419.
- Chen, W., Chen, G., Head, D.L., Mangelsdorf, D.J., Russell, D.W., 2007. Enzymatic reduction of oxysterols impairs LXR signaling in cultured cells and the livers of mice. *Cell Metab.* 5, 73–79.
- Chuu, C.P., 2011. Modulation of liver X receptor signaling as a prevention and therapy for colon cancer. *Med. Hypotheses* 76, 697–699.
- Coffey, D.S., 2001. Similarities of prostate and breast cancer: evolution, diet, and estrogens. *Urology* 57, 31–38.
- Crisafulli, C., Mazzon, E., Paterniti, I., Galuppo, M., Bramanti, P., Cuzzocrea, S., 2010. Effects of liver X receptor agonist treatment on signal transduction pathways in acute lung inflammation. *Respir. Res.* 11, 19.
- Cummins, C.L., Volle, D.H., Zhang, Y., McDonald, J.G., Sion, B., Lefrançois-Martinez, A.M., Cairra, F., Veyssiere, G., Mangelsdorf, D.J., Lobaccaro, J.M., 2006. Liver X receptors regulate adrenal cholesterol balance. *J. Clin. Invest.* 116, 1902–1912.
- Dahlman, I., Nilsson, M., Jiao, H., Hoffstedt, J., Lindgren, C.M., Humphreys, K., Kere, J., Gustafsson, J.A., Arner, P., Dahlman-Wright, K., 2006. Liver X receptor gene polymorphisms and adipose tissue expression levels in obesity. *Pharmacogenet. Genomics* 16, 881–889.
- Dave, V.P., Kaul, D., Sharma, Y., Bhattacharya, R., 2009. Functional genomics of blood cellular LXR-alpha gene in human coronary heart disease. *J. Mol. Cell. Cardiol.* 46, 536–544.
- Demerjian, M., Choi, E.H., Man, M.Q., Chang, S., Elias, P.M., Feingold, K.R., 2009. Activators of PPARs and LXR decrease the adverse effects of exogenous glucocorticoids on the epidermis. *Exp. Dermatol.* 18, 643–649.
- DiBlasio-Smith, E.A., Arai, M., Quinet, E.M., Evans, M.J., Kornaga, T., Basso, M.D., Chen, L., Feingold, I., Halpern, A.R., Liu, Q.Y., Nambi, P., Savio, D., Wang, S., Mounts, W.M., Isler, J.A., Slager, A.M., Burczynski, M.E., Dorner, A.J., LaVallie, E.R., 2008. Discovery and implementation of transcriptional biomarkers of synthetic LXR agonists in peripheral blood cells. *J. Transl. Med.* 6, 59.
- Donkin, J.J., Stukas, S., Hirsch-Reinshagen, V., Namjoshi, D., Wilkinson, A., May, S., Chan, J., Fan, J., Collins, J., Wellington, C.L., 2010. ATP-binding cassette transporter A1 mediates the beneficial effects of the liver X receptor agonist GW3965 on object recognition memory and amyloid burden in amyloid precursor protein/presenilin 1 mice. *J. Biol. Chem.* 285, 34144–34154.



- El-Hajjaji, F.Z., Oumeddour, A., Pommier, A.J., Ouvrier, A., Viennois, E., Dufour, J., Cairra, F., Drevet, J.R., Volle, D.H., Baron, S., Saez, F., Lobaccaro, J.M., 2011. Liver X receptors, lipids and their reproductive secrets in the male. *Biochim. Biophys. Acta* 1812, 974–981.
- Fitz, N.F., Cronican, A., Pham, T., Fogg, A., Fauq, A.H., Chapman, R., Lefterov, I., Koldamova, R., 2010. Liver X receptor agonist treatment ameliorates amyloid pathology and memory deficits caused by high-fat diet in APP23 mice. *J. Neurosci.* 30, 6862–6872.
- Fukuchi, J., Hiipakka, R.A., Kokontis, J.M., Hsu, S., Ko, A.L., Fitzgerald, M.L., Liao, S., 2004a. Androgenic suppression of ATP-binding cassette transporter A1 expression in LNCaP human prostate cancer cells. *Cancer Res.* 64, 7682–7685.
- Fukuchi, J., Kokontis, J.M., Hiipakka, R.A., Chuu, C.P., Liao, S., 2004b. Antiproliferative effect of liver X receptor agonists on LNCaP human prostate cancer cells. *Cancer Res.* 64, 7686–7689.
- Giannarelli, C., Cimmino, G., Connolly, T.M., Ibanez, B., Garcia Ruiz, J.M., Alique, M., Zafar, M.U., Fuster, V., Feuerstein, G., Badimon, J.J., 2011. Synergistic effect of liver X receptor activation and simvastatin on plaque regression and stabilization: an magnetic resonance imaging study in a model of advanced atherosclerosis. *Eur Heart J.* [Epub ahead of print].
- Goldwasser, J., Cohen, P.Y., Yang, E., Balaguer, P., Yarmush, M.L., Nahmias, Y., 2010. Transcriptional regulation of human and rat hepatic lipid metabolism by the grapefruit flavonoid naringenin: role of PPARalpha, PPARgamma and LXRalpha. *PLoS One* 5, e12399.
- Gong, H., Guo, P., Zhai, Y., Zhou, J., Uppal, H., Jarzynka, M.J., Song, W.C., Cheng, S.Y., Xie, W., 2007. Estrogen deprivation and inhibition of breast cancer growth in vivo through activation of the orphan nuclear receptor liver X receptor. *Mol. Endocrinol.* 21, 1781–1790.
- Gronberg, H., 2003. Prostate cancer epidemiology. *Lancet* 361, 859–864.
- Hanley, K., Ng, D.C., He, S.S., Lau, P., Min, K., Elias, P.M., Bikle, D.D., Mangelsdorf, D.J., Williams, M.L., Feingold, K.R., 2000. Oxysterols induce differentiation in human keratinocytes and increase Ap-1-dependent involucrin transcription. *J. Invest. Dermatol.* 114, 545–553.
- Hannedouche, S., Zhang, J., Yi, T., Shen, W., Nguyen, D., Pereira, J.P., Guerini, D., Baumgarten, B.U., Roggo, S., Wen, B., Knochenmuss, R., Noel, S., Gessier, F., Kelly, L.M., Vanek, M., Laurent, S., Preuss, I., Miault, C., Christen, I., Karuna, R., Li, W., Koo, D.I., Suply, T., Schmedt, C., Peters, E.C., Falchetto, R., Katopodis, A., Spanka, C., Roy, M.O., Dethieux, M., Chen, Y.A., Schultz, P.G., Cho, C.Y., Seuwen, K., Cyster, J.G., Sailer, A.W., 2011. Oxysterols direct immune cell migration via EBI2. *Nature* 475, 524–527.
- Hatano, Y., Man, M.Q., Uchida, Y., Crumrine, D., Mauro, T.M., Feingold, K.R., Elias, P.M., Holleran, W.M., 2010. Murine atopic dermatitis responds to peroxisome proliferator-activated receptors alpha and beta/delta (but not gamma) and liver X receptor activators. *J. Allergy Clin. Immunol.* 125, 160 e-1–169 e-5.
- Herath, K., Jayasuriya, H., Ondeyka, J.G., Guan, Z., Borris, R.P., Stijfhoorn, E., Stevenson, D., Wang, J., Sharma, N., Macnaul, K., Menke, J.G., Ali, A., Schulman, M.J., Singh, S.B., 2005. Guttiferone 1, a new prenylated benzophenone from *Garcinia humilis* as a liver X receptor ligand. *J. Nat. Prod.* 68, 617–619.
- Houck, K.A., Borchert, K.M., Hepler, C.D., Thomas, J.S., Bramlett, K.S., Michael, L.F., Burris, T.P., 2004. T0901317 is a dual LXR/FXR agonist. *Mol. Genet. Metab.* 83, 184–187.
- Hu, B., Unwalla, R.J., Goljer, I., Jetter, J.W., Quinet, E.M., Berrodin, T.J., Basso, M.D., Feingold, I.B., Nilsson, A.G., Wilhelmsson, A., Evans, M.J., Wrobel, J.E., 2010. Identification of phenylsulfone-substituted quinoxaline (WYE-672) as a tissue selective liver X-receptor (LXR) agonist. *J. Med. Chem.* 53, 3296–3304.
- Hu, X., Li, S., Wu, J., Xia, C., Lala, D.S., 2003. Liver X receptors interact with corepressors to regulate gene expression. *Mol. Endocrinol.* 17, 1019–1026.
- Huang, T.H., Razmovski-Naumovski, V., Salam, N.K., Duke, R.K., Tran, V.H., Duke, C.C., Roufogalis, B.D., 2005. A novel LXR-alpha activator identified from the natural product *Gynostemma pentaphyllum*. *Biochem. Pharmacol.* 70, 1298–1308.
- Infante, J., Rodriguez-Rodriguez, E., Mateo, I., Llorca, J., Vazquez-Higuera, J.L., Berciano, J., Combarros, O., 2010. Gene-gene interaction between heme oxygenase-1 and liver X receptor-beta and Alzheimer's disease risk. *Neurobiol. Aging* 31, 710–714.
- Janowski, B.A., Willy, P.J., Devi, T.R., Falck, J.R., Mangelsdorf, D.J., 1996. An oxysterol signalling pathway mediated by the nuclear receptor LXR alpha. *Nature* 383, 728–731.
- Janowski, B.A., Grogan, M.J., Jones, S.A., Wisely, G.B., Kliewer, S.A., Corey, E.J., Mangelsdorf, D.J., 1999. Structural requirements of ligands for the oxysterol liver X receptors LXRalpha and LXRbeta. *Proc. Natl. Acad. Sci. USA* 96, 266–271.
- Jayasuriya, H., Herath, K.B., Ondeyka, J.G., Guan, Z., Borris, R.P., Tiwari, S., de Jong, W., Chavez, F., Moss, J., Stevenson, D.W., Beck, H.T., Slattery, M., Zamora, N., Schulman, M., Ali, A., Sharma, N., MacNaul, K., Hayes, N., Menke, J.G., Singh, S.B., 2005. Diterpenoid, steroid, and triterpenoid agonists of liver X receptors from diversified terrestrial plants and marine sources. *J. Nat. Prod.* 68, 1247–1252.
- Jiang, Y.J., Lu, B., Kim, P., Elias, P.M., Feingold, K.R., 2006. Regulation of ABCA1 expression in human keratinocytes and murine epidermis. *J. Lipid Res.* 47, 2248–2258.
- Jiang, Y.J., Lu, B., Tarling, E.J., Kim, P., Man, M.Q., Crumrine, D., Edwards, P.A., Elias, P.M., Feingold, K.R., 2010. Regulation of ABCG1 expression in human keratinocytes and murine epidermis. *J. Lipid Res.* 51, 3185–3195.
- Jick, H., Zornberg, G.L., Jick, S.S., Seshadri, S., Drachman, D.A., 2000. Statins and the risk of dementia. *Lancet* 356, 1627–1631.
- Joseph, S.B., Laffitte, B.A., Patel, P.H., Watson, M.A., Matsukuma, K.E., Walczak, R., Collins, J.L., Osborne, T.F., Tontonoz, P., 2002a. Direct and indirect mechanisms for regulation of fatty acid synthase gene expression by liver X receptors. *J. Biol. Chem.* 277, 11019–11025.
- Joseph, S.B., McKilligan, E., Pei, L., Watson, M.A., Collins, A.R., Laffitte, B.A., Chen, M., Noh, G., Goodman, J., Hagger, G.N., Tran, J., Tipping, T.K., Wang, X., Lusis, A.J., Hsueh, W.A., Law, R.E., Collins, J.L., Willson, T.M., Tontonoz, P., 2002b. Synthetic LXR ligand inhibits the development of atherosclerosis in mice. *Proc. Natl. Acad. Sci. USA* 99, 7604–7609.
- Joseph, S.B., Bradley, M.N., Castrillo, A., Bruhn, K.W., Mak, P.A., Pei, L., Hogenesch, J., O'Connell, R. M., Cheng, G., Saez, E., Miller, J.F., Tontonoz, P., 2004. LXR-dependent gene expression is important for macrophage survival and the innate immune response. *Cell* 119, 299–309.
- Kaneko, E., Matsuda, M., Yamada, Y., Tachibana, Y., Shimomura, I., Makishima, M., 2003. Induction of intestinal ATP-binding cassette transporters by a phytosterol-derived liver X receptor agonist. *J. Biol. Chem.* 278, 36091–36098.
- Kase, E.T., Wensasa, A.J., Aas, V., Hojlund, K., Levin, K., Thoresen, G.H., Beck-Nielsen, H., Rustan, A.C., Gaster, M., 2005. Skeletal muscle lipid accumulation in type 2 diabetes may involve the liver X receptor pathway. *Diabetes* 54, 1108–1115.
- Katz, A., Udata, C., Ott, E., Hickey, L., Burczynski, M.E., Burghart, P., Vesterqvist, O., Meng, X., 2009. Safety, pharmacokinetics, and pharmacodynamics of single doses of LXR-623, a novel liver X-receptor agonist, in healthy participants. *J. Clin. Pharmacol.* 49, 643–649.
- Ketterer, C., Mussig, K., Machicao, F., Stefan, N., Fritsche, A., Haring, H.U., Staiger, H., 2011. Genetic variation within the NR1H2 gene encoding liver X receptor beta associates with insulin secretion in subjects at increased risk for type 2 diabetes. *J. Mol. Med.* 89, 75–81.
- Kim, K.H., Choi, S.H., Lee, T.S., Oh, W.K., Kim, D.S., Kim, J.B., 2006. Selective LXRalpha inhibitory effects observed in plant extracts of MEH184 (*Parthenocissua tricuspidata*) and MEH185 (*Euscaphis japonica*). *Biochem. Biophys. Res. Commun.* 349, 513–518.
- Koldamova, R., Lefterov, I., 2007. Role of LXR and ABCA1 in the pathogenesis of Alzheimer's disease – implications for a new therapeutic approach. *Curr. Alzheimer Res.* 4, 171–178.
- Koldamova, R.P., Lefterov, I.M., Staufenbiel, M., Wolfe, D., Huang, S., Glorioso, J.C., Walter, M., Roth, M.G., Lazo, J.S., 2005. The liver X receptor ligand T0901317 decreases amyloid beta production in vitro and in a mouse model of Alzheimer's disease. *J. Biol. Chem.* 280, 4079–4088.
- Komuves, L.G., Schmuth, M., Fowler, A.J., Elias, P.M., Hanley, K., Man, M.Q., Moser, A.H., Lobaccaro, J.M., Williams, M.L., Mangelsdorf, D.J., Feingold, K.R., 2002. Oxysterol stimulation of epidermal differentiation is mediated by liver X receptor-beta in murine epidermis. *J. Invest. Dermatol.* 118, 25–34.
- Kotani, H., Tanabe, H., Mizukami, H., Makishima, M., Inoue, M., 2010. Identification of a naturally occurring retinoid, honokiol, that activates the retinoid X receptor. *J. Nat. Prod.* 73, 1332–1336.
- Krycer, J.R., Brown, A.J., 2011. Crosstalk between the androgen receptor and the liver X receptor: implications for cholesterol homeostasis. *J. Biol. Chem.* 286, 20637–20647.
- Kumar, N., Solt, L.A., Conkright, J.J., Wang, Y., Istrate, M.A., Busby, S.A., Garcia-Ordenez, R.D., Burris, T.P., Griffin, P.R., 2010. The benzenesulfonamide T0901317 [N-(2,2,2-trifluoroethyl)-N-[4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenyl]-benzenesulfonamide] is a novel retinoic acid receptor-related orphan receptor-alpha/gamma inverse agonist. *Mol. Pharmacol.* 77, 228–236.
- Laffitte, B.A., Chao, L.C., Li, J., Walczak, R., Hummasti, S., Joseph, S.B., Castrillo, A., Wilpitz, D.C., Mangelsdorf, D.J., Collins, J.L., Saez, E., Tontonoz, P., 2003. Activation of liver X receptor improves glucose tolerance through coordinate regulation of glucose metabolism in liver and adipose tissue. *Proc. Natl. Acad. Sci. USA* 100, 5419–5424.
- Lazar, M.A., 2003. Nuclear receptor corepressors. *Nucl. Recept. Signal* 1, e001.
- Lee, S., Lee, J., Lee, S.K., Lee, J.W., 2008. Activating signal cointegrator-2 is an essential adaptor to recruit histone H3 lysine 4 methyltransferases MLL3 and MLL4 to the liver X receptors. *Mol. Endocrinol.* 22, 1312–1319.
- Lee, J.H., Park, S.M., Kim, O.S., Lee, C.S., Woo, J.H., Park, S.J., Joe, E.H., Jou, I., 2009. Differential SUMOylation of LXRalpha and LXRbeta mediates transrepression of STAT1 inflammatory signaling in IFN-gamma-stimulated brain astrocytes. *Mol. Cell.* 35, 806–817.
- Lehmann, J.M., Kliewer, S.A., Moore, L.B., Smith-Oliver, T.A., Oliver, B.B., Su, J.L., Sundseth, S.S., Winegar, D.A., Blanchard, D.E., Spencer, T.A., Willson, T.M., 1997. Activation of the nuclear receptor LXR by oxysterols defines a new hormone response pathway. *J. Biol. Chem.* 272, 3137–3140.
- Leoni, V., Caccia, C., 2011. Oxysterols as biomarkers in neurodegenerative diseases. *Chem. Phys. Lipids* 164, 515–524.
- Li, X., Zhang, S., Blander, G., Tse, J.G., Krieger, M., Guarente, L., 2007. SIRT1 deacetylates and positively regulates the nuclear receptor LXR. *Mol. Cell* 28, 91–106.
- Li, X., Yeh, V., Molteni, V., 2010. Liver X receptor modulators: a review of recently patented compounds (2007–2009). *Expert Opin. Ther. Pat.* 20, 535–562.
- Liu, C., Yang, X.V., Wu, J., Kuei, C., Mani, N.S., Zhang, L., Yu, J., Sutton, S.W., Qin, N., Banie, H., Karlsson, L., Sun, S., Lovenberg, T.W., 2011. Oxysterols direct B-cell migration through EBI2. *Nature* 475, 519–523.
- Lo Sasso, G., Murzilli, S., Salvatore, L., D'Errico, I., Petruzzelli, M., Conca, P., Jiang, Z.Y., Calabresi, L., Parini, P., Moschetta, A., 2010. Intestinal specific LXR activation stimulates reverse cholesterol transport and protects from atherosclerosis. *Cell Metab.* 12, 187–193.
- Magura, L., Blanchard, R., Hope, B., Beal, J.R., Schwartz, G.G., Sahmoun, A.E., 2008. Hypercholesterolemia and prostate cancer: a hospital-based case-control study. *Cancer Causes Control* 19, 1259–1266.

- Makishima, M., Okamoto, A.Y., Repa, J.J., Tu, H., Learned, R.M., Luk, A., Hull, M.V., Lustig, K.D., Mangelsdorf, D.J., Shan, B., 1999. Identification of a nuclear receptor for bile acids. *Science* 284, 1362–1365.
- Misra, A., Khurana, L., 2008. Obesity and the metabolic syndrome in developing countries. *J. Clin. Endocrinol. Metab.* 93, S9–30.
- Mitro, N., Vargas, L., Romeo, R., Koder, A., Saez, E., 2007. T0901317 is a potent PXR ligand: implications for the biology ascribed to LXR. *FEBS Lett.* 581, 1721–1726.
- Morales, J.R., Ballesteros, I., Deniz, J.M., Hurtado, O., Vivancos, J., Nombela, F., Lizasoain, I., Castrillo, A., Moro, M.A., 2008. Activation of liver X receptors promotes neuroprotection and reduces brain inflammation in experimental stroke. *Circulation* 118, 1450–1459.
- Morello, F., de Boer, R.A., Steffensen, K.R., Gnecci, M., Chisholm, J.W., Boomsma, F., Anderson, L.M., Lawn, R.M., Gustafsson, J.A., Lopez-Illasaca, M., Pratt, R.E., Dzau, V.J., 2005. Liver X receptors alpha and beta regulate renin expression in vivo. *J. Clin. Invest.* 115, 1913–1922.
- Motoshima, K., Noguchi-Yachide, T., Sugita, K., Hashimoto, Y., Ishikawa, M., 2009. Separation of alpha-glucosidase-inhibitory and liver X receptor-antagonistic activities of phenethylphenyl phthalimide analogs and generation of LXRalpha-selective antagonists. *Bioorg. Med. Chem.* 17, 5001–5014.
- Noguchi-Yachide, T., Aoyama, A., Makishima, M., Miyachi, H., Hashimoto, Y., 2007. Liver X receptor antagonists with a phthalimide skeleton derived from thalidomide-related glucosidase inhibitors. *Bioorg. Med. Chem. Lett.* 17, 3957–3961.
- Ondeyka, J.G., Jayasuriya, H., Herath, K.B., Guan, Z., Schulman, M., Collado, J., Dombrowski, A.W., Kwon, S.S., McCallum, C., Sharma, N., MacNaul, K., Hayes, N., Menke, J.G., Singh, S.B., 2005. Steroidal and triterpenoidal fungal metabolites as ligands of liver X receptors. *J. Antibiot. (Tokyo)* 58, 559–565.
- Ou, J., Tu, H., Shan, B., Luk, A., DeBose-Boyd, R.A., Bashmakov, Y., Goldstein, J.L., Brown, M.S., 2001. Unsaturated fatty acids inhibit transcription of the sterol regulatory element-binding protein-1c (SREBP-1c) gene by antagonizing ligand-dependent activation of the LXR. *Proc. Natl. Acad. Sci. USA* 98, 6027–6032.
- Park, M.C., Kwon, Y.J., Chung, S.J., Park, Y.B., Lee, S.K., 2010. Liver X receptor agonist prevents the evolution of collagen-induced arthritis in mice. *Rheumatology (Oxford)* 49, 882–890.
- Peet, D.J., Turley, S.D., Ma, W., Janowski, B.A., Lobaccaro, J.M., Hammer, R.E., Mangelsdorf, D.J., 1998. Cholesterol and bile acid metabolism are impaired in mice lacking the nuclear oxysterol receptor LXR alpha. *Cell* 93, 693–704.
- Peng, D., Hiipakka, R.A., Xie, J.T., Dai, Q., Kokontis, J.M., Reardon, C.A., Getz, G.S., Liao, S., 2011. A novel potent synthetic steroidal liver X receptor agonist lowers plasma cholesterol and triglycerides and reduces atherosclerosis in LDLR(−/−) mice. *Br. J. Pharmacol.* 162, 1792–1804.
- Pfeifer, T., Bucheiner, M., Chandak, P.G., Patankar, J., Kratzer, A., Obrowsky, S., Rechberger, G.N., Kadam, R.S., Kompella, U.B., Kostner, G.M., Kratky, D., Levak-Frank, S., 2011. Synthetic LXR agonist suppresses endogenous cholesterol biosynthesis and efficiently lowers plasma cholesterol. *Curr. Pharm. Biotechnol.* 12, 285–292.
- Platz, J., Nichols, J.A., Mensink, R.P., 2005. Plant sterols and stanols: effects on mixed micellar composition and LXR (target gene) activation. *J. Lipid Res.* 46, 2468–2476.
- Platz, E.A., Leitzmann, M.F., Visvanathan, K., Rimm, E.B., Stampfer, M.J., Willett, W.C., Giovannucci, E., 2006. Statin drugs and risk of advanced prostate cancer. *J. Natl. Cancer Inst.* 98, 1819–1825.
- Pommier, A.J., Alves, G., Viennois, E., Bernard, S., Communal, Y., Sion, B., Marceau, G., Damon, C., Mouzat, K., Cairra, F., Baron, S., Lobaccaro, J.M., 2010. Liver X Receptor activation downregulates AKT survival signaling in lipid rafts and induces apoptosis of prostate cancer cells. *Oncogene* 29, 2712–2723.
- Qin, Y., Dalen, K.T., Gustafsson, J.A., Nebb, H.L., 2009. Regulation of hepatic fatty acid elongase 5 by LXRalpha-SREBP-1c. *Biochim. Biophys. Acta* 1791, 140–147.
- Quinet, E.M., Savio, D.A., Halpern, A.R., Chen, L., Miller, C.P., Nambi, P., 2004. Gene-selective modulation by a synthetic oxysterol ligand of the liver X receptor. *J. Lipid Res.* 45, 1929–1942.
- Quinet, E.M., Basso, M.D., Halpern, A.R., Yates, D.W., Steffan, R.J., Clerin, V., Resmini, C., Keith, J.C., Berrodin, T.J., Feingold, I., Zhong, W., Hartman, H.B., Evans, M.J., Gardell, S.J., DiBlasio-Smith, E., Mounds, W.M., LaVallie, E.R., Wrobel, J., Nambi, P., Vlasuk, G.P., 2009. LXR ligand lowers LDL cholesterol in primates, is lipid neutral in hamster, and reduces atherosclerosis in mouse. *J. Lipid Res.* 50, 2358–2370.
- Rau, K.M., Kang, H.Y., Cha, T.L., Miller, S.A., Hung, M.C., 2005. The mechanisms and managements of hormone-therapy resistance in breast and prostate cancers. *Endocr. Relat. Cancer* 12, 511–532.
- Repa, J.J., Mangelsdorf, D.J., 2000. The role of orphan nuclear receptors in the regulation of cholesterol homeostasis. *Annu. Rev. Cell Dev. Biol.* 16, 459–481.
- Rough, J.J., Monroy, M.A., Yerrum, S., Daly, J.M., 2010. Anti-proliferative effect of LXR agonist T0901317 in ovarian carcinoma cells. *J. Ovarian Res.* 3, 13.
- Ruan, B., Wilson, W.K., Schroeffer Jr., G.J., 1998. An alternative synthesis of 4,4-dimethyl-5alpha-cholesta-8,14,24-trien-3beta-ol, an intermediate in sterol biosynthesis and a reported activator of meiosis and of nuclear orphan receptor LXR alpha. *Bioorg. Med. Chem. Lett.* 8, 233–236.
- Schroeffer Jr., G.J., 2000. Oxysterols: modulators of cholesterol metabolism and other processes. *Physiol. Rev.* 80, 361–554.
- Schultz, J.R., Tu, H., Luk, A., Repa, J.J., Medina, J.C., Li, L., Schwendner, S., Wang, S., Thoolen, M., Mangelsdorf, D.J., Lustig, K.D., Shan, B., 2000. Role of LXRs in control of lipogenesis. *Genes Dev.* 14, 2831–2838.
- Shannon, J., Tewoderos, S., Garzotto, M., Beer, T.M., Derenick, R., Palma, A., Farris, P.E., 2005. Statins and prostate cancer risk: a case-control study. *Am. J. Epidemiol.* 162, 318–325.
- Shenoy, S.D., Spencer, T.A., Mercer-Haines, N.A., Alipour, M., Gargano, M.D., Runge-Morris, M., Kocarek, T.A., 2004. CYP3A induction by liver X receptor ligands in primary cultured rat and mouse hepatocytes is mediated by the pregnane X receptor. *Drug Metab. Dispos.* 32, 66–71.
- Shinar, D.M., Endo, N., Rutledge, S.J., Vogel, R., Rodan, G.A., Schmidt, A., 1994. NER, a new member of the gene family encoding the human steroid hormone nuclear receptor. *Gene* 147, 273–276.
- Singh, S.B., Ondeyka, J.G., Liu, W., Chen, S., Chen, T.S., Li, X., Bouffard, A., Dropinski, J., Jones, A.B., McCormick, S., Hayes, N., Wang, J., Sharma, N., MacNaul, K., Hernandez, M., Chao, Y.S., Baffic, J., Lam, M.H., Burton, C., Sparrow, C.P., Menke, J.G., 2005. Discovery and development of dimeric podocarpic acid leads as potent agonists of liver X receptor with HDL cholesterol raising activity in mice and hamsters. *Bioorg. Med. Chem. Lett.* 15, 2824–2828.
- Sironi, L., Mitro, N., Cimino, M., Gelosa, P., Guerrini, U., Tremoli, E., Saez, E., 2008. Treatment with LXR agonists after focal cerebral ischemia prevents brain damage. *FEBS Lett.* 582, 3396–3400.
- Song, C., Kokontis, J.M., Hiipakka, R.A., Liao, S., 1994. Ubiquitous receptor: a receptor that modulates gene activation by retinoic acid and thyroid hormone receptors. *Proc. Natl. Acad. Sci. USA* 91, 10809–10813.
- Song, C., Hiipakka, R.A., Liao, S., 2000. Selective activation of liver X receptor alpha by 6alpha-hydroxy bile acids and analogs. *Steroids* 65, 423–427.
- Song, C., Liao, S., 2000. Cholestenic acid is a naturally occurring ligand for liver X receptor alpha. *Endocrinology* 141, 4180–4184.
- Song, C., Hiipakka, R.A., Liao, S., 2001. Auto-oxidized cholesterol sulfates are antagonistic ligands of liver X receptors: implications for the development and treatment of atherosclerosis. *Steroids* 66, 473–479.
- Spector, D., Anthony, M., Alexander, D., Arab, L., 2003. Soy consumption and colorectal cancer. *Nutr. Cancer* 47, 1–12.
- Spyridon, M., Moraes, L.A., Jones, C.I., Sage, T., Sasikumar, P., Bucci, G., Gibbins, J.M., 2011. LXR as a novel antithrombotic target. *Blood* 117, 5751–5761.
- Stein, S., Matter, C.M., 2011. Protective roles of SIRT1 in atherosclerosis. *Cell Cycle* 10, 640–647.
- Sticozzi, C., Pecorelli, A., Belmonte, G., Valacchi, G., 2010. Cigarette smoke affects ABCA1 expression via liver X receptor nuclear translocation in human keratinocytes. *Int. J. Mol. Sci.* 11, 3375–3386.
- Svensson, S., Ostberg, T., Jacobsson, M., Norstrom, C., Stefansson, K., Hallen, D., Johansson, I.C., Zachrisson, K., Ogg, D., Jendeborg, L., 2003. Crystal structure of the heterodimeric complex of LXRalpha and RXRbeta ligand-binding domains in a fully agonistic conformation. *EMBO J.* 22, 4625–4633.
- Tamehiro, N., Sato, Y., Suzuki, T., Hashimoto, T., Asakawa, Y., Yokoyama, S., Kawanishi, T., Ohno, Y., Inoue, K., Nagao, T., Nishimaki-Mogami, T., 2005. Riccardin C: a natural product that functions as a liver X receptor (LXR)alpha agonist and an RXRbeta antagonist. *FEBS Lett.* 579, 5299–5304.
- Tamura, K., Chen, Y.E., Horiuchi, M., Chen, Q., Daviet, L., Yang, Z., Lopez-Illasaca, M., Mu, H., Pratt, R.E., Dzau, V.J., 2000. LXRalpha functions as a cAMP-responsive transcriptional regulator of gene expression. *Proc. Natl. Acad. Sci. USA* 97, 8513–8518.
- Thomas, J., Bramlett, K.S., Montrose, C., Foxworthy, P., Eacho, P.I., McCann, D., Cao, G., Kiefer, A., McCowan, J., Yu, K.L., Grese, T., Chin, W.W., Burris, T.P., Michael, L.F., 2003. A chemical switch regulates fibrin specificity for peroxisome proliferator-activated receptor alpha (PPARalpha) versus liver X receptor. *J. Biol. Chem.* 278, 2403–2410.
- Traves, P.G., Hortelano, S., Zeini, M., Chao, T.H., Lam, T., Neuteboom, S.T., Theodorakis, E.A., Palladin, M.A., Castrillo, A., Bosca, L., 2007. Selective activation of liver X receptors by acanthoic acid-related diterpenes. *Mol. Pharmacol.* 71, 1545–1553.
- Uno, S., Endo, K., Jeong, Y., Kawana, K., Miyachi, H., Hashimoto, Y., Makishima, M., 2009. Suppression of beta-catenin signaling by liver X receptor ligands. *Biochem. Pharmacol.* 77, 186–195.
- Vedin, L.L., Lewandowski, S.A., Parini, P., Gustafsson, J.A., Steffensen, K.R., 2009. The oxysterol receptor LXR inhibits proliferation of human breast cancer cells. *Carcinogenesis* 30, 575–579.
- Viennois, E., Pommier, A.J., Mouzat, K., Oumeddour, A., El Hajjaji, F.Z., Dufour, J., Cairra, F., Volle, D.H., Baron, S., Lobaccaro, J.M., 2011. Targeting liver X receptors in human health: deadlock or promising trail? *Expert. Opin. Ther. Targets* 15, 219–232.
- Volle, D.H., Repa, J.J., Mazur, A., Cummins, C.L., Val, P., Henry-Berger, J., Cairra, F., Veyssiere, G., Mangelsdorf, D.J., Lobaccaro, J.M., 2004. Regulation of the aldo-keto reductase gene *akr1b7* by the nuclear oxysterol receptor LXRalpha (liver X receptor-alpha) in the mouse intestine: putative role of LXRs in lipid detoxification processes. *Mol. Endocrinol.* 18, 888–898.
- Volle, D.H., Lobaccaro, J.M., 2007. Role of the nuclear receptors for oxysterols LXRs in steroidogenic tissues: beyond the “foie gras”, the steroids and sex? *Mol. Cell. Endocrinol.* 265–266, 183–189.
- Wang, Y., Rogers, P.M., Stayrook, K.R., Su, C., Varga, G., Shen, Q., Nagpal, S., Burris, T.P., 2008a. The selective Alzheimer's disease indicator-1 gene (Seladin-1/DHCR24) is a liver X receptor target gene. *Mol. Pharmacol.* 74, 1716–1721.
- Wang, Y., Rogers, P.M., Su, C., Varga, G., Stayrook, K.R., Burris, T.P., 2008b. Regulation of cholesterologenesis by the oxysterol receptor, LXRalpha. *J. Biol. Chem.* 283, 26332–26339.
- Watanabe, M., Nakayama, T., Shiraishi, T., Stemmermann, G.N., Yatani, R., 2000. Comparative studies of prostate cancer in Japan versus the United States. A review. *Urol. Oncol.* 5, 274–283.
- Willy, P.J., Umesono, K., Ong, E.S., Evans, R.M., Heyman, R.A., Mangelsdorf, D.J., 1995. LXR, a nuclear receptor that defines a distinct retinoid response pathway. *Genes Dev.* 9, 1033–1045.

- Willy, P.J., Mangelsdorf, D.J., 1997. Unique requirements for retinoid-dependent transcriptional activation by the orphan receptor LXR. *Genes Dev.* 11, 289–298.
- Wolozin, B., Kellman, W., Ruosseau, P., Celesia, G.G., Siegel, G., 2000. Decreased prevalence of Alzheimer disease associated with 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitors. *Arch. Neurol.* 57, 1439–1443.
- Yang, C., McDonald, J.G., Patel, A., Zhang, Y., Umetani, M., Xu, F., Westover, E.J., Covey, D.F., Mangelsdorf, D.J., Cohen, J.C., Hobbs, H.H., 2006. Sterol intermediates from cholesterol biosynthetic pathway as liver X receptor ligands. *J. Biol. Chem.* 281, 27816–27826.
- Yasuda, T., Grillot, D., Billheimer, J.T., Briand, F., Delerive, P., Huet, S., Rader, D.J., 2010. Tissue-specific liver X receptor activation promotes macrophage reverse cholesterol transport in vivo. *Arterioscler. Thromb. Vasc. Biol.* 30, 781–786.
- Yoshikawa, T., Shimano, H., Yahagi, N., Ide, T., Amemiya-Kudo, M., Matsuzaka, T., Nakakuki, M., Tomita, S., Okazaki, H., Tamura, Y., Iizuka, Y., Ohashi, K., Takahashi, A., Sone, H., Osuga, J., Gotoda, T., Ishibashi, S., Yamada, N., 2002. Polyunsaturated fatty acids suppress sterol regulatory element-binding protein 1c promoter activity by inhibition of liver X receptor (LXR) binding to LXR response elements. *J. Biol. Chem.* 277, 1705–1711.
- Zelcer, N., Tontonoz, P., 2006. Liver X receptors as integrators of metabolic and inflammatory signaling. *J. Clin. Invest.* 116, 607–614.
- Zelcer, N., Khanlou, N., Clare, R., Jiang, Q., Reed-Geaghan, E.G., Landreth, G.E., Vinters, H.V., Tontonoz, P., 2007. Attenuation of neuroinflammation and Alzheimer's disease pathology by liver X receptors. *Proc. Natl. Acad. Sci. USA* 104, 10601–10606.
- Zuercher, W.J., Buckholz, R.G., Campobasso, N., Collins, J.L., Galardi, C.M., Gampe, R.T., Hyatt, S.M., Merrihew, S.L., Moore, J.T., Oplinger, J.A., Reid, P.R., Spearing, P.K., Stanley, T.B., Stewart, E.L., Willson, T.M., 2010. Discovery of tertiary sulfonamides as potent liver X receptor antagonists. *J. Med. Chem.* 53, 3412–3416.