

Research Report

Busy Making Other Plans: A Simulation Study on the Effects of Deviations from Preregistrations

A. J. Vijlbrief
December 18, 2025

Supervisors

A. M. Scheel
H. I. Oberman

Word Count

2,480

Introduction

Preregistrations were popularized during the 2010s in psychology as a way to increase reproducibility (Lindsay et al., 2018). Preregistrations are documents that are published at the start of the research process, in which decisions concerning data collection and analysis are recorded. This is done with the aim of limiting the extent to which such decisions can be influenced by results.

This practice increases transparency about research choices and has become more popular, with the number of preregistrations increasing annually (Ferguson et al., 2023; Lindsay et al., 2018). However, whether the desired effect of reduced flexibility and increased reproducibility has been achieved is debated. Preregistration should lead to more credible and reproducible results (Lakens, 2019; Nosek et al., 2018; Wagenmakers et al., 2012). However, no difference has been found between preregistered and non-preregistered papers in the number of significant findings (van den Akker, van Assen, et al., 2024), and no reduction in *p*-hacking (Brodeur et al., 2024).

There are many reasons for the uncertainty of the effect of preregistration. Preregistrations are often incomplete or overly vague about their decisions (Claesen et al., 2021; Heirene et al., 2024). Researchers also deviate from their predetermined choices (Claesen et al., 2021; van den Akker, Bakker, et al., 2024). Deviations have been found in as many as 93% of preregistered papers, with up to 89% being incompletely reported (Claesen et al., 2021; Willroth & Atherton, 2024). These high deviation rates could be the reason why we see no reduction in significant findings and *p*-hacking.

The aim of this project is to identify potentially problematic deviations and investigate their effects. I first discuss the goals and problems of preregistration and the potential impact of deviations in more detail. Then, through a simulation study, I examine the effect of said deviations on type I and type II error rates.

The Problem and the Solution

The inability to reliably reproduce findings in the field of psychology is a known problem. The Open Science Collaboration (2015) found that only 36% of significant results were reproducible. One reason for this “replication crisis” is “researcher degrees of freedom” (Simmons et al., 2011). This term describes the flexibility that researchers have in decisions made during their work. This includes decisions such as how many observations to collect, which statistical model to use and which covariates to add. When these decisions are made during data collection or analysis, researchers can make opportunistic choices which steer their results towards desired outcomes. Such opportunistic choices are called *p*-hacking and can dramatically increase false-positive rates (Stefan & Schönbrodt, 2023).

The preregistration is proposed as a way to limit these types of practices. Templates are available to researchers, to inform them about what information is important to preregister. The preregistration is then uploaded to a public repository, with a time stamp of when it was published. Peer reviewers, and others, have access to the preregistration and can evaluate it before the start of the study or compare it to the final research paper.

In theory preregistration works well, if researchers are specific in their preregistration and follow it closely, this should lead to better reproducible findings. In practice, preregistrations are often imprecise and incomplete, making them less effective at restricting researcher degrees of freedom. Another common problem are deviations from preregistration (Claesen et al., 2021; van den Akker, Bakker, et al., 2024). Deviating means that the preregistered plan is not being followed during the execution of the study. Deviations occur most commonly in data collection procedures, statistical models and exclusion criteria (van den Akker, Bakker, et al., 2024). This includes differences in the total sample size, how outliers are selected and which covariates are used. These deviations could

serve to further diminish the effectiveness of preregistrations, as it reintroduces researcher degrees of freedom and leaves room to revert back to p -hacking strategies.

Are all deviations bad?

Deviations from preregistrations occur in all different aspects of the research process, but what is not known is the effect of these deviations. Do changes from original research plans reduce research quality? Lakens (2024) theorized what the possible consequences could be. If deviations occur due to assumptions being violated or data no longer being suitable for the planned analysis, then deviating from the original research plan can have positive effects on the validity of a test. Similarly, adding additional analyses can increase the robustness. On the other hand, changes in testing also mean that a hypothesis is generally less “severely” tested. This means that a theory has not been given the proper opportunity to be falsified which can result in higher type II errors. Consequently, little can be said about whether deviations are bad, especially when the reason for the deviation is unknown.

There is no clear consensus on whether inconsistencies between preregistrations and final publications are a problem. When asked, researchers deemed deviations to be problematic to varying degrees (Willroth & Atherton, 2024). Changes to analyses and hypotheses were considered the least acceptable, whereas changing research platforms or software were deemed relatively justifiable. Whether the deviation was reported transparently and the estimated impact on the results of the study were also taken into consideration. To summarize, even though deviations from preregistration are common, a clear conclusion on their effects is still missing.

The present project

With this simulation study, I intend to close this gap in the literature and explore the effects of typical preregistration deviations on research outcomes and the quality of published results. Specifically, I aim to answer the question: ‘How do deviations from preregistrations affect type I and type II error rates?’ The effect of deviations is studied in three domains: sample size, outlier exclusion criteria, and the statistical model. Different deviations are simulated in each domain and compared to a baseline condition without deviations. This research question is investigated in the context of behavioral psychology and uses a linear regression as the model of interest. The effects are examined across two scenarios: a no-effect scenario, and an effect scenario, in which the main effect parameter is changed. The aim of this study is descriptive and therefore no specific hypotheses are formulated.

As a secondary research question, I address the issue of reasons for deviations being unknown. The reason behind a deviation is important for assessing its effect. A researcher being forced to deviate due to circumstance is very different from a researcher *choosing* to deviate. When a researcher chooses to deviate, this could be because they are picking the results they prefer. Initially, the deviation conditions are simulated and the outcomes are used to assess type I and type II error as compared to the baseline condition. In the second phase, deviations are simulated but the outcomes are selectively chosen from either the baseline or the deviation conditions, based on which condition has a significant p -value. This is done in order to proxy the opportunistic selection of results by the researcher.

With these simulations I hope to shed light on the effect of deviations from preregistration on type I and type II errors, as well as the effect of picking deviations opportunistically.

Methods

Conditions

I reviewed which deviations to investigate by (1) how common they are, (2) their potential impact and (3) their justifiability. Based on this deviations were chosen in the following domains: sample size, outlier exclusion criteria, and statistical model. Within the three chosen domains, multiple conditions were simulated and compared to a baseline condition without any deviations.

Sample size

Deviations in the sample size are some of the most common deviations, with consistency between preregistrations and published papers only being 28% for the exact sample size (van den Akker, Bakker, et al., 2024). Changing the sample size can inflate type II error (by decreasing power) as well as the type I error (Lakens, 2024; Simmons et al., 2011). Small deviations in sample size are common and often due to factors outside of the researcher’s agency. Researchers also admit to stopping collection earlier or continuing collection longer after finding disappointing results (John et al., 2012). However, it is unknown how often this is also used as a reason to deviate from preregistrations. Based on this, the conditions in Table 1 are simulated within the sample size domain.

Outlier exclusion criteria

Literature shows that over half of published papers do not adhere to their pre-specified outlier exclusion criteria (Claesen et al., 2021; van den Akker, Bakker, et al., 2024). Vague or missing criteria in preregistrations make it harder to assess deviations and leave a lot of room for interpretation and ad hoc decisions (Heirene et al., 2024). This could potentially lead researchers to choose how to exclude outliers based on which method provides better results. Researchers themselves, however, deem outlier deviations to be relatively acceptable (Willroth & Atherton, 2024).

Lakens (2024) argues that adding additional exclusion criteria can undermine the severity of a test. This idea is corroborated by Stefan & Schönbrodt (2023), who showed that the type I error increases linearly with the number of outlier detection methods used. Furthermore, 38% of researchers admit to excluding outliers only after examining their impact on the data (John et al., 2012). If this is also the reason why many people deviate from their preregistered outlier criteria, then these deviations are important.

The most common method for identifying outliers in the social sciences is the z -score (Bakker & Wicherts, 2014; Leys et al., 2013). A minimum of three standard deviations from the mean is often maintained as the rule of thumb for excluding cases. Within linear regression, outliers are also often identified based on their influence, commonly using Cook’s distance. Cook’s distance excludes outliers based on how much their exclusion would influence the regression coefficients. In this study, the baseline condition will be to not exclude any datapoints, with deviation conditions based on z -score and Cook’s distance (Table 1).

Statistical model

Amongst all of the aforementioned domains, the least is known about why people deviate from their preregistered statistical model. “Statistical model” is a very broad domain and includes many types of deviations, such as changes in the dependent variable, independent variable, statistical inference

criteria and the actual statistical test. In this paper, “statistical model” refers to the above-mentioned types of changes, not only the statistical test and its specifications.

The broadness of the domain also means that deviations within the domain have been investigated from multiple angles. Claesen et al. (2021) found a deviation rate of 70% within the “analysis” domain, including deviations like examining additional effects, changing which model is tested and performing unregistered robustness checks. The majority of these deviations went unreported. Another study reported inconsistencies in 40% of the statistical models, deviations included the specifications of the variables, which model was tested and how variables were used (van den Akker, Bakker, et al., 2024).

As Lakens (2024) argues, at times it might be necessary to alter the statistical model. This could happen when a variable has been measured at a different level than expected or a test assumption has been violated. However, changing the statistical model can also be done opportunistically. Choosing to add an extra dependent variable because results are not yet satisfying can increase the type I error rate to 9.5%, and the addition of a covariate can increase it to 11.7% (Simmons et al., 2011). The reasoning behind the change thus becomes very important. Therefore, deviations due to failing to properly specify how variables would be operationalized in the preregistration are also seen as a larger shortcoming than model deviations due to unforeseen consequences (Willroth & Atherton, 2024). Within the statistical model, three deviation conditions are simulated: outcome switching, adding a continuous covariate, and adding a dichotomous covariate (Table 1).

Table 1: Parameter Values for Baseline Condition and Deviation Conditions per Domain

Domain	Baseline condition	Deviations conditions
Sample size	200	+5 +30 -5 -30
Outlier exclusion criteria	No outliers excluded	Exclusion based on z -scores Exclusion based on Cook’s distance
Statistical methods: <i>covariates</i>	No covariates	Continuous covariate Dichotomous covariate
Statistical methods: <i>outcome switching</i>	Y	Ya

Data-generating mechanism

Data is generated under two scenarios: “X has an effect on Y” and “X has no effect on Y”. The only difference in data generation between the two scenarios is in the main effect parameter, β_1 . In the effect scenario, β_1 is 0.2, and in the no effect scenario, β_1 is 0. In the baseline condition, the statistical model is defined as

$$Y = \beta_0 + \beta_1 X + \epsilon,$$

where Y is the dependent variable, β_0 the intercept, β_1 the main effect, and ϵ the random error.

The population parameter values are presented in Table 2, together with the parameter values for the deviation conditions. The complete data-generating model is defined as

$$Y = \beta_0 + \beta_1 X + \beta_2 Z + \beta_3 D + \epsilon,$$

$$Y_a \sim Y, \quad \rho(Y, Y_a) = 0.6,$$

where Z and D represent a continuous and categorical covariate respectively and Y_a represents an alternative outcome variable with a correlation of 0.6 with variable Y .

As mentioned, data is generated under two scenarios: an effect scenario and a no-effect scenario. The only difference in data generation between the two scenarios is in the main effect parameter, β_1 . In the effect scenario, β_1 is 0.2, and in the no-effect scenario, β_1 is 0.

In order to assess differences between outlier exclusion criteria, the data needs to include outliers. These are simulated by generating 5% of the data, 2.5% at each end of the distribution, with -3 and 3 as mean values for the normal distribution from which ϵ is drawn.

Table 2: Parameter Values for the Data generating mechanism

Parameter	Value
Intercept	$\beta_0 = 0$
Regression coefficient	$\beta_1 = 0$ or $\beta_1 = 0.20$
Independent variable	$X \sim \mathcal{N}(\mu, \sigma^2)$
Random error	95% = $\epsilon \sim \mathcal{N}(0, 1)$ 2.5% = $\epsilon \sim \mathcal{N}(-3, 1)$ 2.5% = $\epsilon \sim \mathcal{N}(3, 1)$
Continuous covariate regression coefficient	$\beta_2 = 0.06$
Continuous demographic variable	$Z \sim \mathcal{N}(\mu, \sigma^2)$
Dichotomous covariate regression coefficient	$\beta_3 = 0.06$
Dichotomous demographic variable	$D \sim \text{Bernoulli}(0.5)$
Alternative outcome error	$\epsilon_a \sim \mathcal{N}(\mu, \sigma^2)$
Sample size	200

Estimands

The estimand of this study is β_1 , which represents the effect of the independent variable X on dependent variable Y . The coefficient is estimated using the `stats` package in R (R Core Team, 2024), under the baseline and deviation conditions.

Methods


As previously mentioned, there are two scenarios under which data is generated. A no-effect scenario and an effect scenario. Within these scenarios each simulated data set is examined under different conditions. The conditions include a baseline condition, with no deviations, and one condition for each possible deviation (Table 1).

To answer the main research question, the type I and type II error rates are compared between the baseline result and each deviation result. All ten conditions, one baseline and nine deviation conditions, are executed using the same data set. This is done iteratively, where the β_1 regression coefficient, the p-value and the confidence interval are recorded for each condition.

To answer the second research question, the effect of choosing whether or not to deviate is examined. This is done using the same data set as in the first section. For each iteration, the baseline p-value is assessed. When it is significant, the baseline condition is recorded for that iteration, if it is insignificant, the deviations conditions of that iteration are checked. If any of the deviations conditions are significant, one is chosen at random and recorded for that iteration. This provides an ‘optimized’ data set, with the largest number of significant iterations possible in the data set. The error rates will

then be compared between the baseline results and the ‘optimized’ result. An overview can be found in Figure 1.

Figure 1: Condition Comparisons



Note. This figure illustrates the study phases. In phase 1, the baseline result is compared to the deviation result. In phase 2, the baseline result is compared to the best result.

Performance measures

Performance is assessed through the type I and type II error for the estimand β_1 . Specifically, the regression coefficient itself, the p -value (for decision making in phase 2) and the confidence interval (to assess coverage) are collected for each condition in each iteration.

The type I error refers to a false-positive, or rejecting the null-hypothesis when it is true. In this study that would mean detecting an effect in the no-effect scenario. Type I error is generally acceptable at a rate of 5% or below, based on an alpha level of .05. In this case, a rate of higher than 5% is considered an inflated type I error rate.

Type II error refers to a false negative, or failing to reject the null-hypothesis when it is false. In this study that would mean failing to detect an effect in the effect scenario. Type II error is commonly assessed in the form of power. Power is $1 - \beta$, where β is the type II error. The nominal type II error rate of .20 results in a generally accepted power level of 80%. In this study, type II error will be expressed in the form of power rates and consequently, power rates below 80% are considered deflated power.

In order to account for the uncertainty introduced by simulationThe number of iterations is based on the desired precision of the project. A Monte-Carlo standard error (MCSE) of 1% is deemed appropriate. According to the formula by Morris et al. (2019),

$$\begin{aligned}
MCSE &= \sqrt{\frac{\widehat{\text{Power}} (1 - \widehat{\text{Power}})}{n_{\text{sim}}}}, \\
0.01 &= \sqrt{\frac{\widehat{0.8} (1 - \widehat{0.8})}{n_{\text{sim}}}}, \\
n_{\text{sim}} &= \frac{0.8 \times 0.2}{(0.01)^2}, \\
n_{\text{sim}} &= 1600
\end{aligned}$$

the required number of iterations to achieve the desired MCSE is 1600.

With an MCSE of 1%, power rates are assessed at a precision interval of [79, 81] around the nominal power level of 80%. Power levels outside of this range can be stated to have increased or deflated power levels that are not due to simulation noise.

For the type I error, 1600 iterations results in an MCSE of 0.00545.

$$\begin{aligned}
MCSE &= \sqrt{\frac{\widehat{\text{Type I error}} (1 - \widehat{\text{Type I error}})}{n_{\text{sim}}}}, \\
MCSE &= \sqrt{\frac{\widehat{0.05} (1 - \widehat{0.05})}{1600}}, \\
MCSE &= \sqrt{2.96875e-05}, \\
MCSE &= 0.00545
\end{aligned}$$

This means that type I error is assessed with a precision interval of [0.045, 0.055] around the nominal α level of 0.05. Any type I error rates outside of this interval are considered deflated or inflated.

Preregistration, reproducibility and ethics

This project was preregistered on the 31th of March, 2026. The preregistration can be found at [link]. In order to prevent seed hacking, I preregistered a seed which would only become available after the date of preregistration. In order to advance reproducibility, all files related to this project are publicly available at [github link]. This includes data, code scripts and output files. I made use of **renv** to create a reproducible environment in which package versions and r version are recorded. Instructions on how to reproduce the findings can be found under the README file on github. During this project an R package was created under the name **thepack** to store the most important functions. The package is publicly available at [githublink].

Ethics approval was obtained on October 7th, 2025 from the University of Utrecht under case-number #25-1980.

Results

Unplanned deviations

Effects on power


The results for the primary research question show an effect of deviating on power in each of the three deviation domains (Figure 2).

Within the sample size domain, small changes in sample size show no effect on power whereas larger changes in sample size are impactful. It is an established fact that increasing sample size can increase the power of a study [source]. This effect is also observed in the results of this study. A 15% decrease in sample size resulted in a deflated power level of 74.5 %. This means that deviating in the form of collecting fewer participants than planned, is likely to lead to an underpowered study. When deviating by large sample size *increases*, the results fail to show a negative impact. The power level increases to 85.1% in the 15% increase condition.

Deviations in the Outlier exclusion criteria domain also impact power levels. When using a strict z -score (exclusion based on 2 s.d. instead of 3 s.d.) power increases to 83.8 %. When using Cook's distance instead of z -scores, the power level drops to 68.2 %.

In the Statistical model domain, deflated power is only observed in the alternative outcome condition. Deviating by using an alternative, correlated outcome variable results in a power level of 35.2%. This is incredibly low when compared to the nominal power level of 80%. Both the continuous and dichotomous covariate conditions fail to show an effect on power.


Figure 2: Power level for Baseline and Deviation Conditions in the Effect scenario



Effects on Type I error

The Monte Carlo standard error for type I error resulted in a precision range of $[0.045, 0.055]$ around the nominal α level of 0.05. Only two conditions show results outside of this precision range. Small increases in effect size show an increased type I error of 5.6%. Using a strict z -score results in a type I error rate of 5.8%. All other deviation scenarios fall within the prespecified precision range and therefore do not indicate any effects on the type I error (Figure 3).

Figure 3: Type I error rate for Baseline and Deviation scenarios in the no-effect Scenario



Choosing to deviate

The second research question pertains to the effect of *choosing* to deviate when the baseline condition is insignificant. The results fail to show a negative effect of choosing to deviate on power levels. The power level does not deflate and is actually increased to 92.7%. In the no-effect scenario, an inflated type I error rate of 16.3% is observed. This is a large inflation compared to the nominal level of 5% and similarly much higher than the highest type I error rate in a singular deviation condition 5.8% in the Strict z -score condition).

In the effect scenario, 1279 out of 1600 repetitions resulted in a significant baseline. Of the 321 repetitions where the baseline was insignificant, 36.1% also reported no significant deviation conditions. The remaining 64% of repetitions in which the baseline was insignificant had at least one significant deviation, with seven significant deviation conditions at most.

Table 3: Distribution of the number of significant deviations

Significant deviation conditions	Occurences	
	Effect	No-effect
(Significant baseline)	1279 (79.9%)	83 (5.2%)
0	116 (7.2%)	1339 (83.7%)
1	75 (4.7%)	126 (7.9%)
2	61 (3.8%)	27 (1.7%)
3	30 (1.9%)	17 (1.1%)
4	31 (1.9%)	5 (0.3%)
5	4 (0.2%)	1 (0.1%)
6	3 (0.2%)	2 (0.1%)
7	1 (0.1%)	0 (0%)

In the no-effect scenario, the majority (83.7%) of the repetitions showed an insignificant baseline as well as insignificant deviations (Table 3). However of the repetitions in which the baseline was insignificant, 13.3% showed at least one significant result. Neither of the scenarios contained a repetition in which all deviations were significant when the baseline was not.

Discussion

This simulation study demonstrates that type I and type II error are both affected by deviations from a preregistration.

Power can be affected both positively and negatively, depending on the type of deviation. Collecting much fewer participants, using Cook’s distance as outlier criterium and switching to an alternative outcome all deflate power levels. Power is most impacted by switching outcomes with a steep drop off from the nominal 80% to 35.2% power. Small changes in sample size and the addition of a covariate show little effect, whereas collecting many more participants than planned and using a z -score of two standard deviations show slight increases in power.

Only two deviation conditions showed effects on the type I error. Small increases in sample size resulted in a small inflation of type I error and using two standard deviations as outlier criterium showed a slightly larger inflation. For all other conditions, effects on type I error were not strong enough to suggest an effect outside of simulation noise. This shows that there is little risk of inflating the type I error when being forced to deviate from a preregistration except for using a stricter z -score as outlier criterium.

The small inflation when slightly increasing the sample size is a borderline case. Using the MCSE of 0.055 which was preregistered for this study means that it falls outside of the precision range, outside of which I stated that effects were unlikely to be due to simulation noise. Using the empirical MCSE of this condition however shows that the effect might still be due to noise. With a type I error of 0.055625, the empirical MCSE becomes 0.00573. This means that type I error rates higher than 0.055729 would be considered inflated, which this condition just remains beneath. When theoretically considering this effect, it is more likely that this effect is in fact due to simulation error rather than an

actual effect. If this was an effect of the deviation, it would be expected to see an opposite effect when decreasing the sample size slightly as well as a stronger effect when increasing the sample size more extremely. Neither of these effects can be observed in the other sample size conditions, thus making it more likely that the observed effect is indeed due to simulation noise. (In light of this, all other conditions were also evaluated using the empirical MCSE. The slight increase in sample size remained the only condition in which this changed the conclusion.) From these results

A limitation of this study is the fact that the data was simulated based on common structures and numbers found in the social sciences. This is likely to make the results less generalizable to other disciplines. The baseline sample size, effect size and methods investigated are all common among social sciences but might look much different for other statistics oriented disciplines such as medicine. It remains unknown whether the found effects would remain under smaller and larger samples or effect sizes.

Conclusion

-larger false positive rate in published papers -papers with deviations in outlier criterium more likely to be false positive -because opportunistic choosing increases type I that much more than individual deviations it is more important to transparently record the reasons for deviating.

References

- Bakker, M., & Wicherts, J. M. (2014). Outlier Removal and the Relation with Reporting Errors and Quality of Psychological Research. *PLOS ONE*, 9(7), e103360. <https://doi.org/10.1371/journal.pone.0103360>
- Brodeur, A., Cook, N. M., Hartley, J. S., & Heyes, A. (2024). Do Preregistration and Preanalysis Plans Reduce p -Hacking and Publication Bias? Evidence from 15,992 Test Statistics and Suggestions for Improvement. *Journal of Political Economy Microeconomics*, 2(3), 527–561. <https://doi.org/10.1086/730455>
- Claesen, A., Gomes, S., Tuerlinckx, F., & Vanpaemel, W. (2021). Comparing dream to reality: An assessment of adherence of the first generation of preregistered studies. *Royal Society Open Science*, 8(10), 211037. <https://doi.org/10.1098/rsos.211037>
- Ferguson, J., Littman, R., Christensen, G., Paluck, E. L., Swanson, N., Wang, Z., Miguel, E., Birke, D., & Pezzuto, J.-H. (2023). Survey of open science practices and attitudes in the social sciences. *Nature Communications*, 14(1), 5401. <https://doi.org/10.1038/s41467-023-41111-1>
- Heirene, R., LaPlante, D., Louderback, E., Keen, B., Bakker, M., Serafimovska, A., & Gainsbury, S. (2024). Preregistration specificity and adherence: A review of preregistered gambling studies and cross-disciplinary comparison. *Meta-Psychology*, 8. <https://doi.org/10.15626/MP.2021.2909>
- John, L. K., Loewenstein, G., & Prelec, D. (2012). Measuring the Prevalence of Questionable Research Practices With Incentives for Truth Telling. *Psychological Science*, 23(5), 524–532. <https://doi.org/10.1177/0956797611430953>
- Lakens, D. (2019). *The Value of Preregistration for Psychological Science: A Conceptual Analysis*. PsyArXiv. <https://doi.org/10.31234/osf.io/jbh4w>
- Lakens, D. (2024). When and How to Deviate From a Preregistration. *Collabra: Psychology*, 10(1), 117094. <https://doi.org/10.1525/collabra.117094>
- Leys, C., Ley, C., Klein, O., Bernard, P., & Licata, L. (2013). Detecting outliers: Do not use standard deviation around the mean, use absolute deviation around the median. *Journal of Experimental Social Psychology*, 49(4), 764–766. <https://doi.org/10.1016/j.jesp.2013.03.013>
- Lindsay, S. D., Nosek, B. A., & Stephen, D. (2018). Preregistration Becoming the Norm in Psychological Science. *APS Observer*, 31.
- Morris, T. P., White, I. R., & Crowther, M. J. (2019). Using simulation studies to evaluate statistical methods. *Statistics in Medicine*, 38(11), 2074–2102. <https://doi.org/10.1002/sim.8086>
- Nosek, B. A., Ebersole, C. R., DeHaven, A. C., & Mellor, D. T. (2018). The preregistration revolution. *Proceedings of the National Academy of Sciences*, 115(11), 2600–2606. <https://doi.org/10.1073/pnas.1708274114>
- Open Science Collaboration. (2015). Estimating the reproducibility of psychological science. *Science*, 349(6251), aac4716. <https://doi.org/10.1126/science.aac4716>
- R Core Team. (2024). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing.
- Simmons, J. P., Nelson, L. D., & Simonsohn, U. (2011). False-Positive Psychology: Undisclosed Flexibility in Data Collection and Analysis Allows Presenting Anything as Significant. *Psychological Science*, 22(11), 1359–1366. <https://doi.org/10.1177/0956797611417632>
- Stefan, A. M., & Schönbrodt, F. D. (2023). *Big little lies: A compendium and simulation of p-hacking strategies*.
- van den Akker, O. R., Bakker, M., Van Assen, M. A. L. M., Pennington, C. R., Verweij, L., Elsherif, M. M., Claesen, A., Gaillard, S. D. M., Yeung, S. K., Frankenberger, J.-L., Krautter, K., Cockcroft, J. P., Kreuer, K. S., Evans, T. R., Heppel, F. M., Schoch, S. F., Korbacher, M., Yamada,

- Y., Albayrak-Aydemir, N., ... Wicherts, J. M. (2024). The potential of preregistration in psychology: Assessing preregistration producibility and preregistration-study consistency. *Psychological Methods*. <https://doi.org/10.1037/met0000687>
- van den Akker, O. R., van Assen, M. A. L. M., Bakker, M., Elsherif, M., Wong, T. K., & Wicherts, J. M. (2024). Preregistration in practice: A comparison of preregistered and non-preregistered studies in psychology. *Behavior Research Methods*, 56(6), 5424–5433. <https://doi.org/10.3758/s13428-023-02277-0>
- Wagenmakers, E.-J., Wetzels, R., Borsboom, D., van der Maas, H. L. J., & Kievit, R. A. (2012). An Agenda for Purely Confirmatory Research. *Perspectives on Psychological Science*, 7(6), 632–638. <https://doi.org/10.1177/1745691612463078>
- Willroth, E. C., & Atherton, O. E. (2024). *Best Laid Plans: A Guide to Reporting Preregistration Deviations*.