

# Quantitative Risk Management



<http://www.qrmtutorial.org>

29th International Summer School of the SAA

P. Embrechts, R. Frey, M. Hofert, A. J. McNeil

# Course information

- Website: <http://www.qrmtutorial.org>
- Book: A. J. McNeil, R. Frey, P. Embrechts  
*Quantitative Risk Management* (1st edition: 2005; revised edition: 2015)



# Overview

- 1 Risk in perspective
- 2 Basics concepts in risk management
- 3 Empirical properties of financial data
- 4 Financial time series
- 5 Extreme value theory
- 6 Multivariate models
- 7 Copulas and dependence
- 8 Aggregate risk
- 9 Market Risk
- 10 Credit risk

- 11 Portfolio credit risk management**
- 14 Multivariate time series**
- 17 Introduction to counterparty risk**

# 1 Risk in perspective

1.1 Risk

1.2 Why manage financial risk?

1.3 Quantitative Risk Management

# 1.1 Risk

- The Concise Oxford English Dictionary: “hazard, a chance of bad consequences, loss or exposure to mischance”.
- McNeil et al. (2005): “any **event or action** that may adversely **affect an organization's** ability **to achieve its objectives** and execute its strategies”.
- No single one-sentence definition captures all aspects of risk.  
For us: **risk** = **chance of loss**  $\Rightarrow$  uncertainty  $\Rightarrow$  randomness

## 1.1.1 Risk and randomness

- Kolmogorov (1933) introduced the notion of a **probability space**  $(\Omega, \mathcal{F}, \mathbb{P})$ :
  - ▶  $\Omega$  is the **sample space** which contains realizations  $\omega \in \Omega$  (“state of nature”) of an experiment;
  - ▶ the  **$\sigma$ -algebra**  $\mathcal{F}$  contains all sets (“events”) to which we can assign probabilities; and
  - ▶  $\mathbb{P}(\cdot)$  denotes a **probability measure**.

- We will mostly model situations in which an investor holds today an asset with an uncertain future value.
- To this end, we model the value of the asset/risky position as a random variable  $X : \Omega \rightarrow \mathbb{R}$  (or  $L$ , typically representing a loss). Several risky positions are modelled by a random vector  $\mathbf{X} : \Omega \rightarrow \mathbb{R}^d$ .
- Most of this modelling concerns the distribution functions

$$F_X(x) = \mathbb{P}(X \leq x), \quad x \in \mathbb{R}, \quad \text{and} \quad F_{\mathbf{X}}(\mathbf{x}) = \mathbb{P}(\mathbf{X} \leq \mathbf{x}), \quad \mathbf{x} \in \mathbb{R}^d,$$

of  $X$  and  $\mathbf{X}$ , respectively.

- If time matters, one can consider sequences of random variables  $(X_t)_{t \geq 0}$ , so-called stochastic processes.
- Our modelling tools will mainly come from probability and statistics.

## 1.1.2 Financial Risk

There are various **types of risks**. We **focus on**

**Market risk** Risk of loss in a financial position due to **changes** in the **underlying components** (e.g. stock/bond/commodity prices)

**Credit risk** Risk of a **counterparty** failing to meet its obligations (**default**), i.e. the risk of not receiving promised repayments (e.g. loans/bonds).

**Operational risk (OpRisk)** Risk of loss resulting from inadequate or **failed internal processes, people and systems** or from **external events** (e.g. fraud, fat-finger trades, earthquakes).

There are many **other types** of risks such as liquidity risk, underwriting risk, or **model risk** (the risk of using a misspecified or inappropriate **model** for measuring risk; model risk is always present to some degree).



## 1.1.3 Measurement and management

### Risk measurement

- Suppose we hold a **portfolio** of  $d$  investments with weights  $w_1, \dots, w_d$ . Let  $X_j$  denote the change in value of the  $j$ th investment. The **change in value** – *profit and loss (P&L)* – of the portfolio over a given **holding period** is then

$$X = \sum_{j=1}^d w_j X_j.$$

Measuring the risk now consists of determining the *distribution function*  $F$  (or functionals of it, e.g. mean, variance,  $\alpha$ -quantiles  $F^{\leftarrow}(\alpha) = \inf\{x \in \mathbb{R} : F(x) \geq \alpha\}$ ).

- To this end, we **need a properly calibrated joint model** for  $\mathbf{X} = (X_1, \dots, X_d)$ . Statistical estimates of  $F$  or one of its functionals are obtained based on historical observations of this model.

# Risk management

- What is RM? Kloman (1990) writes:

“RM is a discipline for living with the possibility that future events may cause adverse effects.”

⇒ It is about ensuring resilience to future events.
- Note that financial firms are not passive/defensive towards risk, banks and insurers actively/willingly take risks because they seek a return. RM thus belongs to the core competence of a bank or insurance company.
- What does managing risks involve?
  - ▶ Determine the capital to hold to absorb losses, both for regulatory purposes (to comply with regulators) and economic capital purposes (to survive as a company).
  - ▶ Ensuring portfolios are well diversified.
  - ▶ Optimizing portfolios according to risk-return considerations

## 1.2 Why manage financial risk?

- **Society** (single customers and as a whole (systemic risk)) relies on the stability of the banking and insurance
- This is related to *systemic importance* of the company in question (size and connectivity to other firms). Considering some firms as **too big to fail** creates a **moral hazard** (should be avoided!) since the **management of such a firm may take more risk knowing that it would be bailed out in a crisis.**
- **Better risk management can** reduce the risk of company failure and **protect customers and policyholders.** **However, regulation** must be designed with care and **should not promote herding, procyclical behaviour or other forms of endogenous risk** that could result in a systemic crisis.

# 1.3 Quantitative Risk Management

## 1.3.1 The Q in QRM

- We treat QRM as a quantitative science using the language of mathematics in general, and probability and statistics in particular.
- Mathematics and statistics provide us with with a suitable language and with appropriate concepts for describing financial risks.
- We also point out assumptions and limitations of the methodology used.
- The Q in QRM is an essential part of the RM process. We believe it remains (if applied correctly and honestly) a part of the solution to managing risk (not the problem). See also Shreve (2008):  
“Don’t blame the quants. Hire good ones instead and listen to them.”

## 1.3.2 The nature of the challenge

- Our approach to QRM has two main strands:
  - ▶ Put current practice onto a firmer mathematical ground;
  - ▶ Put together techniques and tools which go beyond current practice and address some of the deficiencies.
- In particular, some of the challenges of QRM are:
  - ▶ Extremes matter.
  - ▶ Interdependence and concentration of risks.
  - ▶ The problem of scale (models for all risk factors may not be feasible).
  - ▶ Interdisciplinarity.
  - ▶ Communication and education.

## 2 Basics concepts in risk management

2.1 Modelling value and value change

2.2 Risk measurement

## 2.1 Modelling value and value change

### 2.1.1 Mapping of risks

To start of we set up a general mathematical model for (changes in) value caused by financial risks. For this we work on a *probability space*  $(\Omega, \mathcal{F}, \mathbb{P})$  and consider a risk or loss as a *random variable*  $X : \Omega \rightarrow \mathbb{R}$  (or:  $L$ ).

- Consider a *portfolio* of assets and possibly liabilities. The *value* of the portfolio at time  $t$  (*today*) is denoted by  $V_t$  (a random variable; assumed to be known at  $t$ ; its *df* is typically *not trivial to determine!*).
- We consider a given *time horizon*  $\Delta t$  and *assume*:
  - 1) the *portfolio composition remains fixed* over  $\Delta t$ ;
  - 2) there are *no intermediate payments* during  $\Delta t$ $\Rightarrow$  *Fine for small  $\Delta t$  but unlikely to hold for large  $\Delta$ .*

- The *change* in value of the portfolio is then given by

$$\Delta V_{t+1} = V_{t+1} - V_t$$

and we define the (random) *loss* by the *sign-adjusted* value change

$$L_{t+1} = -\Delta V_{t+1}$$

(as QRM is mainly concerned with losses).

### Remark 2.1

- 1) The *distribution of  $L_{t+1}$*  is called *loss distribution* (df  $F_L$  or simply  $F$ ).
- 2) Practitioners often consider the *profit-and-loss (P&L) distribution* which is the distribution of  $-L_{t+1} = \Delta V_{t+1}$ .
- 3) For longer time intervals,  $\Delta V_{t+1} = V_{t+1}/(1 + r) - V_t$  ( $r$  = *risk-free interest rate*) would be more appropriate, but we will *mostly neglect* this issue.



- $V_t$  is typically modelled as a function  $f$  of time  $t$  and a  $d$ -dimensional random vector  $\mathbf{Z} = (Z_{t,1}, \dots, Z_{t,d})$  of risk factors, that is,

$$V_t = f(t, \mathbf{Z}_t) \quad (\text{mapping of risks})$$

for some measurable  $f : \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}$ . The choice of  $f$  and  $\mathbf{Z}_t$  is problem-specific (but typically known).

- It is often convenient to work with the risk-factor changes

$$\mathbf{X}_{t+1} = \mathbf{Z}_{t+1} - \mathbf{Z}_t.$$

We can rewrite  $L_{t+1}$  in terms of  $\mathbf{X}_t$  via

$$\begin{aligned} L_{t+1} &= -(V_{t+1} - V_t) = -(f(t+1, \mathbf{Z}_{t+1}) - f(t, \mathbf{Z}_t)) \\ &= -(f(t+1, \mathbf{Z}_t + \mathbf{X}_{t+1}) - f(t, \mathbf{Z}_t)). \end{aligned}$$

We see that the loss df is determined by the loss df of  $\mathbf{X}_{t+1}$ . We will thus also write  $L_{t+1} = L(\mathbf{X}_{t+1})$ , where  $L(\mathbf{x}) = -(f(t+1, \mathbf{Z}_t + \mathbf{x}) - f(t, \mathbf{Z}_t))$  is known as loss operator.

- If  $f$  is differentiable, its **first-order (Taylor) approximation** ( $f(\mathbf{y}) \approx f(\mathbf{y}_0) + \nabla f(\mathbf{y}_0)(\mathbf{y} - \mathbf{y}_0)$  for  $\mathbf{y} = (t+1, Z_{t,1} + X_{t+1,1}, \dots, Z_{t,d} + X_{t+1,d})$  and  $\mathbf{y}_0 = (t, Z_{t,1}, \dots, Z_{t,d})$ ) is

$$f(t+1, \mathbf{Z}_t + \mathbf{X}_{t+1}) \approx f(t, \mathbf{Z}_t) + f_t(t, \mathbf{Z}_t) \cdot 1 + \sum_{j=1}^d f_{z_j}(t, \mathbf{Z}_t) \cdot X_{t+1,j}$$

We can thus approximate  $L_{t+1}$  by the **linearized loss**

$$L_{t+1}^{\Delta} = - \left( \underbrace{f_t(t, \mathbf{Z}_t)}_{=: c_t} + \sum_{j=1}^d \underbrace{f_{z_j}(t, \mathbf{Z}_t) X_{t+1,j}}_{=: b_{t,j}} \right) = -(c_t + \mathbf{b}'_t \mathbf{X}_{t+1}),$$

a linear function of  $X_{t+1,1}, \dots, X_{t+1,d}$  (indices denote partial derivatives). The **approximation is best if the risk-factor changes are small in absolute value**.

## Example 2.2 (Stock portfolio)

Consider a **portfolio**  $\mathcal{P}$  of  $d$  **stocks**  $S_{t,1}, \dots, S_{t,d}$  ( $S_{t,j}$  = value of stock  $j$  at time  $t$ ) and denote by  $\lambda_j$  the **number of shares** of stock  $j$  in  $\mathcal{P}$ . In finance and risk management, one typically uses **logarithmic prices as risk factors**, i.e.  $Z_{t,j} = \log S_{t,j}$ ,  $j \in \{1, \dots, d\}$ . Then

$$V_t = f(t, \mathbf{Z}_t) = \sum_{j=1}^d \lambda_j S_{t,j} = \sum_{j=1}^d \lambda_j e^{Z_{t,j}}.$$

- The one-period ahead loss is then given by

$$\begin{aligned} L_{t+1} &= -(V_{t+1} - V_t) = -\sum_{j=1}^d \lambda_j (e^{Z_{t,j} + X_{t+1,j}} - e^{Z_{t,j}}) \\ &= -\sum_{j=1}^d \lambda_j e^{Z_{t,j}} (e^{X_{t+1,j}} - 1) = -\sum_{j=1}^d \underbrace{\lambda_j S_{t,j}}_{=: w_{t,j}} (e^{X_{t+1,j}} - 1) \quad (1) \end{aligned}$$

which is non-linear in  $X_{t+1,j}$  (here:  $L(\mathbf{x}) = -\sum_{j=1}^d w_{t,j} (e^{x_j} - 1)$ ).

- With  $f_{z_j}(t, \mathbf{Z}_t) = \lambda_j e^{Z_{t,j}} = \lambda_j S_{t,j} = w_{t,j}$ , the **linearized loss** is

$$\begin{aligned} L_{t+1}^\Delta &= -\left(f_t(t, \mathbf{Z}_t) + \sum_{j=1}^d f_{z_j}(t, \mathbf{Z}_t) X_{t+1,j}\right) = -\left(0 + \sum_{j=1}^d w_{t,j} X_{t+1,j}\right) \\ &= -\mathbf{w}'_t \mathbf{X}_{t+1}. \end{aligned}$$

- Note that  $L_{t+1}^\Delta = -(c_t + \mathbf{b}'_t \mathbf{X}_{t+1})$  for  $c_t = 0$  and  $\mathbf{b}_t = \mathbf{w}_t$ .
- If  $\boldsymbol{\mu} = \mathbb{E} \mathbf{X}_{t+1}$  and  $\Sigma = \text{cov } \mathbf{X}_{t+1}$  are known, then **expectation** and **variance of the (linearized) one-period ahead loss** are

$$\begin{aligned} \mathbb{E} L_{t+1}^\Delta &= -\sum_{j=1}^d w_{t,j} \mathbb{E}(X_{t+1,j}) = -\mathbf{w}'_t \boldsymbol{\mu}, \\ \text{var } L_{t+1}^\Delta &= \text{var}(\mathbf{w}'_t \mathbf{X}_{t+1}) = \mathbf{w}'_t \text{cov}(\mathbf{X}_{t+1}) \mathbf{w}_t = \mathbf{w}'_t \Sigma \mathbf{w}_t. \end{aligned}$$

## 2.1.2 Loss distributions

Having determined the mapping  $f$  (may involve *valuation models*, e.g. Black–Scholes, or numerical approximation), we can identify the following **key statistical tasks of QRM**:

- 1) Find a statistical **model for  $\mathbf{X}_{t+1}$**  (typically a model for forecasting  $\mathbf{X}_{t+1}$ , estimated based on historical data);
- 2) Compute/derive the **df  $F_{L_{t+1}}$**  (requires the df of  $f(t+1, \mathbf{Z}_t + \mathbf{X}_{t+1})$ );
- 3) Compute a **risk measure from  $F_{L_{t+1}}$** .

There are **three general methods** to approach these challenges.

### 1) Analytical method

**Idea:** Choose  $F_{\mathbf{X}_{t+1}}$  and  $f$  such that  $F_{L_{t+1}}$  can be determined explicitly.

Prime example: *Variance-covariance method*, see RiskMetrics (1996):

**Assumption 1**  $\mathbf{X}_{t+1} \sim N(\boldsymbol{\mu}, \Sigma)$  (e.g. if  $(\mathbf{Z}_t)$  is a Brownian motion,  $(S_t)$  a geometric Brownian motion)

**Assumption 2**  $F_{L_{t+1}^\Delta}$  is a good approximation to  $F_{L_{t+1}}$ .

$$L_{t+1}^\Delta = -(c_t + \mathbf{b}_t' \mathbf{X}_{t+1}) \underset{\text{Ass. 1}}{\Rightarrow} L_{t+1}^\Delta \sim N(-c_t - \mathbf{b}_t' \boldsymbol{\mu}, \mathbf{b}_t' \Sigma \mathbf{b}_t).$$

**Advantages:**

- $F_{L_{t+1}^\Delta}$  explicit ( $\Rightarrow$  typically explicit risk measures)
- Easy to implement

**Drawbacks:** Assumption 1 is unlikely to be realistic for daily (probably also weekly/monthly) data.

## 2) Historical simulation

**Idea:** Estimate  $F_{L_{t+1}}$  by its *empirical distribution function (edf)*

$$\hat{F}_{L_{t+1},n}(x) = \frac{1}{n} \sum_{i=1}^n I_{\{\tilde{L}_{t-i+1} \leq x\}}, \quad x \in \mathbb{R}, \quad (2)$$

based on

$$\tilde{L}_k = L(\mathbf{X}_k) = -(f(t+1, \mathbf{Z}_t + \mathbf{X}_k) - f(t, \mathbf{Z}_t)), \quad (3)$$

$k \in \{t-n+1, \dots, t\}$ .  $\tilde{L}_{t-n+1}, \dots, \tilde{L}_t$  show what would happen to the current portfolio if the past  $n$  risk-factor changes were to recur.

Advantages: ■ Easy to implement

- No estimation of the distribution of  $\mathbf{X}_{t+1}$  required

Drawbacks: ■ Sufficient data for all risk-factor changes required

- Only past losses considered (“driving a car by looking in the back mirror”)

### 3) Monte Carlo method

**Idea:** Take any model for  $X_{t+1}$ , simulate  $X_{t+1}$ , compute the corresponding losses as in (3) and estimate  $F_{L_{t+1}}$  (typically via edf as in (2)).

**Advantages:** ■ Quite general (applicable to any model of  $X_{t+1}$  which is easy to sample)

**Drawbacks:** ■ Unclear how to find an appropriate model for  $X_{t+1}$  (any result is only as good as the chosen  $F_{X_{t+1}}$ )

■ Computational cost (every simulation requires to evaluate the mapping  $f$ ; expensive, e.g. if the latter contains derivatives which are priced via Monte Carlo themselves  $\Rightarrow$  Nested Monte Carlo simulations)

So-called *economic scenario generators* used in insurance also fall under the heading of Monte Carlo methods.



## 2.2 Risk measurement

- A *risk measure* for a financial position with (random) loss  $L$  is a *real number* which measures the “riskiness of  $L$ ”. In the Basel or Solvency context, it is often interpreted as the amount of *capital required to make a position with loss  $L$  acceptable* to an (internal/external) regulator.
- Some *reasons for using risk measures* in practice:
  - ▶ To *determine the amount of capital to hold* as a buffer against unexpected future losses on a portfolio (in order to satisfy a regulator/manager concerned with the institution’s solvency).
  - ▶ As a *tool for limiting* the amount of *risk of a business unit* (e.g. by requiring that the daily 95% value-at-risk (i.e. the 95%-quantile) of a trader’s position should not exceed a given bound).
  - ▶ To determine the *riskiness* (and *thus fair premium*) of an *insurance contract*.

## 2.2.1 Approaches to risk measurement

Existing risk measurement approaches grouped into three categories:

### 1) Notional-amount approach

- oldest approach; “standardized approaches” of Basel II (e.g. OpRisk)
- *risk of a portfolio* = summed notational values of the securities times their riskiness factor

### 2) Risk measures based on loss distributions

- Most modern risk measures are characteristics of the underlying loss distribution over some predetermined time horizon  $\Delta t$ .
- Examples: variance, value-at-risk, expected shortfall (see later)
- Advantages:
  - ▶ Makes sense on all levels (from single portfolios to the overall position of a financial institution).
  - ▶ Loss distributions reflect netting and diversification.

### 3) Scenario-based risk measures

- Typically considered in stress testing.
- One considers possible future risk-factor changes (*scenarios*; e.g. a 20% drop in a market index).
- *Risk of a portfolio* = maximum (weighted) loss under all scenarios.
- If  $\mathcal{X} = \{\mathbf{x}_1, \dots, \mathbf{x}_n\}$  denote the risk-factor changes (*scenarios*) with corresponding weights  $\mathbf{w} = (w_1, \dots, w_n)$ , the risk is

$$\psi_{\mathcal{X}, \mathbf{w}} = \max_{1 \leq i \leq n} \{w_i L(\mathbf{x}_i)\}, \quad (4)$$

where  $L(\mathbf{x})$  denotes the loss the portfolio would suffer if the hypothetical scenario  $\mathbf{x}$  were to occur. Many risk measures are of the form (4); see *CME SPAN: Standard Portfolio Analysis of Risk* (2010).

- Mathematical interpretation of (4):
  - Assume  $L(\mathbf{0}) = 0$  (okay if  $\Delta t$  small) and  $w_i \in [0, 1] \ \forall i$ .

- ▶  $w_i L(\mathbf{x}_i) = w_i L(\mathbf{x}_i) + (1 - w_i) L(\mathbf{0}) = \mathbb{E}_{\mathbb{P}_i}(L(\mathbf{X}_i))$  where  $\mathbf{X}_i \sim \mathbb{P}_i = w_i \delta_{\mathbf{x}_i} + (1 - w_i) \delta_{\mathbf{0}}$  ( $\delta_{\mathbf{x}}$  the Dirac measure at  $\mathbf{x}$ ) is a probability measure on  $\mathbb{R}^d$ .

Therefore,  $\psi_{\mathcal{X}, w} = \max\{\mathbb{E}_{\mathbb{P}}(L(\mathbf{X})) : \mathbf{X} \sim \mathbb{P} \in \{\mathbb{P}_1, \dots, \mathbb{P}_n\}\}$ .  
Such a risk measure is known as a *generalized scenario*.

- **Advantages:**
  - ▶ Useful for portfolios with few risk factors.
  - ▶ Useful complementary information to risk measures based on loss distributions (past data).


**Drawbacks:** ▶ Determining scenarios and weights.

## 2.2.2 Value-at-risk

### Definition 2.3 (Value-at-risk)

For a loss  $L \sim F_L$ , *value-at-risk (VaR)* at confidence level  $\alpha \in (0, 1)$  is defined by  $\text{VaR}_\alpha = \text{VaR}_\alpha(L) = F_L^{\leftarrow}(\alpha) = \inf\{x \in \mathbb{R} : F_L(x) \geq \alpha\}$ .

- $\text{VaR}_\alpha$  is simply the  $\alpha$ -quantile of  $F_L$ . As such,  $F_L(x) < \alpha$  for all  $x < \text{VaR}_\alpha(L)$  and  $F_L(\text{VaR}_\alpha(L)) = F_L(F_L^\leftarrow(\alpha)) \geq \alpha$ .
- Known since 1994: Weatherstone 4<sup>15</sup> report (J.P. Morgan; RiskMetrics)
- VaR is the most widely used risk measure (by Basel II or Solvency II)
- $\text{VaR}_\alpha(L)$  is not a what if risk measure: It does not provide information about the severity of losses which occur with probability  $\leq 1 - \alpha$ .



### Example 2.4 ( $\text{VaR}_\alpha$ for $N(\mu, \sigma^2)$ and $t_\nu(\mu, \sigma^2)$ )

1) Let  $L \sim N(\mu, \sigma^2)$ . Then

$$F_L(x) = \mathbb{P}(L \leq x) = \mathbb{P}((L - \mu)/\sigma \leq (x - \mu)/\sigma) = \Phi((x - \mu)/\sigma).$$

This implies that

$$\text{VaR}_\alpha(L) = F_L^{\leftarrow}(\alpha) = F_L^{-1}(\alpha) = \mu + \sigma\Phi^{-1}(\alpha).$$

Check:  $F_L(\text{VaR}_\alpha(L)) = \Phi(((\mu + \sigma\Phi^{-1}(\alpha)) - \mu)/\sigma) = \alpha$ .

2) Let  $L \sim t_\nu(\mu, \sigma^2)$ , so  $(L - \mu)/\sigma \sim t_\nu$  and thus, as above,

$$\text{VaR}_\alpha(L) = \mu + \sigma t_\nu^{-1}(\alpha).$$

Note that  $X \sim t_\nu = t_\nu(0, 1)$  has density

$$f_X(x) = \frac{\Gamma((\nu + 1)/2)}{\sqrt{\nu\pi}\Gamma(\nu/2)} (1 + x^2/\nu)^{-\frac{\nu+1}{2}}.$$

Furthermore, if  $\nu > 1$ ,  $\mathbb{E}X$  exists and  $\mathbb{E}X = 0$ ; and if  $\nu > 2$ ,  $\text{var } X$  exists and  $\text{var } X = \frac{\nu}{\nu-2}$ .

## Choices of parameters $\Delta t, \alpha$ :

- $\Delta t$  should reflect the time period over which the portfolio is held (unchanged) (e.g. insurance contracts:  $\Delta t = 1 \text{ y}$ )
- $\Delta t$  should be relatively small (more risk-factor change data is available).
- Typical choices:
  - ▶ According to Basel II:
    - Market risk:  $\alpha = 0.99$ ,  $\Delta t = 10 \text{ d}$  (2 trading weeks)
    - Credit risk and operational risk:  $\alpha = 0.999$ ,  $\Delta t = 1 \text{ y}$
  - ▶ According to Solvency II:  $\alpha = 0.995$ ,  $\Delta t = 1 \text{ y}$
- Backtesting often needs to be carried out at lower confidence levels in order to have sufficient statistical power to detect poor models.
- Be cautious with strictly interpreting  $\text{VaR}_\alpha(L)$  (and other risk measure) estimates (considerable model/liquidity risk).

## Interlude: Generalized inverses

$T \nearrow$  means that  $T$  is *increasing*, i.e.  $T(x) \leq T(y)$  for all  $x < y$ .  $T \uparrow$  means that  $T$  is *strictly increasing*, i.e.  $T(x) < T(y)$  for all  $x < y$ .

### Definition 2.5 (Generalized inverse)

For any increasing function  $T : \mathbb{R} \rightarrow \mathbb{R}$ , with  $T(-\infty) = \lim_{x \downarrow -\infty} T(x)$  and  $T(\infty) = \lim_{x \uparrow \infty} T(x)$ , the *generalized inverse*  $T^{\leftarrow} : \mathbb{R} \rightarrow \bar{\mathbb{R}} = [-\infty, \infty]$  of  $T$  is defined by


$$T^{\leftarrow}(y) = \inf\{x \in \mathbb{R} : T(x) \geq y\}, \quad y \in \mathbb{R},$$

with the convention that  $\inf \emptyset = \infty$ . If  $T$  is a df,  $T^{\leftarrow} : [0, 1] \rightarrow \bar{\mathbb{R}}$  is the *quantile function* of  $T$ .

- If  $T$  is *continuous and  $\uparrow$* , then  $T^{\leftarrow} \equiv T^{-1}$  (ordinary inverse).
- There are *rules for working with  $T^{\leftarrow}$*  similar to  $T^{-1}$ .



How to determine  $F_L^{\leftarrow}$  from  $F_L$ :



$\Rightarrow$  Flat parts of  $F_L$  correspond to jumps of  $F_L^{\leftarrow}$ ; Jumps of  $F_L$  correspond to flat parts of  $F_L^{\leftarrow}$ .

## 2.2.3 VaR in risk capital calculations

### 1) VaR in regulatory capital calculations for the trading book

For banks using the *internal model (IM)* approach for market risk in Basel II, the daily risk capital formula is

$$RC^t = \max \left\{ VaR_{0.99}^{t,10}, \frac{k}{60} \sum_{i=1}^{60} VaR_{0.99}^{t-i+1,10} \right\} + c.$$

- $VaR_{\alpha}^{s,10}$  denotes the 10-day  $VaR_{\alpha}$  calculated at day  $s$  ( $t = \text{today}$ ).
- $k \in [3, 4]$  is a multiplier (or *stress factor*).
- $c = \text{stressed VaR charge}$  (calculated from data from a volatile market period) + *incremental risk charge (IRC;  $VaR_{0.999}$ -estimate of the annual distribution of losses due to defaults and downgrades)* + *charges for specific risks*.

The averaging tends to lead to smooth changes in the capital charge over time unless  $VaR_{0.99}^{t,10}$  is very large.

## 2) The Solvency Capital Requirement in Solvency II

The *Solvency Capital Requirement (SCR)* is the amount of capital that enables the insurer to meet its obligations over  $\Delta t = 1$  y with  $\alpha = 0.995$ . Let  $V_t = A_t - B_t$  denote the equity capital. The insurer wants to determine the minimum amount of extra capital  $x_0$  to put aside to be solvent in  $\Delta t$  with probability  $(\geq)\alpha$ . So

$$\begin{aligned}x_0 &= \inf\{x \in \mathbb{R} : \mathbb{P}(V_{t+1} + x(1+r) \geq 0) \geq \alpha\} \\&= \inf\left\{x \in \mathbb{R} : \mathbb{P}\left(-\left(\frac{V_{t+1}}{1+r} - V_t\right) \leq x + V_t\right) \geq \alpha\right\} \\&= \inf\{x \in \mathbb{R} : \mathbb{P}(L_{t+1} \leq x + V_t) \geq \alpha\} \\&= \inf\{x \in \mathbb{R} : F_{L_{t+1}}(x + V_t) \geq \alpha\} \\&= \inf\{z - V_t \in \mathbb{R} : F_{L_{t+1}}(z) \geq \alpha\} = \text{VaR}_\alpha(L_{t+1}) - V_t\end{aligned}$$

and thus  $\text{SCR} = V_t + x_0 = \text{VaR}_\alpha(L_{t+1})$  (available capital now + capital required to be solvent in  $\Delta t$  with probability  $\geq \alpha$ ). If  $x_0 < 0$ , the company is already well capitalized.

## 2.2.4 Expected shortfall

### Definition 2.6 (Expected shortfall)

For a loss  $L \sim F_L$  with  $\mathbb{E}|L| < \infty$ , *expected shortfall (ES)* at confidence level  $\alpha \in (0, 1)$  is defined by

$$\text{ES}_\alpha = \text{ES}_\alpha(L) = \frac{1}{1 - \alpha} \int_\alpha^1 \text{VaR}_u(L) du. \quad (5)$$

- $\text{ES}_\alpha$  is the **average over  $\text{VaR}_u$**  for all  $u \geq \alpha \Rightarrow \text{ES}_\alpha \geq \text{VaR}_\alpha$ .
- Besides VaR, ES is the **most important risk measure** in practice.
- $\text{ES}_\alpha$  **looks further into the tail** of  $F_L$ , it is a “what if” risk measure ( $\text{VaR}_\alpha$  is **frequency**-based;  $\text{ES}_\alpha$  is **severity**-based).
- $\text{ES}_\alpha$  is **more difficult to estimate and backtest** than  $\text{VaR}_\alpha$  (the variance of estimators is typically larger; larger sample size required).
- $\text{ES}_\alpha(L) < \infty$  **requires  $\mathbb{E}|L| < \infty$** .
- **Subadditivity** and **elicitability**. One can show:

- ▶ In contrast to  $\text{VaR}_\alpha$ ,  $\text{ES}_\alpha$  is subadditive.
- ▶ In contrast to  $\text{ES}_\alpha$ ,  $\text{VaR}_\alpha$  exists if  $\mathbb{E}|L| = \infty$  and is elicitable (i.e., minimizes a certain expected functional (expected scoring function); see Gneiting (2011). This can be used for backtesting).


### Example 2.7 (A comparison of VaR and ES for stock returns)

- Consider Example 2.2 with a portfolio consisting of a single stock  $V_t = S_t = 10\,000$ . In this case,  $L_{t+1}^\Delta = -S_t X_{t+1}$ , where  $X_{t+1} = \log(S_{t+1}/S_t)$ .
- Let  $\sigma = 0.2/\sqrt{250}$  (annualized volatility of 20%) and assume
  - 1)  $X_{t+1} \sim N(0, \sigma^2) \Rightarrow L_{t+1}^\Delta \sim N(0, S_t^2 \sigma^2)$ ;
  - 2)  $X_{t+1} \sim t_\nu(0, \sigma^2 \frac{\nu-2}{\nu})$ ,  $\nu > 2$  (so that  $\text{var } X_{t+1}$  is also  $\sigma^2$ ). Then

$$X_{t+1} = \sqrt{\sigma^2 \frac{\nu-2}{\nu}} Y \quad \text{for } Y \sim t_\nu.$$

$$\Rightarrow L_{t+1}^\Delta = -S_t \sqrt{\sigma^2 \frac{\nu-2}{\nu}} Y \sim t_\nu(0, S_t^2 \sigma^2 \frac{\nu-2}{\nu}) \quad (\text{var}(L_{t+1}^\Delta) = S_t^2 \sigma^2).$$

Consider  $\nu = 4$  and note that  $\text{VaR}_\alpha^{t_4} \geq \text{VaR}_\alpha^{\text{normal}}$  and  $\text{ES}_\alpha^{t_4} \geq \text{ES}_\alpha^{\text{normal}}$  only hold for sufficiently large  $\alpha$ .



$\Rightarrow$  The  $t_4$  model is not always “riskier” than the normal model.

**Example 2.8 (Example 2.4 continued;  $\text{ES}_\alpha$  for  $N(\mu, \sigma^2)$  and  $t_\nu(\mu, \sigma^2)$ )**

1) Let  $\tilde{L} \sim N(0, 1)$ . Then  $\text{VaR}_\alpha(\tilde{L}) = 0 + 1 \cdot \Phi^{-1}(\alpha)$  and thus

$$\text{ES}_\alpha(\tilde{L}) = \frac{1}{1-\alpha} \int_\alpha^1 \Phi^{-1}(u) du \stackrel{x=\Phi^{-1}(u)}{=} \frac{1}{1-\alpha} \int_{\Phi^{-1}(\alpha)}^\infty x \varphi(x) dx,$$

where  $\varphi(x) = \Phi'(x) = \exp(-x^2/2)/\sqrt{2\pi}$ . Note that  $x\varphi(x) = -\varphi'(x)$ , so that

$$\text{ES}_\alpha(\tilde{L}) = \frac{-[\varphi(x)]_{\Phi^{-1}(\alpha)}^\infty}{1-\alpha} = \frac{-(0 - \varphi(\Phi^{-1}(\alpha)))}{1-\alpha} = \frac{\varphi(\Phi^{-1}(\alpha))}{1-\alpha}.$$

This implies that  $L \sim N(\mu, \sigma^2)$  has expected shortfall

$$\text{ES}_\alpha(L) = \mu + \sigma \text{ES}_\alpha(\tilde{L}) = \mu + \sigma \frac{\varphi(\Phi^{-1}(\alpha))}{1-\alpha}.$$

2) Let  $L \sim t_\nu(\mu, \sigma^2)$ ,  $\nu > 1$ . Similarly as above, one obtains that

$$\text{ES}_\alpha(L) = \mu + \sigma \frac{f_{t_\nu}(t_\nu^{-1}(\alpha))(\nu + t_\nu^{-1}(\alpha)^2)}{(1-\alpha)(\nu-1)},$$

where  $f_{t_\nu}$  denotes the density of  $t_\nu$ ; see Example 2.4.

By l'Hôpital's Rule (case "0/0"), one can show that

$$1 \leq \lim_{\alpha \uparrow 1} \frac{\text{ES}_\alpha(L)}{\text{VaR}_\alpha(L)} = \frac{\nu}{\nu - 1}.$$

- In finance, often  $\nu \in (3, 5)$ . With  $\nu = 3$ ,  $\text{ES}_\alpha(L)$  is 50% larger than  $\text{VaR}_\alpha(L)$  (in the limit for large  $\alpha$ ).
- For  $\nu \uparrow \infty$ ,  $\lim_{\alpha \uparrow 1} \frac{\text{ES}_\alpha(L)}{\text{VaR}_\alpha(L)} \downarrow 1$ ; for  $\nu \downarrow 1$ ,  $\lim_{\alpha \uparrow 1} \frac{\text{ES}_\alpha(L)}{\text{VaR}_\alpha(L)} \uparrow \infty$ .

### Conclusion:

For losses with *heavy tails* (power-like), the difference between using VaR and ES as risk measures for computing risk capital can be huge (for large  $\alpha$  as required by Basel II).

### Proposition 2.9 ( $\text{ES}_\alpha(L)$ under continuity)

If  $F_L$  is continuous,  $\text{ES}_\alpha(L) = \mathbb{E}(L \mid L > \text{VaR}_\alpha(L))$ .



## 2.2.5 Coherent and convex risk measures

- Artzner et al. (1999) (coherent risk measures) and Föllmer and Schied (2002) (convex risk measures) propose **axioms of a good risk measure**.
- Assume that **risk measures**  $\varrho$  are defined on a **linear space of random variables**  $\mathcal{M}$  (including constants; we can thus add rvs, multiply them with constants etc.), so  $\varrho : \mathcal{M} \rightarrow \mathbb{R}$ .
- There are **two possible interpretations** of elements of  $\mathcal{M}$ :
  - 1) **Elements of  $\mathcal{M}$  are random variables**  $V_{t+1}$ :  $\tilde{\varrho}(V_{t+1})$  denotes the amount of **additional capital** that needs to be added to a position with future value  $V_{t+1}$  to make it acceptable to a regulator.
  - 2) **Elements of  $\mathcal{M}$  are losses**  $L_{t+1} = -(V_{t+1} - V_t)$ :  $\varrho(L_{t+1})$  denotes the **total amount of capital** necessary to back a position with loss  $L$ .
- 1) and 2) are **related via**  $\varrho(L_{t+1}) = V_t + \tilde{\varrho}(V_{t+1})$  (total capital = available capital + additional capital). **We focus on 2) and drop “ $t + 1$ ”.**

## Axioms of coherence

**Axiom 1** (**monotonicity**)  $L_1, L_2 \in \mathcal{M}$ ,  $L_1 \leq L_2$  (a.s., i.e. almost surely)  
 $\Rightarrow \varrho(L_1) \leq \varrho(L_2)$

**Interpr.:** Positions which lead to a higher loss in every state of the world require more risk capital.

**Criticism:** none

**Axiom 2** (**translation invar.**)  $\varrho(L + l) = \varrho(L) + l$  for all  $L \in \mathcal{M}$ ,  $l \in \mathbb{R}$

**Interpr.:**

- By adding  $l \in \mathbb{R}$  to a position with loss  $L$ , we alter the capital requirements accordingly.
- If  $\varrho(L) > 0$ , and  $l = -\varrho(L)$ , then  $\varrho(L - \varrho(L)) = \varrho(L + l) = \varrho(L) + l = 0$  so that adding  $\varrho(L)$  to a position with loss  $L$  makes it acceptable.

**Criticism:** Most people believe this to be reasonable.

**Axiom 3** (**subadditivity**)  $\varrho(L_1 + L_2) \leq \varrho(L_1) + \varrho(L_2)$  for all  $L_1, L_2 \in \mathcal{M}$

- Interpr.:**
- Reflects the idea of **diversification**
  - Using a non-subadditive (that is, a **superadditive**)  $\varrho$  encourages institutions to legally break up into subsidiaries to reduce regulatory capital requirements.
  - Subadditivity makes decentralization possible: Assume  $L = L_1 + L_2$  and that we want to bound  $\varrho(L)$  by  $M$ . Choose  $M_j$  such that  $\varrho(L_j) \leq M_j$ ,  $j \in \{1, 2\}$ , and  $M_1 + M_2 \leq M$ . Then  $\varrho(L) \leq_{\text{subadd.}} \varrho(L_1) + \varrho(L_2) \leq M_1 + M_2 \leq M$ .

**Criticism:** **VaR is ruled out** under certain scenarios (see later).  
**VaR is monotone, translation invariant, and positive homogeneous**, but in general **not subadditive**.

**Axiom 4** (positive homogeneity)  $\varrho(\lambda L) = \lambda \varrho(L)$  for all  $L \in \mathcal{M}$ ,  $\lambda > 0$

Interpr.: (or motivation):  $\lambda = n \in \mathbb{N}$  and subadditivity imply  $\varrho(nL) \leq n\varrho(L)$ , but  $n$  times the same loss  $L$  means no diversification, so equality should hold.

Criticism: If  $\lambda > 0$  is large, liquidity risk plays a role and one should rather have  $\varrho(\lambda L) > \lambda \varrho(L)$  (also to penalize concentration or risk), but this contradicts subadditivity. This has led to convex risk measures, i.e. risk measures  $\varrho$  satisfying

$$\varrho(\lambda L_1 + (1 - \lambda)L_2) \leq \lambda \varrho(L_1) + (1 - \lambda)\varrho(L_2)$$

for all  $L_1, L_2 \in \mathcal{M}$ ,  $0 \leq \lambda \leq 1$ .

### Definition 2.10 (Coherent risk measure)

A risk measure  $\varrho$  which satisfies Axioms 1–4 is called *coherent*.

### Example 2.11 (Generalized scenario risk measures)

Let  $L(x)$  denote the hypothetical loss under scenario  $x$  (risk-factor change).

The generalized scenario risk measure

$$\psi_{\mathcal{X},w}(L) = \max\{\mathbb{E}_{\mathbb{P}}(L(\mathbf{X})) : \mathbf{X} \sim \mathbb{P} \in \{\mathbb{P}_1, \dots, \mathbb{P}_n\}\}$$

is coherent. Monotonicity, translation invariance, positive homogeneity are clear (by monotonicity and linearity of  $\mathbb{E}_{\mathbb{P}}(\cdot)$ ); for subadditivity, note that

$$\begin{aligned}\psi_{\mathcal{X},w}(L_1 + L_2) &= \max\{\underbrace{\mathbb{E}_{\mathbb{P}}(L_1(\mathbf{X}) + L_2(\mathbf{X}))}_{=\mathbb{E}_{\mathbb{P}}(L_1(\mathbf{X})) + \mathbb{E}_{\mathbb{P}}(L_2(\mathbf{X}))} : \mathbf{X} \sim \mathbb{P} \in \{\mathbb{P}_1, \dots, \mathbb{P}_n\}\} \\ &\leq \psi_{\mathcal{X},w}(L_1) + \psi_{\mathcal{X},w}(L_2).\end{aligned}$$

One can show that all coherent risk measures can be represented as generalized scenarios via

$$\varrho(L) = \sup\{\mathbb{E}_{\mathbb{P}}(L) : \mathbb{P} \in \mathcal{P}\}$$

for a suitable set  $\mathcal{P}$  of probability measures.

### Definition 2.12 (Convex risk measure)

A risk measure  $\rho$  which is **monotone**, **translation invariant** and **convex** is called a **convex risk measure**.

- Justification: Again diversification but they don't have to be positive homogeneous.
- Any coherent risk measure is also a convex risk measure. The **converse is not true in general**, but for positive homogeneous risk measures, convexity and subadditivity are equivalent.

### Theorem 2.13 (Coherence of ES)

**ES is a coherent risk measure.**

*Proof.* **Monotonicity**, **translation invariance** and **positive homogeneity** follow from VaR. **Subadditivity** is more involved but can be shown in various ways. □

## Superadditivity scenarios for VaR

Under the following scenarios,  $\text{VaR}_\alpha$  is typically **superadditive**:

- 1)  $L_1, L_2$  have **skewed** distributions;
- 2) **Independent, light-tailed**  $L_1, L_2$  and **small**  $\alpha$ ;
- 3)  $L_1, L_2$  have **special dependence**;
- 4)  $L_1, L_2$  have **heavy tailed** distributions.

### Exercise 2.14 (Skewed loss distributions)

Consider a portfolio of two independently defaultable zero-coupon bonds (maturity  $T = 1y$ , nominal/face value 100, paid interest of 5%, default probability  $p = 0.009$ , no recovery). The loss of bond  $j$  (from the money lender's/investor's perspective) is thus

$$L_j = \begin{cases} -5, & \text{with prob. } 1 - p = 0.991, \\ 100, & \text{with prob. } p = 0.009, \end{cases} \quad j \in \{1, 2\}.$$

Set  $\alpha = 0.99$ . Then  $\text{VaR}_\alpha(L_j) = -5$ ,  $j \in \{1, 2\}$ .

The loss  $L_1 + L_2$  is given by


$$L_1 + L_2 = \begin{cases} -10, & \text{with prob. } (1 - p)^2 = 0.982081, \\ 95, & \text{with prob. } 2p(1 - p) = 0.017838, \\ 200, & \text{with prob. } p^2 = 0.000081. \end{cases}$$

Therefore,  $\text{VaR}_\alpha(L_1 + L_2) = 95 > -10 = \text{VaR}_\alpha(L_1) + \text{VaR}_\alpha(L_2)$ . Hence  $\text{VaR}_\alpha$  is superadditive.



## Exercise 2.15 (Independent, light-tailed $L_1, L_2$ and small $\alpha$ )

If  $L_1, L_2 \stackrel{\text{ind.}}{\sim} \text{Exp}(1)$ ,  $\text{VaR}_\alpha$  is superadditive  $\iff \alpha < 0.71$ .



## Exercise 2.16 (Special dependence)

Let  $\alpha \in (0, 1)$ ,  $L_1 \sim U(0, 1)$  and define  $L_2 \stackrel{\text{a.s.}}{=} \begin{cases} L_1, & \text{if } L_1 < \alpha, \\ 1 + \alpha - L_1, & \text{if } L_1 \geq \alpha. \end{cases}$

One can show that  $L_2 \sim U(0, 1)$ . Also,  $L_1 + L_2 = \begin{cases} 2L_1, & \text{if } L_1 < \alpha, \\ 1 + \alpha, & \text{if } L_1 \geq \alpha, \end{cases}$   
from which one can show that

$$F_{L_1+L_2}(x) = \begin{cases} 0, & \text{if } x < 0, \\ x/2, & \text{if } x \in [0, 2\alpha), \\ \alpha, & \text{if } x \in [2\alpha, 1 + \alpha), \\ 1, & \text{if } x \geq 1 + \alpha. \end{cases}$$

For all  $\varepsilon \in (0, \frac{1-\alpha}{2})$ , we thus obtain that

$$\text{VaR}_{\alpha+\varepsilon}(L_1 + L_2) = 1 + \alpha > \underset{\varepsilon \in (0, \frac{1-\alpha}{2})}{2(\alpha + \varepsilon)} = \text{VaR}_{\alpha+\varepsilon}(L_1) + \text{VaR}_{\alpha+\varepsilon}(L_2).$$

## Remark 2.17 (Special case of comonotone risks; elliptical risks)

- Note that  $L_1 \stackrel{\text{a.s.}}{=} L_2$  does not lead to the largest  $\text{VaR}_\alpha(L_1 + L_2)$  since

$$\text{VaR}_\alpha(L_1 + L_2) \stackrel[\text{hom.}]{\text{pos.}} 2 \text{VaR}_\alpha(L_1) = \text{VaR}_\alpha(L_1) + \text{VaR}_\alpha(L_2),$$

so “only” equality.

- $\text{VaR}_\alpha$  is subadditive and thus coherent for a certain class of multivariate distributions (strictly including the multivariate normal and  $t$ ); see later.

## Exercise 2.18 (Heavy tailed loss distributions)

Let  $L_1, L_2 \stackrel{\text{ind.}}{\sim} F(x) = 1 - x^{-1/2}$ ,  $x \in [1, \infty)$ . By deriving the distribution function

$$F_{L_1+L_2}(x) = 1 - 2\sqrt{x-1}/x, \quad x \geq 2,$$

of  $L_1 + L_2$  (via the density convolution formula; tedious), one can show (via solving a quadratic equation) that  $\text{VaR}_\alpha$  is superadditive for all  $\alpha \in (0, 1)$ .

# 3 Empirical properties of financial data

3.1 Stylized facts of financial return series

3.2 Multivariate stylized facts


## 3.1 Stylized facts of financial return series

- The **stylized facts** are a collection of **empirical observations** and **related inferences**, which apply to many time series of risk-factor changes (e.g. log-returns on equities, indices, exchange rates, commodity prices).
- The best-known stylized facts **apply to daily log-returns** (also to intra-daily, weekly, monthly). Tick-by-tick (**high-frequency**) data **have their own stylized facts** (not discussed here) and annual return (**low-frequency**) data **are more difficult** to investigate (**data sparseness**; non-stationarity).
- Consider **discrete-time risk-factor changes**  $X_t = Z_t - Z_{t-1}$  for a log-price or rate  $Z_t = \log S_t$ . In this case

$$X_t = \log(S_t/S_{t-1}) \approx S_t/S_{t-1} - 1 = (S_t - S_{t-1})/S_{t-1};$$


the former is often called a **(log-)return**, the latter a **relative return**.

## 3.1.1 Volatility Clustering




- (a) Log-returns for the DAX index from 1985-01-02 to 1994-12-30 ( $n = 2608$ ).
- (b) Simulated iid data from a fitted normal with  $\hat{\mu} = \bar{X}_n$ ,  $\hat{\sigma}^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2$  show too few extremes.
- (c) Simulated iid data from a fitted  $t_{3.8}$ . Better range of values but still no volatility clustering (= tendency for extreme returns to be followed by extreme returns).

Estimated *autocorrelation function (ACF)*  $\rho(h) = \text{corr}(X_0, X_h), h \in \mathbb{Z}$



- Estimated  
 (a) ACF of  $(X_t)_{t \in \mathbb{Z}}$   
 (b) ACF of  $(|X_t|)_{t \in \mathbb{Z}}$
- Non-zero ACF at lag 1 implies a tendency for a return to be followed by a return of equal sign; not the case here.  
 $\Rightarrow$  Predicted return  $\approx 0$
- For iid process  $(X_t)_{t \in \mathbb{Z}}$   
 $\rho_X(h) = \rho_{|X|}(h) = I_{\{h=0\}}$ ;  
 not the case here. (Can confirm with Ljung–Box tests.)
- $(Z_t)_{t \in \mathbb{Z}}$  not a random walk  
 $(S_t)_{t \in \mathbb{Z}}$  not GBM.

Concerning clustering of extremes, consider the 100 largest losses of the...



- (a) ... DAX index  
(c) ... simulated fitted  $t_{3.8}$
- (b), (d) Q-Q plots of waiting times between these large losses (should be  $\text{Exp}(\lambda)$  for iid data; see EVT chapter).
- The DAX data shows shorter and longer waiting times than the iid data  
⇒ clustering of extremes.



## 3.1.2 Non-normality and heavy tails

### Formal statistical tests of normality



- For **general univariate** df  $F$ :
  - ▶ Kolmogorov–Smirnov (test statistic  $T_n = \sup_x |\hat{F}_n(x) - F(x)|$ )
  - ▶ Cramér–von Mises ( $T_n = n \int_{-\infty}^{\infty} (\hat{F}_n(x) - F(x))^2 dF(x)$ )
  - ▶ **Anderson–Darling** ( $T_n = n \int_{-\infty}^{\infty} \frac{(\hat{F}_n(x) - F(x))^2}{F(x)(1-F(x))} dF(x)$ ; recommended by D’Agostino and Stephens (1986))
- For  $F = N(\mu, \sigma^2)$ :
  - ▶ Shapiro–Wilk (idea: quantify Q-Q plot in one number)
  - ▶ D’Agostino (based on skewness and kurtosis as Jarque–Bera)
  - ▶ **Jarque–Bera test**: Compares skewness  $\beta = \frac{\mathbb{E}((X-\mu)^3)}{\sigma^3}$  and kurtosis  $\kappa = \frac{\mathbb{E}((X-\mu)^4)}{\sigma^4}$  with sample versions. The test statistic is

$$T_n = \frac{n}{6} \left( \hat{\beta}^2 + \frac{1}{4} (\hat{\kappa} - 3)^2 \right) \overset{H_0}{\underset{n \text{ large}}{\sim}} \chi^2_2.$$

## Graphical tests

- Suppose we want to **graphically test whether  $X_1, \dots, X_n \sim F$  for some df  $F$**  based on realizations of iid  $X_1, \dots, X_n$ .
- Let  $X_{(1)} \leq \dots \leq X_{(n)}$  denote the corresponding **order statistics** and note that  $\hat{F}_n(x) = \frac{1}{n} \sum_{i=1}^n I_{\{X_i \leq x\}} = \frac{1}{n} \sum_{i=1}^n I_{\{X_{(i)} \leq x\}}$ ,  $x \in \mathbb{R}$ , i.e. the order statistics contain all relevant information about  $X_1, \dots, X_n$ .
- Possible graphical tests:
  - ▶ **P-P plot**: Plot  $\{(p_i, F(X_{(i)})) : i = 1, \dots, n\}$ , where  $p_i \stackrel{n > 10}{=} \frac{i-1/2}{n} \approx \frac{i}{n} \approx \hat{F}_n(X_{(i)})$ . If  $F \approx \hat{F}_n$ , the points lie roughly on a **line with slope 1**; this also applies to Q-Q plots.
  - ▶ **Q-Q plot**: Plot  $\{(F^{\leftarrow}(p_i), X_{(i)}) : i = 1, \dots, n\}$  (tail differences better visible).

Interpreting Q-Q plots (**S-shape** hints at **heavier tails** than  $N(\mu, \sigma^2)$ ):



Daily returns typically have kurtosis  $\kappa > 3$  (*leptokurtic*; narrower center, heavier tails than  $N(\mu, \sigma^2)$  for which  $\kappa = 3$ ). They have **power-like tails** rather than exponential.

To summarize, we can infer the following stylized facts about univariate financial return series:

- (U1) Return series are not iid although they show little serial correlation;
- (U2) Series of absolute or squared returns show profound serial correlation;
- (U3) Conditional expected returns are close to zero;
- (U4) Volatility (conditional standard deviation) appears to vary over time;
- (U5) Extreme returns appear in clusters;
- (U6) Return series are leptokurtic or heavy-tailed (power-like tail).


## 3.2 Multivariate stylized facts

Consider **multivariate** log-return data  $X_1, \dots, X_n$ .

### 3.2.1 Correlation between series


- By (U1), the **returns of stock A at  $t$  and  $t + h$  show little correlation**. The same applies to the returns of stock A at  $t$  and stock B at  $t + h$ ,  $h > 0$ . Stock **A and stock B on day  $t$  may be correlated** due to factors that affect the whole market (*contemporaneous dependence*).
- **Correlations of returns at  $t$  vary over time** (difficult to detect whether changes are continual or constant within regimes; fit different models for changing correlation, then make a formal comparison).
- **Periods of high/low volatility** are typically **common to more than one stock**  $\Rightarrow$  Returns of large magnitude in A at  $t$  may be followed by returns of large magnitude in A and B at  $t + h$ .

## Estimated correlations between/within series:



2000 values from period 1985-01-23 to 1994-09-22


## Estimated correlations between/within series of absolute values:



2000 values from period 1985-01-23 to 1994-09-22

### 3.2.2 Tail dependence

(BMW, Siemens) log-returns from 1985-01-23 to 1994-09-22 ( $n = 2000$ )



In **volatile/extreme periods**, dependence is **stronger** (1: 1987-10-19 **Black Monday** (DJ drop by 22%); 2: 1989-10-16 **Monday demonstrations in Leipzig** (Wende); 3: 1991-08-19 **coup against soviet president M. Gorbachev**).



To summarize, we can infer the following **stylized facts** about **multivariate** financial return series:

- (M1) **Multivariate return series** show **little** evidence of **cross-correlation**, **except** for **contemporaneous returns** (i.e. at the same  $t$ );
- (M2) Multivariate series of **absolute returns** show **profound cross-correlation**;
- (M3) **Correlations** between contemporaneous returns **vary over time**;
- (M4) **Extreme returns** in one series **often coincide with extreme returns** in several **other series** (e.g. tail dependence).

## 4 Financial time series

4.1 Fundamentals of time series analysis

4.2 GARCH models for changing volatility

# 4.1 Fundamentals of time series analysis

## 4.1.1 Basic definitions

A *stochastic process* is a family of rvs  $(X_t)_{t \in I}$ ,  $I \subseteq \mathbb{R}$ , defined on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . A *time series* is a discrete-time ( $I \subseteq \mathbb{Z}$ ) stochastic process.

### Definition 4.1 (Mean function, autocovariance function)

Assuming they exist, the *mean function*  $\mu(t)$  and the *autocovariance function*  $\gamma(t, s)$  of  $(X_t)_{t \in \mathbb{Z}}$  are defined by

$$\mu(t) = \mathbb{E}(X_t), \quad t \in \mathbb{Z},$$

$$\gamma(t, s) = \text{cov}(X_t, X_s) = \mathbb{E}((X_t - \mathbb{E}X_t)(X_s - \mathbb{E}X_s)), \quad t, s \in \mathbb{Z}.$$



## Definition 4.2 ((Weak/strict) stationarity)

- 1)  $(X_t)_{t \in \mathbb{Z}}$  is *(weakly/covariance) stationary* if  $\mathbb{E}(X_t^2) < \infty$ ,  $\mu(t) = \mu \in \mathbb{R}$  and  $\gamma(t, s) = \gamma(t + h, s + h)$  for all  $t, s, h \in \mathbb{Z}$ .
- 2)  $(X_t)_{t \in \mathbb{Z}}$  is *strictly stationary* if  $(X_{t_1}, \dots, X_{t_n}) \stackrel{d}{=} (X_{t_1+h}, \dots, X_{t_n+h})$  for all  $t_1, \dots, t_n, h \in \mathbb{Z}$ ,  $n \in \mathbb{N}$ .

## Remark 4.3

- 1) Both types of stationarity formalize the idea that  $(X_t)_{t \in \mathbb{Z}}$  behaves similarly in any time period.
- 2) ■ *Strict stationarity  $\nRightarrow$  stationarity unless we also assume that  $\mathbb{E}(X_t^2)$  exists.*  
■ *Stationarity  $\nRightarrow$  strict stationarity because  $\mathbb{E}(|X_t|^p)$ ,  $p > 2$ , could change.*
- 3)  $\gamma(0, t - s) = \gamma(s, t) = \gamma(t, s) = \gamma(0, s - t)$ , so  $\gamma(t, s)$  only depends on the lag  $h = |t - s|$ . We can thus write  $\gamma(h) := \gamma(0, |h|)$ ,  $h \in \mathbb{Z}$ .

# Stationary?



## Autocorrelation in stationary time series

### Definition 4.4 (ACF)

The *autocorrelation function (ACF)* (or *serial correlation*) of a stationary time series  $(X_t)_{t \in \mathbb{Z}}$  is defined by

$$\rho(h) := \text{corr}(X_0, X_h) = \gamma(h)/\gamma(0), \quad h \in \mathbb{Z}.$$

## White noise processes

### Definition 4.5 ((Strict) white noise)

- 1)  $(X_t)_{t \in \mathbb{Z}}$  is a *white noise* process if  $(X_t)_{t \in \mathbb{Z}}$  is *stationary* with  $\rho(h) = I_{\{h=0\}}$  (*no serial correlation*). If  $\mu(t) = 0$ ,  $\gamma(0) = \sigma^2$ ,  $(X_t)_{t \in \mathbb{Z}}$  is denoted by  $\text{WN}(0, \sigma^2)$ .
- 2)  $(X_t)_{t \in \mathbb{Z}}$  is a *strict white noise* process if  $(X_t)_{t \in \mathbb{Z}}$  is a sequence of *iid rvs* with  $\gamma(0) = \sigma^2 < \infty$ . If  $\mu(t) = 0$ , we write  $\text{SWN}(0, \sigma^2)$ .

For GARCH processes (see later), we need another notion of noise.

Let  $(X_t)_{t \in \mathbb{Z}}$  be a stochastic process on  $(\Omega, \mathcal{F}, \mathbb{P})$ . A sequence  $(\mathcal{F}_t)_{t \in \mathbb{Z}}$  of  $\sigma$ -algebras is called *filtration* if  $\mathcal{F}_t \subseteq \mathcal{F}_{t+1} \subseteq \mathcal{F}$ ,  $t \in \mathbb{Z}$ . If  $\mathcal{F}_t = \sigma(\{X_s : s \leq t\})$ , we call  $(\mathcal{F}_t)_{t \in \mathbb{Z}}$  the *natural filtration* of  $(X_t)_{t \in \mathbb{Z}}$ .  $(X_t)_{t \in \mathbb{Z}}$  is *adapted* to  $(\mathcal{F}_t)_{t \in \mathbb{Z}}$  if  $X_t \in \mathcal{F}_t$ ,  $t \in \mathbb{Z}$  ( $X_t$  is  $\mathcal{F}_t$ -measurable).

## Definition 4.6 (MGDS)

$(X_t)_{t \in \mathbb{Z}}$  is a *martingale-difference sequence (MGDS)* w.r.t.  $(\mathcal{F}_t)_{t \in \mathbb{Z}}$  if

- i)  $\mathbb{E}|X_t| < \infty$  for all  $t$ ;
- ii)  $(X_t)_{t \in \mathbb{Z}}$  is adapted to  $(\mathcal{F}_t)_{t \in \mathbb{Z}}$ ; and
- iii)  $\mathbb{E}(X_{t+1} | \mathcal{F}_t) = 0$  for all  $t \in \mathbb{Z}$ .

- If  $\mathbb{E}(X_{t+1} | \mathcal{F}_t) = X_t$  a.s., then  $(X_t)$  is a (discrete-time) *martingale* and  $\varepsilon_t = X_t - X_{t-1}$  is a MGDS (winnings in rounds of a *fair game*).
- One can show that a MGDS  $(\varepsilon_t)_{t \in \mathbb{Z}}$  with  $\sigma^2 = \mathbb{E}(\varepsilon_t^2) < \infty$  satisfies
  - ▶  $\rho(h) = 0$ ,  $h \neq 0$ , so  $(\varepsilon_t)_{t \in \mathbb{Z}} \sim \text{WN}(0, \sigma^2)$ ;
  - ▶  $\mathbb{E}(\varepsilon_{t+1+k} | \mathcal{F}_t) = \mathbb{E}(\mathbb{E}(\varepsilon_{t+1+k} | \mathcal{F}_{t+k}) | \mathcal{F}_t) = 0$ ,  $k \in \mathbb{N}$ .

## 4.1.2 ARMA processes

### Definition 4.7 (ARMA( $p, q$ ))

Let  $(\varepsilon_t)_{t \in \mathbb{Z}} \sim \text{WN}(0, \sigma^2)$ .  $(X_t)_{t \in \mathbb{Z}}$  is a *zero-mean ARMA( $p, q$ ) process* if it is stationary and satisfies, for all  $t \in \mathbb{Z}$ ,

$$X_t - \phi_1 X_{t-1} - \cdots - \phi_p X_{t-p} = \varepsilon_t + \theta_1 \varepsilon_{t-1} + \cdots + \theta_q \varepsilon_{t-q}. \quad (6)$$

$(X_t)_{t \in \mathbb{Z}}$  is ARMA( $p, q$ ) with *mean  $\mu$*  if  $(X_t - \mu)_{t \in \mathbb{Z}}$  is a zero-mean ARMA( $p, q$ ).

### Remark 4.8

- If the *innovations*  $(\varepsilon_t)_{t \in \mathbb{Z}}$  are  $\text{SWN}(0, \sigma^2)$ , then  $(X_t)_{t \in \mathbb{Z}}$  is strictly stationary (follows from the representation as a linear process below).
- The defining equation (6) can be written as  $\phi(B)X_t = \theta(B)\varepsilon_t$ ,  $t \in \mathbb{Z}$ , where  $B$  denotes the *backshift operator* (such that  $B^k X_t = X_{t-k}$ ) and  $\phi(z) = 1 - \phi_1 z - \cdots - \phi_p z^p$  and  $\theta(z) = 1 + \theta_1 z + \cdots + \theta_q z^q$ .



## Causal processes

For practical purposes, it suffices to consider *causal ARMA processes*, that is, ARMA processes  $(X_t)_{t \in \mathbb{Z}}$  satisfying

$$X_t = \sum_{k=0}^{\infty} \psi_k \varepsilon_{t-k} \quad (\text{depends on the past/present, not the future})$$

for  $\sum_{k=0}^{\infty} |\psi_k| < \infty$  (*absolute summability condition*; guarantees  $\mathbb{E}|X_t| < \infty$ ).

### Proposition 4.9 (ACF for causal processes)

Let  $X_t = \sum_{k=0}^{\infty} \psi_k \varepsilon_{t-k}$  with  $\sum_{k=0}^{\infty} |\psi_k| < \infty$ . This process is stationary with ACF given by

$$\rho(h) = \frac{\sum_{k=0}^{\infty} \psi_k \psi_{k+|h|}}{\sum_{k=0}^{\infty} \psi_k^2}, \quad h \in \mathbb{Z}.$$

## Theorem 4.10 (Stationary and causal ARMA solutions)

Let  $(X_t)_{t \in \mathbb{Z}}$  be an ARMA( $p, q$ ) process for which  $\phi(z), \theta(z)$  have no roots in common. Then

$$(X_t)_{t \in \mathbb{Z}} \text{ is stationary and causal} \quad \Leftrightarrow \quad \phi(z) \neq 0 \quad \forall z \in \mathbb{C} : |z| \leq 1.$$

In this case,  $X_t = \sum_{k=0}^{\infty} \psi_k \varepsilon_{t-k}$  for  $\sum_{k=0}^{\infty} \psi_k z^k = \theta(z)/\phi(z)$ ,  $|z| \leq 1$ .

- If  $\theta(z) \neq 0$ ,  $|z| \leq 1$  (known as *invertibility condition*), we can recover  $\varepsilon_t$  from  $(X_s)_{s \leq t}$  via  $\varepsilon_t = \phi(B)X_t/\theta(B)$ , so  $\varepsilon_t$  is  $\mathcal{F}_t$ -measurable for  $\mathcal{F}_t = \sigma(\{X_s : s \leq t\})$  if  $(X_t)_{t \in \mathbb{Z}}$  is invertible.
- An ARMA( $p, q$ ) process with mean  $\mu$  can be written as  $X_t = \mu_t + \varepsilon_t$  for  $\mu_t = \mu + \sum_{k=1}^p \phi_k (X_{t-k} - \mu) + \sum_{k=1}^q \theta_k \varepsilon_{t-k}$ . If  $(X_t)_{t \in \mathbb{Z}}$  is invertible,  $\mu_t \in \mathcal{F}_{t-1}$ . If  $(\varepsilon_t)_{t \in \mathbb{Z}}$  is a MGDS w.r.t.  $(\mathcal{F}_t)_{t \in \mathbb{Z}}$ , then  $\mu_t = \mathbb{E}(X_t | \mathcal{F}_{t-1})$ . Therefore, ARMA processes put structure on the conditional mean  $\mu_t$  given the past. We will see that GARCH processes put structure on  $\sigma_t^2 = \text{var}(X_t | \mathcal{F}_{t-1})$  (helpful for modeling volatility clustering).

## 4.1.3 Analysis in the time domain

### Correlogram

A *correlogram* is a plot of  $(h, \hat{\rho}(h))_{h \geq 0}$  for the sample ACF

$$\hat{\rho}(h) = \frac{\sum_{t=1}^n (X_{t+h} - \bar{X}_n)(X_t - \bar{X}_n)}{\sum_{t=1}^n (X_t - \bar{X}_n)^2}, \quad h \in \{0, \dots, n\}.$$

### Theorem 4.11

Let  $X_t - \mu = \sum_{k=0}^{\infty} \psi_k Z_{t-k}$  and  $(Z_t) \sim \text{SWN}(0, \sigma^2)$ . Under suitable conditions,

$$\sqrt{n} \left( \begin{pmatrix} \hat{\rho}(1) \\ \vdots \\ \hat{\rho}(h) \end{pmatrix} - \begin{pmatrix} \rho(1) \\ \vdots \\ \rho(h) \end{pmatrix} \right) \xrightarrow[(n \rightarrow \infty)]{d} N_h(\mathbf{0}, W), \quad h \in \mathbb{N},$$

for a matrix  $W$  depending on  $\rho$ ; see McNeil et al. (2015, Theorem 4.13).

If the ARMA process is SWN itself,  $\sqrt{n} \begin{pmatrix} \hat{\rho}(1) \\ \vdots \\ \hat{\rho}(h) \end{pmatrix} \xrightarrow[(n \rightarrow \infty)]{d} N_h(\mathbf{0}, I_h)$ , so that with probability  $1 - \alpha$ ,

$$\hat{\rho}(k) \underset{(n \text{ large})}{\in} \left[ -\frac{q_{1-\alpha/2}}{\sqrt{n}}, \frac{q_{1-\alpha/2}}{\sqrt{n}} \right], \quad k \in \{1, \dots, h\},$$

where  $q_{1-\alpha/2} = \Phi^{-1}(1-\alpha/2)$ . This interval is usually shown in correlogram.

- As a formal test of the SWN hypothesis, one can use the Ljung–Box test with test statistic

$$T = n(n+2) \sum_{k=1}^h \frac{\hat{\rho}(k)^2}{n-k} \underset{n \text{ large}}{\sim} \chi_h^2; \quad \text{reject if } T > \chi_h^{2-1}(1-\alpha).$$

- If  $(X_t)_{t \in \mathbb{Z}}$  is SWN, so is  $(X_t^2)_{t \in \mathbb{Z}}$ . It is a good idea to also apply the correlogram and Ljung–Box tests to  $(|X_t|)_{t \in \mathbb{Z}}$  or  $(X_t^2)_{t \in \mathbb{Z}}$ .

## 4.2 GARCH models for changing volatility

- (G)ARCH = (generalized) autoregressive conditionally heteroscedastic
- They are the most important models for daily risk-factor returns.

### 4.2.1 ARCH processes

#### Definition 4.12 (ARCH( $p$ ))

Let  $(Z_t)_{t \in \mathbb{Z}} \sim \text{SWN}(0, 1)$ .  $(X_t)_{t \in \mathbb{Z}}$  is an ARCH( $p$ ) process if it is strictly stationary and satisfies

$$X_t = \sigma_t Z_t,$$
$$\sigma_t^2 = \alpha_0 + \sum_{k=1}^p \alpha_k X_{t-k}^2,$$

where  $\alpha_0 > 0$ ,  $\alpha_k \geq 0$ ,  $k \in \{1, \dots, p\}$ .

Typical examples:  $Z_t \stackrel{\text{ind.}}{\sim} \text{N}(0, 1)$  or  $Z_t \stackrel{\text{ind.}}{\sim} t_\nu(0, (\nu - 2)/\nu)$ .

## Remark 4.13

- 1)  $\sigma_{t+1}$  is  $\mathcal{F}_t$ -measurable  $\Rightarrow \mathbb{E}(X_{t+1} | \mathcal{F}_t) = \sigma_{t+1} \mathbb{E}(Z_{t+1} | \mathcal{F}_t) = \sigma_{t+1} \mathbb{E}(Z_{t+1}) = 0$ . Thus, ARCH( $p$ ) processes are MGDs w.r.t. the natural filtration  $(\mathcal{F}_t)_{t \in \mathbb{Z}}$ . If they are stationary, they are white noise since



$$\begin{aligned} \gamma(h) &= \mathbb{E}(X_t X_{t+h}) \stackrel{\text{tower property}}{=} \mathbb{E}(\mathbb{E}(X_t X_{t+h} | \mathcal{F}_{t+h-1})) \\ &= \mathbb{E}(X_t \mathbb{E}(X_{t+h} | \mathcal{F}_{t+h-1})) = 0, \quad h \in \mathbb{N}. \end{aligned}$$

This also applies to GARCH processes; see below.

- 2) If  $(X_t)_{t \in \mathbb{Z}}$  is stationary, then  $\text{var}(X_t | \mathcal{F}_{t-1}) = \mathbb{E}((\sigma_t Z_t)^2 | \mathcal{F}_{t-1}) = \sigma_t^2 \mathbb{E}(Z_t^2 | \mathcal{F}_{t-1}) = \sigma_t^2 \mathbb{E}(Z_t^2) = \sigma_t^2$ .  
 $\Rightarrow$  Volatility  $\sigma_t$  (conditional standard deviation) is changing in time, depending on past values of the process. ARCH models can thus capture volatility clustering (if one of  $|X_{t-1}|, \dots, |X_{t-p}|$  is large,  $X_t$  is drawn from a distribution with large variance). This is where “autoregressive conditionally heteroscedastic” comes from.



### Example 4.14 (ARCH(1))

- One can show that an ARCH(1) process  $(X_t)_{t \in \mathbb{Z}}$  is strictly stationary  $\Leftrightarrow \mathbb{E}(\log(\alpha_1 Z_t^2)) < 0$ . In this case,  $X_t^2 = \alpha_0 \sum_{k=0}^{\infty} \alpha_1^k \prod_{j=0}^k Z_{t-j}^2$ .
- $(X_t)_{t \in \mathbb{Z}}$  is stationary  $\Leftrightarrow \alpha_1 < 1$ . In this case,  $\text{var}(X_t) = \alpha_0 / (1 - \alpha_1)$ .
- Provided that  $\mathbb{E}(Z_t^4) < \infty$  and  $\alpha_1 < (\mathbb{E}(Z_t^4))^{-1/2}$ , one can show that  $\kappa(X_t) = \frac{\mathbb{E}(X_t^4)}{\mathbb{E}(X_t^2)^2} = \frac{\kappa(Z_t)(1-\alpha_1^2)}{(1-\alpha_1^2\kappa(Z_t))}$ . If  $\kappa(Z_t) > 1$ ,  $\kappa(X_t) > \kappa(Z_t)$ . For Gaussian or  $t$  innovations,  $\kappa(X_t) > 3$  (leptokurtic).
- Parallels with the AR(1) process: If  $\mathbb{E}(X_t^4) < \infty$ ,  $\alpha_1 < 1$  and  $\varepsilon_t = \sigma_t^2(Z_t^2 - 1)$ , one can show that  $(X_t^2)_{t \in \mathbb{Z}}$  is an AR(1) of the form  $X_t^2 - \frac{\alpha_0}{1-\alpha_1} = \alpha_1(X_{t-1}^2 - \frac{\alpha_0}{1-\alpha_1}) + \varepsilon_t$ .



a)  $n = 1000$  realizations of an ARCH(1) process with  $\alpha_0 = 0.5$ ,  $\alpha_1 = 0.5$  and Gaussian innovations;

b) Realization of the volatility  $(\sigma_t)_{t \in \mathbb{Z}}$ ;



c) Correlogram of  $(X_t)_{t \in \mathbb{Z}}$ , compare with Remark 4.13 1);

d) Correlogram of  $(X_t^2)_{t \in \mathbb{Z}}$  (AR(1)); dashed line = true ACF



## 4.2.2 GARCH processes

### Definition 4.15 (GARCH( $p, q$ ))

Let  $(Z_t)_{t \in \mathbb{Z}} \sim \text{SWN}(0, 1)$ .  $(X_t)_{t \in \mathbb{Z}}$  is a **GARCH( $p, q$ ) process** if it is strictly stationary and satisfies


$$X_t = \sigma_t Z_t,$$
$$\sigma_t^2 = \alpha_0 + \sum_{k=1}^p \alpha_k X_{t-k}^2 + \sum_{k=1}^q \beta_k \sigma_{t-k}^2,$$

where  $\alpha_0 > 0$ ,  $\alpha_k \geq 0$ ,  $k \in \{1, \dots, p\}$ ,  $\beta_k \geq 0$ ,  $k \in \{1, \dots, q\}$ .

If one of  $|X_{t-1}|, \dots, |X_{t-p}|$  or  $\sigma_{t-1}, \dots, \sigma_{t-q}$  is large,  $X_t$  is drawn from a distribution with (persistently) large variance. Periods of high volatility tend to be more persistent.

### Example 4.16 (GARCH(1,1))

- One can show (via stoch. recurrence relations) that a GARCH(1,1) process  $(X_t)_{t \in \mathbb{Z}}$  is strictly stationary if  $\mathbb{E}(\log(\alpha_1 Z_t^2 + \beta_1)) < \infty$ . In this case,  $X_t = Z_t \sqrt{\alpha_0 (1 + \sum_{k=1}^{\infty} \prod_{j=1}^k (\alpha_1 Z_{t-j}^2 + \beta_1))}$ .
- $(X_t)_{t \in \mathbb{Z}}$  is stationary  $\Leftrightarrow \alpha_1 + \beta_1 < 1$ . In this case,  $\text{var}(X_t) = \frac{\alpha_0}{1 - \alpha_1 - \beta_1}$ .
- Provided that  $\mathbb{E}((\alpha_1 Z_t^2 + \beta_1)^2) < 1$  (or  $(\alpha_1 + \beta_1)^2 < 1 - (\kappa(Z_t) - 1)\alpha_1^2$ ), one can show that  $\kappa(X_t) = \frac{\kappa(Z_t)(1 - (\alpha_1 + \beta_1)^2)}{1 - (\alpha_1 + \beta_1)^2 - (\kappa(Z_t) - 1)\alpha_1^2}$ . If  $\kappa(Z_t) > 1$  (Gaussian, scaled  $t$  innovations),  $\kappa(X_t) > \kappa(Z_t)$ .
- Parallels with the ARMA(1,1) process: If  $\mathbb{E}(X_t^4) < \infty$ ,  $\alpha_1 + \beta_1 < 1$  and  $\varepsilon_t = \sigma_t^2(Z_t^2 - 1)$ , one can show that  $(X_t^2)_{t \in \mathbb{Z}}$  is an ARMA(1,1) of the form  $X_t^2 - \frac{\alpha_0}{1 - \alpha_1 - \beta_1} = (\alpha_1 + \beta_1)(X_{t-1}^2 - \frac{\alpha_0}{1 - \alpha_1 - \beta_1}) + \varepsilon_t - \beta_1 \varepsilon_{t-1}$ .



- a)  $n = 1000$  realization of a GARCH(1,1) process with  $\alpha_0 = 0.5$ ,  $\alpha_1 = 0.1$ ,  $\beta_1 = 0.85$  and Gaussian innovations;
- b) Realization of the volatility  $(\sigma_t)_{t \in \mathbb{Z}}$ ;
- c) Correlogram of  $(X_t)_{t \in \mathbb{Z}}$ , compare with Remark 4.13 1);
- d) Correlogram of  $(X_t^2)_{t \in \mathbb{Z}}$  (ARMA(1,1)); dashed line = true ACF

## The GARCH(p,q) model

- Higher-order GARCH models have the same general behaviour as ARCH(1) and GARCH(1,1) models, but their mathematical analysis becomes more tedious.
- One can show that  $(X_t)_{t \in \mathbb{Z}}$  is stationary  $\Leftrightarrow \sum_{k=1}^p \alpha_k + \sum_{k=1}^q \beta_k < 1$ .
- A squared GARCH(p,q) process has the structure

$$X_t^2 = \alpha_0 + \sum_{k=1}^{\max(p,q)} (\alpha_k + \beta_k) X_{t-k}^2 + \varepsilon_t - \sum_{k=1}^q \beta_k \varepsilon_{t-k},$$

where  $\varepsilon_t = \sigma_t^2(Z_t^2 - 1)$ ,  $\alpha_k = 0$ ,  $k \in \{p+1, \dots, q\}$  if  $q > p$ , or  $\beta_k = 0$  for  $k \in \{q+1, \dots, p\}$  if  $p > q$ . This resembles the ARMA(max(p,q), q) process and is formally such a process provided  $\mathbb{E}(X_t^4) < \infty$ .

- There are also IGARCH models (i.e. non-stationary GARCH(p,q) models with  $\sum_{k=1}^p \alpha_k + \sum_{k=1}^q \beta_k = 1$ ; infinite variance).

## 4.2.3 Simple extensions of the GARCH model

Consider stationary GARCH processes as white noise for ARMA processes.

### **Definition 4.17** (ARMA( $p_1, q_1$ ) with GARCH( $p_2, q_2$ ) errors)

Let  $(Z_t)_{t \in \mathbb{Z}} \sim \text{SWN}(0, 1)$ .  $(X_t)_{t \in \mathbb{Z}}$  is an **ARMA( $p_1, q_1$ ) process with GARCH( $p_2, q_2$ ) errors** if it is stationary and satisfies

$$X_t = \mu_t + \varepsilon_t \quad \text{for} \quad \varepsilon_t = \sigma_t Z_t \quad (\text{so } X_t = \mu_t + \sigma_t Z_t),$$

$$\mu_t = \mu + \sum_{k=1}^{p_1} \phi_k (X_{t-k} - \mu) + \sum_{k=1}^{q_1} \theta_k (X_{t-k} - \mu_{t-k}),$$

$$\sigma_t^2 = \alpha_0 + \sum_{k=1}^{p_2} \alpha_k (X_{t-k} - \mu_{t-k})^2 + \sum_{k=1}^{q_2} \beta_k \sigma_{t-k}^2,$$

where  $\alpha_0 > 0$ ,  $\alpha_k \geq 0$ ,  $k \in \{1, \dots, p_2\}$ ,  $\beta_k \geq 0$ ,  $k \in \{1, \dots, q_2\}$ ,  
 $\sum_{k=1}^{p_2} \alpha_k + \sum_{k=1}^{q_2} \beta_k < 1$ .

- ARMA models with GARCH errors are quite flexible models. It is easy to see that the conditional mean of  $(X_t)_{t \in \mathbb{Z}}$  is  $\mu_t = \mathbb{E}(X_t | \mathcal{F}_{t-1})$  and that the conditional variance of  $(X_t)_{t \in \mathbb{Z}}$  is  $\sigma_t^2 = \text{var}(X_t | \mathcal{F}_{t-1})$ .
- Other extensions not further discussed here:
  - ▶ *GJR-GARCH*. These models introduce a parameter in the volatility equation in order for the volatility to react asymmetrically to recent returns (bad news leading to a fall in the equity value of a company tends to increase volatility, the so-called *leverage effect*).
  - ▶ *Threshold GARCH (TGARCH)*. More general models (than GJR-GARCH) in which the dynamics at time  $t$  depend on whether  $X_{t-1}$  (or  $Z_{t-1}$ ; sometimes even a coefficient) was below/above a threshold.
  - ▶ Note that one could also use an asymmetric innovation distribution with mean 0 and variance 1, e.g. from the generalized hyperbolic family or skewed  $t$  distribution.

## 4.2.4 Fitting GARCH models to data

- The most widely used approach is **maximum likelihood**.
- After model fitting, **we check residuals**. Consider an **ARMA model with GARCH errors**  $X_t = \mu_t + \varepsilon_t = \mu_t + \sigma_t Z_t$ ; see Definition 4.17.
- We distinguish **two kinds of residuals**:
  - 1) **Unstandardized residuals**. These are the **residuals**  $\hat{\varepsilon}_1, \dots, \hat{\varepsilon}_n$  and should behave like a realization of a GARCH process.
  - 2) **Standardized residuals**. These are reconstructed **realizations of the SWN** which drives the GARCH process. They are **calculated from the unstandardized residuals** via

$$\hat{Z}_t = \hat{\varepsilon}_t / \hat{\sigma}_t, \quad \hat{\sigma}_t^2 = \hat{\alpha}_0 + \sum_{k=1}^{p_2} \hat{\alpha}_k \hat{\varepsilon}_{t-k}^2 + \sum_{k=1}^{q_2} \hat{\beta}_k \hat{\sigma}_{t-k}^2; \quad (7)$$

starting values for  $\hat{\varepsilon}_t$  are taken as 0 and starting values for  $\hat{\sigma}_t$  are taken as the sample variance (or 0); ignore the first few values then.

- The standardized residuals should behave like SWN. Check this via correlograms of  $(\hat{Z}_t)$  and  $(|\hat{Z}_t|)$  and by applying the Ljung–Box test of strict white noise. In case of no rejection (the dynamics have been satisfactorily captured), the validity of the innovation distribution can also be assessed (e.g. via Q-Q plots or goodness-of-fit tests).

⇒ *Two-stage analysis possible*: First estimate the dynamics via QMLE (known as *pre-whitening* of the data), then model the innovation distribution using the standardized residuals.

- Advantages:
- ▶ More transparency in model building;
  - ▶ Separating of volatility modelling and modelling of shocks that drive the process;
  - ▶ Practical in higher dimensions.

Drawbacks: ARMA fitting errors propagate through to the fitting of innovations (overall error hard to quantify).



## 4.2.5 Volatility forecasting and risk measure estimation

- Consider a weakly and strictly stationary time series  $(X_t)_{t \in \mathbb{Z}}$  of the form

$$X_t = \mu_t + \sigma_t Z_t$$

adapted to a filtration  $(\mathcal{F}_t)_{t \in \mathbb{Z}}$ , where  $\mu_t, \sigma_t \in \mathcal{F}_{t-1}$  and  $\mathbb{E}Z_t = 0$ ,  $\text{var } Z_t = 1$ , independent of  $\mathcal{F}_{t-1}$  (e.g.  $(X_t)_{t \in \mathbb{Z}}$  could be a GARCH model or ARMA model with GARCH errors).

- Assume we know  $X_{t-n+1}, \dots, X_t$  and want to compute  $P_t \sigma_{t+h}$ ,  $h \geq 1$ , a forecast of volatility based on these data.
- Since  $\mathbb{E}(\sigma_{t+h}^2 | \mathcal{F}_t) = \mathbb{E}((X_{t+h} - \mu_{t+h})^2 | \mathcal{F}_t)$  our forecasting problem is related to the problem of predicting  $(X_{t+h} - \mu_{t+h})^2$ .
- We consider two approaches: (1) calculating conditional expectations (optimal squared error forecasts) using model of GARCH type; (2) the more ad hoc exponentially weighted moving average (EWMA) approach.

## Conditional expectation

The general procedure becomes clear from examples.

### Example 4.18 (Prediction in the GARCH(1,1) model)

- A GARCH(1,1) model is of type  $X_t = \mu_t + \sigma_t Z_t$  for  $\mu_t = 0$ . Since  $\mathbb{E}(X_{t+h} | \mathcal{F}_t) = 0$ ,  $\hat{\mu}_{t+h} = P_t X_{t+h} = 0$  for all  $h \in \mathbb{N}$ .
- A natural prediction of  $X_{t+1}^2$  based on  $\mathcal{F}_t$  is its conditional mean

$$\mathbb{E}(X_{t+1}^2 | \mathcal{F}_t) = \sigma_{t+1}^2 = \alpha_0 + \alpha_1 X_t^2 + \beta_1 \sigma_t^2.$$

If  $\mathbb{E}(X_t^4) < \infty$ , this is the optimal squared error prediction.

- We thus obtain the one-step-ahead forecast

$$\hat{\sigma}_{t+1}^2 = \mathbb{E}(\widehat{X_{t+1}^2} | \mathcal{F}_t) = \alpha_0 + \alpha_1 X_t^2 + \beta_1 \hat{\sigma}_t^2.$$

- If  $h > 1$ ,  $\sigma_{t+h}^2$  and  $X_{t+h}^2$  are rvs. Their predictions (coincide and) are

$$\begin{aligned}\mathbb{E}(\sigma_{t+h}^2 | \mathcal{F}_t) &= \alpha_0 + \alpha_1 \mathbb{E}(X_{t+h-1}^2 | \mathcal{F}_t) + \beta_1 \mathbb{E}(\sigma_{t+h-1}^2 | \mathcal{F}_t) \\ &= \alpha_0 + (\alpha_1 + \beta_1) \mathbb{E}(\sigma_{t+h-1}^2 | \mathcal{F}_t)\end{aligned}$$

so that a **general formula** is

$$\mathbb{E}(\sigma_{t+h}^2 | \mathcal{F}_t) = \alpha_0 \sum_{k=0}^{h-1} (\alpha_1 + \beta_1)^k + (\alpha_1 + \beta_1)^{h-1} (\alpha_1 X_t^2 + \beta_1 \sigma_t^2).$$

Note that for  $h \rightarrow \infty$ ,  $\mathbb{E}(\sigma_{t+h}^2 | \mathcal{F}_t) \xrightarrow{\text{a.s.}} \frac{\alpha_0}{1-\alpha_1-\beta_1}$ , so the prediction of squared volatility converges to the unconditional variance of the process.

## Exponentially weighted moving averages

- A **one-period ahead forecast**  $P_t X_{t+1}$  of  $X_{t+1}$  based on  $\mathcal{F}_t$  is given by

$$P_t X_{t+1} = \alpha X_t + (1 - \alpha) P_{t-1} X_t. \quad (8)$$

Applied to  $(X_{t+1} - \mu_{t+1})^2$  leads to

$$P_t (X_{t+1} - \mu_{t+1})^2 = \alpha (X_t - \mu_t)^2 + (1 - \alpha) P_{t-1} (X_t - \mu_t)^2. \quad (9)$$

- Since  $\sigma_{t+1}^2 = \mathbb{E}((X_{t+1} - \mu_{t+1})^2 | \mathcal{F}_t)$ , we can **use (9) as exponential smoothing scheme** for the unobserved squared volatility  $\sigma_{t+1}^2$ . **This yields**

a recursive scheme for the one-step-ahead volatility forecast given by

$$\hat{\sigma}_{t+1}^2 = \alpha(X_t - \hat{\mu}_t)^2 + (1 - \alpha)\hat{\sigma}_t^2,$$

which is then iterated.

- $\alpha$  is typically **small** (e.g. RiskMetrics:  $\alpha = 0.06$ );  $\hat{\mu}_t$  is usually set to 0 (see Chapter 3).

## Estimators of $\text{VaR}_\alpha$ and $\text{ES}_\alpha$

- Suppose we now **want to estimate**  $\text{VaR}_\alpha^t$ ,  $\text{ES}_\alpha^t$ , risk measures based on  $F_{X_{t+1}|\mathcal{F}_t}$ . If  $Z_t \stackrel{\text{ind.}}{\sim} F_Z$ , the  $\mathcal{F}_t$ -measurability of  $\mu_{t+1}$  and  $\sigma_{t+1}$ , and  $X_t = \mu_t + \sigma_t Z_t$  imply that

$$F_{X_{t+1}|\mathcal{F}_t}(x) = \mathbb{P}(\mu_{t+1} + \sigma_{t+1}Z_{t+1} \leq x | \mathcal{F}_t) = F_Z\left(\frac{x - \mu_{t+1}}{\sigma_{t+1}}\right).$$

- Then  $\text{VaR}_\alpha^t = \mu_{t+1} + \sigma_{t+1}F_Z^{\leftarrow}(\alpha)$  and  $\text{ES}_\alpha^t = \mu_{t+1} + \sigma_{t+1}\text{ES}_\alpha(Z)$ .
- If we have **estimated**  $\sigma_{t+1}$  (and  $\mu_{t+1}$ ) it remains to **estimate**  $F_Z^{\leftarrow}(\alpha)$  and  $\text{ES}_\alpha(Z)$ .

- ▶ For GARCH-type models it is easy to calculate  $F_Z^{\leftarrow}(\alpha)$  and  $\text{ES}_\alpha(Z)$ .
- ▶ If we use exponential smoothing to estimate  $\mu_{t+1}$ ,  $\sigma_{t+1}$ , we can use the residuals

$$\hat{Z}_s = (X_s - \hat{\mu}_s) / \hat{\sigma}_s, \quad s \in \{t - n + 1, \dots, n\},$$

to estimate  $F_Z^{\leftarrow}(\alpha)$  and  $\text{ES}_\alpha(Z)$ .

# 5 Extreme value theory

## 5.1 Maxima

## 5.2 Threshold exceedances

## 5.1 Maxima

Consider a series of financial losses  $(X_k)_{k \in \mathbb{N}}$ .

### 5.1.1 Generalized extreme value distribution

#### Convergence of sums

Let  $(X_k)_{k \in \mathbb{N}}$  be iid with  $\mathbb{E}(X_1^2) < \infty$  (mean  $\mu$ , variance  $\sigma^2$ ) and  $S_n = \sum_{k=1}^n X_k$ . As  $n \rightarrow \infty$ ,  $\bar{X}_n \xrightarrow{\text{a.s.}} \mu$  by the Strong Law of Large Numbers (SLLN), so  $(\bar{X}_n - \mu)/\sigma \xrightarrow{\text{a.s.}} 0$ . By the CLT,

$$\sqrt{n} \frac{\bar{X}_n - \mu}{\sigma} = \frac{S_n - n\mu}{\sqrt{n}\sigma} \xrightarrow[n \uparrow \infty]{d} N(0, 1) \text{ or } \lim_{n \rightarrow \infty} \mathbb{P}\left(\frac{S_n - d_n}{c_n} \leq x\right) = \Phi(x),$$

where the sequences  $c_n = \sqrt{n}\sigma$  and  $d_n = n\mu$  give normalization and where  $\Phi(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-z^2/2} dz$ .

## Convergence of maxima

QRM is concerned with maximal losses (worst-case losses). Let  $(X_i)_{i \in \mathbb{N}} \stackrel{\text{ind.}}{\sim} F$  (can be relaxed to a strictly stationary time series) and  $F$  continuous. Then the *block maximum* is given by

$$M_n = \max\{X_1, \dots, X_n\}.$$

One can show that, for  $n \rightarrow \infty$ ,  $M_n \xrightarrow{\text{a.s.}} x_F$  (similar as in the SLLN) where  $x_F := \sup\{x \in \mathbb{R} : F(x) < 1\} = F^{\leftarrow}(1) \leq \infty$  denotes the *right endpoint of  $F$*  (similar to the SLLN).

**Question:** Is there a “CLT” for block maxima?



**Idea CLT:** What about **linear transformations** (the simplest possible)?

**Definition 5.1 (Maximum domain of attraction)**

Suppose we find *normalizing sequences* of real numbers  $(c_n) > 0$  and  $(d_n)$  such that  $(M_n - d_n)/c_n$  converges in distribution, i.e.

$$\mathbb{P}((M_n - d_n)/c_n \leq x) = \mathbb{P}(M_n \leq c_n x + d_n) = F^n(c_n x + d_n) \xrightarrow[n \uparrow \infty]{} H(x),$$

for some *non-degenerate* df  $H$  (not a unit jump). Then  $F$  is in the *maximum domain of attraction of  $H$*  ( $F \in \text{MDA}(H)$ ).

One can show that  $H$  is determined up to location/scale, i.e.  $H$  specifies a *unique type* of distribution. This is guaranteed by the **convergence to types theorem**.

**Question:** What does  $H$  look like?

## Definition 5.2 (Generalized extreme value (GEV) distribution)

The (standard) *generalized extreme value (GEV) distribution* is given by

$$H_{\xi}(x) = \begin{cases} \exp(-(1 + \xi x)^{-1/\xi}), & \text{if } \xi \neq 0, \\ \exp(-e^{-x}), & \text{if } \xi = 0, \end{cases}$$

where  $1 + \xi x > 0$  (MLE!). A three-parameter family is obtained by a location-scale transform  $H_{\xi, \mu, \sigma}(x) = H_{\xi}((x - \mu)/\sigma)$ ,  $\mu \in \mathbb{R}$ ,  $\sigma > 0$ .

- The parameterization is continuous in  $\xi$  (simplifies statistical modelling).
- The larger  $\xi$ , the heavier tailed  $H_{\xi}$  (if  $\xi > 0$ ,  $\mathbb{E}(X^k) = \infty$  iff  $k \geq \frac{1}{\xi}$ ).
- $\xi$  is the *shape* (determines moments, tail). Special cases:
  - 1)  $\xi < 0$ : the Weibull df, short-tailed,  $x_{H_{\xi}} < \infty$ ;
  - 2)  $\xi = 0$ : the Gumbel df,  $x_{H_0} = \infty$ , decays exponentially;
  - 3)  $\xi > 0$ : the Fréchet df,  $x_{H_{\xi}} = \infty$ , heavy-tailed ( $\bar{H}_{\xi}(x) \approx (\xi x)^{-1/\xi}$ ), most important case for practice

### Theorem 5.3 (Fisher–Tippett–Gnedenko)

If  $F \in \text{MDA}(H)$  for some non-degenerate  $H$ , then  $H$  must be of GEV type, i.e.  $H = H_\xi$  for some  $\xi \in \mathbb{R}$ .

*Proof.* Non-trivial. For a sketch, see Embrechts et al. (1997, p. 122).  $\square$

- **Interpretation:** If location-scale transformed maxima converge in distribution to a non-degenerate limit, the limiting distribution must be a location-scale transformed GEV distribution (that is, of GEV type).
- We can always choose normalizing sequences  $(c_n) > 0$ ,  $(d_n)$  such that  $H_\xi$  appears in standard form.
- All commonly encountered continuous distributions are in the MDA of a GEV distribution.

### Example 5.4 (Pareto distribution)

For  $(X_i)_{i \in \mathbb{N}} \stackrel{\text{ind.}}{\sim} \text{Par}(\theta, \kappa)$  with  $F(x) = 1 - (\frac{\kappa}{\kappa+x})^\theta$ ,  $x \geq 0$ ,  $\theta, \kappa > 0$ , choosing  $c_n = \kappa n^{1/\theta}/\theta$ ,  $d_n = \kappa(n^{1/\theta} - 1)$ ,  $F^n(c_n x + d_n)$  equals

$$\begin{aligned} & \left(1 - \left(\frac{\kappa}{\kappa + x\kappa n^{1/\theta}/\theta + \kappa(n^{1/\theta} - 1)}\right)^\theta\right)^n \\ &= \left(1 - \left(\frac{1}{1 + xn^{1/\theta}/\theta + n^{1/\theta} - 1}\right)^\theta\right)^n = \left(1 - \left(\frac{1}{n^{1/\theta}(1 + x/\theta)}\right)^\theta\right)^n \\ &= \left(1 - \frac{(1 + x/\theta)^{-\theta}}{n}\right)^n \xrightarrow{n \uparrow \infty} \exp(-(1 + x/\theta)^{-\theta}) = H_{1/\theta}(x) \text{ (Fréchet)} \end{aligned}$$

Therefore,  $F \in \text{MDA}(H_{1/\theta})$ .

## 5.1.2 Maximum domains of attraction

All commonly applied continuous  $F$  belong to  $\text{MDA}(H_\xi)$  for some  $\xi \in \mathbb{R}$ .  $\mu, \sigma$  can be estimated, but how can we characterize/determine  $\xi$ ? All  $F \in \text{MDA}(H_\xi)$  for  $\xi > 0$  have an elegant characterization involving the following notions.

### Definition 5.5 (Slowly/regularly varying functions)

- 1) A positive, Lebesgue-measurable function  $L$  on  $(0, \infty)$  is *slowly varying at  $\infty$*  if  $\lim_{x \rightarrow \infty} \frac{L(tx)}{L(x)} = 1, t > 0$ . The class of all such functions is denoted by  $\mathcal{R}_0$ ; e.g.  $c, \log \in \mathcal{R}_0$ .
- 2) A positive, Lebesgue-measurable function  $h$  on  $(0, \infty)$  is *regularly varying at  $\infty$  with index  $\alpha \in \mathbb{R}$*  if  $\lim_{x \rightarrow \infty} \frac{h(tx)}{h(x)} = t^\alpha, t > 0$ . The class of all such functions is denoted by  $\mathcal{R}_\alpha$ ; e.g.  $x^\alpha L(x) \in \mathcal{R}_\alpha$ .

If  $\bar{F} \in \mathcal{R}_{-\alpha}, \alpha > 0$ , the tail of  $F$  decays like a power function (Pareto like).

## The Fréchet case

### Theorem 5.6 (Fréchet MDA, Gnedenko (1943))

$F \in \text{MDA}(H_\xi)$  for  $\xi > 0$  if and only if  $\bar{F}(x) = x^{-1/\xi}L(x)$  for some  $L \in \mathcal{R}_0$ . If  $F \in \text{MDA}(H_\xi)$ ,  $\xi > 0$ , the normalizing sequences can be chosen as  $c_n = F^\leftarrow(1 - 1/n)$  and  $d_n = 0$ ,  $n \in \mathbb{N}$ .


- **Interpretation:** Distributions in  $\text{MDA}(H_\xi)$ ,  $\xi > 0$ , are those whose tails decay like power functions;  $\alpha = 1/\xi$  is known as *tail index*.
- **Examples in  $\text{MDA}(H_\xi)$ ,  $\xi > 0$ :** Inverse gamma, Student  $t$ , log-gamma,  $F$ , Cauchy,  $\alpha$ -stable with  $0 < \alpha < 2$ , Burr and Pareto

### Example 5.7 (Pareto distribution)

For  $F = \text{Par}(\theta, \kappa)$ ,  $\bar{F}(x) = (\kappa/(\kappa + x))^\theta = (1 + x/\kappa)^{-\theta} = x^{-\theta}L(x)$ ,  $x \geq 0$ ,  $\theta, \kappa > 0$ , where  $L(x) = (\kappa^{-1} + x^{-1})^{-\theta} \in \mathcal{R}_0$ . We (again) see that  $F \in \text{MDA}(H_\xi)$ ,  $\xi > 0$ .

### 5.1.3 The block maxima method (BMM)

The basic idea in a picture based on losses  $X_1, \dots, X_{12}$ :



Consider the maximal loss from each block and fit  $H_{\xi, \mu, \sigma}$  to them.

## Fitting the GEV distribution

- Suppose  $(x_i)_{i \in \mathbb{N}}$  are realizations of  $(X_i)_{i \in \mathbb{N}} \stackrel{\text{ind.}}{\sim} F \in \text{MDA}(H_\xi)$ ,  $\xi \in \mathbb{R}$ . The Fisher–Tippett–Gnedenko Theorem implies that

$$\mathbb{P}(M_n \leq x) = \mathbb{P}((M_n - d_n)/c_n \leq (x - d_n)/c_n) \underset{n \text{ large}}{\approx} H_{\xi, \mu=d_n, \sigma=c_n}(x).$$

- For fitting  $\theta = (\xi, \mu, \sigma)$ , divide the realizations into  $m$  blocks of size  $n$  denoted by  $M_{n1}, \dots, M_{nm}$  (e.g. daily log-returns  $\Rightarrow$  monthly maxima)
- Assume the block size  $n$  to be sufficiently large so that (regardless of whether the underlying data are dependent or not), the block maxima can be considered independent.
- The density  $h_\xi$  of  $H_\xi$  is

$$h_\xi(x) = \begin{cases} (1 + \xi x)^{-1/\xi-1} H_\xi(x) I_{\{1+\xi x > 0\}}, & \text{if } \xi \neq 0, \\ e^{-x} H_0(x), & \text{if } \xi = 0. \end{cases}$$



The **log-likelihood** is thus

$$\ell(\boldsymbol{\theta}; M_{n1}, \dots, M_{nm}) = \sum_{i=1}^m \log\left(\frac{1}{\sigma} h_{\xi}\left(\frac{M_{ni} - \mu}{\sigma}\right)\right).$$

Maximize w.r.t.  $\boldsymbol{\theta} = (\xi, \mu, \sigma)$  to get  $\hat{\boldsymbol{\theta}} = (\hat{\xi}, \hat{\mu}, \hat{\sigma})$ .

### Remark 5.8

- 1) Sufficiently many/large blocks **require large amounts of data**.
- 2) Bias and variance must be traded off (**bias-variance tradeoff**):
  - Block size  $n \uparrow \Rightarrow$  GEV approximation more accurate  $\Rightarrow$  **bias**  $\downarrow$
  - Number of blocks  $m \uparrow \Rightarrow$  more data for MLE  $\Rightarrow$  **variance**  $\downarrow$
- 3) There is **no general best strategy** known to find the **optimal block size**.

## Return levels and stress losses (exceedances)

Let  $M_n \sim H$  (exact or estimated).  $H$  can be used to estimate the...

- 1) ... size of an event with prescribed frequency (*return-level problem*)
  - The level  $r_{n,k}$  which is expected to be exceeded in one out of every  $k$  blocks of size  $n$  satisfies  $\mathbb{P}(M_n > r_{n,k}) = 1/k$  (e.g. 10-year return level  $r_{260,10}$  = level exceeded in one out of every 10y; 260d  $\approx$  1y).
  - $r_{n,k} = H^{\leftarrow}(1 - 1/k)$  is known as *k n-block return level* with parametric estimator  $\hat{r}_{n,k} = H_{\hat{\xi}, \hat{\mu}, \hat{\sigma}}^{\leftarrow}(1 - 1/k) = \hat{\mu} + \frac{\hat{\sigma}}{\hat{\xi}}((-\log(1 - 1/k))^{-\hat{\xi}} - 1)$ .
- 2) ... frequency of an event with prescribed size (*return-period problem*)
  - The smallest number  $k_{n,u}$  of  $n$ -blocks for which we expect to see at least one  $n$ -block exceeding  $u$  satisfies  $r_{n,k_{n,u}} = u$  (so one has  $\mathbb{P}(M_n > u) = 1/k_{n,u}$ ).
  - $k_{n,u} = 1/\bar{H}(u)$  is known as *return period* of the event  $\{M_n > u\}$  with parametric estimator  $\hat{k}_{n,u} = 1/\bar{H}_{\hat{\xi}, \hat{\mu}, \hat{\sigma}}(u)$ .

### Example 5.9 (Block maxima analysis of S&P500)

Suppose it is Friday 1987-10-16; the Friday before Black Monday (1987-10-19). The S&P 500 index fell by 9.12% this week. On that Friday alone the index is down 5.16%. We fit a GEV distribution to (bi)annual maxima of daily negative log-returns  $X_t = -\log(S_t/S_{t-1})$  since 1960-01-01.

**Analysis 1:** Annual maxima ( $m = 28$ ; including the latest from the incomplete year 1987):  $\hat{\theta} = (0.30, 0.02, 0.007) \Rightarrow$  Heavy-tailed Fréchet distribution (infinite fourth moment). The corresponding standard errors are  $(0.21, 0.002, 0.001) \Rightarrow$  High uncertainty ( $m$  small) for estimating  $\xi$ .

**Analysis 2:** Biannual maxima ( $m = 56$ ):  $\hat{\theta} = (0.34, 0.02, 0.006)$  with standard errors  $(0.14, 0.0009, 0.0008) \Rightarrow$  Even heavier tails. In what follows we work with the annual maxima.

- What is the probability that next year's maximal risk-factor change exceeds all previous ones?  $1 - H_{\hat{\xi}, \hat{\mu}, \hat{\sigma}}(\text{"previous maxima"})$

- Was a risk-factor change as on **Black Monday** foreseeable?
    - ▶ Based on data up to and including Friday 1987-10-16, the 10-year return level  $r_{260,10}$  is estimated as  $\hat{r}_{260,10} = 4.42\%$ .
    - ▶ Index drop Black Monday:  $20.47\% \Rightarrow X_{t+1} = 22.9\% \gg \hat{r}_{260,10}$ .
    - ▶ One can show that  $22.9\%$  is in the 95% confidence interval of  $r_{260,50}$  (estimated as  $\hat{r}_{260,50} = 7.49\%$ ), but the 28 maxima are too few to get a reliable estimate of a once-in-50-years event.
  - Based on the available data, what is the (estimated) **return period of a risk-factor change** at least as large as on **Black Monday**?
    - ▶ The estimated return period  $k_{260,0.229}$  is  $\hat{k}_{260,0.229} = 1877$  years.
    - ▶ One can show that the 95% confidence interval encompasses everything from 45y to essentially never!  $\Rightarrow$  **Very high uncertainty!**
- $\Rightarrow$  On 1987-10-16 we did not have enough data to say anything meaningful about such an event. Quantifying such events is difficult.

## 5.2 Threshold exceedances

The **BMM** is **wasteful of data** (only the maxima of large blocks are used). It has been largely superseded in practice by methods based on **threshold exceedances** (*peaks-over-threshold (POT) approach*), where **all data above** a designated high **threshold**  $u$  are used.

### 5.2.1 Generalized Pareto distribution

#### Definition 5.10 (Generalized Pareto distribution (GPD))

The *generalized Pareto distribution (GPD)* is given by

$$G_{\xi,\beta}(x) = \begin{cases} 1 - (1 + \xi x/\beta)^{-1/\xi}, & \text{if } \xi \neq 0, \\ 1 - \exp(-x/\beta), & \text{if } \xi = 0, \end{cases}$$

where  $\beta > 0$ , and the support is  $x \geq 0$  when  $\xi \geq 0$  and  $x \in [0, -\beta/\xi]$  when  $\xi < 0$ .

- The larger  $\xi$ , the heavier tailed  $G_{\xi,\beta}$  (if  $\xi > 0$ ,  $\mathbb{E}(X^k) = \infty$  iff  $k \geq \frac{1}{\xi}$ ; if  $\xi < 1$ , then  $\mathbb{E}X = \beta/(1 - \xi)$ ).
- $\xi$  is known as *shape*;  $\beta$  as *scale*. Special cases:
  - 1)  $\xi > 0$ :  $\text{Par}(1/\xi, \beta/\xi)$
  - 2)  $\xi = 0$ :  $\text{Exp}(1/\beta)$
  - 3)  $\xi < 0$ : short-tailed Pareto type II distribution
- The density  $g_{\xi,\beta}$  of  $G_{\xi,\beta}$  is given by

$$g_{\xi,\beta}(x) = \begin{cases} \frac{1}{\beta}(1 + \xi x/\beta)^{-1/\xi-1}, & \text{if } \xi \neq 0, \\ \frac{1}{\beta} \exp(-x/\beta), & \text{if } \xi = 0, \end{cases}$$

where  $x \geq 0$  when  $\xi \geq 0$  and  $x \in [0, -\beta/\xi)$  when  $\xi < 0$  (MLE!).

- $G_{\xi,\beta} \in \text{MDA}(H_\xi)$ ,  $\xi \in \mathbb{R}$ .

### Definition 5.11 (Excess distribution over $u$ , mean excess function)

Let  $X \sim F$ . The *excess distribution over the threshold  $u$*  is defined by

$$F_u(x) = \mathbb{P}(X - u \leq x \mid X > u) = \frac{F(x + u) - F(u)}{1 - F(u)}, \quad x \in [0, x_F - u).$$

If  $\mathbb{E}|X| < \infty$ , the *mean excess function* is defined by

$$e(u) = \mathbb{E}(X - u \mid X > u) \quad (\text{i.e. the mean w.r.t. } F_u)$$

### Interpretation

$F_u$  describes the distribution of the excess loss  $X - u$  over  $u$ , given that  $X$  exceeds  $u$ .  $e(u)$  is the mean of  $F_u$  as a function in  $u$ .

### Example 5.12 ( $F_u$ , $e(u)$ for $\text{Exp}(\lambda)$ , $G_{\xi, \beta}$ )

- 1) If  $F$  is  $\text{Exp}(\lambda)$ , then  $F_u(x) = 1 - e^{-\lambda x}$  and  $e(u) = 1/\lambda = \mathbb{E}X$ .
- 2) If  $F$  is  $G_{\xi, \beta}$ , then  $F_u(x) = G_{\xi, \beta + \xi u}(x)$  and  $e(u) = \frac{\beta + \xi u}{1 - \xi}$  for all  $u : \beta + \xi u > 0$ , which is linear in  $u$ .

### Theorem 5.13 (Pickands–Balkema–de Haan (1974/75))

There exists a positive, measurable function  $\beta(u)$ , such that

$$\lim_{u \uparrow x_F} \sup_{0 \leq x < x_F - u} |F_u(x) - G_{\xi, \beta(u)}(x)| = 0.$$

if and only if  $F \in \text{MDA}(H_\xi)$ ,  $\xi \in \mathbb{R}$ .

*Proof.* Non-trivial; see, e.g. Pickands (1975) and Balkema and de Haan (1974). □


### Interpretation

- The GPD is the canonical df for excess losses over high  $u$ .
- The result is also a characterization of  $\text{MDA}(H_\xi)$ ,  $\xi \in \mathbb{R}$ . All  $F \in \text{MDA}(H_\xi)$  form a set of df for which the excess distribution converges to the GPD  $G_{\xi, \beta}$  with the same  $\xi$  as in  $H_\xi$  when  $u$  is raised.



## 5.2.2 Modelling excess losses

The basic idea in a picture based on losses  $X_1, \dots, X_{12}$ .



Consider all **excesses over  $u$**  and fit  $G_{\xi, \beta}$  to them.

## The peaks-over-threshold (POT) method

- Given losses  $X_1, \dots, X_n \sim F \in \text{MDA}(H_\xi)$ ,  $\xi \in \mathbb{R}$ , let
  - ▶  $N_u = |\{i \in \{1, \dots, n\} : X_i > u\}|$  denote the *number of exceedances* over the (given; see later) threshold  $u$ ;
  - ▶  $\tilde{X}_1, \dots, \tilde{X}_{N_u}$  denote the *exceedances*; and
  - ▶  $Y_k = \tilde{X}_k - u$ ,  $k \in \{1, \dots, N_u\}$ , the corresponding *excesses*.
- If  $Y_1, \dots, Y_{N_u}$  are iid and (roughly) distributed as  $G_{\xi, \beta}$ , the *log-likelihood* is given by

$$\begin{aligned}\ell(\xi, \beta; Y_1, \dots, Y_{N_u}) &= \sum_{k=1}^{N_u} \log g_{\xi, \beta}(Y_k) \\ &= -N_u \log(\beta) - (1 + 1/\xi) \sum_{k=1}^{N_u} \log(1 + \xi Y_k / \beta)\end{aligned}$$

$\Rightarrow$  Maximize w.r.t.  $\beta > 0$  and  $1 + \xi Y_k / \beta > 0$  for all  $k \in \{1, \dots, N_u\}$ .

## Excesses over higher thresholds

Once a model is fitted to  $F_u$ , we can infer a model for  $F_v$ ,  $v \geq u$ .

### Lemma 5.14

Assume, for some  $u$ ,  $F_u(x) = G_{\xi, \beta}(x)$  for  $0 \leq x < x_F - u$ . Then  $F_v(x) = G_{\xi, \beta + \xi(v-u)}(x)$  for all  $v \geq u$ .

$\Rightarrow$  The excess distribution over  $v \geq u$  remains GPD with the same  $\xi$  (and  $\beta$  growing linearly in  $v$ ); makes sense for a limiting distribution for  $u \uparrow$ .

If  $\xi < 1$  (so if it exists), the mean excess function is given by

$$e(v) = \frac{\xi}{1 - \xi}v + \frac{\beta - \xi u}{1 - \xi}, \quad v \in [u, \infty) \text{ if } \xi \in [0, 1), \quad (10)$$

and  $v \in [u, u - \beta/\xi]$  if  $\xi < 0$ . This forms the basis for a graphical method for choosing  $u$ .

## Sample mean excess plot and choice of the threshold

### Definition 5.15 (Sample mean excess function, mean excess plot)

For  $X_1, \dots, X_n > 0$ , the *sample mean excess function* is defined by

$$e_n(v) = \frac{\sum_{i=1}^n (X_i - v) I_{\{X_i > v\}}}{\sum_{i=1}^n I_{\{X_i > v\}}}, \quad v < X_{(n)}.$$


The *mean excess plot* is the plot of  $\{(X_{(i)}, e_n(X_{(i)})) : 1 \leq i \leq n-1\}$ , where  $X_{(i)}$  denotes the  $i$ th order statistic.

- If the data supports the GPD model over  $u$ ,  $e_n(v)$  should become increasingly “linear” for higher values of  $v \geq u$ . Select  $u$  as the smallest point where  $e_n(v)$ ,  $v \geq u$ , becomes linear.
- $e_n(v)$  is rarely perfectly linear. The choice of a good threshold  $u$  is as difficult as finding an adequate block size for the Block Maxima method. One should always analyze the data for several  $u$ .

### Example 5.16 (Danish fire loss data)


- 2156 fire insurance losses over 1M Danish kroner from 1980-01-03 to 1990-12-31; combined loss for a building and its contents, in some cases also a loss of business earnings. The losses are inflation adjusted to reflect values as of 1985.
- The mean excess function shows a “kink” below 10; “straightening out” above 10  $\Rightarrow$  Our choice is  $u = 10$  (so 10M Danish kroner).
- MLE  $(\hat{\xi}, \hat{\beta}) = (0.50, 7.0)$  (with standard errors  $(0.14, 1.1)$ )  
 $\Rightarrow$  very heavy-tailed, infinite-variance model
- We can then estimate the expected loss given exceedance of 10M kroner or any higher threshold (via  $e(v)$  in (10) based on  $\hat{\xi}, \hat{\beta}$  and the chosen  $u$ ), even beyond the data.  
 $\Rightarrow$  EVT allows us to estimate “in the data” and then “scale up”.

(a): Losses ( $> 1M$ ; in  $M$ ); (b):  $e_n(u)$  ( $\uparrow$ ); (c)  $\hat{F}_{u,n}(x - u)$ ,  $G_{\hat{\xi}, \hat{\beta}}(x - u)$



$\Rightarrow$  Choose the threshold  $u = 10$ .

Sensitivity of the estimated shape parameter  $\hat{\xi}$  to changes in  $u$ :



⇒ The higher  $u$ , the wider the confidence intervals (also support  $u = 10$ ).

## 5.2.3 Modelling tails and measures of tail risk

- How can the fitted GPD model be used to estimate the tail of the loss distribution  $F$  and associated risk measures?
- Assume  $F_u(x) = G_{\xi, \beta}(x)$  for  $0 \leq x < x_F - u$ ,  $\xi \neq 0$  and some  $u$ .
- We obtain the following GPD-based formula for tail probabilities:

$$\begin{aligned}\bar{F}(x) &= \mathbb{P}(X > x) = \mathbb{P}(X > u) \mathbb{P}(X > x \mid X > u) \\ &= \bar{F}(u) \mathbb{P}(X - u > x - u \mid X > u) = \bar{F}(u) \bar{F}_u(x - u) \\ &= \bar{F}(u) \left(1 + \xi \frac{x - u}{\beta}\right)^{-1/\xi}, \quad x \geq u.\end{aligned}\tag{11}$$

- Assuming we know  $\bar{F}(u)$ , inverting this formula for  $\alpha \geq F(u)$  leads to

$$\text{VaR}_\alpha = F^{\leftarrow}(\alpha) = u + \frac{\beta}{\xi} \left( \left( \frac{1 - \alpha}{\bar{F}(u)} \right)^{-\xi} - 1 \right),\tag{12}$$

$$\text{ES}_\alpha = \frac{\text{VaR}_\alpha}{1 - \xi} + \frac{\beta - \xi u}{1 - \xi}, \quad \xi < 1.\tag{13}$$



- $\bar{F}(x)$ ,  $\text{VaR}_\alpha$  and  $\text{ES}_\alpha$  are all of the form  $g(\xi, \beta, \bar{F}(u))$ . If we have sufficient samples above  $u$ , we obtain semi-parametric plug-in estimators via  $g(\hat{\xi}, \hat{\beta}, N_u/n)$ .
- We hope to gain over empirical estimators by using a kind of extrapolation based on the GPD for more extreme tail probabilities and risk measures.
- In this spirit, Smith (1987) proposed the *tail estimator*


$$\hat{\bar{F}}(x) = \frac{N_u}{n} \left( 1 + \hat{\xi} \frac{x - u}{\hat{\beta}} \right)^{-1/\hat{\xi}}, \quad x \geq u \quad (\text{see (11)});$$

also known as the *Smith estimator* (note that it is only valid for  $x \geq u$ ). It faces a **bias-variance tradeoff**: If  $u$  is increased, the bias of parametrically estimating  $\bar{F}_u(x - u)$  decreases, but the variance of it and the nonparametrically estimated  $\bar{F}(u)$  increases.

- Similarly, semi-parametric GPD-based  $\widehat{\text{VaR}}_\alpha$ ,  $\widehat{\text{ES}}_\alpha$  for  $\alpha \geq 1 - N_u/n$  can be obtained from (12), (13).

### Example 5.17 (Danish fire loss data (continued))

Here are  $\hat{\bar{F}}(x)$ ,  $x \geq u$ ,  $\widehat{\text{VaR}}_{0.99}$ ,  $\widehat{\text{ES}}_{0.99}$  including confidence intervals.



Log-log scale often helpful: If  $\bar{F}(x) = x^{-\alpha}L(x)$ ,  $\log \bar{F}(x) = -\alpha \log(x) + \log L(x)$  which is approximately linear in  $\log x$ .

## 5.2.4 Conditional EVT for financial time series

- Assume  $X_{t-n+1}, \dots, X_t$  are negative log-returns generated by a strictly stationary time series process  $(X_t)$  of the form

$$X_t = \mu_t + \sigma_t Z_t,$$

where  $\mu_t$  and  $\sigma_t$  are  $\mathcal{F}_{t-1}$ -measurable and  $Z_t \stackrel{\text{ind.}}{\sim} F_Z$ ; e.g. ARMA model with GARCH errors. Furthermore, let  $Z \sim F_Z$ .

- $\text{VaR}_\alpha^t$  and  $\text{ES}_\alpha^t$  based on  $F_{X_{t+1}|\mathcal{F}_t}$  are given by

$$\text{VaR}_\alpha^t(X_{t+1}) = \mu_{t+1} + \sigma_{t+1} \text{VaR}_\alpha(Z),$$

$$\text{ES}_\alpha^t(X_{t+1}) = \mu_{t+1} + \sigma_{t+1} \text{ES}_\alpha(Z).$$

- To obtain estimates  $\widehat{\text{VaR}}_\alpha^t(X_{t+1})$  and  $\widehat{\text{ES}}_\alpha^t(X_{t+1})$ , proceed as follows:
  - Fit an ARMA-GARCH model  $\Rightarrow$  Estimates of  $\mu_{t+1}$  and  $\sigma_{t+1}$ .
  - Fit a GPD to  $F_Z \Rightarrow$  GPD-based estimates of  $\text{VaR}_\alpha(Z)$  (see (12)) and  $\text{ES}_\alpha(Z)$  (see (13)).

# 6 Multivariate models

- 6.1 Basics of multivariate modelling
- 6.2 Normal mixture distributions
- 6.3 Spherical and elliptical distributions
- 6.4 Dimension reduction techniques

# 6.1 Basics of multivariate modelling

## 6.1.1 Random vectors and their distributions

### Joint and marginal distributions

- Let  $\mathbf{X} = (X_1, \dots, X_d) : \Omega \rightarrow \mathbb{R}^d$  be a  $d$ -dimensional *random vector* (representing risk-factor changes, risks, etc.).
- The *(joint) distribution function (df)  $F$  of  $\mathbf{X}$*  is

$$F(\mathbf{x}) = F_{\mathbf{X}}(\mathbf{x}) = \mathbb{P}(\mathbf{X} \leq \mathbf{x}) = \mathbb{P}(X_1 \leq x_1, \dots, X_d \leq x_d), \quad \mathbf{x} \in \mathbb{R}^d.$$

- The  *$j$ th margin or marginal df  $F_j$  of  $\mathbf{X}$*  is

$$\begin{aligned} F_j(x_j) &= \mathbb{P}(X_j \leq x_j) \\ &= \mathbb{P}(X_1 \leq \infty, \dots, X_{j-1} \leq \infty, X_j \leq x_j, X_{j+1} \leq \infty, \dots, X_d \leq \infty) \\ &= F(\infty, \dots, \infty, x_j, \infty, \dots, \infty), \quad x_j \in \mathbb{R}, \quad j \in \{1, \dots, d\}. \end{aligned}$$

(interpreted as a *limit*).

- Similarly for *k-dimensional margins*. Suppose we partition  $\mathbf{X}$  into  $(\mathbf{X}'_1, \mathbf{X}'_2)'$ , where  $\mathbf{X}_1 = (X_1, \dots, X_k)'$  and  $\mathbf{X}_2 = (X_{k+1}, \dots, X_d)'$ , then the marginal distribution function of  $\mathbf{X}_1$  is

$$F_{\mathbf{X}_1}(\mathbf{x}_1) = \mathbb{P}(\mathbf{X}_1 \leq \mathbf{x}_1) = F(x_1, \dots, x_k, \infty, \dots, \infty).$$

- *F is absolutely continuous* if

$$F(\mathbf{x}) = \int_{-\infty}^{x_d} \dots \int_{-\infty}^{x_1} f(z_1, \dots, z_d) dz_1 \dots dz_d = \int_{(-\infty, \mathbf{x}]} f(\mathbf{z}) d\mathbf{z}$$

for some  $f \geq 0$  known as the *(joint) density of  $\mathbf{X}$  (or  $F$ )*. Similarly, the *jth marginal df  $F_j$  is absolutely continuous* if  $F_j(x) = \int_{-\infty}^x f_j(z) dz$  for some  $f_j \geq 0$  known as the *density of  $X_j$  (or  $F_j$ )*.

- Existence of a *joint density*  $\Rightarrow$  Existence of *marginal densities* for all  $k$ -dimensional marginals,  $1 \leq k \leq d - 1$ . The *converse is false in general* (counter-examples can be constructed with singular *copulas*; see Chapter 7).
- We sometimes work with the *survival function  $\bar{F}$  of  $\mathbf{X}$* ,

$$\bar{F}(\mathbf{x}) = \bar{F}_{\mathbf{X}}(\mathbf{x}) = \mathbb{P}(\mathbf{X} > \mathbf{x}) = \mathbb{P}(X_1 > x_1, \dots, X_d > x_d), \quad \mathbf{x} \in \mathbb{R}^d,$$

with corresponding *j*th marginal survival function  $\bar{F}_j$

$$\begin{aligned}\bar{F}_j(x_j) &= \mathbb{P}(X_j > x_j) \\ &= \bar{F}(-\infty, \dots, -\infty, x_j, -\infty, \dots, -\infty), \quad x_j \in \mathbb{R}, \quad j \in \{1, \dots, d\}.\end{aligned}$$

- Note that  $\bar{F}(\mathbf{x}) \neq 1 - F(\mathbf{x})$  in general (unless  $d = 1$ ).

## Independence

- A **multivariate model** for risks in the form of a joint df, survival function or density, **implicitly describes** their **dependence structure**.
- $\mathbf{X}_1, \mathbf{X}_2$  are **independent** if  $F(\mathbf{x}_1, \mathbf{x}_2) = F_{\mathbf{X}_1}(\mathbf{x}_1)F_{\mathbf{X}_2}(\mathbf{x}_2)$  for all  $\mathbf{x}_1, \mathbf{x}_2$ .
- If  $F$  has density  $f$ , then  $\mathbf{X}_1, \mathbf{X}_2$  are independent if  $f(\mathbf{x}_1, \mathbf{x}_2) = f_{\mathbf{X}_1}(\mathbf{x}_1)f_{\mathbf{X}_2}(\mathbf{x}_2)$  for all  $\mathbf{x}_1, \mathbf{x}_2$ .
- The components  $X_1, \dots, X_d$  of  $\mathbf{X}$  are **(mutually) independent** if  $F(\mathbf{x}) = \prod_{j=1}^d F_j(x_j)$  for all  $\mathbf{x}$  or, if  $F$  has density  $f$ , if  $f(\mathbf{x}) = \prod_{j=1}^d f_j(x_j)$  for all  $\mathbf{x}$ .

## Moments and characteristic function

- If  $\mathbb{E}|X_j| < \infty$ ,  $j \in \{1, \dots, d\}$ , the *mean vector of  $\mathbf{X}$*  is defined by

$$\mathbb{E}\mathbf{X} = (\mathbb{E}X_1, \dots, \mathbb{E}X_d).$$

One can show:  $X_1, \dots, X_d$  independent  $\Rightarrow \mathbb{E}(X_1 \cdots X_d) = \prod_{j=1}^d \mathbb{E}(X_j)$

- If  $\mathbb{E}(X_j^2) < \infty$  for all  $j$ , the *covariance matrix of  $\mathbf{X}$*  is defined by

$$\text{cov}(\mathbf{X}) = \mathbb{E}((\mathbf{X} - \mathbb{E}\mathbf{X})(\mathbf{X} - \mathbb{E}\mathbf{X})').$$

If we write  $\Sigma = \text{cov}(\mathbf{X})$ , its  $(i, j)$ th element is

$$\begin{aligned}\sigma_{ij} &= \Sigma_{ij} = \text{cov}(X_i, X_j) = \mathbb{E}((X_i - \mathbb{E}X_i)(X_j - \mathbb{E}X_j)) \\ &= \mathbb{E}(X_i X_j) - \mathbb{E}(X_i)\mathbb{E}(X_j);\end{aligned}$$

the diagonal elements are  $\sigma_{jj} = \text{var}(X_j)$ ,  $j \in \{1, \dots, d\}$ .

- $X_1, X_2$  independent  $\not\Rightarrow \text{cov}(X_1, X_2) = 0$  (counter-examples can be constructed with *copulas*; see Chapter 7).



- If  $\mathbb{E}(X_j^2) < \infty$ ,  $j \in \{1, \dots, d\}$ , the *correlation matrix* of  $\mathbf{X}$  is defined by the matrix  $\text{corr}(\mathbf{X})$  with  $(i, j)$ th element

$$\text{corr}(X_i, X_j) = \frac{\text{cov}(X_i, X_j)}{\sqrt{\text{var}(X_i) \text{var}(X_j)}}, \quad i, j \in \{1, \dots, d\},$$

which is in  $[-1, 1]$  with  $\text{corr}(X_i, X_j) = \pm 1$  if and only if  $X_j \stackrel{\text{a.s.}}{=} aX_i + b$  for some  $a \neq 0$  and  $b \in \mathbb{R}$ .

- **Some properties of  $\mathbb{E}()$  and  $\text{cov}()$ :**

1) For all  $A \in \mathbb{R}^{k \times d}$ ,  $\mathbf{b} \in \mathbb{R}^k$ :

- ▶  $\mathbb{E}(A\mathbf{X} + \mathbf{b}) = A\mathbb{E}\mathbf{X} + \mathbf{b} = A\boldsymbol{\mu} + \mathbf{b}$ ;
- ▶  $\text{cov}(A\mathbf{X} + \mathbf{b}) = A \text{cov}(\mathbf{X}) A' = A\Sigma A'$ ; if  $k = 1$  ( $A = \mathbf{a}'$ ),

$$\mathbf{a}'\Sigma\mathbf{a} = \text{cov}(\mathbf{a}'\mathbf{X}) = \text{var}(\mathbf{a}'\mathbf{X}) \geq 0, \quad \mathbf{a} \in \mathbb{R}^d, \quad (14)$$

i.e. *covariance matrices are positive semidefinite*.

2) If  $\Sigma$  is a *positive definite matrix* (i.e.  $\mathbf{a}'\Sigma\mathbf{a} > 0$  for all  $\mathbf{a} \in \mathbb{R}^d \setminus \{\mathbf{0}\}$ ), one can show that  $\Sigma$  is invertible.

3) A symmetric, positive (semi)definite  $\Sigma$  can be written as

$$\Sigma = AA' \quad \text{Cholesky decomposition} \quad (15)$$

for a lower triangular matrix  $A$  with  $A_{jj} > 0$  ( $A_{jj} \geq 0$ ) for all  $j$ .  $A$  is known as *Cholesky factor* (and is also denoted by  $\Sigma^{1/2}$ ).

- Properties of  $\mathbf{X}$  can often be shown with the *characteristic function* (cf)

$$\phi_{\mathbf{X}}(\mathbf{t}) = \mathbb{E}(\exp(i\mathbf{t}'\mathbf{X})), \quad \mathbf{t} \in \mathbb{R}^d.$$

$X_1, \dots, X_d$  are independent  $\Leftrightarrow \phi_{\mathbf{X}}(\mathbf{t}) = \prod_{j=1}^d \phi_{X_j}(t_j)$  for all  $\mathbf{t}$ .

### Proposition 6.1 (Characterization of covariance matrices)

A symmetric matrix  $\Sigma$  is a covariance matrix if and only if it is positive semidefinite.

## 6.1.2 Standard estimators of covariance and correlation

- Assume  $\mathbf{X}_1, \dots, \mathbf{X}_n \sim F$  (daily/weekly/monthly/yearly risk-factor changes) are **serially uncorrelated** (i.e. multivariate white noise) with  $\boldsymbol{\mu} := \mathbb{E}\mathbf{X}_1$ ,  $\Sigma := \text{cov } \mathbf{X}_1$  and  $P = \text{corr}(\mathbf{X}_1)$ .
- Standard estimators of  $\boldsymbol{\mu}, \Sigma, P$  are

$$\bar{\mathbf{X}} = \frac{1}{n} \sum_{i=1}^n \mathbf{X}_i \quad (\text{sample mean})$$

$$S = \frac{1}{n} \sum_{i=1}^n (\mathbf{X}_i - \bar{\mathbf{X}})(\mathbf{X}_i - \bar{\mathbf{X}})' \quad (\text{sample covariance matrix})$$

$$R = (R_{ij}) \text{ for } R_{ij} = \frac{S_{ij}}{\sqrt{S_{ii}S_{jj}}} \quad (\text{sample correlation matrix})$$

- Under joint normality ( $F$  multivariate normal),  $\bar{\mathbf{X}}$ ,  $S$  and  $R$  are also **MLEs**.  $S$  is biased, but an unbiased version can be obtained by

$$S_n = \frac{n}{n-1} S.$$

## 6.1.3 The multivariate normal distribution

### Definition 6.2 (Multivariate normal distribution)

$\mathbf{X} = (X_1, \dots, X_d)$  has a *multivariate normal* (or *Gaussian*) *distribution* if

$$\mathbf{X} \stackrel{d}{=} \boldsymbol{\mu} + \mathbf{A}\mathbf{Z}, \quad (16)$$

where  $\mathbf{Z} = (Z_1, \dots, Z_k)$ ,  $Z_l \stackrel{\text{ind.}}{\sim} \mathcal{N}(0, 1)$ ,  $\mathbf{A} \in \mathbb{R}^{d \times k}$ ,  $\boldsymbol{\mu} \in \mathbb{R}^d$ .

- $\mathbb{E}\mathbf{X} = \boldsymbol{\mu} + \mathbf{A}\mathbb{E}\mathbf{Z} = \boldsymbol{\mu}$
- $\text{cov}(\mathbf{X}) = \text{cov}(\boldsymbol{\mu} + \mathbf{A}\mathbf{Z}) = \mathbf{A} \text{cov}(\mathbf{Z}) \mathbf{A}' = \mathbf{A}\mathbf{A}' =: \Sigma$

### Proposition 6.3 (Cf of the multivariate normal distribution)

Let  $\mathbf{X}$  be as in (16) and  $\Sigma = \mathbf{A}\mathbf{A}'$ . Then the cf of  $\mathbf{X}$  is

$$\phi_{\mathbf{X}}(\mathbf{t}) = \mathbb{E}(\exp(it'\mathbf{X})) = \exp\left(it'\boldsymbol{\mu} - \frac{1}{2}\mathbf{t}'\Sigma\mathbf{t}\right), \quad \mathbf{t} \in \mathbb{R}^d.$$

- We see that the multivariate normal distribution is characterized by  $\mu$  and  $\Sigma$ , hence the notation  $\mathbf{X} \sim N_d(\mu, \Sigma)$ .





### Proposition 6.4 (Density)

Let  $\mathbf{X} \sim N_d(\mu, \Sigma)$  with  $\text{rank } \Sigma = d = k$  ( $\Rightarrow \Sigma$  pos. definite, invertible).  $\mathbf{X}$  has density

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{1}{(2\pi)^{d/2} \sqrt{\det \Sigma}} \exp\left(-\frac{1}{2}(\mathbf{x} - \mu)' \Sigma^{-1}(\mathbf{x} - \mu)\right), \quad \mathbf{x} \in \mathbb{R}^d.$$

### Consequences:

- Sets of the form  $S_c = \{\mathbf{x} \in \mathbb{R}^d : (\mathbf{x} - \mu)' \Sigma^{-1}(\mathbf{x} - \mu) = c\}$ ,  $c > 0$ , describe points of equal density. Contours of equal density are thus ellipsoids.
- The components of  $\mathbf{X} \sim N_d(\mu, \Sigma)$  are mutually independent if and only if  $\Sigma$  is diagonal, i.e. if and only if the components of  $\mathbf{X}$  are uncorrelated.



Left:  $N_d(\boldsymbol{\mu}, \Sigma)$  for  $\boldsymbol{\mu} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$ ,  $\Sigma = \begin{pmatrix} 1 & -0.7 \\ -0.7 & 1 \end{pmatrix}$ ; Right:  $t_\nu(\boldsymbol{\mu}, \frac{\nu-2}{\nu}\Sigma)$ ,  $\nu = 4$ ,  
 (same mean and covariance matrix as on the left-hand side)

The definition of  $N_d(\boldsymbol{\mu}, \Sigma)$  in terms of a **stochastic representation** ( $\mathbf{X} \stackrel{d}{=} \boldsymbol{\mu} + A\mathbf{Z}$ ) **directly justifies the following sampling algorithm**.

### Algorithm 6.5 (Sampling $N_d(\boldsymbol{\mu}, \Sigma)$ )

Let  $\mathbf{X} \sim N_d(\boldsymbol{\mu}, \Sigma)$  with  $\Sigma$  symmetric and positive definite.

- 1) Compute the Cholesky factor  $A$  of  $\Sigma$ ; see, e.g. Press et al. (1992).
- 2) Generate  $Z_j \stackrel{\text{ind.}}{\sim} N(0, 1)$ ,  $j \in \{1, \dots, d\}$ .
- 3) Return  $\mathbf{X} = \boldsymbol{\mu} + A\mathbf{Z}$ , where  $\mathbf{Z} = (Z_1, \dots, Z_d)$ .

## Further useful properties of multivariate normal distributions

### ■ Linear combinations

If  $\mathbf{X} \sim N_d(\boldsymbol{\mu}, \Sigma)$  and  $B \in \mathbb{R}^{k \times d}$ ,  $\mathbf{b} \in \mathbb{R}^k$ , then

$$\begin{aligned} B\mathbf{X} + \mathbf{b} &= B(\boldsymbol{\mu} + A\mathbf{Z}) + \mathbf{b} = (B\boldsymbol{\mu} + \mathbf{b}) + BA\mathbf{Z} \\ &\sim N_k(B\boldsymbol{\mu} + \mathbf{b}, BA(BA)') = N_k(B\boldsymbol{\mu} + \mathbf{b}, B\Sigma B'). \end{aligned}$$

Special case :  $\mathbf{b}'\mathbf{X} \sim N(\mathbf{b}'\boldsymbol{\mu}, \mathbf{b}'\Sigma\mathbf{b})$

## ■ Marginal dfs

Let  $\mathbf{X} \sim N_d(\boldsymbol{\mu}, \Sigma)$  and write  $\mathbf{X} = (\mathbf{X}'_1, \mathbf{X}'_2)$ , where  $\mathbf{X}_1 \in \mathbb{R}^k$ ,  $\mathbf{X}_2 \in \mathbb{R}^{d-k}$ , and  $\boldsymbol{\mu} = (\boldsymbol{\mu}'_1, \boldsymbol{\mu}'_2)$ ,  $\Sigma = \begin{pmatrix} \Sigma_{11} & \Sigma_{12} \\ \Sigma_{21} & \Sigma_{22} \end{pmatrix}$ . Then

$$\mathbf{X}_1 \sim N_k(\boldsymbol{\mu}_1, \Sigma_{11}) \quad \text{and} \quad \mathbf{X}_2 \sim N_{d-k}(\boldsymbol{\mu}_2, \Sigma_{22}).$$

## ■ Quadratic forms

Let  $\mathbf{X} \sim N_d(\boldsymbol{\mu}, \Sigma)$  and  $\Sigma$  be positive definite with Cholesky factor  $A$ . Furthermore, let  $\mathbf{Z} = A^{-1}(\mathbf{X} - \boldsymbol{\mu})$ . Then  $\mathbf{Z} \sim N_d(\mathbf{0}, I_d)$ . Moreover,

$$(\mathbf{X} - \boldsymbol{\mu})' \Sigma^{-1} (\mathbf{X} - \boldsymbol{\mu}) = \mathbf{Z}' \mathbf{Z} \sim \chi_d^2, \quad (17)$$

which is useful for (goodness-of-fit) testing of  $N_d(\boldsymbol{\mu}, \Sigma)$ .

## 6.1.4 Testing multivariate normality

- For testing **univariate normality**, all tests of Section 3.1.2 can be applied.
- Now consider **multivariate normality**.

$$\mathbf{X}_1, \dots, \mathbf{X}_n \stackrel{\text{ind.}}{\sim} N_d(\boldsymbol{\mu}, \Sigma) \Rightarrow \mathbf{a}' \mathbf{X}_1, \dots, \mathbf{a}' \mathbf{X}_n \stackrel{\text{ind.}}{\sim} N(\mathbf{a}' \boldsymbol{\mu}, \mathbf{a}' \Sigma \mathbf{a}).$$



This can be tested statistically (for some  $\alpha$ ) with various goodness-of-fit tests (e.g. Q-Q plots) used for univariate normality. Alternatively, (17) can be used to test joint normality.

- Multivariate Shapiro–Wilk

- Mardia's test

- ▶ According to (17), if  $\mathbf{X} \sim N_d(\boldsymbol{\mu}, \Sigma)$  with  $\Sigma$  positive definite, then  $(\mathbf{X} - \boldsymbol{\mu})' \Sigma^{-1} (\mathbf{X} - \boldsymbol{\mu}) \sim \chi_d^2$ .
- ▶ Let  $D_i^2 = (\mathbf{X}_i - \bar{\mathbf{X}})' S^{-1} (\mathbf{X}_i - \bar{\mathbf{X}})$  denote the *squared Mahalanobis distances* and  $D_{ij} = (\mathbf{X}_i - \bar{\mathbf{X}})' S^{-1} (\mathbf{X}_j - \bar{\mathbf{X}})$  the *Mahalanobis angles*.
- ▶ Let  $b_d = \frac{1}{n^2} \sum_{i=1}^n \sum_{j=1}^n D_{ij}^3$  and  $k_d = \frac{1}{n} \sum_{i=1}^n D_i^4$ . Under the null hypothesis one can show that asymptotically for  $n \rightarrow \infty$ ,

$$\frac{n}{6} b_d \sim \chi_{d(d+1)(d+2)/6}^2, \quad \frac{k_d - d(d+2)}{\sqrt{8d(d+2)/n}} \sim N(0, 1),$$


which can be used for testing; see Joensuu and Vogel (2014).

- We can also compare  $D_i^2$  data to a  $\chi_{10}^2$  graphically using a Q-Q plot.


### Example 6.6 (Simulated data vs BMW–Siemens)

Is the BMW–Siemens data (see Section 3.2.2) jointly normal?

Simulated data (fitted multivariate normal)




Real risk-factor changes




Considering the **first margin** only:

**Q-Q plot for margin 1 (simulated data)**




**Q-Q plot for margin 1 (real data)**




Considering the **second margin** only:

**Q-Q plot for margin 2 (simulated data)**




**Q-Q plot for margin 2 (real data)**




Q-Q plot of the simulated (left) or real (right)  $D_i^2$ 's against a  $\chi_2^2$ :

Q-Q plot of  $D_i^2$  (simulated data)



Q-Q plot of  $D_i^2$  (real data)



## Advantages of $N_d(\mu, \Sigma)$

- Inference “easy”.
- Distribution is determined by  $\mu$  and  $\Sigma$ .
- Linear combinations are normal ( $\Rightarrow$   $\text{VaR}_\alpha$  and  $\text{ES}_\alpha$  calculations for portfolios are easy).
- Marginal distributions are normal.
- Conditional distributions are normal.
- Quadratic forms are known.
- Convolutions are normal.
- Sampling is straightforward.
- Independence and uncorrelatedness are equivalent.

## Drawbacks of $N_d(\boldsymbol{\mu}, \Sigma)$ for modelling risk-factor changes

- 1) **Tails of univariate** (normal) **margins are too thin** (generate too few extreme events).
- 2) **Joint tails are too thin** (generate too few joint extreme events).  
 $N_d(\boldsymbol{\mu}, \Sigma)$  **cannot capture** the notion of **tail dependence** (see Chapter 7).
- 3) Very strong symmetry known as **radial symmetry**:  $\mathbf{X}$  is called **radially symmetric about  $\boldsymbol{\mu}$**  if  $\mathbf{X} - \boldsymbol{\mu} \stackrel{d}{=} \boldsymbol{\mu} - \mathbf{X}$ . This is true for  $N_d(\boldsymbol{\mu}, \Sigma)$ .

### Short outlook:

- **Normal variance mixtures** (or, more general, **elliptical distributions** **can address 1) and 2)** while sharing many of the desirable properties of  $N_d(\boldsymbol{\mu}, \Sigma)$ .
- **Normal mean-variance mixtures** **can also address 3)** (**but at the expense of tractability** in comparison to  $N_d(\boldsymbol{\mu}, \Sigma)$ ).

## 6.2 Normal mixture distributions

**Idea:** Randomize  $\Sigma$  (and  $\mu$ ) with a non-negative rv  $W$ .

### 6.2.1 Normal variance mixtures

#### Definition 6.7 (Multivariate normal variance mixtures)

The random vector  $\mathbf{X}$  has a (multivariate) *normal variance mixture distribution* if

$$\mathbf{X} \stackrel{d}{=} \boldsymbol{\mu} + \sqrt{W} \mathbf{A} \mathbf{Z}, \quad (18)$$

where  $\mathbf{Z} \sim N_k(\mathbf{0}, I_k)$ ,  $W \geq 0$  is a rv independent of  $\mathbf{Z}$ ,  $\mathbf{A} \in \mathbb{R}^{d \times k}$ , and  $\boldsymbol{\mu} \in \mathbb{R}^d$ .  $\boldsymbol{\mu}$  is called *location vector* and  $\Sigma = \mathbf{A} \mathbf{A}'$  *scale* (or *dispersion matrix*).

Observe that  $(\mathbf{X} | W = w) \stackrel{d}{=} \boldsymbol{\mu} + \sqrt{w} \mathbf{A} \mathbf{Z} = N_d(\boldsymbol{\mu}, w \mathbf{A} \mathbf{A}') = N_d(\boldsymbol{\mu}, w \Sigma)$ ; or  $(\mathbf{X} | W) \stackrel{d}{=} N_d(\boldsymbol{\mu}, W \Sigma)$ .  $W$  can be interpreted as a *shock* affecting the variances of all risk factors.



## Properties of multivariate normal variance mixtures

Let  $\mathbf{X} = \boldsymbol{\mu} + \sqrt{W}\mathbf{A}\mathbf{Z}$  and  $\mathbf{Y} = \boldsymbol{\mu} + \mathbf{A}\mathbf{Z}$ . Assume that  $\text{rank}(\mathbf{A}) = d \leq k$  and that  $\Sigma$  is positive definite.

■ If  $\mathbb{E}\sqrt{W} < \infty$ , then  $\mathbb{E}(\mathbf{X}) \stackrel{\text{ind.}}{=} \boldsymbol{\mu} + \mathbb{E}(\sqrt{W})\mathbf{A}\mathbb{E}(\mathbf{Z}) = \boldsymbol{\mu} + \mathbf{0} = \boldsymbol{\mu} = \mathbb{E}\mathbf{Y}$

■ If  $\mathbb{E}W < \infty$ , then

$$\begin{aligned}\text{cov}(\mathbf{X}) &= \text{cov}(\sqrt{W}\mathbf{A}\mathbf{Z}) = \mathbb{E}((\sqrt{W}\mathbf{A}\mathbf{Z})(\sqrt{W}\mathbf{A}\mathbf{Z})') \\ &\stackrel{\text{ind.}}{=} \mathbb{E}(W) \cdot \mathbb{E}(\mathbf{A}\mathbf{Z}\mathbf{Z}'\mathbf{A}') = \mathbb{E}(W) \cdot \mathbf{A}\mathbb{E}(\mathbf{Z}\mathbf{Z}')\mathbf{A}' \\ &= \mathbb{E}(W)\mathbf{A}\mathbf{I}_k\mathbf{A}' = \mathbb{E}(W)\Sigma \neq \Sigma \quad (\text{in general} \quad (= \text{cov}(\mathbf{Y})))\end{aligned}$$

■ However, if they exist (i.e. if  $\mathbb{E}W < \infty$ )  $\text{corr}(\mathbf{X}) = \text{corr}(\mathbf{Y})$  since

$$\begin{aligned}\text{corr}(X_i, X_j) &= \frac{\text{cov}(X_i, X_j)}{\sqrt{\text{var}(X_i) \text{var}(X_j)}} = \frac{\mathbb{E}(W)\Sigma_{ij}}{\sqrt{\mathbb{E}(W)\Sigma_{ii}\mathbb{E}(W)\Sigma_{jj}}} \\ &= \frac{\Sigma_{ij}}{\sqrt{\Sigma_{ii}\Sigma_{jj}}} = \text{corr}(Y_i, Y_j), \quad i, j \in \{1, \dots, d\}.\end{aligned}$$

**Recall:** If  $\mathbf{X} \sim N_d(\boldsymbol{\mu}, \Sigma)$ , then  $\phi_{\mathbf{X}}(\mathbf{t}) = \exp(i\mathbf{t}'\boldsymbol{\mu} - \frac{1}{2}\mathbf{t}'\Sigma\mathbf{t})$ .

Furthermore,  $\mathbf{X} \mid W = w \sim N_d(\boldsymbol{\mu}, w\Sigma)$

- **Characteristic function:** The cf of a multivariate normal variance mixtures is

$$\begin{aligned}\phi_{\mathbf{X}}(\mathbf{t}) &= \mathbb{E}(\exp(i\mathbf{t}'\mathbf{X})) = \mathbb{E}(\mathbb{E}(\exp(i\mathbf{t}'\mathbf{X}) \mid W)) \\ &= \mathbb{E}(\exp(i\mathbf{t}'\boldsymbol{\mu} - \frac{1}{2}W\mathbf{t}'\Sigma\mathbf{t})) = \exp(i\mathbf{t}'\boldsymbol{\mu})\mathbb{E}(\exp(-W\frac{1}{2}\mathbf{t}'\Sigma\mathbf{t})).\end{aligned}$$

This depends on the *Laplace-Stieltjes transform*  $\hat{F}_W$  of  $F_W$ . We thus introduce the notation  $\mathbf{X} \sim M_d(\boldsymbol{\mu}, \Sigma, \hat{F}_W)$  for a  $d$ -dimensional multivariate normal variance mixture.

- **Density:** If  $\Sigma$  is positive definite,  $\mathbb{P}(W = 0) = 0$ , the density of  $\mathbf{X}$  is

$$\begin{aligned}f_{\mathbf{X}}(\mathbf{x}) &= \int_0^\infty f_{\mathbf{X} \mid W}(\mathbf{x} \mid w) dF_W(w) \\ &= \int_0^\infty \frac{1}{(2\pi)^{d/2}w^{d/2}|\Sigma|^{1/2}} \exp\left(-\frac{(\mathbf{x} - \boldsymbol{\mu})'\Sigma^{-1}(\mathbf{x} - \boldsymbol{\mu})}{2w}\right) dF_W(w).\end{aligned}$$

⇒ Only depends on  $x$  through  $(x - \mu)' \Sigma^{-1} (x - \mu)$ .

⇒ Multivariate normal variance mixtures are **elliptical distributions**.

If  $\Sigma$  is diagonal and  $\mathbb{E}W < \infty$ ,  $\mathbf{X}$  is **uncorrelated** (as  $\text{cov}(\mathbf{X}) = \mathbb{E}(W)\Sigma$ )  
**but not independent** unless  $W$  is constant a.s.

- **Linear combinations:** For  $\mathbf{X} \sim M_d(\mu, \Sigma, \hat{F}_W)$  and  $\mathbf{Y} = B\mathbf{X} + \mathbf{b}$ , where  $B \in \mathbb{R}^{k \times d}$  and  $\mathbf{b} \in \mathbb{R}^k$ , we have  $\mathbf{Y} \sim M_k(B\mu + \mathbf{b}, B\Sigma B', \hat{F}_W)$ ; this can be shown via cfs.

- **Sampling:**

**Algorithm 6.8 (Simulation of  $\mathbf{X} = \mu + \sqrt{W}AZ \sim M_d(\mu, \Sigma, \hat{F}_W)$ )**

- 1) Generate  $\mathbf{Z} \sim N_d(\mathbf{0}, I_d)$ .
- 2) Generate  $W \sim F_W$  (with LS transform  $\hat{F}_W$ ), independent of  $\mathbf{Z}$ .
- 3) Compute the Cholesky factor  $A$  (such that  $AA' = \Sigma$ ).
- 4) Return  $\mathbf{X} = \mu + \sqrt{W}AZ$ .

## Examples of multivariate normal variance mixtures

- **Multivariate normal distribution**

$W = 1$  a.s. (degenerate case)

- **Two point mixture**

$$W = \begin{cases} w_1 & \text{with probability } p, \\ w_2 & \text{with probability } 1 - p \end{cases} \quad w_1, w_2 > 0, w_1 \neq w_2.$$

Can be used to model **ordinary and stress regimes**; extends to  $k$  regimes.

- **Symmetric generalised hyperbolic distribution**

$W$  has a generalised inverse Gaussian distribution (GIG); see McNeil et al. (2015, p. 187)

- **Multivariate  $t$  distribution**

$W$  has an inverse gamma distribution  $W = 1/V$  for  $V \sim \Gamma(\nu/2, \nu/2)$ .




- ▶  $\mathbb{E}(W) = \frac{\nu}{\nu-2} \Rightarrow \text{cov}(\mathbf{X}) = \frac{\nu}{\nu-2} \Sigma$ . For finite variances/correlations,  $\nu > 2$  is required. For finite mean,  $\nu > 1$  is required.

- The density of the multivariate  $t$  distribution is given by

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{\Gamma((\nu + d)/2)}{\Gamma(\nu/2)(\nu\pi)^{d/2}|\Sigma|^{1/2}} \left( 1 + \frac{(\mathbf{x} - \boldsymbol{\mu})'\Sigma^{-1}(\mathbf{x} - \boldsymbol{\mu})}{\nu} \right)^{-\frac{\nu+d}{2}},$$

where  $\boldsymbol{\mu} \in \mathbb{R}^d$ ,  $\Sigma \in \mathbb{R}^{d \times d}$  is a positive definite matrix, and  $\nu$  is the degrees of freedom. Notation:  $\mathbf{X} \sim t_d(\boldsymbol{\mu}, \Sigma)$ .

- $t_d(\nu, \boldsymbol{\mu}, \Sigma)$  has heavier marginal and joint tails than  $N_d(\boldsymbol{\mu}, \Sigma)$ .
- BMW–Siemens data; simulations from fitted  $N_d(\boldsymbol{\mu}, \Sigma)$  and  $t_d(3, \boldsymbol{\mu}, \Sigma)$ :



## 6.2.2 Normal mean-variance mixtures

- Radial symmetry implies that **all one-dimensional margins of normal variance mixtures are symmetric**.
- Often visible in data: **joint losses have heavier tails** than joint gains.

**Idea:** Introduce **asymmetry by mixing** normal distributions **with different means and variances**.

$\mathbf{X}$  has a (multivariate) *normal mean-variance mixture distribution* if

$$\mathbf{X} \stackrel{d}{=} \mathbf{m}(W) + \sqrt{W} \mathbf{A} \mathbf{Z}, \quad (19)$$

where

- $\mathbf{Z} \sim N_k(\mathbf{0}, I_k)$ ;
- $W \geq 0$  is a scalar random variable which is independent of  $\mathbf{Z}$ ;
- $\mathbf{A} \in \mathbb{R}^{d \times k}$  is a matrix of constants;
- $\mathbf{m} : [0, \infty) \rightarrow \mathbb{R}^d$  is a measurable function.

- Normal mean-variance mixtures add **skewness**: Let  $\Sigma = AA'$  and observe that  $\mathbf{X} | W = w \sim N_d(\mathbf{m}(w), w\Sigma)$ . In general, **they are no longer elliptical** (see later).

### Example 6.9

- Suppose we have  $\mathbf{m}(W) = \boldsymbol{\mu} + W\boldsymbol{\gamma}$ . Since

$$\begin{aligned}\mathbb{E}(\mathbf{X} | W) &= \boldsymbol{\mu} + W\boldsymbol{\gamma}, \\ \text{cov}(\mathbf{X} | W) &= W\Sigma\end{aligned}$$

we have

$$\begin{aligned}\mathbb{E}\mathbf{X} &= \mathbb{E}(\mathbb{E}(\mathbf{X} | W)) = \boldsymbol{\mu} + \mathbb{E}(W)\boldsymbol{\gamma} \quad \text{if } \mathbb{E}W < \infty, \\ \text{cov}(\mathbf{X}) &= \mathbb{E}(\text{cov}(\mathbf{X} | W)) + \text{cov}(\mathbb{E}(\mathbf{X} | W)) \\ &= \mathbb{E}(W)\Sigma + \text{var}(W)\boldsymbol{\gamma}\boldsymbol{\gamma}' \quad \text{if } \mathbb{E}(W^2) < \infty.\end{aligned}$$

- If  $W$  has a GIG distribution, then  $\mathbf{X}$  follows a *generalised hyperbolic distribution*.  $\boldsymbol{\gamma} = \mathbf{0}$  leads to (elliptical) normal variance mixtures; see McNeil et al. (2015, Sections 6.2.3) for details.

## 6.3 Spherical and elliptical distributions

Empirical examples (see McNeil et al. (2015, Sections 6.2.4)) show that

- 1)  $M_d(\boldsymbol{\mu}, \Sigma, \hat{F}_W)$  (e.g. multivariate  $t$ , NIG) provide superior models to  $N_d(\boldsymbol{\mu}, \Sigma)$  for daily/weekly US stock-return data;
- 2) the more general skewed normal mean-variance mixture distributions offer only a modest improvement.

We study elliptical distributions, a generalization of  $M_d(\boldsymbol{\mu}, \Sigma, \hat{F}_W)$ .

### 6.3.1 Spherical distributions

#### Definition 6.10 (Spherical distribution)

A random vector  $\mathbf{Y} = (Y_1, \dots, Y_d)$  has a spherical distribution if for every orthogonal  $U \in \mathbb{R}^{d \times d}$  (i.e.  $U \in \mathbb{R}^{d \times d}$  with  $UU' = U'U = I_d$ )

$\mathbf{Y} \stackrel{d}{=} U\mathbf{Y}$  (distributionally invariant under rotations and reflections)



### Theorem 6.11 (Characterization of spherical distributions)

Let  $\|\mathbf{t}\| = (t_1^2 + \dots + t_d^2)^{1/2}$ ,  $\mathbf{t} \in \mathbb{R}^d$ . The following are equivalent:

- 1)  $\mathbf{Y}$  is spherical (notation:  $\mathbf{Y} \sim S_d(\psi)$  for  $\psi$  as below).
- 2)  $\exists$  a characteristic generator  $\psi : [0, \infty) \rightarrow \mathbb{R}$ , such that  $\phi_{\mathbf{Y}}(\mathbf{t}) = \mathbb{E}(e^{i\mathbf{t}'\mathbf{Y}}) = \psi(\|\mathbf{t}\|^2)$ ,  $\forall \mathbf{t} \in \mathbb{R}^d$ .
- 3) For every  $\mathbf{a} \in \mathbb{R}^d$ ,  $\mathbf{a}'\mathbf{Y} \stackrel{d}{=} \|\mathbf{a}\|Y_1$  (lin. comb. are of the same type).  
 $\Rightarrow$  Subadditivity of  $\text{VaR}_\alpha$  for jointly elliptical losses



### Theorem 6.12 (Stochastic representation)

$\mathbf{Y} \sim S_d(\psi)$  if and only if  $\mathbf{Y} \stackrel{d}{=} R\mathbf{S}$  for an independent radial part  $R \geq 0$  and  $\mathbf{S} \sim U(\{\mathbf{x} \in \mathbb{R}^d : \|\mathbf{x}\| = 1\})$ .

This is the key to understanding the structure of spherical distributions.

### Example 6.13 (Understanding spherical distributions)

$n = 500$  realizations of  $\mathbf{S}$  (left) and  $\mathbf{Y} = R\mathbf{S}$  (right) for  $R \sim \sqrt{dF(d, \nu)}$ ,  $d = 2$ ,  $\nu = 4$  (as for the multivariate  $t$  distribution with  $\nu = 4$ ).



## 6.3.2 Elliptical distributions

### Definition 6.14 (Elliptical distribution)

A random vector  $\mathbf{X} = (X_1, \dots, X_d)$  has an *elliptical distribution* if



$$\mathbf{X} \stackrel{d}{=} \boldsymbol{\mu} + A\mathbf{Y}, \quad (\text{multivariate affine transformation})$$

where  $\mathbf{Y} \sim S_k(\psi)$ ,  $A \in \mathbb{R}^{d \times k}$  (*scale matrix*  $\Sigma = AA'$ ), and (*location vector*)  $\boldsymbol{\mu} \in \mathbb{R}^d$ .

- By Theorem 6.12, an elliptical random vector **admits the stochastic representation**  $\mathbf{X} \stackrel{d}{=} \boldsymbol{\mu} + RAS$ , with  $R$  and  $S$  as before.
- Notation:  $\mathbf{X} \sim E_d(\boldsymbol{\mu}, \Sigma, \psi)$
- **Normal variance mixture distributions are elliptical** (most useful examples) since  $\mathbf{X} \stackrel{d}{=} \boldsymbol{\mu} + \sqrt{W}A\mathbf{Z} = \boldsymbol{\mu} + \sqrt{W}\|\mathbf{Z}\|A\mathbf{Z}/\|\mathbf{Z}\| = \boldsymbol{\mu} + RAS$  with  $R = \sqrt{W}\|\mathbf{Z}\|$  and  $S = \mathbf{Z}/\|\mathbf{Z}\|$ .

### Example 6.15 (Understanding elliptical distributions)

$n = 500$  realizations of  $\mathbf{X} = \mathbf{RAS}$  (left) and  $\mathbf{X} = \boldsymbol{\mu} + \mathbf{RAS}$  (right) for  $R \sim \sqrt{dF(d, \nu)}$ ,  $d = 2$ ,  $\nu = 4$ ; based on the same samples as in Example 6.13.



### 6.3.3 Properties of elliptical distributions

- **Density:** Let  $\Sigma$  be positive definite and  $\mathbf{Y} \sim S_d(\psi)$  have density generator  $g$ .  $\mathbf{X} = \boldsymbol{\mu} + A\mathbf{Y}$  has density

$$f_{\mathbf{X}}(\mathbf{x}) = \frac{1}{\sqrt{\det \Sigma}} g((\mathbf{x} - \boldsymbol{\mu})' \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu})),$$

which depends on  $\mathbf{x}$  only through  $(\mathbf{x} - \boldsymbol{\mu})' \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu})$ , i.e. is constant on ellipsoids (hence the name “elliptical”).

- **Linear combinations:** For  $\mathbf{X} \sim E_d(\boldsymbol{\mu}, \Sigma, \psi)$ ,  $B \in \mathbb{R}^{k \times d}$  and  $\mathbf{b} \in \mathbb{R}^k$ ,

$$B\mathbf{X} + \mathbf{b} \sim E_k(B\boldsymbol{\mu} + \mathbf{b}, B\Sigma B', \psi) \quad (\text{via cfs}).$$

If  $\mathbf{a} \in \mathbb{R}^d$  (take  $\mathbf{b} = \mathbf{0}$  and  $B = \mathbf{a}' \in \mathbb{R}^{1 \times d}$ ),

$$\mathbf{a}'\mathbf{X} \sim E_1(\mathbf{a}'\boldsymbol{\mu}, \mathbf{a}'\Sigma\mathbf{a}, \psi) \quad (\text{as for } N(\boldsymbol{\mu}, \Sigma)). \quad (20)$$

From  $\mathbf{a} = \mathbf{e}_j = (0, \dots, 0, 1, 0, \dots, 0)$  we see that all marginal distributions are of the same type.

- **Marginal dfs:** As for  $N_d(\boldsymbol{\mu}, \Sigma)$ , it immediately follows that  $\mathbf{X} = (\mathbf{X}'_1, \mathbf{X}'_2)' \sim E_d(\boldsymbol{\mu}, \Sigma, \psi)$  satisfies  $\mathbf{X}_1 \sim E_k(\boldsymbol{\mu}_1, \Sigma_{11}, \psi)$  and that  $\mathbf{X}_2 \sim E_{d-k}(\boldsymbol{\mu}_2, \Sigma_{22}, \psi)$ ; i.e. **margins of elliptical distributions are elliptical**.
- **Quadratic forms:**  $(\mathbf{X} - \boldsymbol{\mu})' \Sigma^{-1} (\mathbf{X} - \boldsymbol{\mu}) \stackrel{d}{=} R^2$ . If  $\mathbf{X} \sim N_d(\boldsymbol{\mu}, \Sigma)$ ,  $R^2 \sim \chi_d^2$ ; and if  $\mathbf{X} \sim t_d(\nu, \boldsymbol{\mu}, \Sigma)$ ,  $R^2/d \sim F(d, \nu)$ .

## 6.4 Dimension reduction techniques

### 6.4.1 Factor models

Explain the variability of  $\mathbf{X}$  in terms of common factors.

#### Definition 6.16 ( $p$ -factor model)

$\mathbf{X}$  follows a  *$p$ -factor model* if

$$\mathbf{X} = \mathbf{a} + B\mathbf{F} + \boldsymbol{\varepsilon}, \quad (21)$$

where

- 1)  $B \in \mathbb{R}^{d \times p}$  is a *matrix of factor loadings* and  $\mathbf{a} \in \mathbb{R}^d$ ;
- 2)  $\mathbf{F} = (F_1, \dots, F_p)$  is the random vector of *(common) factors* with  $p < d$  and  $\boldsymbol{\Omega} := \text{cov}(\mathbf{F})$ , (*systematic risk*);
- 3)  $\boldsymbol{\varepsilon} = (\varepsilon_1, \dots, \varepsilon_d)$  is the random vector of *idiosyncratic error terms* with  $\mathbb{E}(\boldsymbol{\varepsilon}) = \mathbf{0}$ ,  $\boldsymbol{\Upsilon} := \text{cov}(\boldsymbol{\varepsilon})$  diag.,  $\text{cov}(\mathbf{F}, \boldsymbol{\varepsilon}) = (0)$  (*idiosync. risk*).

- **Goals:** Identify or estimate  $\mathbf{F}_t$ ,  $t \in \{1, \dots, n\}$ , then model the distribution/dynamics of the (lower-dimensional) factors (instead of  $\mathbf{X}_t$ ,  $t \in \{1, \dots, n\}$ ).
- Factor models imply that  $\Sigma := \text{cov}(\mathbf{X}) = B\Omega B' + \Upsilon$ .
- With  $B^* = B\Omega^{1/2}$  and  $\mathbf{F}^* = \Omega^{-1/2}(\mathbf{F} - \mathbb{E}(\mathbf{F}))$ , we have

$$\mathbf{X} = \boldsymbol{\mu} + B^* \mathbf{F}^* + \boldsymbol{\varepsilon},$$

where  $\boldsymbol{\mu} = \mathbb{E}(\mathbf{X})$ . We have  $\Sigma = B^*(B^*)' + \Upsilon$ . Conversely, if  $\text{cov}(\mathbf{X}) = BB' + \Upsilon$  for some  $B \in \mathbb{R}^{d \times p}$  with  $\text{rank}(B) = p < d$  and diagonal matrix  $\Upsilon$ , then  $\mathbf{X}$  has a factor-model representation for a  $p$ -dimensional  $\mathbf{F}$  and  $d$ -dimensional  $\boldsymbol{\varepsilon}$ .



## 6.4.2 Statistical estimation strategies

Consider  $\mathbf{X}_t = \mathbf{a} + B\mathbf{F}_t + \varepsilon_t$ ,  $t \in \{1, \dots, n\}$ . Three types of factor model are commonly used:

- 1) *Macroeconomic factor models*: Here we assume that  $\mathbf{F}_t$  is observable,  $t \in \{1, \dots, n\}$ . Estimation of  $B, \mathbf{a}$  is accomplished by time series regression.
- 2) *Fundamental factor models*: Here we assume that the matrix of factor loadings  $B$  is known but the factors  $\mathbf{F}_t$  are unobserved (and have to be estimated from  $\mathbf{X}_t$ ,  $t \in \{1, \dots, n\}$ , using cross-sectional regression at each  $t$ ).
- 3) *Fundamental factor models*: Here we assume that neither the factors  $\mathbf{F}_t$  nor the factor loadings  $B$  are observed (both have to be estimated from  $\mathbf{X}_t$ ,  $t \in \{1, \dots, n\}$ ). The factors can be found with principal component analysis.

## 6.4.3 Estimating macroeconomic factor models

This is achieved by [time series regression](#).

### Univariate regression

- Consider the (univariate) [time series regression](#) model

$$X_{t,j} = a_j + \mathbf{b}'_j \mathbf{F}_t + \varepsilon_{t,j}, \quad t \in \{1, \dots, n\}.$$

- To justify the use of the [ordinary least-squares \(OLS\)](#) method to derive statistical properties of the method it is usually [assumed that](#), conditional on the factors, the errors  $\varepsilon_{1,j}, \dots, \varepsilon_{n,j}$  [form a white noise process](#) (i.e. are identically distributed and serially uncorrelated).
- $\hat{a}_j$  estimates  $a_j$ ,  $\hat{\mathbf{b}}_j$  estimates the  $j$ th row of  $B$ .

Models can also be estimated simultaneously using [multivariate regression](#); see McNeil et al. (2015).

## 6.4.4 Estimating fundamental factor models

- Consider the cross-sectional regression model  $\mathbf{X}_t = B\mathbf{F}_t + \boldsymbol{\varepsilon}_t$  ( $B$  known;  $\mathbf{F}_t$  to be estimated;  $\text{cov}(\boldsymbol{\varepsilon}) = \Upsilon$ ); note that  $\mathbf{a}$  can be absorbed into  $\mathbf{F}_t$ . To obtain precision in estimating  $\mathbf{F}_t$ , we need  $d \gg p$ .
- First estimate  $\mathbf{F}_t$  via OLS by  $\hat{\mathbf{F}}_t^{\text{OLS}} = (B'B)^{-1}B'\mathbf{X}_t$ . This is the best linear unbiased estimator if the  $\boldsymbol{\varepsilon}$  is homoskedastic. However, it is possible to obtain linear unbiased estimates with a smaller covariance matrix via generalized least squares (GLS).
- To this end, estimate  $\Upsilon$  by  $\hat{\Upsilon}$  via the diagonal of the sample covariance matrix of the residuals  $\hat{\boldsymbol{\varepsilon}}_t = \mathbf{X}_t - B\hat{\mathbf{F}}_t^{\text{OLS}}$ ,  $t \in \{1, \dots, n\}$ .
- Then estimate  $\mathbf{F}_t$  via  $\hat{\mathbf{F}}_t = (B'\hat{\Upsilon}^{-1}B)^{-1}B'\hat{\Upsilon}^{-1}\mathbf{X}_t$ .

## 6.4.5 Principal component analysis

- **Goal:** Reduce the dimensionality of highly correlated data by finding a small number of uncorrelated linear combinations which account for most of the variance in the data; this can be used for finding factors.
- **Key:** Any symmetric  $A$  admits a *spectral decomposition*

$$A = \Gamma \Lambda \Gamma',$$

where

- 1)  $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_d)$  is the diagonal matrix of eigenvalues of  $A$  which, w.l.o.g., are ordered so that  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_d$ ; and
  - 2)  $\Gamma$  is an orthogonal matrix whose columns are eigenvectors of  $A$  standardized to have length 1.
- Let  $\Sigma = \Gamma \Lambda \Gamma'$  with  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_d \geq 0$  (positive semidefiniteness  $\Rightarrow$  all eigenvalues  $\geq 0$ ) and  $\mathbf{Y} = \Gamma'(\mathbf{X} - \boldsymbol{\mu})$  (the so-called *principal component transform*). The  $j$ th component  $Y_j = \boldsymbol{\gamma}_j'(\mathbf{X} - \boldsymbol{\mu})$  is the  $j$ th *principal component of  $\mathbf{X}$*  (where  $\boldsymbol{\gamma}_j$  is the  $j$ th column of  $\Gamma$ ).

- We have  $\mathbb{E}\mathbf{Y} = \mathbf{0}$  and  $\text{cov}(\mathbf{Y}) = \Gamma'\Sigma\Gamma = \Gamma'\Gamma\Lambda\Gamma'\Gamma = \Lambda$ , so the principal components are uncorrelated and  $\text{var}(Y_j) = \lambda_j$ ,  $j \in \{1, \dots, d\}$ . The principal components are thus ordered by decreasing variance.
- One can show:
  - ▶ The first principal component is that standardized linear combination of  $\mathbf{X}$  which has maximal variance among all such combinations, i.e.  $\text{var}(\gamma_1'\mathbf{X}) = \max\{\text{var}(\mathbf{a}'\mathbf{X}) : \mathbf{a}'\mathbf{a} = 1\}$ .
  - ▶ For  $j \in \{2, \dots, d\}$ , the  $j$ th principal component is that standardized linear combination of  $\mathbf{X}$  which has maximal variance among all such linear combinations which are orthogonal to (and hence uncorrelated with) the first  $j - 1$ -many linear combinations.
- $\sum_{j=1}^d \text{var}(Y_j) = \sum_{j=1}^d \lambda_j = \text{trace}(\Sigma) = \sum_{j=1}^d \text{var}(X_j)$ , so we can interpret  $\sum_{j=1}^k \lambda_j / \sum_{j=1}^d \lambda_j$  as the fraction of total variance explained by the first  $k$  principal components.

## Principal components as factors

- Inverting the principal component transform  $\mathbf{Y} = \Gamma'(\mathbf{X} - \boldsymbol{\mu})$ , we have

$$\mathbf{X} = \boldsymbol{\mu} + \Gamma \mathbf{Y} = \boldsymbol{\mu} + \Gamma_1 \mathbf{Y}_1 + \Gamma_2 \mathbf{Y}_2 =: \boldsymbol{\mu} + \Gamma_1 \mathbf{Y}_1 + \boldsymbol{\varepsilon}$$

where  $\mathbf{Y}_1 \in \mathbb{R}^k$  contains the first  $k$  principal components. This is reminiscent of the basic factor model.

- Although  $\varepsilon_1, \dots, \varepsilon_d$  will tend to have small variances, the assumptions of the factor model are generally violated (since they need not have a diagonal covariance matrix and need not be uncorrelated with  $\mathbf{Y}_1$ ). Nevertheless, principal components are often interpreted as factors.
- In principle, the same can be applied to the sample covariance matrix to obtain the sample principal components.

# 7 Copulas and dependence

## 7.1 Copulas

## 7.2 Dependence concepts and measures

# 7.1 Copulas

## 7.1.1 Basic properties

### Definition 7.1 (Copula)

A *copula*  $C$  is a df with  $U(0, 1)$  margins.

### Characterization

$C : [0, 1]^d \rightarrow [0, 1]$  is a copula if and only if

1)  $C$  is *grounded*, that is,

$C(u_1, \dots, u_d) = 0$  if  $u_j = 0$  for at least one  $j \in \{1, \dots, d\}$ .

2)  $C$  has standard *uniform* univariate *margins*, that is,


$C(1, \dots, 1, u_j, 1, \dots, 1) = u_j$  for all  $u_j \in [0, 1]$  and  $j \in \{1, \dots, d\}$ .

3)  $C$  is *d-increasing*, that is, for all  $\mathbf{a}, \mathbf{b} \in [0, 1]^d$ ,  $\mathbf{a} \leq \mathbf{b}$ ,  $\Delta_{(\mathbf{a}, \mathbf{b})}C = \sum_{\mathbf{i} \in \{0, 1\}^d} (-1)^{\sum_{j=1}^d i_j} C(a_1^{i_1} b_1^{1-i_1}, \dots, a_d^{i_d} b_d^{1-i_d}) \geq 0$ . Equivalently, if it exists, the density  $c$  of  $C$  satisfies  $c(\mathbf{u}) \geq 0$  for all  $\mathbf{u} \in (0, 1)^d$ .



2-increasingness explained in a picture:

$$\begin{aligned}\Delta_{(a,b]}C &= C(b_1, b_2) - C(b_1, a_2) - C(a_1, b_2) + C(a_1, a_2) \\ &= \mathbb{P}(U \in (a, b]) \geq 0\end{aligned}$$



$\Rightarrow \Delta_{(a,b]}C$  is the **probability** of a random vector  $U \sim C$  to be in  $(a, b]$ .

## Preliminaries

### Lemma 7.2 (Probability transformation)

Let  $X \sim F$ ,  $F$  continuous. Then  $F(X) \sim U(0, 1)$ .

*Idea of the proof.*  $\mathbb{P}(F(X) \leq u) = \mathbb{P}(F^{\leftarrow}(F(X)) \leq F^{\leftarrow}(u)) = \mathbb{P}(X \leq F^{\leftarrow}(u)) = F(F^{\leftarrow}(u)) = u, u \in [0, 1]$ .  $\square$

Note that  $F$  needs to be continuous (otherwise  $F(X)$  would not reach all intervals  $\subseteq [0, 1]$ ).

### Lemma 7.3 (Quantile transformation)

Let  $U \sim U(0, 1)$  and  $F$  be any df. Then  $X = F^{\leftarrow}(U) \sim F$ .

*Proof.*  $\mathbb{P}(F^{\leftarrow}(U) \leq x) = \mathbb{P}(U \leq F(x)) = F(x), x \in \mathbb{R}$ .  $\square$

Probability and quantile transformations are the key to all applications involving copulas. They allow us to go from  $\mathbb{R}^d$  to  $[0, 1]^d$  and back.

# Sklar's Theorem

## Theorem 7.4 (Sklar's Theorem)

- 1) For any df  $F$  with margins  $F_1, \dots, F_d$ , there exists a copula  $C$  such that

$$F(x_1, \dots, x_d) = C(F_1(x_1), \dots, F_d(x_d)), \quad \mathbf{x} \in \mathbb{R}^d. \quad (22)$$

$C$  is uniquely defined on  $\prod_{j=1}^d \text{ran } F_j$  and given by

$$C(u_1, \dots, u_d) = F(F_1^{\leftarrow}(u_1), \dots, F_d^{\leftarrow}(u_d)), \quad \mathbf{u} \in \prod_{j=1}^d \text{ran } F_j,$$

where  $\text{ran } F_j = \{F_j(x) : x \in \mathbb{R}\}$  denotes the *range* of  $F_j$ .

- 2) Conversely, given any copula  $C$  and univariate dfs  $F_1, \dots, F_d$ ,  $F$  defined by (22) is a df with margins  $F_1, \dots, F_d$ .

*Proof.*

1) **Proof for continuous  $F_1, \dots, F_d$  only.** Let  $\mathbf{X} \sim F$  and define  $U_j = F_j(X_j)$ ,  $j \in \{1, \dots, d\}$ . By the probability transformation,  $U_j \sim U(0, 1)$  (continuity!),  $j \in \{1, \dots, d\}$ , so the df  $C$  of  $\mathbf{U}$  is a copula. Since  $F_j \uparrow$  on  $\text{ran } X_j$ ,  $\mathbf{X}_j = F_j^{\leftarrow}(F_j(X_j)) = F_j^{\leftarrow}(U_j)$ ,  $j \in \{1, \dots, d\}$ . Therefore,

$$\begin{aligned} F(\mathbf{x}) &= \mathbb{P}(X_j \leq x_j \ \forall j) = \mathbb{P}(F_j^{\leftarrow}(U_j) \leq x_j \ \forall j) = \mathbb{P}(U_j \leq F_j(x_j) \ \forall j) \\ &= C(F_1(x_1), \dots, F_d(x_d)), \quad \mathbf{x} \in \mathbb{R}^d. \end{aligned}$$

Hence  $C$  is a copula and satisfies (22).

Since  $F_j(F_j^{\leftarrow}(u_j)) = u_j$  for all  $u_j \in \text{ran } F_j$ , so

$$\begin{aligned} C(u_1, \dots, u_d) &= C(F_1(F_1^{\leftarrow}(u_1)), \dots, F_d(F_d^{\leftarrow}(u_d))) \\ &= F(F_1^{\leftarrow}(u_1), \dots, F_d^{\leftarrow}(u_d)), \quad \mathbf{u} \in \prod_{j=1}^d \text{ran } F_j. \end{aligned}$$

2) For  $\mathbf{U} \sim C$ , define  $\mathbf{X} = (F_1^{\leftarrow}(U_1), \dots, F_d^{\leftarrow}(U_d))$ . Then

$$\begin{aligned}\mathbb{P}(\mathbf{X} \leq \mathbf{x}) &= \mathbb{P}(F_j^{\leftarrow}(U_j) \leq x_j \ \forall j) = \mathbb{P}(U_j \leq F_j(x_j) \ \forall j) \\ &= C(F_1(x_1), \dots, F_d(x_d)), \quad \mathbf{x} \in \mathbb{R}^d.\end{aligned}$$

Therefore,  $F$  defined by (22) is a df (that of  $\mathbf{X}$ ), with margins  $F_1, \dots, F_d$  (obtained by the quantile transformation).  $\square$

- A *copula model* for  $\mathbf{X}$  means  $F(\mathbf{x}) = C(F_1(x_1), \dots, F_d(x_d))$  for some (parametric) copula  $C$  and (parametric) marginals  $F_1, \dots, F_d$ .
- $\mathbf{X}$  (or  $F$ ) with margins  $F_1, \dots, F_d$  *has copula*  $C$  if (22) holds.

## Invariance principle

### Theorem 7.5 (Invariance principle)

Let  $\mathbf{X} \sim F$  with continuous margins  $F_1, \dots, F_d$  and copula  $C$ . If  $T_j \uparrow$  on  $\text{ran } X_j$  for all  $j$ , then  $(T_1(X_1), \dots, T_d(X_d))$  (also) has copula  $C$ .

### Interpretation of Sklar's Theorem (and the invariance principle)


- 1) Part 1) of Sklar's Theorem allows one to **decompose any df  $F$  into its margins and a copula**. This, together with the invariance principle, allows one to **study dependence independently of the margins via the margin-free  $\mathbf{U} = (F_1(X_1), \dots, F_d(X_d))$  instead of  $\mathbf{X} = (X_1, \dots, X_d)$**  (they both have the same copula!). This is interesting for statistical applications, e.g. **parameter estimation** or **goodness-of-fit**.
- 2) Part 2) allows one to **construct flexible multivariate distributions** for particular applications.

## Visualizing Part 1) of Sklar's Theorem


**Left:** Scatter plot of  $n = 1000$  samples from  $(X_1, X_2) \sim N_2(\mathbf{0}, P)$ , where  $P = \begin{pmatrix} 1 & 0.7 \\ 0.7 & 1 \end{pmatrix}$ . We mark three points A, B, C.

**Right:** Scatter plot of the corresponding Gauss copula (after applying the df  $\Phi$  of  $N(0, 1)$ ). Note how A, B, C change.

1000 realizations of  $(X, Y)$  for a joint normal distribution with  $\rho = 0.7$





$(U, V) = (F(X), F(Y))$  for  $(X, Y)$  a joint normal distribution with  $\rho = 0.7$



## Visualizing Part 2) of Sklar's Theorem

**Left:** Same Gauss copula scatter plot as before. Apply marginal Exp(2)-quantile functions ( $F_j^{-1}(u) = -\log(1-u)/2$ ,  $j \in \{1, 2\}$ ).

**Right:** The corresponding transformed random variates. Again, note the three points A, B, C.







## Visualizing Part 1) of Sklar's Theorem

**Left:** Scatter plot of  $n = 1000$  samples from  $(X_1, X_2) \sim t_2(4, \mathbf{0}, P)$ , where  $P = \begin{pmatrix} 1 & 0.7 \\ 0.7 & 1 \end{pmatrix}$ . We mark three points I, J, K.



**Right:** Scatter plot of the corresponding  $t_4$  copula (after applying the df  $t_4$ ). Note how A, B, C change.



## Visualizing Part 2) of Sklar's Theorem

**Left:** Same  $t_4$  copula scatter plot as before. Apply marginal Exp(2)-quantile functions ( $F_j^{-1}(u) = -\log(1-u)/2$ ,  $j \in \{1, 2\}$ ).

**Right:** The corresponding transformed random variates. Again, note the three points I, J, K.



## Fréchet–Höfding bounds

### Theorem 7.6 (Fréchet–Höfding bounds)

Let  $W(\mathbf{u}) = \max\{\sum_{j=1}^d u_j - d + 1, 0\}$  and  $M(\mathbf{u}) = \min_{1 \leq j \leq d} \{u_j\}$ .

1) For any  $d$ -dimensional copula  $C$ ,

$$W(\mathbf{u}) \leq C(\mathbf{u}) \leq M(\mathbf{u}), \quad \mathbf{u} \in [0, 1]^d.$$

2)  $W$  is a copula if and only if  $d = 2$ .



3)  $M$  is a copula for all  $d \geq 2$ .

■ It is easy to verify that, for  $U \sim U(0, 1)$ ,

▶  $(U, \dots, U) \sim M$ ;

▶  $(U, 1 - U) \sim W$ .

- Plot of  $W, M$  for  $d = 2$  (compare with  $(U, 1 - U) \sim W$ ,  $(U, U) \sim M$ )



- The Fréchet–Höfding bounds correspond to perfect dependence (negative for  $W$ ; positive for  $M$ ); see Proposition 7.10 later.
- The Fréchet–Höfding bounds lead to bounds for any df  $F$ , via

$$\max\left\{\sum_{j=1}^d F_j(x_j) - d + 1, 0\right\} \leq F(\mathbf{x}) \leq \min_{1 \leq j \leq d} \{F_j(x_j)\}.$$

We will use them later to derive bounds for the correlation coefficient.

## 7.1.2 Examples of copulas

- *Fundamental copulas*: important special copulas;
- *Implicit copulas*: extracted from known  $F$  via Sklar's Theorem;
- *Explicit copulas*: have simple closed-form expressions and follow construction principles of copulas.

### Fundamental copulas

- $\Pi(\mathbf{u}) = \prod_{j=1}^d u_j$  is the *independence copula* since  $C(F_1(x_1), \dots, F_d(x_d)) = F(\mathbf{x}) \stackrel{\text{Sklar}}{=} \prod_{j=1}^d F_j(x_j) \stackrel{\text{ind.}}{=} \prod_{j=1}^d F_j^{\leftarrow}(u_j)$  if and only if  $C(\mathbf{u}) = \Pi(\mathbf{u})$  (replace  $x_j$  by  $F_j^{\leftarrow}(u_j)$ ). Therefore,  $X_1, \dots, X_d$  are independent if and only if their copula is  $\Pi$ ; the density is thus  $c(\mathbf{u}) = 1, \mathbf{u} \in [0, 1]^d$ .
- The Fréchet–Höfding bound  $W$  is the *countermonotonicity copula*. It is the df of  $(U, 1 - U)$ . If  $X_1, X_2$  are perfectly negatively dependent ( $X_2$  is a.s. a strictly decreasing function in  $X_1$ ), their copula is  $W$ .

- The Fréchet–Höfdding bound  $M$  is the *comonotonicity copula*. It is the df of  $(U, \dots, U)$ . If  $X_1, \dots, X_d$  are perfectly positively dependent ( $X_2, \dots, X_{d-1}$  are a.s. strictly increasing functions in  $X_1$ ), their copula is  $M$ .

## Implicit copulas

*Elliptical copulas* are implicit copulas arising from elliptical distributions via Sklar's Theorem. The two most prominent parametric families are the *Gauss copula* and the *t copula* (stemming from normal variance mixtures).

## Gauss copulas

- Consider (w.l.o.g.)  $\mathbf{X} \sim N_d(\mathbf{0}, P)$ . The *Gauss copula* (family) is given by

$$\begin{aligned} C_P^{\text{Ga}}(\mathbf{u}) &= \mathbb{P}(\Phi(X_1) \leq u_1, \dots, \Phi(X_d) \leq u_d) \\ &= \Phi_P(\Phi^{-1}(u_1), \dots, \Phi^{-1}(u_d)) \end{aligned}$$

where  $\Phi_P$  is the df of  $N_d(\mathbf{0}, P)$  and  $\Phi$  the df of  $N(0, 1)$ .

- Special cases: If  $P = I_d$  then  $C = \Pi$ , and if  $P = J_d = \mathbf{1}\mathbf{1}'$  then  $C = M$ .  
If  $d = 2$  and  $\rho = P_{12} = -1$  then  $C = W$ .
- Sklar's Theorem  $\Rightarrow$  The density of  $C(\mathbf{u}) = F(F_1^{\leftarrow}(u_1), \dots, F_d^{\leftarrow}(u_d))$  is

$$c(\mathbf{u}) = \frac{f(F_1^{\leftarrow}(u_1), \dots, F_d^{\leftarrow}(u_d))}{\prod_{j=1}^d f_j(F_j^{\leftarrow}(u_j))}, \quad \mathbf{u} \in (0, 1)^d.$$

In particular, the density of  $C_P^{\text{Ga}}$  is

$$c_P^{\text{Ga}}(\mathbf{u}) = \frac{1}{\sqrt{\det P}} \exp\left(-\frac{1}{2} \mathbf{x}'(P^{-1} - I_d) \mathbf{x}\right), \quad (23)$$

where  $\mathbf{x} = (\Phi^{-1}(u_1), \dots, \Phi^{-1}(u_d))$ .

## $t$ copulas

- Consider (w.l.o.g.)  $\mathbf{X} \sim t_d(\nu, \mathbf{0}, P)$ . The  $t$  copula (family) is given by

$$\begin{aligned} C_{\nu, P}^t(\mathbf{u}) &= \mathbb{P}(t_{\nu}(X_1) \leq u_1, \dots, t_{\nu}(X_d) \leq u_d) \\ &= t_{\nu, P}(t_{\nu}^{-1}(u_1), \dots, t_{\nu}^{-1}(u_d)) \end{aligned}$$

where  $t_{\nu,P}$  is the df of  $t_d(\nu, \mathbf{0}, P)$  and  $t_\nu$  the df of the univariate  $t$  distribution with  $\nu$  degrees of freedom.

- Special cases:  $P = J_d = \mathbf{1}\mathbf{1}'$  then  $C = M$ . However, if  $P = I_d$  then  $C \neq \Pi$  (unless  $\nu = \infty$  in which case  $C_{\nu,P}^t = C_P^{\text{Ga}}$ ). If  $d = 2$  and  $\rho = P_{12} = -1$  then  $C = W$ .
- Sklar's Theorem  $\Rightarrow$  The density of  $C_{\nu,P}^t$  is



$$c_{\nu,P}^t(\mathbf{u}) = \frac{\Gamma((\nu + d)/2)}{\Gamma(\nu/2)\sqrt{\det P}} \left( \frac{\Gamma(\nu/2)}{\Gamma((\nu + 1)/2)} \right)^d \frac{(1 + \mathbf{x}'P^{-1}\mathbf{x}/\nu)^{-(\nu+d)/2}}{\prod_{j=1}^d (1 + x_j^2/\nu)^{-(\nu+1)/2}},$$

for  $\mathbf{x} = (t_\nu^{-1}(u_1), \dots, t_\nu^{-1}(u_d))$ .

- For more details, see Demarta and McNeil (2005).
- For scatter plots, see the visualization of Sklar's Theorem above. Note the difference in the tails: The smaller  $\nu$ , the more mass is concentrated in the joint tails.



Perspective plots of the densities of  $C_{\rho=0.3}^{\text{Ga}}$  (left) and  $C_{4,\rho=0.3}^t(\mathbf{u})$  (right).



Advantages and drawbacks of elliptical copulas:

### Advantages:

- Modelling pairwise dependencies (comparably flexible)
- Density available
- Sampling (typically) simple

### Drawbacks:

- Typically,  $C$  is not explicit
- Radially symmetric (so the same lower/upper tail behaviour)

## Explicit copulas

*Archimedean copulas* are copulas of the form

$$C(\mathbf{u}) = \psi(\psi^{-1}(u_1) + \cdots + \psi^{-1}(u_d)), \quad \mathbf{u} \in [0, 1]^d,$$



where the (*Archimedean*) *generator*  $\psi : [0, \infty) \rightarrow [0, 1]$  is  $\downarrow$  on  $[0, \inf\{t : \psi(t) = 0\}]$  and satisfies  $\psi(0) = 1$ ,  $\psi(\infty) = \lim_{t \rightarrow \infty} \psi(t) = 0$ ; we set  $\psi^{-1}(0) = \inf\{t : \psi(t) = 0\}$ . The set of all generators is denoted by  $\Psi$ . If  $\psi(t) > 0$ ,  $t \in [0, \infty)$ , we call  $\psi$  *strict*.

## Examples

- **Clayton copula:** Obtained for  $\psi(t) = (1+t)^{-1/\theta}$ ,  $t \in [0, \infty)$ ,  $\theta \in (0, \infty) \Rightarrow C_\theta^c(\mathbf{u}) = (u_1^{-\theta} + \cdots + u_d^{-\theta} - d + 1)^{-1/\theta}$ . For  $\theta \downarrow 0$ ,  $C \rightarrow \Pi$ ; and for  $\theta \uparrow \infty$ ,  $C \rightarrow M$ .
- **Gumbel copula:** Obtained for  $\psi(t) = \exp(-t^{1/\theta})$ ,  $t \in [0, \infty)$ ,  $\theta \in [1, \infty) \Rightarrow C_\theta^g(\mathbf{u}) = \exp(-((- \log u_1)^\theta + \cdots + (- \log u_d)^\theta)^{1/\theta})$ . For  $\theta = 1$ ,  $C = \Pi$ ; and for  $\theta \rightarrow \infty$ ,  $C \rightarrow M$ .



**Left:** Plot of a bivariate **Clayton copula** (Kendall's tau 0.5; see later).

**Right:** Corresponding **scatter plot** (sample size  $n = 1000$ )



**Left:** Plot of the **corresponding density**.


**Right:** **Level plot** of the density (with heat colors).




**Left:** Cloud plot of a trivariate Clayton copula (sample size  $n = 1000$ ; Kendall's tau 0.5).

**Right:** Corresponding scatter plot matrix.

Clayton copula cloud plot ( $n = 1000$ ,  $\tau = 0.5$ )





Scatter plot matrix of a Clayton copula ( $n = 1000$ ,  $\tau = 0.5$ )



**Left:** Plot of a bivariate **Gumbel copula** (Kendall's tau 0.5).


**Right:** Corresponding **scatter plot** (sample size  $n = 1000$ )




**Left:** Plot of the **corresponding density**.

**Right:** **Level plot** of the density (with heat colors).

Gumbel copula density ( $\tau = 0.5$ )




Level plot Gumbel copula density ( $\tau = 0.5$ )




**Left:** Cloud plot of a trivariate Gumbel copula (sample size  $n = 1000$ ; Kendall's tau 0.5).

**Right:** Corresponding scatter plot matrix.

Gumbel copula cloud plot ( $n = 1000$ ,  $\tau = 0.5$ )



Scatter plot matrix of a Gumbel copula ( $n = 1000$ ,  $\tau = 0.5$ )





## Advantages and drawbacks of Archimedean copulas:

### Advantages:

- Typically **explicit** (if  $\psi^{-1}$  is available)
- Useful in calculations:  
**Properties** can typically be expressed **in terms of  $\psi$**
- **Densities** of various examples available
- **Sampling** often simple
- **Not restricted to radial symmetry**

### Drawbacks:

- All margins of the same dimension are equal (symmetry or **exchangeability**)
- Often used only with a small **number of parameters** (some extensions available, but still less than  $d(d-1)/2$ )

### 7.1.3 Meta distributions

- *Fréchet class*: Class of all dfs  $F$  with given marginal dfs  $F_1, \dots, F_d$ ;  
*Meta- $C$  models*: All dfs  $F$  with the same given copula  $C$ .
- **Example**: A *meta- $t$  model* is a multivariate df  $F$  with  $t$  copula  $C$  and some margins  $F_1, \dots, F_d$ .

### 7.1.4 Simulation of copulas and meta distributions

Copulas are typically sampled via specific stochastic representations.

#### Sampling implicit copulas

##### Algorithm 7.7 (Simulation of implicit copulas)

- 1) Sample  $\mathbf{X} \sim F$ , where  $F$  is a df with continuous margins  $F_1, \dots, F_d$ .
- 2) Return  $\mathbf{U} = (F_1(X_1), \dots, F_d(X_d))$  (**probability transformation**).

## Example 7.8

- Sampling **Gauss copulas**  $C_P^{\text{Ga}}$ :

- 1) Sample  $\mathbf{X} \sim N_d(\mathbf{0}, P)$  ( $\mathbf{X} \stackrel{d}{=} A\mathbf{Z}$  for  $AA' = P$ ,  $\mathbf{Z} \sim N_d(\mathbf{0}, I_d)$ ).
- 2) Return  $\mathbf{U} = (\Phi(X_1), \dots, \Phi(X_d))$ .

- Sampling  **$t_\nu$  copulas**  $C_{\nu, P}^t$ :

- 1) Sample  $\mathbf{X} \sim t_d(\nu, \mathbf{0}, P)$  ( $\mathbf{X} \stackrel{d}{=} \sqrt{W}A\mathbf{Z}$  for  $W = \frac{1}{V}$ ,  $V \sim \Gamma(\frac{\nu}{2}, \frac{\nu}{2})$ ).
- 2) Return  $\mathbf{U} = (t_\nu(X_1), \dots, t_\nu(X_d))$ .


## Sampling meta distributions

Meta- $C$  distributions can be sampled via Sklar's Theorem, Part 2).


### Algorithm 7.9 (Sampling meta- $C$ models)

- 1) Sample  $\mathbf{U} \sim C$ .
- 2) Return  $\mathbf{X} = (F_1^{\leftarrow}(U_1), \dots, F_d^{\leftarrow}(U_d))$  (**quantile transformation**).

2000 samples from (a):  $C_{\rho=0.7}^{\text{Ga}}$ ; (b):  $C_{\theta=2}^{\text{G}}$ ; (c):  $C_{\theta=2.2}^{\text{C}}$ ; (d):  $C_{\nu=4, \rho=0.71}^t$



... transformed to  $N(0, 1)$  margins; all have linear correlation  $\approx 0.7$ !



## 7.1.5 Further properties of copulas

### Copula densities

- By [Sklar's Theorem](#), if  $F_j$  has density  $f_j$ ,  $j \in \{1, \dots, d\}$ , and  $C$  has density  $c$ , then the density  $f$  of  $F$  satisfies

$$f(\mathbf{x}) = c(F_1(x_1), \dots, F_d(x_d)) \prod_{j=1}^d f_j(x_j) \quad (24)$$

As seen before, we can recover  $c$  via

$$c(\mathbf{u}) = \frac{f(F_1^{-1}(u_1), \dots, F_d^{-1}(u_d))}{f_1(F_1^{-1}(u_1)) \cdots f_d(F_d^{-1}(u_d))}.$$

- It follows from (24) that the [log-density](#) splits into

$$\log f(\mathbf{x}) = \log c(F_1(x_1), \dots, F_d(x_d)) + \sum_{j=1}^d \log f_j(x_j).$$

which [allows for a two-stage estimation](#) ([marginal](#) and [copula parameters](#)).

## 7.2 Dependence concepts and measures

*Measures of association/dependence* are scalar measures which summarize the dependence in terms of a single number. There are better and worse examples of such measures, which we will study in this section.

### 7.2.1 Perfect dependence

$X_1, X_2$  are *countermonotone* if  $(X_1, X_2)$  has copula  $W$ .

$X_1, \dots, X_d$  are *comonotone* if  $(X_1, \dots, X_d)$  has copula  $M$ .

#### Proposition 7.10 (Perfect dependence)

- 1)  $X_2 = T(X_1)$  a.s. with decreasing  $T(x) = F_2^{\leftarrow}(1 - F_1(x))$  (countermonotone) if and only if  $C(u_1, u_2) = W(u_1, u_2)$ ,  $u_1, u_2 \in [0, 1]$ .
- 2)  $X_j = T_j(X_1)$  a.s. with increasing  $T_j(x) = F_j^{\leftarrow}(F_1(x))$ ,  $j \in \{2, \dots, d\}$  (comonotone), if and only if  $C(\mathbf{u}) = M(\mathbf{u})$ ,  $\mathbf{u} \in [0, 1]^d$ .

## 7.2.2 Linear correlation

For two random variables  $X_1$  and  $X_2$  with  $\mathbb{E}(X_j^2) < \infty$ ,  $j \in \{1, 2\}$ , the (*linear* or *Pearson's*) *correlation coefficient*  $\rho$  is defined by

$$\rho(X_1, X_2) = \frac{\text{cov}(X_1, X_2)}{\sqrt{\text{var } X_1} \sqrt{\text{var } X_2}} = \frac{\mathbb{E}((X_1 - \mathbb{E}X_1)(X_2 - \mathbb{E}X_2))}{\sqrt{\mathbb{E}((X_1 - \mathbb{E}X_1)^2)} \sqrt{\mathbb{E}((X_2 - \mathbb{E}X_2)^2)}}.$$

### Proposition 7.11 (Höfdding's formula)

Let  $X_j \sim F_j$ ,  $j \in \{1, 2\}$ , be two random variables with  $\mathbb{E}(X_j^2) < \infty$ ,  $j \in \{1, 2\}$ , and joint distribution function  $F$ . Then

$$\text{cov}(X_1, X_2) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (F(x_1, x_2) - F_1(x_1)F_2(x_2)) dx_1 dx_2.$$



## Classical properties and drawbacks of linear correlation

Let  $X_1$  and  $X_2$  be two random variables with  $\mathbb{E}(X_j^2) < \infty$ ,  $j \in \{1, 2\}$ .


Note that  $\rho$  depends on the marginal distributions! In particular, second moments have to exist (not the case, e.g. for  $X_1, X_2 \stackrel{\text{ind.}}{\sim} F(x) = 1 - x^{-3}!$ )

- $|\rho| \leq 1$ . Furthermore,  $|\rho| = 1$  if and only if there are constants  $a \in \mathbb{R} \setminus \{0\}, b \in \mathbb{R}$  with  $X_2 = aX_1 + b$  a.s. with  $a \geq 0$  if and only if  $\rho = \pm 1$ . This discards other strong functional dependence such as  $X_2 = X_1^2$ , for example.
- If  $X_1$  and  $X_2$  are independent, then  $\rho = 0$ . However, the converse is not true in general; see Example 7.12 below.
- $\rho$  is invariant under strictly increasing linear transformations on  $\text{ran } X_1 \times \text{ran } X_2$  but not invariant under strictly increasing functions in general. To see this, consider  $(X_1, X_2) \sim N_2(\mathbf{0}, P)$  with  $P_{12} = \rho$ . Then  $\rho(X_1, X_2) = \rho$ , but  $\rho(F_1(X_1), F_2(X_2)) = \frac{6}{\pi} \arcsin(\rho/2)$ .

## Correlation fallacies

### Fallacy 1: $F_1$ , $F_2$ , and $\rho$ uniquely determine $F$

This is true for bivariate elliptical distributions, but wrong in general. The following samples both have  $N(0, 1)$  margins and correlation  $\rho = 0.7$ , yet come from different (copula) models:



Another example is this.

### Example 7.12 (Uncorrelated $\nRightarrow$ independent)

- Consider the two risks

$$X_1 = Z \quad (\text{Profit \& Loss Country A}),$$

$$X_2 = ZV \quad (\text{Profit \& Loss Country B}),$$

where  $V, Z$  are independent with  $Z \sim N(0, 1)$  and  $\mathbb{P}(V = -1) = \mathbb{P}(V = 1) = 1/2$ . Then  $X_2 \sim N(0, 1)$  and  $\rho(X_1, X_2) = \text{cov}(X_1, X_2) = \mathbb{E}(X_1 X_2) \underset{\text{ind.}}{=} \mathbb{E}(V)\mathbb{E}(Z^2) = 0$ , but  $X_1$  and  $X_2$  are not independent (in fact,  $V$  switches between counter- and comonotonicity).

- Consider  $(X'_1, X'_2) \sim N_2(\mathbf{0}, I_2)$ . Both  $(X'_1, X'_2)$  and  $(X_1, X_2)$  have  $N(0, 1)$  margins and  $\rho = 0$ , but the copula of  $(X'_1, X'_2)$  is  $\Pi$  and the copula of  $(X_1, X_2)$  is the convex combination  $C(\mathbf{u}) = \lambda M(\mathbf{u}) + (1 - \lambda)W(\mathbf{u})$  for  $\lambda = 0.5$ .

## Fallacy 2: Given $F_1, F_2$ , any $\rho \in [-1, 1]$ is attainable

This is true for elliptically distributed  $(X_1, X_2)$  with  $\mathbb{E}(R^2) < \infty$  (as then  $\text{corr } \mathbf{X} = \mathbf{P}$ ), but wrong in general:

- If  $F_1$  and  $F_2$  are not of the same type (no linearity),  $\rho(X_1, X_2) = 1$  is not attainable (recall that  $|\rho| = 1$  if and only if there are constants  $a \in \mathbb{R} \setminus \{0\}, b \in \mathbb{R}$  with  $X_2 = aX_1 + b$  a.s.).
- What is the attainable range then? Höfding's formula



$$\text{cov}(X_1, X_2) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} (C(F_1(x_1), F_2(x_2)) - F_1(x_1)F_2(x_2)) dx_1 dx_2.$$

implies bounds on attainable  $\rho$ :

$$\rho \in [\rho_{\min}, \rho_{\max}] \quad (\rho_{\min} \text{ is attained for } C = W, \rho_{\max} \text{ for } C = M).$$

### Example 7.13 (Bounds for a model with $\text{LN}(0, \sigma_j^2)$ margins)

Let  $X_j \sim \text{LN}(0, \sigma_j^2)$ ,  $j \in \{1, 2\}$ . One can show that minimal ( $\rho_{\min}$ ; left) and maximal ( $\rho_{\max}$ ; right) correlations are given as follows.



For  $\sigma_1^2 = 1$ ,  $\sigma_2^2 = 16$  one has  $\rho \in [-0.0003, 0.0137]!$

### Fallacy 3: $\rho$ maximal (i.e. $C = M$ ) $\Rightarrow \text{VaR}_\alpha(X_1 + X_2)$ maximal

- This is true if  $(X_1, X_2)$  is elliptically distributed since the maximal  $\rho = 1$  implies that  $X_1, X_2$  are comonotone, so  $\text{VaR}_\alpha$  can be shown to be additive and additivity provides the largest possible bound in this case as  $\text{VaR}_\alpha$  can be shown to be subadditive in this case.
- Any superadditivity example  $\text{VaR}_\alpha(X_1 + X_2) > \text{VaR}_\alpha(X_1) + \text{VaR}_\alpha(X_2)$  under comonotonicity (under comonotonicity, so maximal correlation qrm, the right-hand side is  $\text{VaR}_\alpha(X_1 + X_2)$ ) serves as a counterexample; see Section 2.2.5.

## 7.2.3 Rank correlation

Rank correlation coefficients are...

- ... always defined;
- ... invariant under strictly increasing transformations of the random variables (hence only depend on the underlying copula).

## Kendall's tau and Spearman's rho

### Definition 7.14 (Kendall's tau)

Let  $X_j \sim F_j$  with  $F_j$  continuous,  $j \in \{1, 2\}$ . Let  $(X'_1, X'_2)$  be an independent copy of  $(X_1, X_2)$ . *Kendall's tau* is defined by

$$\begin{aligned}\rho_\tau &= \mathbb{E}(\text{sign}((X_1 - X'_1)(X_2 - X'_2))) \\ &= \mathbb{P}((X_1 - X'_1)(X_2 - X'_2) > 0) - \mathbb{P}((X_1 - X'_1)(X_2 - X'_2) < 0),\end{aligned}$$

where  $\text{sign}(x) = I_{(0, \infty)}(x) - I_{(-\infty, 0)}(x)$  (so  $-1$  for  $x < 0$ ,  $0$  for  $x = 0$  and  $1$  for  $x > 0$ ).

By definition, Kendall's tau is *the probability of concordance* ( $\mathbb{P}((X_1 - X'_1)(X_2 - X'_2) > 0)$ ; probability of two independent points from  $F$  to have a positive slope) *minus the probability of discordance* ( $\mathbb{P}((X_1 - X'_1)(X_2 - X'_2) < 0)$ ; probability of two independent points from  $F$  to have a negative slope).

### Proposition 7.15 (Formula for Kendall's tau)

Let  $X_j \sim F_j$  with  $F_j$  continuous,  $j \in \{1, 2\}$ , and copula  $C$ . Then

$$\rho_\tau = 4 \int_0^1 \int_0^1 C(u_1, u_2) dC(u_1, u_2) - 1 = 4\mathbb{E}(C(U_1, U_2)) - 1,$$

where  $(U_1, U_2) \sim C$ .

### Definition 7.16 (Spearman's rho)

Let  $X_j \sim F_j$  with  $F_j$  continuous,  $j \in \{1, 2\}$ . *Spearman's rho* is defined by  $\rho_S = \rho(F_1(X_1), F_2(X_2))$ .

### Proposition 7.17 (Formula for Spearman's rho)

Let  $X_j \sim F_j$  with  $F_j$  continuous,  $j \in \{1, 2\}$ , and copula  $C$ . Then

$$\rho_S = 12 \int_0^1 \int_0^1 C(u_1, u_2) du_1 du_2 - 3 = 12\mathbb{E}(C(U'_1, U'_2)) - 3,$$

where  $(U'_1, U'_2) \sim \Pi$ .



- **Fallacy 1** ( $F_1, F_2, \rho$  uniquely determine  $F$ ) is **not solved by** replacing  $\rho$  by **rank correlation coefficients  $\kappa$**  (it is easy to construct several copulas with the same Kendall's tau, e.g. via Archimedean copulas).
- **Fallacy 2** (For  $F_1, F_2$ , any  $\rho \in [-1, 1]$  is attainable) is **solved** when  $\rho$  is  $\rho_\tau$  or  $\rho_S$ . **Take**

$$F(x_1, x_2) = \lambda M(F_1(x_1), F_2(x_2)) + (1 - \lambda) W(F_1(x_1), F_2(x_2)).$$

This is a model with  $\rho_\tau = \rho_S = 2\lambda - 1$  (choose  $\lambda \in [0, 1]$  as desired).

- **Fallacy 3** ( $C = M$  implies  $\text{VaR}_\alpha(X_1 + X_2)$  maximal) is **also not solved by rank correlation coefficients  $\kappa = 1$** : Although  $\kappa = 1$  corresponds to  $C = M$ , this copula does not necessarily provide the largest  $\text{VaR}_\alpha(X_1 + X_2)$ ; see Fallacy 3 earlier.
- **Nevertheless**, rank correlations are useful **to summarize dependence**, to parameterize copula families to **make dependence comparable** and for copula parameter calibration or estimation.

## 7.2.4 Coefficients of tail dependence

**Goal:** Measure *extremal dependence*, i.e. dependence in the *joint tails*.

### Definition 7.18 (Tail dependence)

Let  $X_j \sim F_j$ ,  $j \in \{1, 2\}$ , be continuously distributed random variables. Provided that the limits exist, the *lower tail-dependence coefficient*  $\lambda_l$  and *upper tail-dependence coefficient*  $\lambda_u$  of  $X_1$  and  $X_2$  are defined by

$$\lambda_l = \lim_{u \downarrow 0} \mathbb{P}(X_2 \leq F_2^{\leftarrow}(u) \mid X_1 \leq F_1^{\leftarrow}(u)),$$

$$\lambda_u = \lim_{u \uparrow 1} \mathbb{P}(X_2 > F_2^{\leftarrow}(u) \mid X_1 > F_1^{\leftarrow}(u)).$$

If  $\lambda_l \in (0, 1]$  ( $\lambda_u \in (0, 1]$ ), then  $(X_1, X_2)$  is *lower (upper) tail dependent*.  
If  $\lambda_l = 0$  ( $\lambda_u = 0$ ), then  $(X_1, X_2)$  is *lower (upper) tail independent*.

As (conditional) probabilities, we clearly have  $\lambda_l, \lambda_u \in [0, 1]$ .

- Tail dependence is a copula property, since

$$\begin{aligned}\mathbb{P}(X_2 \leq F_2^{\leftarrow}(u) \mid X_1 \leq F_1^{\leftarrow}(u)) &= \frac{\mathbb{P}(X_1 \leq F_1^{\leftarrow}(u), X_2 \leq F_2^{\leftarrow}(u))}{\mathbb{P}(X_1 \leq F_1^{\leftarrow}(u))} \\ &= \frac{F(F_1^{\leftarrow}(u), F_2^{\leftarrow}(u))}{F_1(F_1^{\leftarrow}(u))} \stackrel{\text{Sklar}}{=} \frac{C(u, u)}{u}, \quad u \in (0, 1), \text{ so } \lambda_1 = \lim_{u \downarrow 0} \frac{C(u, u)}{u}.\end{aligned}$$

- If  $u \mapsto C(u, u)$  is differentiable in a neighborhood of 0 and the limit exists, then  $\lambda_1 = \lim_{u \downarrow 0} \frac{d}{du} C(u, u)$  (l'Hôpital's Rule).
- If  $C$  is totally differentiable in a neighborhood of 0 and the limit exists, then  $\lambda_1 = \lim_{u \downarrow 0} (D_1 C(u, u) + D_2 C(u, u))$  (Chain Rule). If  $C$  is symmetric,  $\lambda_1 = 2 \lim_{u \downarrow 0} D_1 C(u, u) = 2 \lim_{u \downarrow 0} C_{2|1}(u \mid u) = 2 \lim_{u \downarrow 0} \mathbb{P}(U_2 \leq u \mid U_1 = u)$  for  $(U_1, U_2) \sim C$ . Combined with any continuous df  $F$  and  $(X_1, X_2) = (F^{\leftarrow}(U_1), F^{\leftarrow}(U_2))$ , one has

$$\lambda_1 = 2 \lim_{x \downarrow -\infty} \mathbb{P}(X_2 \leq x \mid X_1 = x) \stackrel{\text{if density}}{=} 2 \lim_{x \downarrow -\infty} \int_{-\infty}^x f_{X_2|X_1=x}(x_2) dx_2. \quad (25)$$

- Similarly as above, for the upper tail-dependence coefficient,

$$\begin{aligned}\lambda_u &= \lim_{u \uparrow 1} \frac{1 - 2u + C(u, u)}{1 - u} \\ &= \lim_{u \uparrow 1} \frac{2(1 - u) - (1 - C(u, u))}{1 - u} = 2 - \lim_{u \uparrow 1} \frac{1 - C(u, u)}{1 - u}.\end{aligned}$$

- For all **radially symmetric copulas** (e.g. the bivariate  $C_P^{\text{Ga}}$  and  $C_{\nu, P}^t$  copulas), we have  $\lambda_l = \lambda_u =: \lambda$ .
- For **Archimedean copulas with strict  $\psi$** , a substitution and l'Hôpital's Rule show:

$$\begin{aligned}\lambda_l &= \lim_{u \downarrow 0} \frac{\psi(2\psi^{-1}(u))}{u} = \lim_{t \rightarrow \infty} \frac{\psi(2t)}{\psi(t)} = 2 \lim_{t \rightarrow \infty} \frac{\psi'(2t)}{\psi'(t)}, \\ \lambda_u &= 2 - \lim_{u \uparrow 1} \frac{1 - \psi(2\psi^{-1}(u))}{1 - u} = 2 - \lim_{t \downarrow 0} \frac{1 - \psi(2t)}{1 - \psi(t)} = 2 - 2 \lim_{t \downarrow 0} \frac{\psi'(2t)}{\psi'(t)}.\end{aligned}$$

**Clayton:**  $\lambda_l = 2^{-1/\theta}$ ,  $\lambda_u = 0$ ; **Gumbel:**  $\lambda_l = 0$ ,  $\lambda_u = 2 - 2^{1/\theta}$

### Example 7.19 ( $\lambda$ for the Gauss and $t$ copula)


- Considering the bivariate  $N(\mathbf{0}, P)$  density, one can show (via  $f_{X_2|X_1}(x_2 | x_1) = \frac{f_{X_1, X_2}(x_1, x_2)}{f_{X_1}(x_1)}$ ) that  $X_2 | X_1 = x \sim N(\rho x, 1 - \rho^2)$ . This implies that  $\lambda = 2 \lim_{x \downarrow -\infty} \mathbb{P}(X_2 \leq x | X_1 = x) = 2 \lim_{x \downarrow -\infty} \Phi\left(\frac{x(1-\rho)}{\sqrt{1-\rho^2}}\right) = I_{\{\rho=1\}}$  (essentially **no tail dependence**).

- For  $C_{\nu, P}^t$ , one can show that  $X_2 | X_1 = x \sim t_{\nu+1}\left(\rho x, \frac{(1-\rho^2)(\nu+x^2)}{\nu+1}\right)$  and thus  $\mathbb{P}(X_2 \leq x | X_1 = x) = t_{\nu+1}\left(\frac{x-\rho x}{\sqrt{\frac{(1-\rho^2)(\nu+x^2)}{\nu+1}}}\right)$ . Hence

$$\lambda = 2t_{\nu+1}\left(-\sqrt{\frac{(\nu+1)(1-\rho)}{1+\rho}}\right) \quad (\text{tail dependence; } \lambda \uparrow \text{ in } \rho \uparrow \text{ and } \nu \downarrow).$$

$\nu$	$\rho = -0.5$	$\rho = 0$	$\rho = 0.5$	$\rho = 0.9$	$\rho = 1$
$\infty$	0	0	0	0	1
10	0.00	0.01	0.08	0.46	1
4	0.01	0.08	0.25	0.63	1
2	0.06	0.18	0.39	0.72	1

# Joint quantile exceedance probabilities



5000 samples from

- (a)  $N_2(\mathbf{0}, P = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix})$ ,  $\rho = 0.5$ ;
- (b)  $C_{\rho}^{\text{Ga}}$  with  $t_4$  margins (same dependence as in (a));
- (c)  $C_{4,\rho}^t$  with  $N(0, 1)$  margins;
- (d)  $t_2(4, \mathbf{0}, P)$  (same dependence as in (c)).

Lines denote the true 0.005- and 0.995-quantiles.

Note the different number of points in the bivariate tails (all models have the same Kendall's tau!)

# 8 Aggregate risk

8.1 Risk measures for linear portfolios

8.2 Risk aggregation

8.3 Capital allocation

## 8.1 Risk measures for linear portfolios

We now consider **linear portfolios** in

$$\mathcal{M} = \{L : L = m + \boldsymbol{\lambda}'\mathbf{X}, m \in \mathbb{R}, \boldsymbol{\lambda} \in \mathbb{R}^d\}, \quad (26)$$

for a fixed  $d$ -dimensional random vector  $\mathbf{X}$ .

- **Many standard approaches** to risk aggregation and capital allocation are based on the **assumption** that losses have a linear relationship to underlying risk factors.
- It is **common to use linear approximations** for losses due to market risks over short time horizons.



## 8.1.1 Elliptically distributed risk factors

### Theorem 8.1 (Risk measurement for elliptical risk factors)

Let  $\mathbf{X} \sim E_d(\boldsymbol{\mu}, \Sigma, \psi)$  and  $\varrho$  be any positive-homogeneous, translation-invariant and law-invariant risk measure on  $\mathcal{M}$ . Then:

- 1) For any  $L = m + \boldsymbol{\lambda}'\mathbf{X} \in \mathcal{M}$ ,  $\varrho(L) = m + \boldsymbol{\lambda}'\boldsymbol{\mu} + \sqrt{\boldsymbol{\lambda}'\Sigma\boldsymbol{\lambda}}\varrho(Y_1)$  for  $Y_1 \sim S_1(\psi)$ .
- 2) If  $\varrho(Y_1) \geq 0$ ,  $\varrho$  is subadditive on  $\mathcal{M}$  (e.g.,  $\text{VaR}_\alpha$  for  $\alpha \geq 0.5$ ).
- 3) If  $\mathbb{E}\mathbf{X}$  exists then, for any  $L = m + \boldsymbol{\lambda}'\mathbf{X} \in \mathcal{M}$  and  $\rho_{ij} = \wp(\Sigma)_{ij}$ ,

$$\varrho(L - \mathbb{E}L) = \sqrt{\sum_{i=1}^d \sum_{j=1}^d \rho_{ij} \lambda_i \lambda_j \varrho(X_i - \mathbb{E}X_i) \varrho(X_j - \mathbb{E}X_j)}.$$

- 4) If  $\text{cov}(\mathbf{X})$  exists and  $\varrho(Y_1) > 0$  then, for every  $L \in \mathcal{M}$ ,  
 $\varrho(L) = \mathbb{E}(L) + k_\varrho \sqrt{\text{var}(L)}$  for some  $k_\varrho > 0$  depending on  $\varrho$ .

*Proof.*

- 1) Let  $\mathbf{Y} \sim S_k(\psi)$ ,  $AA' = \Sigma$ .  $L = m + \boldsymbol{\lambda}'\mathbf{X} \stackrel{d}{=} m + \boldsymbol{\lambda}'\boldsymbol{\mu} + \boldsymbol{\lambda}'A\mathbf{Y}$ . By Theorem 6.11 3),  $L \stackrel{d}{=} m + \boldsymbol{\lambda}'\boldsymbol{\mu} + \|A'\boldsymbol{\lambda}\|Y_1$ . Thus  $\varrho(L) = m + \boldsymbol{\lambda}'\boldsymbol{\mu} + \|A'\boldsymbol{\lambda}\|\varrho(Y_1) = m + \boldsymbol{\lambda}'\boldsymbol{\mu} + \sqrt{\boldsymbol{\lambda}'\Sigma\boldsymbol{\lambda}}\varrho(Y_1)$ .
- 2) Set  $L_1 = m_1 + \boldsymbol{\lambda}'_1\mathbf{X}$  and  $L_2 = m_2 + \boldsymbol{\lambda}'_2\mathbf{X}$ . Subadditivity follows from 1) and  $\|A'(\boldsymbol{\lambda}_1 + \boldsymbol{\lambda}_2)\| \leq \|A'\boldsymbol{\lambda}_1\| + \|A'\boldsymbol{\lambda}_2\|$  and  $\varrho(Y_1) \geq 0$ .
- 3)  $\varrho(L - \mathbb{E}L) = \varrho(L) - \mathbb{E}(L) = \varrho(L) - (m + \boldsymbol{\lambda}'\boldsymbol{\mu}) = \sqrt{\boldsymbol{\lambda}'\Sigma\boldsymbol{\lambda}}\varrho(Y_1)$ , so

$$\varrho(L - \mathbb{E}L) = \sqrt{\sum_{i=1}^d \sum_{j=1}^d \rho_{ij} \lambda_i \lambda_j \sigma_i \sigma_j \varrho(Y_1)},$$

where  $\sigma_j = \sqrt{\Sigma_{jj}}$  for  $j \in \{1, \dots, d\}$ . For  $\boldsymbol{\lambda} = \mathbf{e}_j$ ,  $\varrho(X_j - \mathbb{E}X_j) = \varrho(\mathbf{e}'_j\mathbf{X} - \mathbb{E}(\mathbf{e}'_j\mathbf{X})) = \sigma_j\varrho(Y_1)$ , from which the result follows.

- 4)  $\text{cov}(\mathbf{X}) = c\Sigma$  for some  $c > 0$ . Since  $\text{var}(L) = \text{var}(\boldsymbol{\lambda}'\mathbf{X}) = \boldsymbol{\lambda}'c\Sigma\boldsymbol{\lambda}$ , 3) implies that  $\varrho(L) = \mathbb{E}(L) + \sqrt{\boldsymbol{\lambda}'\Sigma\boldsymbol{\lambda}}\varrho(Y_1) = \mathbb{E}(L) + \sqrt{\text{var}(L)}\varrho(Y_1)/\sqrt{c}$ .

□

- 2) gives a special case where VaR is subadditive and thus coherent. In particular, if  $(L_1, \dots, L_d)$  is jointly elliptical,  $\text{VaR}_\alpha$  is subadditive for  $\alpha \geq 0.5$ .
- 3) provides a useful interpretation of risk measures on  $\mathcal{M}$  in terms of the aggregation of stress tests.
- 4) is relevant to portfolio optimization. If we consider losses  $L \in \mathcal{M}$  for which  $\mathbb{E}(L)$  is fixed, the weights that minimize  $\varrho$  also minimize the variance. The portfolio minimizing  $\varrho$  is thus the same as the Markowitz variance-minimizing portfolio.

## 8.2 Risk aggregation

- A *risk aggregation rule* is a mapping

$$f(\text{EC}_1, \dots, \text{EC}_d) = \text{EC}$$

which maps the individual capital amounts  $\text{EC}_1, \dots, \text{EC}_d$  to the aggregate capital  $\text{EC}$  (economic capital). Examples are:

- ▶ *Simple summation*  $\text{EC} = \text{EC}_1 + \dots + \text{EC}_d$  (a special case of and upper bound for correlation adjusted summation)
- ▶ *Correlation adjusted summation*

$$\text{EC} = \sqrt{\sum_{i=1}^d \sum_{j=1}^d \rho_{ij} \text{EC}_i \text{EC}_j}, \quad (27)$$

where  $\rho_{ij} \in [0, 1]$  are parameters (referred to as *correlations*).

- In what follows we show that correlation adjusted summation is justified as a risk aggregation rule under various setups.

## 8.2.1 Aggregation based on loss distributions

- Suppose that the overall loss is  $L = L_1 + \dots + L_d$  where  $L_1, \dots, L_d$  are the losses arising from sub-units (e.g., business units, asset classes). Consider a translation-invariant  $\varrho$  and define

$$\varrho^{\text{mean}}(\cdot) = \varrho(\cdot - \mathbb{E}(\cdot)) = \varrho(\cdot) - \mathbb{E}(\cdot),$$

that is, the capital required to cover unexpected losses.

- The capital requirements for the sub-units are

$$\text{EC}_j = \varrho^{\text{mean}}(L_j), \quad j \in \{1, \dots, d\},$$

and the aggregate capital should be

$$\text{EC} = \varrho^{\text{mean}}(L).$$

- We require an aggregation rule  $f$  such that  $f(\text{EC}_1, \dots, \text{EC}_d) = \text{EC}$ .

- If  $\varrho(L) = \mathbb{E}(L) + k \text{sd}(L)$ ,  $k > 0$ , and  $\mathbb{E}(L^2) < \infty$  then

$$\text{sd}(L) = \sqrt{\text{var}(\mathbf{1}'\mathbf{L})} = \sqrt{\mathbf{1}'\text{cov}(\mathbf{L})\mathbf{1}} = \sqrt{\sum_{i=1}^d \sum_{j=1}^d \rho_{ij} \text{sd}(L_i) \text{sd}(L_j)},$$

where  $(\rho_{ij})_{i,j} = \text{corr}(\mathbf{L})$ , so correlation adjusted summation follows by noting that  $\text{sd}(L) = \varrho^{\text{mean}}(L)/k = \text{EC}/k$  (and  $\text{sd}(L_j) = \text{EC}_j/k$ ).

- If  $L_j = m_j + \boldsymbol{\lambda}'_j \mathbf{X}$  for  $\mathbf{X} \sim E_d(\boldsymbol{\mu}, \Sigma, \psi)$  with existing  $\text{cov}(\mathbf{X})$ , then this formula and Theorem 8.1 4) imply that correlation adjusted summation is justified for any positive-homogeneous, translation-invariant and law-invariant risk measure  $\varrho$ .
- The assumption on  $\text{cov}(\mathbf{X})$  can be dropped.

## 8.2.2 Aggregation based on stressing risk factors

- Correlation adjusted summation is used in the aggregation of capital contributions  $EC_1, \dots, EC_d$  computed by stressing individual risk factors. An example is the standard formula approach to Solvency II.
- Let  $x = X(\omega)$  be a scenario defined in terms of changes in risk factors and  $L(x)$  the corresponding loss. Assume  $L(x)$  is known and componentwise increasing.
- The  $d$  risk factors are stressed individually by amounts  $k_1, \dots, k_d$ . Capital contributions for each risk factor are computed by

$$EC_j = L(k_j e_j) - L(\mathbb{E}(X_j) e_j)$$

where  $k_j > \mathbb{E}(X_j)$  so that  $EC_j > 0$  (interpreted as the loss incurred by stressing risk factor  $j$  by  $k_j$  relative to the impact of stressing it by its expected change); an example is  $k_j = q_\alpha(X_j)$  for large  $\alpha$ .

- The following justifies correlation adjusted summation as a risk aggregation rule if  $k_j = \varrho(X_j)$  for elliptical  $X$  and  $L(X) = m + \lambda' X$ .

## Proposition 8.2 (Justification for correlation adjusted summation)

Let  $\mathbf{X} \sim E_d(\boldsymbol{\mu}, \Sigma, \psi)$  with  $\mathbb{E}(\mathbf{X}) = \boldsymbol{\mu}$ . Let  $\mathcal{M}$  be the space of linear portfolios (26) and  $\varrho$  be a pos. hom., translation- and law-invariant risk measure on  $\mathcal{M}$ . Then, for any  $L = L(\mathbf{X}) = m + \boldsymbol{\lambda}'\mathbf{X} \in \mathcal{M}$ ,

$$\text{EC} = \varrho(L - \mathbb{E}(L)) = \sqrt{\sum_{i=1}^d \sum_{j=1}^d \rho_{ij} \text{EC}_i \text{EC}_j}$$

where  $\text{EC}_j = L(\varrho(X_j)\mathbf{e}_j) - L(\mathbb{E}(X_j)\mathbf{e}_j)$  and  $\rho_{ij} = \varrho(\Sigma)_{i,j}$ .

*Proof.* Note that  $\text{EC}_j = m + \lambda_j \varrho(X_j) - (m + \lambda_j \mathbb{E}X_j) = \lambda_j \varrho(X_j - \mathbb{E}X_j)$  and plug this into Theorem 8.1 3) to see that the claim holds.  $\square$

- Thus under linearity of the losses in jointly elliptical risk-factor changes, we can aggregate the effects of single-risk-factor stresses to an aggregate capital; this applies to VaR, ES. This idea underscores correlation adjusted summation in Solvency II.
- The  $\rho_{ij}$ s are either estimated (if possible) or set by expert judgement.



## 8.2.3 Risk aggregation and Fréchet problems

- Consider the *margins-plus-copula approach* where  $L_j \sim F_j$ ,  $j \in \{1, \dots, d\}$ , are *treated as known* (estimated or postulated) and  $C$  is *unknown*.
- Consider  $L = L_1 + \dots + L_d$ . Due to the unknown  $C$  (*dependence uncertainty*), risk measures can no longer be computed explicitly.
- Our goal is to find bounds on  $\text{VaR}_\alpha$  and  $\text{ES}_\alpha$  under all possible  $C$ . Let

$$\mathcal{S}_d := \mathcal{S}_d(F_1, \dots, F_d) := \left\{ L = \sum_{j=1}^d L_j : L_j \sim F_j, j = 1, \dots, d \right\}$$

and consider

$$\overline{\varrho}(L) := \overline{\varrho}(\mathcal{S}_d) := \sup\{\varrho(L) : L \in \mathcal{S}_d(F_1, \dots, F_d)\} \quad (\text{worst } \varrho)$$

$$\underline{\varrho}(L) := \underline{\varrho}(\mathcal{S}_d) := \inf\{\varrho(L) : L \in \mathcal{S}_d(F_1, \dots, F_d)\} \quad (\text{best } \varrho)$$

- If  $\varrho = \text{ES}_\alpha$ ,  $\overline{\text{ES}}_\alpha(L) = \sum_{j=1}^d \text{ES}_\alpha(L_j)$  (subadditivity, com. additivity).  
 $\underline{\text{ES}}_\alpha$ ,  $\underline{\text{VaR}}_\alpha$ ,  $\overline{\text{VaR}}_\alpha$  depend on whether the portfolio is *homogeneous* (that is,  $F_1 = \dots = F_d$ ).

## Summary of existing results

$d = 2$ : Fully solved analytically


$d \geq 3$ : Here we distinguish:

- ▶ **Homogeneous case** ( $F_1 = \dots = F_d$ ):
  - $\underline{\text{ES}}_\alpha(L)$  solved analytically for decreasing densities (e.g. Pareto, Exponential)
  - $\underline{\text{VaR}}_\alpha(L)$ ,  $\overline{\text{VaR}}_\alpha(L)$  solved analytically for tail-decreasing densities (e.g. Pareto, Log-normal, Gamma)
- ▶ **Inhomogeneous case:**
  - Few **analytical** results: current research
  - **Numerical** methods: (Adaptive/Block) Rearrangement Algorithm

## The general problem

- We have one-period risks  $L_1 \sim F_1, \dots, L_d \sim F_d$  with **given** (estimated or postulated)  $F_1, \dots, F_d$ . The copula **C** is **unknown**.
- We need to find the **dependence uncertainty gaps** ( $\underline{\text{VaR}}_\alpha(L), \overline{\text{VaR}}_\alpha(L)$ ) (or ( $\underline{\text{ES}}_\alpha(L), \overline{\text{ES}}_\alpha(L)$ )) for  $L = L_1 + \dots + L_d$ .
- The **dependence uncertainty spreads** can be visualized as follows.

- ▶ For  $\text{VaR}_\alpha$ :



- ▶ For  $\text{ES}_\alpha$ :



The Rearrangement Algorithm (RA) can find approximate solutions to the “ $\nless$ ” cases.


## The Rearrangement Algorithm (RA)

Two columns  $\mathbf{a}, \mathbf{b}$  are *oppositely ordered* if  $(a_i - a_j)(b_i - b_j) \leq 0 \ \forall i, j$ .  
Introduce the *minimum row-sum operator*  $s(\mathbf{X}) = \min_{1 \leq i \leq N} \sum_{1 \leq j \leq d} x_{ij}$ .

### Algorithm 8.3 (RA for computing $\overline{\text{VaR}}_\alpha(L)$ )

- 1) Fix  $\alpha \in (0, 1)$ ,  $F_1^\leftarrow, \dots, F_d^\leftarrow$ ,  $N \in \mathbb{N}$  (# of discr. points),  $\varepsilon \geq 0$  (tol.)
- 2) Compute the lower bound  $\underline{s}_N$ :
  - 2.1) Define the  $(N, d)$ -matrix  $\underline{X}^\alpha = \left( F_j^\leftarrow \left( \alpha + \frac{(1-\alpha)(i-1)}{N} \right) \right)_{i,j}$ .
  - 2.2) Permute randomly each column of  $\underline{X}^\alpha$  (to avoid  $\bar{s}_N - \underline{s}_N \nrightarrow 0$ )
  - 2.3) Set  $\underline{Y}^\alpha = \underline{X}^\alpha$ . For  $1 \leq j \leq d$ , rearrange the  $j$ th column of  $\underline{Y}^\alpha$  so that it becomes oppositely ordered to the sum of all others.
  - 2.4) While  $s(\underline{Y}^\alpha) - s(\underline{X}^\alpha) > \varepsilon$ , set  $\underline{X}^\alpha$  to  $\underline{Y}^\alpha$  and repeat Step 2.3).
  - 2.5) Set  $\underline{s}_N = s(\underline{Y}^\alpha)$ .
- 3) Similarly, compute  $\bar{s}_N = s(\bar{Y}^\alpha)$  based on  $\bar{X}^\alpha = \left( F_j^\leftarrow \left( \alpha + \frac{(1-\alpha)i}{N} \right) \right)_{i,j}$ .
- 4) Return  $(\underline{s}_N, \bar{s}_N)$  (*rearrangement range*; taken as bounds on  $\overline{\text{VaR}}_\alpha(L)$ )

**Intuition:** The RA is based on the idea of joint mixability. The dfs  $F_1, \dots, F_d$  are *jointly mixable* if there exists a  $c \in \mathbb{R}$  such that  $\mathbb{P}(L_1 + \dots + L_d = c) = 1$  where  $L_1 \sim F_1, \dots, L_d \sim F_d$ . This is a notion of negative dependence and can be illustrated as follows.



$\Rightarrow$  Minimizing the variance of  $L | L > F_L^{\leftarrow}(\alpha)$  increases  $\text{VaR}_\alpha(L)$ .

## Example 8.4 (How the RA works)

1) Where it works (to compute the optimum of the maximin problem):

$$\begin{array}{ccc}
 \begin{pmatrix} 1 & 1 & 1 \\ 2 & 3 & 2 \\ 3 & 5 & 4 \\ 4 & 7 & 8 \end{pmatrix} & \xRightarrow{\Sigma_{-1} = \begin{pmatrix} 2 \\ 5 \\ 9 \\ 15 \end{pmatrix}} & \begin{pmatrix} 4 & 1 & 1 \\ 3 & 3 & 2 \\ 2 & 5 & 4 \\ 1 & 7 & 8 \end{pmatrix} \xRightarrow{\text{stable sort}} \begin{pmatrix} 4 & 5 & 1 \\ 3 & 7 & 2 \\ 2 & 3 & 4 \\ 1 & 1 & 8 \end{pmatrix} \xRightarrow{\Sigma_{-3} = \begin{pmatrix} 9 \\ 10 \\ 5 \\ 2 \end{pmatrix}} \\
 \begin{pmatrix} 4 & 5 & 2 \\ 3 & 7 & 1 \\ 2 & 3 & 4 \\ 1 & 1 & 8 \end{pmatrix} & \xRightarrow{\Sigma_{-1} = \begin{pmatrix} 7 \\ 8 \\ 7 \\ 9 \end{pmatrix}} & \begin{pmatrix} 3 & 5 & 2 \\ 2 & 7 & 1 \\ 4 & 3 & 4 \\ 1 & 1 & 8 \end{pmatrix} \checkmark \xRightarrow{\Sigma = \begin{pmatrix} 10 \\ 10 \\ 11 \\ 10 \end{pmatrix}} \widehat{\text{VaR}}_{\alpha}(L) \approx 10
 \end{array}$$

2) Where it fails (to compute the optimum of the maximin problem):

$$\begin{pmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \\ 3 & 3 & 3 \end{pmatrix} \xRightarrow{\Sigma_{-1} = \begin{pmatrix} 2 \\ 4 \\ 6 \end{pmatrix}} \begin{pmatrix} 3 & 1 & 1 \\ 2 & 2 & 2 \\ 1 & 3 & 3 \end{pmatrix} \checkmark \xRightarrow{\Sigma = \begin{pmatrix} 5 \\ 6 \\ 7 \end{pmatrix}} \widehat{\text{VaR}}_{\alpha}(L) \approx 5 < 6$$

### Example 8.5 (Par(2) margins)

Let  $L_j \sim \text{Par}(2)$  with  $\bar{F}_j(x) = (1+x)^{-2}$ ,  $j \in \{1, \dots, d\}$  (homogeneous case) and  $\alpha = 0.999$ . One obtains:

	$d = 8$		$d = 56$	
	$\theta = 2$	$\theta = 0.8$	$\theta = 2$	$\theta = 0.8$
$\overline{\text{VaR}}_\alpha(L)$	465	300 182	3454	4 683 172
$\text{VaR}_\alpha^+(L) = d \text{VaR}_\alpha(L_1)$	245	44 979	1715	314 855
$\text{VaR}_\alpha^\perp(L)$	96	75 877	293	862 855
$\underline{\text{VaR}}_\alpha(L)$	31	5622	53	5622
$\overline{\text{ES}}_\alpha(L) = d \text{ES}_\alpha(L_1)$	498	–	3486	–
$\text{ES}_\alpha^\perp(L)$	184	–	518	–
$\underline{\text{ES}}_\alpha(L)$	178	–	472	–

- The “+” and “ $\perp$ ” denote the comonotonic and independent case, resp.
- $\frac{\overline{\text{ES}}_\alpha(L)}{\overline{\text{VaR}}_\alpha(L)} \approx 1$  can be explained; see McNeil et al. (2015, Prop. 8.36).
- The dependence uncertainty spread  $\overline{\text{VaR}}_\alpha(L) - \underline{\text{VaR}}_\alpha(L) \geq \overline{\text{ES}}_\alpha(L) - \underline{\text{ES}}_\alpha(L)$  can be explained; see McNeil et al. (2015, Prop. 8.37).

## Remark 8.6

- The RA finds approximate solutions to *maximin* (for  $\overline{\text{VaR}}_\alpha(L)$ ) and *minimax* (for  $\underline{\text{VaR}}_\alpha(L)$ ) *problems* and is thus of wider interest (e.g., in Operations Research).
- For  $\underline{\text{ES}}_\alpha(L)$ , discretize the whole support of each margin, rearrange, and approximate  $\underline{\text{ES}}_\alpha(L)$  by the nonparametric  $\text{ES}_\alpha$  estimate of the row sums.
- The *Adaptive Rearrangement Algorithm (ARA)*
  - ▶ uses relative (instead of absolute) individual tolerances;
  - ▶ uses a relative joint tolerance to guarantee that  $\underline{s}_N$  and  $\overline{s}_N$  are close;
  - ▶ chooses  $N$  adaptively to reach the joint tolerance; and
  - ▶ determines convergence after each rearranged column.
- The *Block Rearrangement Algorithm* rearranges blocks of columns.



## 8.3 Capital allocation

We now study how the overall capital requirement may be disaggregated into additive contributions/units/investments. Motivation: How can we measure the risk-adjusted performance of different investments?

### 8.3.1 The allocation problem

- The performance of investments is usually measured using a *RORAC (return on risk-adjusted capital) approach* by considering 
$$\frac{\text{expected profit of investment } j}{\text{risk capital for investment } j}.$$
- The risk capital of investment  $j$  with loss  $L_j$  can be computed as follows: Compute  $\varrho(L) = \varrho(L_1 + \dots + L_d)$ . Then allocate  $\varrho(L)$  to the investments according to a capital allocation principle such that

$$\varrho(L) = \sum_{j=1}^d AC_j,$$

where the *risk contribution*  $AC_j$  is the capital allocated to investment  $j$ .

## The formal set-up

- Consider an open set  $\mathbf{1} \in \Lambda \subseteq \mathbb{R}^d \setminus \{\mathbf{0}\}$  of portfolio weights and define

$$L(\boldsymbol{\lambda}) = \boldsymbol{\lambda}' \mathbf{L} = \sum_{j=1}^d \lambda_j L_j, \quad \boldsymbol{\lambda} \in \Lambda.$$

- For a risk measure  $\varrho$ , define the *associated risk-measure function*

$$r_{\varrho}(\boldsymbol{\lambda}) = \varrho(L(\boldsymbol{\lambda})),$$

so that  $r_{\varrho}(\mathbf{1}) = \varrho(L)$ .

### 8.3.2 The Euler principle and examples

- If  $r_{\varrho}$  is positive homogeneous and differentiable at  $\boldsymbol{\lambda} \in \Lambda$ , Euler's rule implies that

$$r_{\varrho}(\boldsymbol{\lambda}) = \sum_{i=1}^d \lambda_i \frac{\partial r_{\varrho}}{\partial \lambda_i}(\boldsymbol{\lambda}) \quad \text{so} \quad \varrho(L) = r_{\varrho}(\mathbf{1}) = \sum_{j=1}^d \frac{\partial r_{\varrho}}{\partial \lambda_j}(\mathbf{1}).$$

Note that  $r_{\varrho}$  is positive homogeneous if  $\varrho$  is.

### Definition 8.7 (Euler capital allocation principle)

If  $r_\varrho$  is a pos.-hom. risk-measure function and differentiable at  $\lambda = \mathbf{1}$ , then the *Euler capital allocation principle* has risk contributions

$$\text{AC}_j = \text{AC}_j^\varrho := \frac{\partial r_\varrho}{\partial \lambda_j}(\mathbf{1}), \quad j \in \{1, \dots, d\}.$$

### Example 8.8 (Covariance principle)

Consider  $r_{\text{SD}}(\lambda) := \sqrt{\text{var}(L(\lambda))} = \sqrt{\lambda' \Sigma \lambda}$  where  $\Sigma = \text{cov}(\mathbf{L})$ . Thus

$$\text{AC}_j^\varrho = \frac{\partial r_{\text{SD}}}{\partial \lambda_j}(\mathbf{1}) = \frac{(\Sigma \mathbf{1})_j}{r_{\text{SD}}(\mathbf{1})} = \frac{\sum_{k=1}^d \text{cov}(L_j, L_k)}{r_{\text{SD}}(\mathbf{1})} = \frac{\text{cov}(L_j, L)}{\sqrt{\text{var}(L)}}.$$

### Corollary 8.9 (Euler allocation under ellipticity)

Assume that  $r_\varrho$  is the risk-measure function of a positive-homogeneous and law invariant  $\varrho$ . Let  $\mathbf{L} \sim E_d(\mathbf{0}, \Sigma, \psi)$ . Then, under an Euler allocation,

$$\frac{\text{AC}_j^\varrho}{\text{AC}_k^\varrho} = \frac{\sum_{l=1}^d \Sigma_{jl}}{\sum_{l=1}^d \Sigma_{kl}}, \quad j, k \in \{1, \dots, d\}.$$

# 9 Market Risk

9.1 Risk factors and mapping

9.2 Market risk measurement

9.3 Backtesting

## 9.1 Risk factors and mapping

### 9.1.1 The loss operator

- The key idea in this section is that of a **loss operator** for expressing the change in value of a portfolio in terms of **risk-factor changes**.
- Let the current time be  $t$  and assume the current value  $V_t$  of an asset portfolio is known, or can be computed with appropriate valuation models.
- We are interested in value changes or losses over a relatively **short time period**  $[t, t + 1]$ , for example one day, two weeks or month.
- Scaling may be applied to derive capital requirements for longer periods.
- We assume there is **no change to the composition of the portfolio** over the time period.
- The future value  $V_{t+1}$  is modelled as a random variable.

- We want to determine the distribution of the loss distribution of  $L_{t+1} = -(V_{t+1} - V_t)$ .
- We map the value at time  $t$  using the formula

$$V_t = g(\tau_t, \mathbf{Z}_t)$$

where  $\tau_t$  is time  $t$  expressed in units of **valuation time**.

## The issue of time

- We will be quite precise about the modelling of time.
- The natural time unit for valuation of positions might be yearly.
- In Black-Scholes valuation the volatility is expressed in annualized terms.
- On the other hand the **risk modelling time horizon**  $[t, t + 1]$  is typically shorter.
- Let  $\Delta t$  be the length of the time horizon in **valuation time**.

- For example, suppose that valuation time is yearly. Then a monthly time horizon would be  $\Delta t = 1/12$  and a trading day  $\Delta t = 1/250$ .
- We set  $\tau_t = t(\Delta t)$  for all  $t$  so that  $\tau_{t+1} - \tau_t = \Delta t$ .

## From the mapping to the loss operator

- The **risk factor changes** over the time horizon are

$$\mathbf{X}_{t+1} = \mathbf{Z}_{t+1} - \mathbf{Z}_t.$$

- Typically, **historical** risk factor data are available as a time series  $\mathbf{X}_{t-n}, \dots, \mathbf{X}_{t-1}, \mathbf{X}_t$  and these are used to model the behaviour of  $\mathbf{X}_{t+1}$ .
- We have

$$\begin{aligned} L_{t+1} &= -(V_{t+1} - V_t) \\ &= -(g(\tau_{t+1}, \mathbf{Z}_{t+1}) - g(\tau_t, \mathbf{Z}_t)) \\ &= -(g(\tau_t + \Delta t, \mathbf{Z}_t + \mathbf{X}_{t+1}) - g(\tau_t, \mathbf{Z}_t)). \end{aligned} \quad (28)$$

- Since the risk factor values  $\mathbf{Z}_t$  are known at time  $t$  the loss  $L_{t+1}$  is determined by the risk factor changes  $\mathbf{X}_{t+1}$ .
- Given a realization  $\mathbf{z}_t$  of  $\mathbf{Z}_t$ , the **loss operator** at time  $t$  is defined to be

$$l_{[t]}(\mathbf{x}) = -(g(\tau_t + \Delta t, \mathbf{z}_t + \mathbf{x}) - g(\tau_t, \mathbf{z}_t)), \quad (29)$$

so that

$$L_{t+1} = l_{[t]}(\mathbf{X}_{t+1}).$$

- The loss operator embodies the idea of **full revaluation**.
- From the perspective of time  $t$  the loss distribution of  $L_{t+1}$  is determined by the multivariate distribution of  $\mathbf{X}_{t+1}$ .
- Generally we consider the **conditional** distribution of  $L_{t+1}$  given history  $\mathcal{F}_t$  up to and including time  $t$ .
- Alternatively we can consider the **unconditional** distribution under assumption that  $(\mathbf{X}_t)$  form stationary time series.



## 9.1.2 Delta and delta–gamma approximations

- If the mapping function  $g$  is differentiable and  $\Delta t$  is relatively small we can approximate  $g$  with a first-order Taylor series approximation

$$g(\tau_t + \Delta t, \mathbf{z}_t + \mathbf{x}) \approx g(\tau_t, \mathbf{z}_t) + g_\tau(\tau_t, \mathbf{z}_t)\Delta t + \sum_{i=1}^d g_{z_i}(\tau_t, \mathbf{z}_t)x_i, \quad (30)$$

where the  $\tau$ -subscript and  $z_i$ -subscript denote partial derivatives with respect to (valuation) time and the risk factors respectively.

- This allows us to approximate the loss operator in (29) by the linear loss operator at time  $t$  given by

$$l_{[t]}^\Delta(\mathbf{x}) := -\left(g_\tau(\tau_t, \mathbf{z}_t)\Delta t + \sum_{i=1}^d g_{z_i}(\tau_t, \mathbf{z}_t)x_i\right). \quad (31)$$

- Note that, when working with a short time horizon  $\Delta t$ , the term  $g_\tau(\tau_t, \mathbf{z}_t)\Delta t$  is typically small and is sometimes omitted in practice.

## Example 9.1 (European call option)

- Consider portfolio consisting of one standard European call on a non-dividend paying stock  $S$  with maturity  $T$  and exercise price  $K$ .
- The Black-Scholes value of this asset at time  $t$  is  $C^{BS}(t, S_t, r, \sigma)$  where

$$C^{BS}(t, S; r, \sigma) = S\Phi(d_1) - Ke^{-r(T-t)}\Phi(d_2),$$

$\Phi$  is standard normal df,  $r$  represents risk-free interest rate,  $\sigma$  the volatility of underlying stock, and where

$$d_1 = \frac{\log(S/K) + (r + \sigma^2/2)(T - t)}{\sigma\sqrt{T - t}} \text{ and } d_2 = d_1 - \sigma\sqrt{T - t}.$$

- While in the BS model, it is assumed that interest rates and volatilities are constant, in reality they tend to fluctuate over time; they should be added to our set of risk factors.

- The risk factors:  $\mathbf{Z}_t = (\log S_t, r_t, \sigma_t)'$ .
- The risk factor changes:  $\mathbf{X}_t = (\log(S_t/S_{t-1}), r_t - r_{t-1}, \sigma_t - \sigma_{t-1})'$ .
- The mapping:

$$V_t = C^{BS}(\tau_t, S_t; r_t, \sigma_t) = g(\tau_t, \mathbf{Z}_t)$$

- For derivative positions it is quite common to use the linear loss operator

$$L_{t+1}^\Delta = l_{[t]}^\Delta(\mathbf{X}_{t+1}) = - \left( g_\tau(\tau_t, \mathbf{z}_t) \Delta t + \sum_{i=1}^3 g_{z_i}(\tau_t, \mathbf{z}_t) X_{t+1,i} \right),$$

where  $g_\tau$ ,  $g_{z_i}$  denote partial derivatives.

- $\Delta t$  is the length of the time interval expressed in years since Black-Scholes parameters relate to units of one year.

- It is more common to write the linear loss operator as

$$l_{[t]}^{\Delta}(\mathbf{x}) = - \left( C_t^{BS} + C_S^{BS} S_t x_1 + C_r^{BS} x_2 + C_{\sigma}^{BS} x_3 \right),$$

in terms of the derivatives of the BS formula or [the Greeks](#).

- ▶  $C_S^{BS}$  is known as the [delta](#) of the option.
- ▶  $C_{\sigma}^{BS}$  is the [vega](#).
- ▶  $C_r^{BS}$  is the [rho](#).
- ▶  $C_t^{BS}$  is the [theta](#).

Note the appearance of  $S_t$  in the  $C_S^{BS}$  term. This is because the risk factor is  $\ln S_t$  rather than  $S_t$  and  $C_{\ln S}^{BS} = C_S^{BS} S_t$ .

## Quadratic loss operator

- Recall the first-order Taylor series approximation of mapping in (30).
- Let  $\delta(\tau_t, \mathbf{z}_t) = (g_{z_1}(\tau_t, \mathbf{z}_t), \dots, g_{z_d}(\tau_t, \mathbf{z}_t))'$  be the first-order partial derivatives of the mapping with respect to the risk factors.

- Let  $\omega(\tau_t, \mathbf{z}_t) = (g_{z_1\tau}(\tau_t, \mathbf{z}_t), \dots, g_{z_d\tau}(\tau_t, \mathbf{z}_t))'$  denote the mixed partial derivatives with respect to time and the risk factors.
- Let  $\Gamma(\tau_t, \mathbf{z}_t)$  denote the matrix with  $(i, j)$ th element given by  $g_{z_i z_j}(\tau_t, \mathbf{z}_t)$ ; this matrix contains **gamma sensitivities** to individual risk factors on the diagonal and **cross gamma sensitivities** to pairs of risk factors off the diagonal.
- The full second-order approximation of the mapping function is  $g$  is

$$g(\tau_t + \Delta t, \mathbf{z}_t + \mathbf{x}) \approx g(\tau_t, \mathbf{z}_t) + g_\tau(\tau_t, \mathbf{z}_t)\Delta t + \boldsymbol{\delta}(\tau_t, \mathbf{z}_t)'\mathbf{x} + \frac{1}{2}(g_{\tau\tau}(\tau_t, \mathbf{z}_t)(\Delta t)^2 + 2\boldsymbol{\omega}(\tau_t, \mathbf{z}_t)'\mathbf{x}\Delta t + \mathbf{x}'\Gamma(\tau_t, \mathbf{z}_t)\mathbf{x}) .$$

- In practice, we would usually omit terms of order  $o(\Delta_t)$  (terms that tend to zero faster than  $\Delta_t$ ). In standard continuous-time financial models like Black-Scholes the risk-factor changes  $\mathbf{x}$  are of order  $\sqrt{\Delta_t}$ .

- This leaves us with the **quadratic loss operator**

$$l_{[t]}^{\Delta\Gamma}(\mathbf{x}) = -(g_{\tau}(\tau_t, \mathbf{z}_t)\Delta t + \boldsymbol{\delta}(\tau_t, \mathbf{z}_t)' \mathbf{x} + \frac{1}{2} \mathbf{x}' \Gamma(\tau_t, \mathbf{z}_t) \mathbf{x}) \quad (32)$$

which is more accurate than the linear loss operator (31).

### Example 9.2 (European call option)

The quadratic loss operator is

$$\begin{aligned} l_{[t]}^{\Delta\Gamma}(\mathbf{x}) = & l_{[t]}^{\Delta}(\mathbf{x}) - 0.5 \left( C_{SS}^{BS} S_t^2 x_1^2 + C_{rr}^{BS} x_2^2 + C_{\sigma\sigma}^{BS} x_3^2 \right) \\ & - \left( C_{Sr}^{BS} S_t x_1 x_2 + C_{S\sigma}^{BS} S_t x_1 x_3 + C_{r\sigma}^{BS} x_2 x_3 \right). \end{aligned}$$

The names of the second-order Greeks (with the exception of gamma) are rather obscure. Here are some of them:

- $C_{SS}^{BS}$  is known as the **gamma** of the option;
- $C_{\sigma\sigma}^{BS}$  is the **vomma**;
- $C_{S\sigma}^{BS}$  is the **vanna**.

## 9.1.3 Mapping bond portfolios

### Basic definitions for bond pricing

- Let  $p(t, T)$  denote the price at time  $t$  of a default-free zero-coupon bond paying one at time  $T$  (also called a discount factor).
- Time is measured in years.
- Many other fixed-income instruments such as coupon bonds or standard swaps can be viewed as portfolios of zero-coupon bonds.
- The mapping  $T \rightarrow p(t, T)$  for different maturities is one way of describing the so-called term structure of interest rates at time  $t$ . An alternative description is based on yields.
- The term structure  $T \rightarrow p(t, T)$  is known at time  $t$ .
- However the future term structure  $T \rightarrow p(t + x, T)$  for  $x > 0$  is not known at time  $t$  and must be modelled stochastically.

- The **continuously compounded yield** of a zero-coupon bond is

$$y(t, T) = -\frac{\ln p(t, T)}{T - t}. \quad (33)$$

- We have the relation

$$p(t, T) = \exp(-(T - t)y(t, T)).$$

- The yield is the constant, annualized rate implied by the price  $p(t, T)$ . Also known as spot rate.
- The mapping  $T \rightarrow y(t, T)$  is referred to as the continuously compounded **yield curve** at time  $t$ .
- Yields are comparable across different times to maturity.

## Detailed mapping of a bond portfolio

- Consider a portfolio of  $d$  default-free zero-coupon bonds with maturities  $T_i$  and prices  $p(t, T_i)$  for  $i = 1, \dots, d$ . Assume  $p(T_i, T_i) = 1$  for all  $i$ .



- By  $\lambda_i$  we denote the number of bonds with maturity  $T_i$  in the portfolio.
- The **portfolio value** at time  $t$  is given by

$$V(t) := \sum_{i=1}^d \lambda_i p(t, T_i) = \sum_{i=1}^d \lambda_i \exp(-(T_i - t)y(t, T_i)).$$

- In a detailed analysis of the change in value one takes all yields  $y(t, T_i)$ ,  $1 \leq i \leq d$ , as risk factors.
- We want to put this in the general discrete-time framework of the mapping

$$V_t = g(\tau_t, \mathbf{Z}_t).$$

- We set

$$\tau_t = t(\Delta t), \quad V_t = V(\tau_t), \quad Z_{t,i} = y(\tau_t, T_i)$$

where  $\Delta t$  is risk management time horizon in years.

- We obtain a mapping of the form

$$V_t = V(\tau_t) = g(\tau_t, \mathbf{Z}_t) = \sum_{i=1}^d \lambda_i \exp(-(T_i - \tau_t)Z_{t,i}). \quad (34)$$

## The loss operator and its approximations

- The portfolio loss is

$$\begin{aligned} L_{t+1} &= -(V_{t+1} - V_t) \\ &= -\sum_{i=1}^d \lambda_i e^{-(T_i - \tau_t)Z_{t,i}} \left( \exp(Z_{t,i}\Delta t - (T_i - \tau_{t+1})X_{t+1,i}) - 1 \right). \end{aligned}$$

- Reverting to standard bond pricing notation the loss operator is

$$l_{[t]}(\mathbf{x}) = -\sum_{i=1}^d \lambda_i p(\tau_t, T_i) \left( \exp(y(\tau_t, T_i)\Delta t - (T_i - \tau_{t+1})x_i) - 1 \right),$$

where  $x_i$  represents the change in yield of the  $i$ th bond.

- The first derivatives of the mapping function (34) are

$$g_{\tau}(\tau_t, \mathbf{z}_t) = \sum_{i=1}^d \lambda_i p(\tau_t, T_i) z_{t,i}$$

$$g_{z_i}(\tau_t, \mathbf{z}_t) = -\lambda_i (T_i - \tau_t) \exp(-(T_i - \tau_t) z_{t,i}).$$

- Inserting these in (31) and reverting to standard bond pricing notation we obtain

$$l_{[t]}^{\Delta}(\mathbf{x}) = - \sum_{i=1}^d \lambda_i p(\tau_t, T_i) \left( y(\tau_t, T_i) \Delta t - (T_i - \tau_t) x_i \right), \quad (35)$$

- For the second-order approximation we need the second derivatives with respect to yields which are

$$g_{z_i z_i}(\tau_t, \mathbf{z}_t) = \lambda_i (T_i - \tau_t)^2 \exp(-(T_i - \tau_t) z_{t,i})$$

and  $g_{z_i z_j}(\tau_t, \mathbf{z}_t) = 0$  for  $i \neq j$ .

- The quadratic loss operator (32) is

$$l_{[t]}^{\Delta\Gamma}(\mathbf{x}) = - \sum_{i=1}^d \lambda_i p(\tau_t, T_i) \left( y(\tau_t, T_i) \Delta t - (T_i - \tau_t) x_i + \frac{1}{2} (T_i - \tau_t)^2 x_i^2 \right). \quad (36)$$

## Relationship of linear operator to duration

- Consider a very simple model for the yield curve at time  $t$  in which

$$y(\tau_{t+1}, T_i) = y(\tau_t, T_i) + x$$

for all maturities  $T_i$ .

- In our mapping notation

$$Z_{t+1,i} = Z_{t,i} + X_{t+1}, \quad \forall i.$$

- In this model we assume that a **parallel shift in level** takes place along the entire yield curve.

- This is **unrealistic but frequently assumed in practice**.
- In this model the loss operator and its linear and quadratic approximations are functions of a scalar variable  $x$ , the change in level.
- Under the parallel shift model we can write

$$l_{[t]}^{\Delta}(x) = -V_t \left( A_t \Delta t - D_t x \right), \quad (37)$$

where

$$D_t := \sum_{i=1}^d \frac{\lambda_i p(\tau_t, T_i)}{v_t} (T_i - \tau_t), \quad A_t := \sum_{i=1}^d \frac{\lambda_i p(\tau_t, T_i)}{V_t} y(\tau_t, T_i).$$

- $D_t$  is usually called the (Macaulay) **duration** of the bond portfolio.
- It is a weighted sum of the times to maturity of the different cash flows in the portfolio, the weights being proportional to the discounted values of the cash flows.

## Interpreting duration

- Over short time intervals losses of value in the bond portfolio will be determined by  $l_{[t]}(x) \approx V_t D_t x$ .
- **Increases** in level of yields lead to **losses**; **decreases** lead to **gains**.
- The duration  $D_t$  is the bond pricing **analogue of the delta of an option**.
- Any two bond portfolios with equal value and duration will be subject to similar losses when there is a small parallel shift of the yield curve.
- Duration is an important tool in traditional bond-portfolio or asset-liability management.
- An asset manager, who invests in various bonds to cover promised cash flows in the future, invests in such a way that the duration of the overall portfolio of bonds and liability cash flows is equal to zero.
- Portfolios are **immunized** against small parallel shifts in yield curve, but not changes of slope and curvature.

## Relationship of quadratic operator to convexity

- It is possible to get more accurate approximations for the loss in a bond portfolio by considering second-order effects.
- The **analogue of the gamma** of an option is **convexity**. Under the parallel shift model, the quadratic loss operator (36) becomes

$$l_{[t]}^{\Delta\Gamma}(x) = -V_t \left( A_t \Delta t - D_t x + \frac{1}{2} C_t x^2 \right), \quad (38)$$

where

$$C_t := \sum_{i=1}^d \frac{\lambda_i p(\tau_t, T_i)}{V_t} (T_i - \tau_t)^2$$

is the convexity of the bond portfolio.

- The convexity is a weighted average of the squared times to maturity and is (minus) the derivative of the duration with respect to yield.

## Interpreting convexity

- Consider two portfolios (1) and (2) with identical durations  $D_t^{(1)} = D_t^{(2)}$  but differing convexities satisfying  $C_t^{(1)} > C_t^{(2)}$ .
- Ignoring terms in  $\Delta t$ , the difference in loss operators satisfies

$$l_{[t]}^{\Delta\Gamma,1}(x) - l_{[t]}^{\Delta\Gamma,2}(x) \approx -\frac{1}{2}V_t(C_t^{(1)} - C_t^{(2)})x^2 < 0.$$

- Since  $l_{[t]}^{\Delta\Gamma,1}(x) < l_{[t]}^{\Delta\Gamma,2}(x)$  an increase in the level of yields ( $x > 0$ ) will lead to smaller losses for portfolio (1)
- Since  $-l_{[t]}^{\Delta\Gamma,1}(x) > -l_{[t]}^{\Delta\Gamma,2}(x)$  a decrease in the level of yields ( $x < 0$ ) will lead to larger gains.
- For this reason higher convexity is considered a desirable attribute of a bond portfolio in risk management.



## 9.1.4 Factor models for bond portfolios

### The need for factor models

- The parallel shift model is unrealistic in practice.
- For large portfolios of fixed-income instruments, such as the overall fixed-income position of a major bank, modelling changes in the yield for every cash flow maturity date becomes impractical.
- Moreover, the statistical task of estimating a distribution for  $X_{t+1}$  is difficult because the **yields are highly dependent for different times to maturity**.
- A pragmatic approach is therefore to build a factor model for yields that captures the main features of the yield curve.
- Three-factor models of the yield curve in which the factors typically represent **level**, **slope** and **curvature** are often used in practice.

## The approach based on the Nelson and Siegel (1987) model


- We assume that at time  $t$  the yield curve can be modelled by

$$y(\tau_t, T) = Z_{t,1} + k_2(T - \tau_t, \eta_t)Z_{t,2} + k_3(T - \tau_t, \eta_t)Z_{t,3}, \quad (39)$$

where the functions  $k_2$  and  $k_3$  are given by

$$k_2(s, \eta) = \frac{1 - \exp(-\eta s)}{\eta s}, \quad k_3(s, \eta) = k_2(s, \eta) - \exp(-\eta s).$$

- Nelson-Siegel functions  $k_2(s, \eta)$  and  $k_3(s, \eta)$  for an  $\eta$  value of 0.623:



- $\eta$  is an extra tuning parameter to improve fit.
- There are other simple factor models including the [Svensson model](#).
- Clearly  $\lim_{s \rightarrow \infty} k_2(s, \eta) = \lim_{s \rightarrow \infty} k_3(s, \eta) = 0$  while  $\lim_{s \rightarrow 0} k_2(s, \eta) = 1$  and  $\lim_{s \rightarrow 0} k_3(s, \eta) = 0$ .
- It follows that

$$\lim_{T \rightarrow \infty} y(\tau_t, T) = Z_{t,1},$$

so that the first factor is usually interpreted as a [long-term level factor](#).

- $Z_{t,2}$  is interpreted as a [slope factor](#) because the difference in short-term and long-term yields satisfies

$$\lim_{T \rightarrow \tau_t} y(\tau_t, T) - \lim_{T \rightarrow \infty} y(\tau_t, T) = Z_{t,2}.$$

- $Z_{t,3}$  has an interpretation as a [curvature factor](#).

- Using (39), the portfolio mapping (34) becomes

$$V_t = g(\tau_t, \mathbf{Z}_t) = \sum_{i=1}^d \lambda_i \exp \left( - (T_i - \tau_t) \sum_{j=1}^3 k_j (T_i - \tau_t, \eta_t) Z_{t,j} \right),$$

where  $k_1(s, \eta) = 1$ .

- It is then straightforward to derive the loss operator  $l_{[t]}(\mathbf{x})$  or its linear version  $l_{[t]}^{\Delta}(\mathbf{x})$  which are functions on  $\mathbb{R}^3$  rather than  $\mathbb{R}^d$ .
- To use this method to evaluate the loss operator at time  $t$  we require realized values  $z_t$  for the risk factors  $\mathbf{Z}_t$ . We have to overcome the fact that **the Nelson-Siegel factors  $\mathbf{Z}_t$  are not directly observed** at time  $t$ . Instead **they have to be estimated** from observable yield curve data.
- Let  $\mathbf{Y}_t = (y(\tau_t, \tau_t + s_1), \dots, y(\tau_t, \tau_t + s_m))'$  denote the data vector at time  $t$ , containing the yields for  $m$  different times to maturity,  $s_1, \dots, s_m$ , where  $m$  is large.

- This is assumed to follow the factor model

$$\mathbf{Y}_t = B_t \mathbf{Z}_t + \boldsymbol{\varepsilon}_t,$$


where  $B_t \in \mathbb{R}^{m \times 3}$  is the matrix with  $i$ th row  $(1, k_2(s_i, \eta_t), k_3(s_i, \eta_t))$  and  $\boldsymbol{\varepsilon}_t \in \mathbb{R}^m$  is an error vector.

- For a given value of  $\eta_t$  the estimation of  $\mathbf{Z}_t$  can be carried out as a cross-sectional regression using weighted least squares. It is a **fundamental** factor model where the loading matrix  $B_t$  is known.
- To estimate  $\eta_t$  a more complicated optimization is carried out.

### Example 9.3

- The data are daily Canadian zero-coupon bond yields for 120 different quarterly maturities ranging from 0.25 years to 30 years.
- They have been generated using pricing data for Government of Canada bonds and treasury bills.

- We model the yield curve on the 8th August 2011.
- The estimated values are  $z_{t,1} = 3.82$ ,  $z_{t,2} = -2.75$ ,  $z_{t,3} = -5.22$  and  $\hat{\eta}_t = 0.623$ . Thus the curves  $k_2(s, \eta)$  and  $k_3(s, \eta)$  are as shown earlier.
- The fitted Nelson-Siegel curve and the data are shown below:



## The approach based on PCA

- The key difference to the Nelson-Siegel approach is that here the dimension reduction via factor modelling is **applied at the level of the risk factor changes**  $\mathbf{X}_{t+1}$  rather than the risk factors  $\mathbf{Z}_t$ .
- We recall that PCA can be used to construct factor models of the form

$$\mathbf{X}_{t+1} = \boldsymbol{\mu} + \Gamma_1 \mathbf{F}_{t+1} + \boldsymbol{\varepsilon}, \quad (40)$$

where  $\mathbf{F}_{t+1}$  is a  $p$ -dimensional vector of principal component factors ( $p < d$ ),  $\Gamma_1 \in \mathbb{R}_{d \times p}$  contains the corresponding loading matrix,  $\boldsymbol{\mu}$  is the mean vector of  $\mathbf{X}_{t+1}$  and  $\boldsymbol{\varepsilon}$  is an error vector.

- Typically, the error term is neglected and  $\boldsymbol{\mu} \approx \mathbf{0}$ , so that we make the approximation  $\mathbf{X}_{t+1} \approx \Gamma_1 \mathbf{F}_{t+1}$ .
- In the case of the **linear loss operator for the bond portfolio** in (35) we

basically replace  $l_{[t]}^{\Delta}(\mathbf{X}_{t+1})$  by

$$l_{[t]}^{\Delta}(\mathbf{F}_{t+1}) = - \sum_{i=1}^d \lambda_i p(\tau_t, T_i) (y(\tau_t, T_i) \Delta t - (T_i - \tau_t)(\Gamma_1 \mathbf{F}_{t+1})_i), \quad (41)$$

so that a  $p$ -dimensional function replaces a  $d$ -dimensional function.

- To calibrate this function, **we require an estimate for the matrix  $\Gamma_1$** . This can be obtained from historical time-series data on yield changes by estimating sample principle components.


### Example 9.4

- To estimate the  $\Gamma_1$  matrix of principal component loadings we require longitudinal (time-series) data rather than the cross-sectional data.
- We again analyse Canadian data. Recall that we have data vectors  $\mathbf{Y}_t = (y(\tau_t, \tau_t + s_1), \dots, y(\tau_t, \tau_t + s_d))$  of yields for different maturities.



- For simplicity assume that the times-to-maturity  $T_1 - \tau_t, \dots, T_d - \tau_t$  of the bonds in the portfolio correspond exactly to the times to maturity  $s_1, \dots, s_d$  available in the historical dataset.
- Assume also that the risk management horizon  $\Delta t$  is one day.
- We analyse the first differences of the data  $\mathbf{X}_t = \mathbf{Y}_t - \mathbf{Y}_{t-1}$  using PCA under an assumption of stationarity.
- In the Canadian dataset we have 2488 days of data from 2 January 2002 to 30 December 2011.
- (Note that a small error is incurred by analysing daily returns of yields with fixed times-to-maturity rather than fixed maturity date.)
- The first principle component explains 87.0% of the variance of the data, the first two components explain 95.9% and the first three components explain 97.5%.

- We choose to work with the first three principal components, meaning that we set  $p = 3$  and set the columns of  $\Gamma_1$  equal to the first three principal component loading vectors.
- These vectors are shown graphically below and lend themselves to a standard interpretation.
- The first principal component has negative loadings for all maturities; the second has negative loadings up to 10 years and positive loadings thereafter; the third has positive loadings for very short maturities (less than 2.5 years) and very long maturities (greater than 15 years) but negative loadings otherwise.
- This suggests that the first principal component can be thought of as inducing a change in the level of all yields, the second induces a change of slope and the third a change in the curvature of the yield curve.



## 9.2 Market risk measurement

The goal in this section is to estimate the distribution of

$$L_{t+1} = l_{[t]}(\mathbf{X}_{t+1})$$

or a linear or quadratic approximation thereof, where

- $\mathbf{X}_{t+1}$  is the vector of risk-factor changes from time  $t$  to time  $t + 1$ ;
- $l_{[t]}$  is the known loss operator function at time  $t$ .

The problem comprises two tasks:

- 1) on the one hand we have the **statistical problem** of estimating the distribution of  $\mathbf{X}_{t+1}$ ;
- 2) on the other hand we have the **computational or numerical problem** of evaluating the distribution of  $L_{t+1} = l_{[t]}(\mathbf{X}_{t+1})$ .

## 9.2.1 Conditional and unconditional loss distributions

- Generally, we want to compute conditional measures of risk based on the most recent information about financial markets.
- In this case, the task is to estimate  $F_{\mathbf{X}_{t+1}|\mathcal{F}_t}$ , the conditional distribution of risk-factor changes, given  $\mathcal{F}_t$ , the sigma field representing the available information at time  $t$ .

- The **conditional loss distribution** is the distribution of the loss operator  $l_{[t]}(\cdot)$  under  $F_{\mathbf{X}_{t+1}|\mathcal{F}_t}$ , that is, the distribution with distribution function

$$F_{L_{t+1}|\mathcal{F}_t} = P(l_{[t]}(\mathbf{X}_{t+1}) \leq l \mid \mathcal{F}_t).$$

- In the unconditional approach we assume that the process of risk-factor changes  $(\mathbf{X}_s)_{s \leq t}$  forms a stationary time series, at least in the recent past.

- In this case we can estimate the stationary distribution  $F_{\mathbf{X}}$  and then evaluate the unconditional loss distribution of  $l_{[t]}(\mathbf{X})$  where  $\mathbf{X}$  represents a generic random vector in  $\mathbb{R}^d$  with distribution function  $F_{\mathbf{X}}$ .
- The unconditional loss distribution is thus the distribution of the loss operator  $l_{[t]}(\cdot)$  under  $F_{\mathbf{X}}$ .
- The unconditional approach may be appropriate for longer time intervals, or for stress testing during quieter periods.
- If the risk-factor changes form an iid series, we obviously have  $F_{\mathbf{X}_{t+1}|\mathcal{F}_t} = F_{\mathbf{X}}$ , so that the conditional and unconditional approaches coincide.

## 9.2.2 Variance-covariance method

- The variance–covariance method is an **analytical** method in which strong assumptions of (conditional) normality and linearity are made.
- We assume that the **conditional distribution of risk-factor changes**  $F_{\mathbf{X}_{t+1}|\mathcal{F}_t}$  **is a multivariate normal distribution.**
- In other words, we assume that  $\mathbf{X}_{t+1} \mid \mathcal{F}_t \sim N_d(\boldsymbol{\mu}_{t+1}, \Sigma_{t+1})$ .
- The estimation of  $F_{\mathbf{X}_{t+1}|\mathcal{F}_t}$  can be carried out in a number of ways:
  - ▶ Fit **multivariate ARMA-GARCH model with multivariate normal innovations**; use model to derive estimates of  $\boldsymbol{\mu}_{t+1}$  and  $\Sigma_{t+1}$ .
  - ▶ Alternatively use the **exponentially weighted moving-average (EWMA)** procedure;  $\Sigma_{t+1}$  estimated recursively by

$$\hat{\Sigma}_{t+1} = \theta \mathbf{X}_t \mathbf{X}_t' + (1 - \theta) \hat{\Sigma}_t$$

where  $\theta$  is a small positive number (typically  $\theta \approx 0.04$ ).

- The **second critical assumption** in the variance–covariance method is that the **linear loss operator is sufficiently accurate**.
- The linear loss operator is a function of the form

$$l_{[t]}^{\Delta}(\mathbf{x}) = -(c_t + \mathbf{b}_t' \mathbf{x})$$

for some constant  $c_t$  and constant vector  $\mathbf{b}_t$ , known at time  $t$ .

- We have seen examples, including derivative and bond portfolios.
- We infer that, conditional on  $\mathcal{F}_t$ ,

$$L_{t+1}^{\Delta} = l_{[t]}^{\Delta}(\mathbf{X}_{t+1}) \sim N(-c_t - \mathbf{b}_t' \boldsymbol{\mu}_{t+1}, \mathbf{b}_t' \Sigma_{t+1} \mathbf{b}_t). \quad (42)$$

- Value-at-Risk and expected shortfall may be easily calculated for a normal loss distribution:

- ▶  $\widehat{\text{VaR}}_{\alpha} = -c_t - \mathbf{b}_t' \hat{\boldsymbol{\mu}}_{t+1} + \sqrt{\mathbf{b}_t' \hat{\Sigma}_{t+1} \mathbf{b}_t} \Phi^{-1}(\alpha).$
- ▶  $\widehat{\text{ES}}_{\alpha} = -c_t - \mathbf{b}_t' \hat{\boldsymbol{\mu}}_{t+1} + \sqrt{\mathbf{b}_t' \hat{\Sigma}_{t+1} \mathbf{b}_t} \frac{\phi(\Phi^{-1}(\alpha))}{1-\alpha}.$



## Pros and cons, extensions

- Pros:** In **contrast** to the methods that follow, variance-covariance offers **analytical solution** with no simulation.
- Cons:**
- ▶ **Linearization** may be a crude approximation.
  - ▶ Assumption of **multivariate normality** may seriously underestimate the **tail** of the loss distribution.
- Extensions:** Instead of assuming normal risk factors, the method **could be** easily **adapted to** use **multivariate Student  $t$**  or multivariate hyperbolic risk-factor changes without sacrificing tractability (the method **works for all elliptical distributions** but linearization is crucial here).

## 9.2.3 Historical simulation

- Historical simulation is by far the most popular method used by banks for the trading book.

- **The Idea**

Instead of estimating the distribution of  $l_{[t]}(\mathbf{X}_{t+1})$  under an explicit parametric model for  $\mathbf{X}_{t+1}$ , the historical-simulation method can be thought of as estimating the distribution of the loss operator under the *empirical distribution* of historical data  $\mathbf{X}_{t-n+1}, \dots, \mathbf{X}_t$ .

- We construct a univariate dataset by applying the loss operator to historical observations of the risk-factor change vector to get a set of historically simulated losses:

$$\{\tilde{L}_s = l_{[t]}(\mathbf{X}_s) : s = t - n + 1, \dots, t\}. \quad (43)$$

- To avoid full revaluation, we may apply linear/quadratic loss operator.

- The values  $\tilde{L}_s$  show what would happen to the current portfolio if the risk-factor changes on day  $s$  were to recur.
- We make inferences about the loss distribution and risk measures using these historically simulated loss data.
- **Inference about the loss distribution**
  - ▶ Use empirical quantile estimation to estimate the VaR directly from the simulated data.  
But: What about precision (sample size; confidence intervals)?
  - ▶ Fit a parametric distribution to the historical losses  $L_{t-n+1}, \dots, L_t$  and calculate risk measures from this distribution.  
But: Which distribution to fit (body or tail)?
  - ▶ Powerful: Use techniques of extreme value theory to estimate the tail of the loss distribution and related risk measures based on the historical losses  $L_{t-n+1}, \dots, L_t$ .

## Theoretical justification

If  $\mathbf{X}_{t-n+1}, \dots, \mathbf{X}_t$  are iid or, more generally, stationary, convergence of the empirical distribution to the true distribution is ensured by a suitable version of the Law of Large Numbers (e.g. Glivenko–Cantelli theorem).

## Pros and Cons

**Pros:** ▶ Easy to **implement**.

- ▶ **No** statistical **estimation** of the distribution of  $\mathbf{X}$  necessary (the **empirical df** of  $\mathbf{X}$  is used implicitly).

**Cons:** ▶ It may be difficult to collect **sufficient** quantities of relevant, synchronized **data** for all risk factors.

- ▶ Historical data may **not contain** examples of **extreme scenarios** (“driving a car by only looking in the back mirror”).

**Note:** ▶ The **dependence** here is given by the **empirical df** of  $\mathbf{X}$ .

- ▶ “Historical simulation method” is actually a misnomer; there is no simulation in the sense of random number generation.

**Extensions:** In its standard form HS is an unconditional method. There are a number of ways of extending historical simulation to take account of volatility dynamics (filtered HS).

## 9.2.4 Dynamic Historical Simulation

### A univariate approach:

- Assume that the historical simulation data  $\{\tilde{L}_s = l_{[t]}(\mathbf{X}_s) : s = t - n + 1, \dots, t\}$  are realizations from a stationary process  $(\tilde{L}_s)$ .
- To predict  $L_{t+1} = l_{[t]}(\mathbf{X}_{t+1})$ , the next random variable in this process, we assume  $(\tilde{L}_s)$  follows a model of the form  $\tilde{L}_s = \mu_s + \sigma_s Z_s$ , where
  - ▶  $\mu_s$  and  $\sigma_s$  are  $\mathcal{F}_{s-1}$ -measurable;
  - ▶  $(Z_s)$  are SWN(0, 1) innovations with distribution function  $F_Z$ .

- An example would be a **GARCH model with ARMA mean structure**.
- Writing  $\text{VaR}_\alpha^t$  for the  $\alpha$ -quantile of  $F_{L_{t+1}|\mathcal{F}_t}$ ,  $\text{ES}_\alpha^t$  for the corresponding expected shortfall, we can obtain formulas:

$$\begin{aligned}\text{VaR}_\alpha^t &= \mu_{t+1} + \sigma_{t+1} \text{VaR}_\alpha(Z), \\ \text{ES}_\alpha^t &= \mu_{t+1} + \sigma_{t+1} \text{ES}_\alpha(Z),\end{aligned}$$

where  $Z$  is a random variable with distribution function  $F_Z$ .

### ■ Estimation Options:

- ▶ Formal **parametric time series modelling** to estimate  $\mu_{t+1}$ ,  $\sigma_{t+1}$ ,  $\text{VaR}_\alpha(Z)$  and  $\text{ES}_\alpha(Z)$ .
- ▶ Often  $\mu_{t+1} \approx 0$  and can be neglected. We can use **EWMA** to estimate  $\sigma_{t-n+1}, \dots, \sigma_t, \sigma_{t+1}$  and carry out separate analysis of the **residuals**  $\{\hat{Z}_s = \tilde{L}_s / \hat{\sigma}_s, s = t - n + 1, \dots, t\}$  to estimate  $\text{VaR}_\alpha(Z)$  and  $\text{ES}_\alpha(Z)$ .

## A multivariate approach:

- We (implicitly) assume risk-factor change data  $\mathbf{X}_{t-n+1}, \dots, \mathbf{X}_t$  are realizations from process  $(\mathbf{X}_s)$  which satisfies

$$\mathbf{X}_s = \boldsymbol{\mu}_s + \Delta_s \mathbf{Z}_s, \quad \Delta_s = \text{diag}(\sigma_{s,1}, \dots, \sigma_{s,d}),$$

where  $(\boldsymbol{\mu}_s)$  is a process of vectors and  $(\Delta_s)$  a process of diagonal matrices (all assumed  $\mathcal{F}_{s-1}$ -measurable) and  $(\mathbf{Z}_s) \sim \text{SWN}(\mathbf{0}, P)$  for some correlation matrix  $P$ .

- The vector  $\boldsymbol{\mu}_s$  contains the conditional means and the matrix  $\Delta_s$  contains the volatilities of the component series at time  $s$ .
- An example of a model that fits into this framework is the [CCC-GARCH \(constant conditional correlation\)](#) process.
- The key idea of the method is to apply historical simulation to the unobserved innovations  $(\mathbf{Z}_s)$ .

- The first step is to compute estimates  $\{\hat{\boldsymbol{\mu}}_s : s = t - n + 1, \dots, t\}$  and  $\{\hat{\Delta}_s : s = t - n + 1, \dots, t\}$ .
- This can be achieved by fitting univariate time series models of ARMA-GARCH type to each of the component series in turn; alternatively we can use the univariate EWMA approach for each series.
- In the second step we construct residuals

$$\{\hat{\mathbf{Z}}_s = \hat{\Delta}_s^{-1}(\mathbf{X}_s - \hat{\boldsymbol{\mu}}_s) : s = t - n + 1, \dots, t\}$$

and treat these as “observations” of the unobserved innovations.

- We then construct the dataset

$$\{\tilde{L}_s = l_{[t]}(\hat{\boldsymbol{\mu}}_{t+1} + \hat{\Delta}_{t+1}\hat{\mathbf{Z}}_s) : s = t - n + 1, \dots, t\} \quad (44)$$

and treat these as observations of  $L_{t+1} = l_{[t]}(\mathbf{X}_{t+1})$ .

- To estimate VaR (or expected shortfall) we can apply simple empirical estimators directly to these data.



## 9.2.5 Monte Carlo

### Idea

- We estimate the distribution of  $L = \ell_{[t]}(\mathbf{X}_{t+1})$  under some **explicit parametric model** for  $\mathbf{X}_{t+1}$ .
- In contrast to the variance-covariance approach we do **not necessarily** make the problem **analytically tractable** by linearizing the loss and making an assumption of normality for the risk factors.
- Instead, make inference about  $L$  using **Monte Carlo** methods, which involves **simulating** new risk factor data.

### The method

- 1) Based on the historical risk-factor data  $\mathbf{X}_{t-n+1}, \dots, \mathbf{X}_t$ , **estimate** a suitable statistical **model** for the **risk-factor changes**.
- 2) Simulate  $N$  new data vectors  $\mathbf{X}_{t+1}^{(1)}, \dots, \mathbf{X}_{t+1}^{(N)}$  from this model.

- 3) Construct the Monte Carlo simulated losses  $L_k = \ell_{[t]}(\mathbf{X}_{t+1}^{(k)})$ ,  $k \in \{1, \dots, N\}$ .
- 4) Make inference about the loss distribution  $F_L$  and risk measures using  $L_k$ ,  $k \in \{1, \dots, N\}$  (similar possibilities as for the historical simulation method: non-parametric/parametric/EVT).

## Pros and Cons

**Pros:** ▶ Any distribution for  $\mathbf{X}_{t+1}$  can be taken  $\Rightarrow$  general

**Cons:** ▶ Can be time consuming if loss operator is difficult to evaluate, which depends on size and complexity of the portfolio.

▶ Note that MC approach does not address the problem of determining the distribution of  $\mathbf{X}_{t+1}$ .

## 9.2.6 Estimating risk measures

**Aim:** In both the historical simulation and Monte Carlo methods we estimate risk measures using simulated loss data. Let us suppose that we have data  $L_1, \dots, L_n$  from an underlying loss distribution  $F_L$  and the aim is to estimate  $\text{VaR}_\alpha = q_\alpha(F_L) = F_L^{\leftarrow}(\alpha)$  or  $\text{ES}_\alpha = (1 - \alpha)^{-1} \int_\alpha^1 q_\theta(F_L) d\theta$ . In the book we consider two possibilities:

- **L-estimators.** These are **linear** combinations of sample order statistics. Easiest to use notation for lower order statistics  $L_{(1)} \leq \dots \leq L_{(n)}$ .
- **GPD-based estimators.** These are semi-parametric estimators based on GPD approximations described in EVT chapter.

### L-estimators:

**VaR:**  $\text{VaR}_\alpha(L) = \inf\{x \in \mathbb{R} : F_L(x) \geq \alpha\}$ . Replacing  $F_L$  by  $\hat{F}_L$  we obtain an L-estimator.

$$\begin{aligned}
\widehat{\text{VaR}}_{\alpha}(L) &= \inf\{x \in \mathbb{R} : \hat{F}_L(x) \geq \alpha\} \\
&= \inf\left\{x \in \mathbb{R} : \sum_{i=1}^n I_{\{L_i \leq x\}} \geq \lceil n\alpha \rceil\right\} \\
&= \inf\left\{x \in \mathbb{R} : \sum_{i=1}^n I_{\{L_{(i)} \leq x\}} \geq \lceil n\alpha \rceil\right\} = L_{(\lceil n\alpha \rceil)}.
\end{aligned}$$

In practice, most software uses an average of two order statistics.

**ES:** Assume  $F_L$  is continuous so that

$$\text{ES}_{\alpha}(L) = \frac{\mathbb{E}(LI_{\{L > F_L^{\leftarrow}(\alpha)\}})}{1 - \alpha} = \frac{\mathbb{E}(LI_{\{L > F_L^{\leftarrow}(\alpha)\}})}{\mathbb{E}(I_{\{L > F_L^{\leftarrow}(\alpha)\}})}.$$

Replacing  $F_L$  by  $\hat{F}_L$  leads to the canonical estimator

$$\widehat{\text{ES}}_{\alpha}(L) = \frac{\sum_{i=1}^n L_i I_{\{L_i > \widehat{\text{VaR}}_{\alpha}(L)\}}}{\sum_{i=1}^n I_{\{L_i > \widehat{\text{VaR}}_{\alpha}(L)\}}}.$$

### GPD-based estimators:

We set a high threshold  $u = L_{(n-k)}$  at an order statistic and fit a GPD distribution to the  $k$  excess losses over  $u$  to obtain maximum likelihood estimates  $\hat{\xi}$  and  $\hat{\beta}$ .

For  $k/n > 1 - \alpha$  we can form the risk measure estimates:

$$\widehat{\text{VaR}}_{\alpha} = u + \frac{\hat{\beta}}{\hat{\xi}} \left( \left( \frac{1 - \alpha}{k/n} \right)^{-\hat{\xi}} - 1 \right)$$
$$\widehat{\text{ES}}_{\alpha} = \frac{\widehat{\text{VaR}}_{\alpha}}{1 - \hat{\xi}} + \frac{\hat{\beta} - \hat{\xi}u}{1 - \hat{\xi}}.$$

## 9.3 Backtesting

- Backtesting is the practice of evaluating risk measurement procedures by comparing *ex ante estimates/forecasts of risk measures* with *ex post realized losses and gains*.
- It allows us to evaluate whether a model and estimation procedure produce *credible risk measure estimates*.

### 9.3.1 Violation-based tests for VaR

- Let  $\text{VaR}_\alpha^t$  denote the  $\alpha$ -quantile of the conditional loss distribution  $F_{L_{t+1}|\mathcal{F}_t}$  and consider the event indicator variable  $I_{t+1} = I_{\{L_{t+1} > \text{VaR}_\alpha^t\}}$ .
- The event  $\{L_{t+1} > \text{VaR}_\alpha^t\}$  is a VaR *violation* or *exception*.
- Assuming a continuous loss distribution, we have, by definition of the quantile,

$$E(I_{t+1} \mid \mathcal{F}_t) = P(L_{t+1} > \text{VaR}_\alpha^t \mid \mathcal{F}_t) = 1 - \alpha, \quad (45)$$

- $I_{t+1}$  is a **Bernoulli variable** with event probability  $(1 - \alpha)$ .
- Moreover, the sequence of VaR exception indicators  $(I_t)$  is an **iid sequence**.
- The sum of exception indicators is **binomially distributed**:

$$M = \sum_{t=1}^m I_t \sim B(m, 1 - \alpha).$$

- Assume exceptions occur at times  $1 \leq T_1 < \dots < T_M \leq m$  and set  $T_0 = 0$ . The spacings  $S_j = T_j - T_{j-1}$  will be independent **geometrically distributed** rvs with mean  $1/(1 - \alpha)$ , so that

$$P(S_j = k) = \alpha^{k-1}(1 - \alpha), \quad k \in \mathbb{N}.$$

- Both of these properties are **testable in empirical data**.
- For small event probability  $1 - \alpha$ , the Bernoulli Trials Process may be well approximated by a **Poisson process**.

- Also for small  $1 - \alpha$  the geometric distribution may be approximated by an **exponential distribution**.
- Suppose we **estimate**  $\text{VaR}_\alpha^t$  at time point  $t$  by  $\widehat{\text{VaR}}_\alpha^t$ .
- In a backtest we consider empirical indicator variables

$$\hat{I}_{t+1} = I_{\{L_{t+1} > \widehat{\text{VaR}}_\alpha^t\}}.$$

- The sequence  $(\hat{I}_t)_{1 \leq t \leq m}$  **should behave** like a realization from a Bernoulli trials process with event probability  $(1 - \alpha)$ .
- To test binomial behaviour for number of violations we compute a **score test statistic**

$$Z_m = \frac{\sum_{t=1}^m \hat{I}_t - m(1 - \alpha)}{\sqrt{m\alpha(1 - \alpha)}}$$

and reject Bernoulli hypothesis at 5% level if  $Z_m > \Phi^{-1}(0.95)$ .

- Exponential spacings can be tested numerically or with a Q-Qplot.



### 9.3.2 Violation-based tests of expected shortfall

- Let  $ES_{\alpha}^t$  denote the one-period expected shortfall and  $\widehat{ES}_{\alpha}^t$  its estimate.
- Assume  $(L_t)$  follows a model of the form  $L_t = \sigma_t Z_t$ , where  $\sigma_t$  is a function of  $\mathcal{F}_{t-1}$  and the  $(Z_t)$  are  $\text{SWN}(0, 1)$  innovations.
- Then we can define a process  $(K_t)$  by

$$K_{t+1} = \frac{(L_{t+1} - ES_{\alpha}^t)}{ES_{\alpha}^t} I_{\{L_{t+1} > \text{VaR}_{\alpha}^t\}} = \frac{Z_{t+1} - ES_{\alpha}(Z)}{ES_{\alpha}(Z)} I_{\{Z_{t+1} > q_{\alpha}(Z)\}},$$

and note that it is a **zero-mean iid sequence**.

- This suggests we form **violation residuals** of the form

$$\widehat{K}_{t+1} = \frac{(L_{t+1} - \widehat{ES}_{\alpha}^t)}{\widehat{ES}_{\alpha}^t} \widehat{I}_{t+1}. \quad (46)$$

- We test for mean-zero behaviour using a bootstrap test on the non-zero violation residuals (McNeil and Frey (2000)).


### 9.3.3 Empirical comparison of methods using backtesting concepts

- We apply various VaR estimation methods to the portfolio of a hypothetical investor in international equity indexes.
- The investor is assumed to have domestic currency sterling (GBP) and to invest in the Financial Times 100 Shares Index (FTSE 100), the Standard & Poor's 500 (S&P 500) and the Swiss Market Index (SMI).
- The portfolio is **influenced by five risk factors**.
- On any day  $t$  we standardize the total portfolio value  $V_t$  in sterling to be one and assume portfolio weights are 30%, 40% and 30%, respectively.
- The loss operator and linear loss operator are:

$$l_{[t]}(\mathbf{x}) = 1 - (0.3e^{x_1} + 0.4e^{x_2+x_4} + 0.3e^{x_3+x_5})$$

$$l_{[t]}^{\Delta}(\mathbf{x}) = -(0.3x_1 + 0.4(x_2 + x_4) + 0.3(x_3 + x_5))$$

- $x_1$ ,  $x_2$  and  $x_3$  represent log-returns on the three indexes and  $x_4$  and  $x_5$  are log-returns on the GBP/USD and GBP/CHF exchange rates.



- The final picture shows the corresponding historical simulation data. The vertical dashed line is Lehman Brothers bankruptcy.

## Estimation methods:

**VC.** The [variance–covariance method](#) assuming multivariate Gaussian risk-factor changes and using the multivariate EWMA method to estimate the conditional covariance matrix of risk-factor changes.

**HS.** The standard [unconditional historical simulation](#) method.

**HS-GARCH.** The univariate [dynamic approach to historical simulation](#) in which a GARCH(1, 1) model with a constant conditional mean term and Gaussian innovations is fitted to the historically simulated losses to estimate the volatility of the next day's loss.

**HS-GARCH- $t$ .** A similar method to HS-GARCH but Student  $t$  innovations are assumed in the GARCH model.

**HS-MGARCH.** The [multivariate dynamic approach to historical simulation](#) in which GARCH(1, 1) models with constant conditional mean terms are fitted to each time series of risk-factor changes to estimate volatilities.

Year	2005	2006	2007	2008	2009	2010	2011	2012	All
Trading days	258	257	258	259	258	259	258	258	2065


### Results for 95% VaR

Expected no. of violations	13	13	13	13	13	13	13	13	103
VC	8	16	17	19	13	15	14	14	116
HS	0	6	28	49	19	6	10	1	119
HS-GARCH	9	13	22	22	13	14	9	15	117
HS-GARCH- $t$	9	14	23	22	14	15	10	15	122
HS-MGARCH	5	14	21	19	12	9	11	12	103

### Results for 99% VaR


Expected no. of violations	2.6	2.6	2.6	2.6	2.6	2.6	2.6	2.6	21
VC	2	8	8	8	2	4	5	6	43
HS	0	0	10	22	2	0	2	0	36
HS-GARCH	2	8	8	10	5	4	3	3	43
HS-GARCH- $t$	2	8	6	8	1	4	2	1	32
HS-MGARCH	0	4	4	5	0	1	2	1	17

- The HS method **does not react to changing volatility:**



- Dotted line is HS; dashed line is HS-MGARCH; vertical line is Lehmann.
- Circle is VaR violation for HS; cross is VaR violation for HS-MGARCH.

- QQ plot of spacings between exceptions:



	Violation residual test			
	95% ES	( <i>n</i> )	99% ES	( <i>n</i> )
VC	0.00	116	0.05	43
HS	0.02	119	0.25	36
HS-GARCH	0.00	117	0.05	43
HS-GARCH- <i>t</i>	0.12	122	0.68	32
HS-MGARCH	0.99	103	0.55	17

# 10 Credit risk

10.1 Credit risky instruments

10.2 Measuring credit quality

10.3 Structural models of default

10.4 Bond and CDS pricing in hazard rate models

10.5 Pricing with stochastic hazard rates



## What is credit risk?

*“Credit risk is the risk of a loss arising from the failure of a counterparty to honour its contractual obligations. This subsumes both default risk (the risk of losses due to the default of a borrower or a trading partner) and downgrade risk (the risk of losses caused by a deterioration in the credit quality of a counterparty that translates into a downgrading in some rating system). ”*

- **Obligor** = a counterparty who has a **financial obligation** to us; for example, a debtor who owes us money, a bond issuer who promises interest, or a counterparty in a derivatives transaction.
- **Default** = failure to fulfil that obligation, for example, failure to repay loan or pay interest/coupon on a loan/bond; generally due to **lack of liquidity** or **insolvency**; may entail **bankruptcy**.

## A crucial risk category

- A **portfolio of loans** or **(corporate) bonds** is obviously affected by credit risk.
- Credit risk accompanies any OTC (over-the-counter) **derivative transaction** such as a swap, because the default of one of the parties involved may substantially affect the actual pay-off of the transaction.
- There is a specialized market for **credit derivatives**, such as credit default swaps.
- Credit risk relates to the core activities of most **banks** but is also highly relevant to **insurance companies**: Insurers are exposed to substantial credit risk in their investment portfolios and counterparty default risk in their reinsurance treaties.

## Credit risk management: A range of tasks

- An enterprise needs to determine the **capital** it requires to absorb losses due to credit risk.
- Portfolios of credit-risky instruments should be well **diversified** and **optimized** according to risk-return considerations.
- Institutions need to manage their portfolio of traded credit derivatives, which involves **pricing**, **hedging** and **managing collateral** for such trades.
- Financial institutions need to control the **counterparty credit risk** in their trades and contracts with other institutions. This has particularly been the case since the 2007–2009 financial crisis.

### 10.1 Credit risky instruments

This comprises loans, bonds, derivatives subject to counterparty risk and credit derivatives such as CDSs.


## 10.1.1 Credit default swaps and related credit derivatives

- Credit derivatives are securities which are primarily used for the hedging and trading of credit risk.
- The promised pay-off of a credit derivative is related to credit events affecting one or more firms.
- Major participants in the market for credit derivatives are banks, insurance companies and investment funds.
- Retail banks are typically net buyers of protection against credit events; other investors such as hedge funds and investment banks often act as both sellers and buyers of credit protection.
- Credit default swaps (CDSs) are the workhorses of the credit derivatives market and the market for CDSs written on larger corporations is fairly liquid.

## Structure of CDS

Consider contract with maturity  $T$  and ignore counterparty credit risk. Three parties are involved (only **two directly**):

- **C (reference entity)**; default at time  $\tau_C < T$  triggers default payment.
- **A (protection buyer)**; pays premiums to B until  $\min(\tau_C, T)$ .
- **B (protection seller)**; makes default payment to A if  $\tau_C < T$ .



## CDS: Payment flows

- If the reference entity experiences a default before the maturity date  $T$  of the contract, the protection seller makes a default payment to the protection buyer, which mimics the loss due to the default of a bond issued by the reference entity (the reference asset); this part of a CDS is called the **default payment leg**.
- As compensation the protection buyer makes periodic premium payments (typically quarterly or semiannually) to the protection seller (the **premium payment leg**); after the default of the reference entity, premium payments stop. There is no initial payment.
- The premium payments are quoted in the form of an annualized percentage  $x^*$  of the notional value of the reference asset;  $x^*$  is termed the (fair or market quoted) **CDS spread**.

## Use of CDS

Investors enter into CDS contracts for various reasons.

- Bond investors with a large credit exposure to the reference entity may buy CDS protection to insure themselves against losses due to default of a bond (easier than reducing the original bond position as CDS contracts are more liquid).
- CDS contracts are also held for speculative reasons: so-called **naked** CDS positions, where the protection buyer does not own the bond are often assumed by investors who are speculating on the widening of the credit spread of the reference entity (similar to short-selling bonds issued by the reference entity.)
- Note that, in contrast to insurance, there is no requirement for the protection buyer to have **insurable interest**, that is, to actually own a bond issued by the reference entity.

## 10.2 Measuring credit quality

### Scores, ratings & measures inferred from prices

There are **two philosophies** for quantifying the credit quality or default probability of an obligor.

- 1) Credit quality can be described by a credit **rating or score** that is based on **empirical** data and expert judgement
- 2) For obligors whose equity is traded on financial markets, **prices** can be used to infer the **market's view** of the credit quality of the obligor.

Credit ratings and scores fulfill a similar function—they allow us to order obligors by their credit risk and map that risk to an estimate of the PD.

Credit ratings tend to be expressed on an ordered categorical scale whereas credit scores are often expressed in points on a metric scale.



## Rating and scoring

- The task of rating obligors is often outsourced to a rating agency such as Moody's or Standard & Poor's (S&P).
- In the S&P rating system there are seven pre-default rating categories labelled AAA, AA, A, BBB, BB, B, CCC, with AAA being the highest and CCC the lowest rating.
- Moody's uses nine pre-default rating categories labelled Aaa, Aa, A, Baa, Ba, B, Caa, Ca, C.
- A finer alpha-numeric system is also used by both agencies.
- Credit scores are traditionally used for retail customers and are based on so-called *scorecards*. Historical data is used to model default risk as a function of demographic, behavioural and financial covariates using techniques like logistic regression. The covariates are weighted and combined into a score.

## 10.2.1 Credit rating migration

- In the credit-migration approach each firm is assigned to a credit-rating category at any given time point.
- We assume that the current credit rating completely determines the default probability.
- The probability of moving from one credit rating to another over a given risk horizon (typically one year) is then specified.
- These probabilities, known as **transition probabilities**, are typically presented in the form of a matrix. They are estimated from historical data on **empirical transition rates**.
- The following example is taken from Ou (2013), (Exhibit 26). It gives average transition rates from one rating to another within one year. WR stands for withdrawn rating.

Initial rating	Rating at year-end (%)									WR
	Aaa	Aa	A	Baa	Ba	B	Caa	Ca-C	Default	
Aaa	87.20	8.20	0.63	0.00	0.03	0.00	0.00	0.00	0.00	3.93
Aa	0.91	84.57	8.43	0.49	0.06	0.02	0.01	0.00	0.02	5.48
A	0.06	2.48	86.07	5.47	0.57	0.11	0.03	0.00	0.06	5.13
Baa	0.039	0.17	4.11	84.84	4.05	7.55	1.63	0.02	0.17	5.65
Ba	0.01	0.05	0.35	5.52	75.75	7.22	0.58	0.07	1.06	9.39
B	0.01	0.03	0.11	0.32	4.58	73.53	5.81	0.59	3.85	11.16
Caa	0.01	0.02	0.02	0.12	0.38	8.70	61.71	3.72	13.34	12.00
Ca-C	0.00	0.00	0.00	0.00	0.40	2.03	9.38	35.46	37.93	14.80

- 1-year default probability for an A-rated company is estimated to be 0.06%, whereas for a Caa-rated company it is 13.3%.
- In practice a correction to the figures would probably be undertaken to account for rating withdrawals

- Rating agencies also publish **cumulative default rates** over longer time horizons.
- These provide estimates of cumulative default probabilities over several years. Alternative estimates of multi-year default probabilities can be inferred from one-year transition matrices as explained later.
- The data are taken from Ou (2013), (Exhibit 33).

Initial rating	Term						
	1	2	3	4	5	10	15
Aaa	0.00	0.01	0.01	0.04	0.11	0.50	0.93
Aa	0.02	0.07	0.14	0.26	0.38	0.92	1.75
A	0.06	0.20	0.41	0.63	0.87	2.48	4.26
Baa	0.18	0.50	0.89	1.37	1.88	4.70	8.62
Ba	1.11	3.08	5.42	7.93	10.18	19.70	29.17
B	4.05	9.60	15.22	20.13	24.61	41.94	52.22
Caa-C	16.45	27.87	36.91	44.13	50.37	69.48	79.18

## 10.2.2 Rating transitions as a Markov chain

- Let  $(R_t)$  denote a discrete-time stochastic process taking values in  $S = \{0, 1, \dots, n\}$  at times  $t = 0, 1, \dots$
- The set  $S$  defines rating states of increasing creditworthiness with 0 representing default.  $(R_t)$  models an obligor's rating over time.
- We will assume that  $(R_t)$  is a **Markov chain**. This means that it has the Markov property that

$$\mathbb{P}(R_t = k \mid R_0 = r_0, R_1 = r_1, \dots, R_{t-1} = j) = \mathbb{P}(R_t = k \mid R_{t-1} = j)$$

for all  $t \geq 1$  and all  $j, r_0, r_1, r_{t-2}, k \in S$ .

- Conditional probabilities of rating transitions given an obligors's rating history depend only on the previous rating  $R_{t-1} = j$  at the last time point and not the more distant history.
- There is evidence that rating histories show **momentum** and **stickiness** which violates the Markov assumption (Lando and Skodeberg (2002)).

## Properties of Markov chains

- The Markov chain is **stationary** if, for all  $t \geq 1$  and rating states  $j, k$ ,

$$\mathbb{P}(R_t = k \mid R_{t-1} = j) = \mathbb{P}(R_1 = k \mid R_0 = j).$$

- In this case we can define the transition matrix  $\mathbf{P} = (p_{jk})$  with elements  $p_{jk} = \mathbb{P}(R_t = k \mid R_{t-1} = j)$ , for any  $t \geq 1$ .
- The Chapman-Kolmogorov equations say that

$$\mathbb{P}(R_t = k \mid R_{t-2} = j) = \sum_{l \in S} p_{jl} p_{lk}.$$

- An implication of this is that the matrix of transition probabilities over two time steps is given by  $\mathbf{P}^2 = \mathbf{P} \times \mathbf{P}$ .
- It is not clear how a matrix of transition probabilities for a fraction of a time period can be computed (one would need continuous-time chains).
- Estimators of transition probabilities from rating history are available, see Section 10.2.2 of the book.

## 10.3 Structural models of default

### 10.3.1 The Merton model

- Merton's model (1974) is the prototype of all firm value models.
- Consider firm with stochastic asset-value ( $V_t$ ), financing itself by **equity** (i.e. by issuing shares) and **debt**.
- Assume that debt consists of single zero coupon bond with face or nominal value  $B$  and maturity  $T$ .
- Denote by  $S_t$  and  $B_t$  the value at time  $t \leq T$  of equity and debt so that

$$V_t = S_t + B_t, \quad 0 \leq t \leq T.$$

- Assume that default occurs if the firm misses a payment to its debt holders and hence only at  $T$ .

## Equity and debt as contingent claims on assets

- At  $T$  we have two possible cases:
  - 1)  $V_T > B$ . In that case the debtholders receive  $B$ ; shareholders receive residual value  $S_T = V_T - B$ , and there is no default.
  - 2)  $V_T \leq B$ . In that case the firm cannot meet its financial obligations, and shareholders hand over control to the bondholders, who liquidate the firm; hence we have  $B_T = V_T$ ,  $S_T = 0$ .
- In summary we obtain

$$S_T = (V_T - B)^+$$

$$B_T = \min(V_T, B) = B - (B - V_T)^+.$$

- The **value of equity** at  $T$  equals the pay-off of a European call option on  $V_T$  with exercise price equal to  $B$ .
- The **value of the debt** at  $T$  equals the nominal value of debt minus the pay-off of a European put option on  $V_T$ .



- The option interpretation explains certain **conflicts of interest** between shareholders and bondholders.
- For example, shareholders have more interest in the firm taking on risky projects/investments since the value of an option increases with the volatility of the underlying security.
- Bondholders have a short position on the firm's assets and would like to see the volatility reduced.

## The asset value process

It is assumed that asset value ( $V_t$ ) follows a diffusion of the form

$$dV_t = \mu_V V_t dt + \sigma_V V_t dW_t$$

for constants  $\mu_V \in \mathbb{R}$ ,  $\sigma_V > 0$ , and a Brownian motion  $(W_t)_{t \geq 0}$ , so that


$$V_T = V_0 \exp\left((\mu_V - \frac{1}{2}\sigma_V^2)T + \sigma_V W_T\right);$$

in particular  $\ln V_T \sim N\left(\ln V_0 + (\mu_V - \frac{1}{2}\sigma_V^2)T, \sigma_V^2 T\right)$ . The **default probability** is thus


$$\mathbb{P}(V_T \leq B) = \mathbb{P}(\ln V_T \leq \ln B) = \Phi\left(\frac{\ln \frac{B}{V_0} - (\mu_V - \frac{1}{2}\sigma_V^2)T}{\sigma_V \sqrt{T}}\right); \quad (47)$$

it is increasing in  $B$  and  $\sigma_V$  (for  $V_0 > B$ ) and decreasing in  $V_0$  and  $\mu_V$ .

## A default path



## A non-default path



## 10.3.2 Pricing in Merton's model

- Under some technical assumptions we can price equity and debt using the [Black–Scholes formula](#).
- The assumptions are that:
  - 1) The risk-free rate is deterministic and equal to  $r \geq 0$ .
  - 2) The asset-value process  $(V_t)$  is independent of the debt level  $B$ .
  - 3) The asset value  $(V_t)$  can be traded on a frictionless market.
- Recall that equity is a call option on the asset value  $(V_t)$ . Hence Black–Scholes formula yields

$$S_t = C^{\text{BS}}(t, V_t; \sigma_V, r, T, B) := V_t \Phi(d_{t,1}) - B e^{-r(T-t)} \Phi(d_{t,2}),$$

where the arguments are given by

$$d_{t,1} = \frac{\ln \frac{V_t}{B} + (r + \frac{1}{2}\sigma_V^2)(T-t)}{\sigma_V \sqrt{T-t}}, \quad d_{t,2} = d_{t,1} - \sigma_V \sqrt{T-t}.$$

## Pricing of debt

- The price at  $t \leq T$  of a default-free zero-coupon bond with maturity  $T$  and a face value of one equals

$$p_0(t, T) = \exp(-r(T - t)).$$

- The value of the firm's debt equals the difference between the value of default-free debt and a put option on  $(V_t)$  with strike  $B$ , i.e.

$$B_t = Bp_0(t, T) - P^{\text{BS}}(t, V_t; r, \sigma_V, B, T).$$

- The Black–Scholes formula for European puts now yields

$$B_t = p_0(t, T)B\Phi(d_{t,2}) + V_t\Phi(-d_{t,1}). \quad (48)$$

- The **path of  $(B_t)$**  is shown on the previous plots. The value of default-free debt  $Bp_0(t, T)$  is shown as a green curve.

## Risk-neutral and physical default probabilities

- Under the risk-neutral measure  $\mathbb{Q}$  the process  $(V_t)$  satisfies the SDE  $dV_t = rV_t dt + \sigma_V V_t d\tilde{W}_t$  for a standard  $\mathbb{Q}$ -Brownian motion  $\tilde{W}$ .
- The drift  $\mu_V$  is replaced by the risk-free interest rate  $r$ .
- Hence the **risk-neutral default probability** is given by

$$q = \mathbb{Q}(V_T \leq B) = \Phi\left(\frac{\ln B - \ln V_0 - (r - \frac{1}{2}\sigma_V^2)T}{\sigma_V \sqrt{T}}\right).$$

- Comparison with **physical default probability**  $p = \mathbb{P}(V_T \leq B)$  yields

$$q = \Phi\left(\Phi^{-1}(p) + \frac{\mu_V - r}{\sigma_V} \sqrt{T}\right). \quad (49)$$

- The correction term  $(\mu_V - r)/\sigma_V$  equals the **Sharpe ratio** of the firm's assets (a popular measure of the risk premium earned by the firm).
- The formula is sometimes applied in practice to go from physical to risk-neutral default probabilities.

## Credit spreads in Merton's model

- The **credit spread** measures the difference between the (continuously compounded) **yield** of a default-free zero coupon bond  $p_0(t, T)$  and a defaultable zero coupon bond  $p_1(t, T)$ , i.e.

$$\begin{aligned}c(t, T) &= \frac{-1}{T-t} (\ln p_1(t, T) - \ln p_0(t, T)) \\&= \frac{-1}{T-t} \ln \frac{p_1(t, T)}{p_0(t, T)}.\end{aligned}$$


- In Merton's model we have  $p_1(t, T) = \frac{1}{B} B_t$  and hence

$$c(t, T) = \frac{-1}{(T-t)} \ln \left( \Phi(d_{t,2}) + \frac{V_t}{B p_0(t, T)} \Phi(-d_{t,1}) \right). \quad (50)$$

- For a fixed time to maturity  $c(t, T)$  depends only on  $\sigma_V$  and on the ratio  $B p_0(t, T)/V_t$  (a measure of **indebtedness** of the firm).
- In line with economic intuition it is **increasing** in both quantities.



## Illustration of credit spreads in Merton's model



Credit spread  $c(t, T)$  (%) as function of  $\sigma_V$  (top) and time to maturity  $T - t$  (bottom) for fixed debt to firm value ratio 0.6. In upper picture  $T - t = 2$ ; in lower picture  $\sigma_V = 0.25$ .

## 10.4 Bond and CDS pricing in hazard rate models

### 10.4.1 Hazard rate models

- These are the simplest **reduced-form** credit risk models.
- A hazard rate model is a model in which the distribution of the default time of an obligor is directly specified by a hazard function without modelling the mechanism by which default occurs.
- To set up a hazard rate model we consider a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$  and a **random default time**  $\tau$  defined on this space, i.e. an  $\mathcal{F}$ -measurable rv taking values in  $[0, \infty]$ .
- We denote the df of  $\tau$  by  $F(t) = \mathbb{P}(\tau \leq t)$  and the tail or survival function by  $\bar{F}(t) = 1 - F(t)$ ; we assume that  $\mathbb{P}(\tau = 0) = F(0) = 0$ , and that  $\bar{F}(t) > 0$  for all  $t < \infty$ .

- The **jump** or **default indicator process** ( $Y_t$ ) associated with  $\tau$  is

$$Y_t = I_{\{\tau \leq t\}}, \quad t \geq 0. \quad (51)$$

- ( $Y_t$ ) is a right-continuous process which jumps from 0 to 1 at the default time  $\tau$ .
- $1 - Y_t = I_{\{\tau > t\}}$  is the **survival indicator** of the firm at time  $t$ .

### Definition 10.1 (cumulative hazard and hazard function)

The function  $\Gamma(t) = -\ln(\bar{F}(t))$  is called the **cumulative hazard function** of the random time  $\tau$ . If  $F$  is absolutely continuous with density  $f$ , the function

$$\gamma(t) = \frac{f(t)}{1 - F(t)} = \frac{f(t)}{\bar{F}(t)} = -\frac{d}{dt} \ln(\bar{F}(t))$$

is called the **hazard function** of  $\tau$ .

- The hazard function  $\gamma(t)$  gives the **hazard rate** at  $t$ , which is a measure of the instantaneous risk of default at  $t$ , given survival up to time  $t$ .

- We can represent the survival function of  $\tau$  by

$$\bar{F}(t) = \exp \left( - \int_0^t \gamma(s) \, ds \right). \quad (52)$$

- We may show that

$$\lim_{h \rightarrow 0} \frac{1}{h} \mathbb{P}(\tau \leq t + h \mid \tau > t) = \frac{1}{\bar{F}(t)} \lim_{h \rightarrow 0} \frac{F(t + h) - F(t)}{h} = \gamma(t).$$

### Example 10.2 (Weibull distribution)

For illustrative purposes we determine the hazard function for the Weibull distribution with df  $F(t) = 1 - \exp(-\lambda t^\alpha)$  for parameters  $\lambda, \alpha > 0$ . Differentiation yields

$$f(t) = \lambda \alpha t^{\alpha-1} \exp(-\lambda t^\alpha) \quad \text{and} \quad \gamma(t) = \lambda \alpha t^{\alpha-1}.$$

In particular,  $\gamma$  is decreasing in  $t$  if  $\alpha < 1$  and increasing if  $\alpha > 1$ . For  $\alpha = 1$  (exponential distribution) the hazard rate equals the constant  $\lambda$ .

## Introducing filtrations

- Filtrations model information available to investors over time.
- A **filtration**  $(\mathcal{F}_t)$  on  $(\Omega, \mathcal{F})$  is an increasing family  $\{\mathcal{F}_t : t \geq 0\}$  of sub- $\sigma$ -algebras of  $\mathcal{F} : \mathcal{F}_t \subset \mathcal{F}_s \subset \mathcal{F}$  for  $0 \leq t \leq s < \infty$ .
- $\mathcal{F}_t$  represents the state of knowledge of an observer at time  $t$ .  $A \in \mathcal{F}_t$  means that at time  $t$  we can determine if  $A$  has occurred.
- In this section we assume that only observable quantity is the default indicator  $(Y_t)$  associated with  $\tau$ . The appropriate filtration is  $(\mathcal{H}_t)$  with

$$\mathcal{H}_t = \sigma(\{Y_u : u \leq t\}), \quad (53)$$

the **default history** up to and including time  $t$ .

- $\tau$  is a  **$(\mathcal{H}_t)$ -stopping time**, since  $\{\tau \leq t\} = \{Y_t = 1\} \in \mathcal{H}_t$  for all  $t \geq 0$ .
- In order to study bond and CDS pricing in hazard rate models we need to compute conditional expectations with respect to the  $\sigma$ -algebra  $\mathcal{H}_t$ .

## A useful result

### Lemma 10.3

Let  $\tau$  be a default time with jump indicator process  $Y_t = I_{\{\tau \leq t\}}$  and natural filtration  $(\mathcal{H}_t)$ . Then, for any integrable rv  $X$  and any  $t \geq 0$ , we have

$$\mathbb{E}(I_{\{\tau > t\}}X \mid \mathcal{H}_t) = I_{\{\tau > t\}} \frac{\mathbb{E}(I_{\{\tau > t\}}X)}{\mathbb{P}(\tau > t)}. \quad (54)$$

This result can be used to determine conditional survival probabilities. For  $t < T$ , applying (54) with  $X := I_{\{\tau > T\}}$  we get

$$\mathbb{P}(\tau > T \mid \mathcal{H}_t) = I_{\{\tau > t\}} \exp\left(-\int_t^T \gamma(s)ds\right), \quad t < T. \quad (55)$$

## Martingale property of jump indicator process

### Proposition 10.4

The process  $(M_t)$  defined as

$$M_t = Y_t - \int_0^t I_{\{\tau > u\}} \gamma(u) \, du, \quad t \geq 0$$

is an  $(\mathcal{H}_t)$ -martingale, that is  $\mathbb{E}(M_s \mid \mathcal{H}_t) = M_t$  for all  $0 \leq t \leq s < \infty$ .

### 10.4.2 Risk-neutral pricing revisited

- According to the first fundamental theorem of asset pricing, a model for security prices is arbitrage free if and (essentially) only if it admits at least one equivalent martingale measure  $\mathbb{Q}$ .
- When building a model for pricing derivatives it is a natural shortcut to model the objects of interest—such as interest rates and default times—directly, under a martingale measure  $\mathbb{Q}$ .

- So-called **martingale modelling** is particularly convenient if the value  $H$  of the underlying assets at some maturity date  $T$  is exogenously given, as in the case of zero-coupon bonds.
- The underlying asset at time  $t < T$  can be computed as the conditional expectation under  $\mathbb{Q}$  of the discounted value at maturity via the risk-neutral pricing rule

$$V_t = \mathbb{E}^{\mathbb{Q}}(e^{-\int_t^T r_s ds} H \mid \mathcal{F}_t). \quad (56)$$

- Model parameters are determined using the requirement that at time  $t = 0$  the model price should coincide with the market price of the security; this is known as **calibration** to market data.



## Pros and cons of Martingale modelling

- Martingale modelling ensures that the resulting model is arbitrage free, which is important for pricing many different securities simultaneously.
- The approach is frequently adopted in default-free term structure models and in reduced-form models for credit-risky securities.
- Martingale modelling has two drawbacks.
  - 1) Historical information is largely useless in estimating model parameters.
  - 2) Realistic models for pricing credit derivatives are typically **incomplete**.
- An arbitrage-free market is complete if and only if there is exactly one equivalent martingale measure.
- In incomplete markets there may be more than one equivalent martingale measure. It will generally not be possible to find a replicating strategy for a derivative (one cannot eliminate all risk by dynamic hedging).

### 10.4.3 Bond pricing

- It suffices to consider **zero-coupon bonds**.
- We use martingale modelling and work directly under some martingale measure  $\mathbb{Q}$ .
- We assume that under  $\mathbb{Q}$  the default time  $\tau$  is a random time with deterministic risk-neutral hazard function  $\gamma^{\mathbb{Q}}(t)$ .
- The information available to investors at time  $t$  is given by the sigma algebra  $\mathcal{H}_t = \sigma(\{Y_u : u \leq t\})$ .
- We take **interest rates** and **recovery rates** to be **deterministic**.
- The percentage loss given default is denoted by  $\delta \in (0, 1)$ .
- The continuously compounded interest rate is denoted by  $r(t) \geq 0$ .
- The price of the default-free zero-coupon bond with maturity  $T \geq t$  is  $p_0(t, T) = \exp(-\int_t^T r(s) ds)$ .

## Analysing the payments

- The payments of a defaultable zero-coupon bond can be represented as a combination of a **survival claim** that pays one unit at the maturity date  $T$  and a **recovery payment** in case of default.
- The survival claim has pay-off  $I_{\{\tau > T\}}$ .
- Recall from (55) that

$$\mathbb{Q}(\tau > T \mid \mathcal{H}_t) = I_{\{\tau > t\}} \exp \left( - \int_t^T \gamma^{\mathbb{Q}}(s) ds \right)$$

and define  $R(t) = r(t) + \gamma^{\mathbb{Q}}(t)$ .

- Then the price of a survival claim at time  $t$  equals

$$\begin{aligned} \mathbb{E}^{\mathbb{Q}}(p_0(t, T) I_{\{\tau > T\}} \mid \mathcal{H}_t) &= \exp \left( - \int_t^T r(s) ds \right) \mathbb{Q}(\tau > T \mid \mathcal{H}_t) \\ &= I_{\{\tau > t\}} \exp \left( - \int_t^T R(s) ds \right). \end{aligned} \quad (57)$$

- Note that for  $\tau > t$ , this can be viewed as the price of a default-free zero-coupon bond with adjusted interest rate  $R(t) > r(t)$ .
- A similar relationship between defaultable and default-free bond prices can be established in many reduced-form credit risk models.

## Recovery models

### 1) Recovery of Treasury (RT).

- The RT model was proposed by Jarrow and Turnbull (1995).
- If default occurs at some point in time  $\tau \leq T$ , the owner of the defaulted bond receives  $(1 - \delta_\tau)$  units of the default-free zero-coupon bond  $p_0(\cdot, T)$  at time  $\tau$ , where  $\delta_\tau \in [0, 1]$  models the percentage loss given default.
- At maturity  $T$  the holder of the defaultable bond therefore receives the payment  $I_{\{\tau > T\}} + (1 - \delta_\tau)I_{\{\tau \leq T\}}$ .

### 2) Recovery of Face Value (RF).

- Under RF, if default occurs at  $\tau \leq T$ , the holder of the bond receives a recovery payment of size  $(1 - \delta_\tau)$  immediately at the default time  $\tau$ .
- Note that even with deterministic loss given default and deterministic interest rates, the **value at maturity of the recovery payment is random** as it depends on the exact timing of default.

RF is slightly more realistic; RT is slightly easier to analyse.

## Pricing recovery payment under RT

- The value of the recovery payment at the maturity date  $T$  is

$$(1 - \delta)I_{\{\tau \leq T\}} = (1 - \delta) - (1 - \delta)I_{\{\tau > T\}}.$$

- Using (57), the value of the recovery payment at time  $t < T$  is hence

$$(1 - \delta)p_0(t, T) - (1 - \delta)I_{\{\tau > t\}} \exp\left(-\int_t^T R(s) ds\right).$$

- Hence the value of the bond is

$$p_1(t, T) = (1 - \delta)p_0(t, T) + \delta I_{\{\tau > t\}} \exp\left(-\int_t^T R(s) ds\right).$$

## Pricing recovery payment under RF

- Under the RF-hypothesis the recovery payment takes the form  $(1 - \delta)I_{\{\tau \leq T\}}$  where the payment occurs directly at time  $\tau$ .
- A payments of this form is a **payment-at-default claim**.
- The value of the recovery payment at time  $t \leq T$  equals

$$\mathbb{E}^{\mathbb{Q}}\left((1 - \delta)I_{\{t < \tau \leq T\}} \exp\left(-\int_t^{\tau} r(s)ds\right) \middle| \mathcal{H}_t\right).$$

- Using (54) we may show that

$$\begin{aligned} & \mathbb{E}^{\mathbb{Q}}\left((1 - \delta)I_{\{t < \tau \leq T\}} \exp\left(-\int_t^{\tau} r(s)ds\right) \middle| \mathcal{H}_t\right) \\ &= (1 - \delta)I_{\{\tau > t\}} \int_t^T \gamma^{\mathbb{Q}}(s) \exp\left(-\int_t^s R(u)du\right) ds. \end{aligned}$$

## 10.4.4 CDS pricing

First we recall payment flows. We write  $\tau = \tau_C$  and consider the following contract:

- Premium payments.

- ▶ These are due at times  $0 < t_1 < \dots < t_N$  measured in years.
- ▶ If  $\tau > t_k$ , A pays a premium of size  $x^*(t_k - t_{k-1})$  at  $t_k$ , where  $x^*$  denotes the fair swap spread.
- ▶ After  $\tau$  premium payments stop, no initial payment.

- Default payment.

- ▶ If  $\tau < t_N = T$ , B makes a default payment  $\delta$  at  $\tau$ .
- ▶ Sometimes B receives an accrued premium payment of size  $x^*(\tau - t_k)$  for  $\tau \in (t_k, t_{k+1})$ . We ignore this feature for simplicity.

## Valuing the premium leg

- The premium leg consists of a set of **survival claims**.
- Introduce a function of  $x$  given by

$$\begin{aligned} & V_t^{\text{prem}}(x; \gamma^{\mathbb{Q}}) \\ &= \mathbb{E}^{\mathbb{Q}} \left( \sum_{k: t_k > t} \exp \left( - \int_t^{t_k} r(u) \, du \right) x(t_k - t_{k-1}) I_{\{\tau > t_k\}} \mid \mathcal{H}_t \right) \\ &= x \sum_{k: t_k > t} p_0(t, t_k) (t_k - t_{k-1}) \mathbb{Q}(\tau > t_k \mid \mathcal{H}_t), \end{aligned}$$

which is easily computed using  $\mathbb{Q}(\tau > t_k) = \exp(-\int_0^{t_k} \gamma^{\mathbb{Q}}(s) ds)$ .

- We obtain

$$V_t^{\text{prem}}(x; \gamma^{\mathbb{Q}}) = I_{\{\tau > t\}} x \sum_{k: t_k > t} (t_k - t_{k-1}) \exp \left( - \int_t^{t_k} R(u) \, du \right).$$



## Valuing the default leg

- The default payment leg is a typical payment-at-default claim.
- We obtain

$$\begin{aligned} & V_t^{\text{def}}(\gamma^{\mathbb{Q}}) \\ &= \mathbb{E}^{\mathbb{Q}}\left(\delta I_{\{t < \tau \leq t_N\}} \exp\left(-\int_t^{\tau} r(s) \, ds\right) \mid \mathcal{H}_t\right) \\ &= I_{\{\tau > t\}} \delta \int_t^{t_N} \gamma^{\mathbb{Q}}(s) \exp\left(-\int_t^s R(u) \, du\right) \, ds. \end{aligned}$$

## The fair CDS spread

- The fair CDS spread  $x_t^*$  quoted for the contract at time  $t$  is chosen such that the value of the contract is equal to zero.

- The equation  $V_t^{\text{prem}}(x_t^*; \gamma^{\mathbb{Q}}) = V_t^{\text{def}}(\gamma^{\mathbb{Q}})$  yields

$$x_t^* = I_{\{\tau > t\}} \frac{\delta \int_t^{t_N} \gamma^{\mathbb{Q}}(s) \exp\left(-\int_t^s R(u) du\right) ds}{\sum_{k: t_k > t} (t_k - t_{k-1}) \exp\left(-\int_t^{t_k} R(s) ds\right)}. \quad (58)$$

## Model calibration

- We have to calibrate our model to the available market information. Hence we have to determine the implied risk-neutral hazard function  $\gamma^{\mathbb{Q}}(t)$ , which ensures that the fair CDS spreads implied by the model equal the spreads quoted in the market.
- Suppose that the market information at time  $t = 0$  consists of the fair spread  $x^*$  of one CDS with maturity  $t_N$ .
- In that case  $\gamma^{\mathbb{Q}}(s)$  is taken constant: for all  $s \geq 0$ ,  $\gamma^{\mathbb{Q}}(s) = \bar{\gamma}^{\mathbb{Q}}$  for some  $\bar{\gamma}^{\mathbb{Q}} > 0$ .

- $\bar{\gamma}^{\mathbb{Q}}$  has to solve the equation

$$x^* \sum_{k=1}^N p_0(0, t_k)(t_k - t_{k-1})e^{-\bar{\gamma}^{\mathbb{Q}}t_k} = \delta \bar{\gamma}^{\mathbb{Q}} \int_0^{t_N} p_0(0, t)e^{-\bar{\gamma}^{\mathbb{Q}}t} dt.$$

- There is a unique solution.
- If we observe spreads for several CDSs on the same reference entity but with different maturities, a constant function is not sufficient. Instead one typically uses piecewise constant or linear hazard functions.
- A first approximation to the implied hazard rate is given by  $\bar{\gamma}^{\mathbb{Q}} \approx x^*/\delta$ .
- This approximation implies that the one-year default probability satisfies  $\mathbb{Q}(\tau \leq 1) = 1 - e^{-\bar{\gamma}^{\mathbb{Q}}} \approx \bar{\gamma}^{\mathbb{Q}} \approx x^*/\delta$ .

## 10.5 Pricing with stochastic hazard rates

### Why stochastic hazard rates?

- In hazard rate models the only risk factor is default risk  $\Rightarrow$  Credit spreads evolve deterministically prior to default, which is clearly unrealistic.
- Moreover, it is not possible to price options on bonds or CDSs or to do risk management for bond portfolios in such models.
- Hence it is of interest to consider models where hazard rate is a stochastic process  $(\gamma_t)_{t \geq 0}$ ; typically hazard rate is driven by a second stochastic process  $\Psi$ , that is  $\gamma_t = \gamma(\Psi_t)$ .
- Simplest such model class are doubly-stochastic random times.

## 10.5.1 Doubly stochastic random times

**Setup.** We work on  $(\Omega, \mathcal{F}, \mathbb{P})$  with **background filtration**  $(\mathcal{F}_t)$  containing information about all other economic events except the default event.

Consider a random time  $\tau$ , that is a measurable rv with values in  $(0, \infty)$ .

- $Y_t = I_{\{\tau \leq t\}}$  is the associated **default indicator** and  $(\mathcal{H}_t) = \sigma\{Y_s, s \leq t\}$  is the **default history** up to  $t$ .
- Define new filtration  $\mathcal{G}_t = \mathcal{F}_t \vee \mathcal{H}_t$ ,  $t \geq 0$ , i.e.  $\mathcal{G}_t$  contains background info  $\mathcal{F}_t$  and default history up to  $t$  (this is the information available to investors).

### Definition 10.5

$\tau$  is called *doubly stochastic* if there is a positive  $(\mathcal{F}_t)$ -adapted process  $(\gamma_t)$  (the hazard rate process) such that for all  $t \geq 0$

$$\mathbb{P}(\tau > t \mid \mathcal{F}^\infty) = \exp\left(-\int_0^t \gamma_s ds\right). \quad (59)$$

### Comments.

- Here  $\mathcal{F}_\infty = \sigma(\bigcup_{t \geq 0} \mathcal{F}_t)$ . Conditioning on  $\mathcal{F}_\infty$  thus means that we know the past and future economic environment and in particular the entire trajectory  $(\gamma_s(\omega))_{s \geq 0}$  of the hazard rate.
- Relation (59) implies that, given the economic environment  $\mathcal{F}^\infty$ ,  $\tau$  is a random time with deterministic hazard function  $s \mapsto \gamma_s(\omega)$ .
- In the literature doubly stochastic random times are also known as **conditional Poisson** or **Cox** random times.

## Sampling doubly stochastic random times

A simple algorithm is based on the following result

### Lemma 10.6

Let  $E$  be a standard exponentially distributed rv independent of  $\mathcal{F}^\infty$ , that is  $\mathbb{P}(\tau > t \mid \mathcal{F}^\infty) = e^{-t}$ . Let  $(\gamma_t)$  be a positive  $\mathcal{F}_t$ -adapted process with  $\int_0^t \gamma_s ds < \infty$  for all  $t$ . Define  $\tau$  by


$$\tau := \inf \left\{ t \geq 0 : \int_0^t \gamma_s ds \geq E \right\}. \quad (60)$$

Then  $\tau$  is doubly stochastic with hazard-rate process  $(\gamma_t)$ .

**Algorithm.** (threshold simulation)

- 1) Generate  $E \sim \text{Exp}(1)$ .
- 2) Generate a trajectory  $(\gamma_s)_{s=0}^\infty$  of hazard rate process.
- 3) Return  $\tau := \inf \{ t \geq 0 : \int_0^t \gamma_s ds \geq E \}$

## Graphical illustration.



A graphical illustration of threshold simulation;  $E \approx 0.44$ ,  $\tau \approx 6.59$ .



## 10.5.2 Pricing formulas

### Setup.

- Consider arbitrage-free security market model on  $(\Omega, \mathcal{F}, (\mathcal{F}_t), \mathbb{Q})$  where  $\mathbb{Q}$  is **equivalent martingale measure**. Prices of default-free securities  $(\mathcal{F}_t)$ -adapted;  $B_t = \exp(\int_0^t r_s \, ds)$  models default-free savings account.
- Let  $\tau$  be the default time of some company. As before we set  $\mathcal{H}_t = \sigma(\{Y_s : s \leq t\})$  and  $\mathcal{G}_t = \mathcal{F}_t \vee \mathcal{H}_t$ ; this is the information available to investors at time  $t$ .
- We use **martingale-modelling**. Hence price at  $t$  of an  $\mathcal{G}_T$ -measurable contingent claim  $H$  is given by

$$H_t = \mathbb{E}^{\mathbb{Q}}\left(\exp\left(-\int_t^T r_s \, ds\right) H \mid \mathcal{G}_t\right). \quad (61)$$

- Under  $\mathbb{Q}$ ,  $\tau$  is a doubly stochastic random time with background filtration  $(\mathcal{F}_t)$  and hazard rate process  $(\gamma_t)$ .

## Key building blocks

The pricing of bonds and CDSs can be reduced to the pricing of the following building blocks:

- A **survival claim**, i.e. a promised  $\mathcal{F}_T$ -measurable payment  $X$  which is made at time  $T$  if there is no default; the actual payment of the survival claim equals  $XI_{\{\tau > T\}}$ .
- A **payment-at-default claim** of the form  $Z_\tau I_{\{\tau \leq T\}}$ , where  $Z = (Z_t)_{t \geq 0}$  is an  $(\mathcal{F}_t)$  adapted stochastic process and where  $Z_\tau$  is short for  $Z_{\tau(\omega)}(\omega)$ . Note that the payment is made directly at  $\tau$ , provided that  $\tau \leq T$  where  $T$  is the maturity date of the claim.

**Example.** Defaultable bond is a combination of a survival claim and a payment at default claim (the recovery payment).

## Pricing the building blocks

Next result shows that pricing of building blocks can be reduced to pricing problem for default-free claims with adjusted interest rate.

### Theorem 10.7

Define adjusted interest rate  $R_t = r_t + \gamma_t$ . Under the above assumptions (in particular for  $\tau$  doubly stochastic) it holds that

$$\begin{aligned}\mathbb{E}^{\mathbb{Q}}\left(\exp\left(-\int_t^T r_s ds\right)I_{\{\tau>T\}}X \mid \mathcal{G}_t\right) &= I_{\{\tau>t\}}\mathbb{E}^{\mathbb{Q}}\left(\exp\left(-\int_t^T R_s ds\right)X \mid \mathcal{F}_t\right), \\ \mathbb{E}^{\mathbb{Q}}\left(I_{\{t<\tau\leq T\}}\exp\left(-\int_t^{\tau} r_s ds\right)Z_{\tau} \mid \mathcal{G}_t\right) \\ &= I_{\{\tau>t\}}\mathbb{E}^{\mathbb{Q}}\left(\int_t^T Z_s\gamma_s \exp\left(-\int_t^s R_u du\right) ds \mid \mathcal{F}_t\right).\end{aligned}$$

## 10.5.3 Applications

**Corporate bonds and RF.** The price at  $t$  of a defaultable zero-coupon bond with maturity  $T \geq t$  is  $p_1(t, T)$ ; price of corresponding default-free bond is  $p_0(t, T)$ .

Consider the **recovery of face value (RF)** recovery model: if default occurs at  $\tau \leq T$ , bondholder receives  $(1 - \delta_\tau)$  immediately at  $\tau$ . Bond price is thus sum of a survival claim and a payment at default claim:

$$p_1(t, T) = I_{\{\tau > t\}} \left( \mathbb{E}^{\mathbb{Q}} \left( \exp \left( - \int_t^T R_s ds \right) \mid \mathcal{F}_t \right) \right. \\ \left. + (1 - \delta) \mathbb{E}^{\mathbb{Q}} \left( \int_t^T \gamma_s \exp \left( - \int_t^s R_u du \right) ds \mid \mathcal{F}_t \right) \right)$$

Recall that the **credit spread** of a defaultable bond is

$$c(t, T) = -\frac{1}{T - t} (\ln p_1(t, T) - \ln p_0(t, T)).$$

We consider the **instantaneous credit spread**

$$c(t, t) = \lim_{T \rightarrow t} c(t, T) = -\frac{\partial}{\partial T} \Big|_{T=t} (\ln p_1(t, T) - \ln p_0(t, T)). \quad (62)$$

It holds that  $c(t, t) = \delta \gamma_t^{\mathbb{Q}} > 0$ .

**CDS contracts.** Here premium payments constitute a sequence of survival claims; default payment is a payment-at-default claim. This gives the following formula for the **fair CDS spread**  $x^*$  at  $t$ :

$$x^* = \frac{\delta \mathbb{E}^{\mathbb{Q}}(\int_t^T \gamma_s e^{-\int_t^s R_u du} ds \mid \mathcal{F}_t)}{\sum_{T_k > t} (t_k - t_{k-1}) \mathbb{E}^{\mathbb{Q}}(\exp(-\int_t^{t_k} R_s ds \mid \mathcal{F}_t))}. \quad (63)$$

- For  $T \rightarrow t$  we get that  $x^*$  converges to  $\delta \gamma_t$ .
- The formula is a generalization of (58).

## 10.5.4 Evaluation of pricing formulas

In most models with doubly stochastic default time used in practice it is assumed that  $(r_t)$  and  $(\gamma_t)$  are functions of some **Markov process**  $(\Psi_t)$

- Natural background filtration is  $(\mathcal{F}_t) = \sigma(\{\Psi_s : s \leq t\})$ .
- $R_t := r_t + \gamma_t$  is of the form  $R_t = R(\Psi_t)$

To evaluate general pricing formulas we hence have to compute conditional expectations of the form

$$\mathbb{E}\left(e^{-\int_t^T R(\Psi_s) ds} g(\Psi_T) \mid \mathcal{F}_t\right) \quad (64)$$

for generic  $g : D \rightarrow \mathbb{R}_+$ . Since  $(\Psi_t)$  is Markov, (64) is a function  $f(t, \Psi_t)$  of time and of  $\Psi_t$ .

A case where this can be done explicitly is the CIR model.

## The CIR square-root diffusion

The **CIR** or **square-root** diffusion model due to Cox et al. (1985) is a popular model. **CIR dynamics**.

$$d\Psi_t = \kappa(\bar{\theta} - \Psi_t) dt + \sigma\sqrt{\Psi_t} dW_t, \quad \Psi_0 = \psi > 0, \quad (65)$$

for parameters  $\kappa, \bar{\theta}, \sigma > 0$  and state space  $D = [0, \infty)$ .

### Properties.

- (65) implies that  $(\Psi_t)$  is *mean reverting*.
- Mean reversion sufficiently strong  $\Rightarrow$  trajectories never reach zero: Let  $\tau_0(\Psi) := \inf\{t \geq 0 : \Psi_t = 0\}$ . For  $\kappa\bar{\theta} \geq \frac{1}{2}\sigma^2$ ,  $\mathbb{P}(\tau_0(\Psi) < \infty) = 0$ ; for  $\kappa\bar{\theta} < \frac{1}{2}\sigma^2$ ,  $\mathbb{P}(\tau_0(\Psi) < \infty) = 1$ .

## CIR term structure

### Theorem 10.8

Suppose that the factor  $\Psi$  follows the CIR model and that adjusted interest rate is an affine function of the state,  $R(\psi) = \rho^0 + \rho^1\psi$ . Then it holds that

$$\mathbb{E}\left(\exp\left(-\int_t^T (\rho^0 + \rho^1\Psi_s) ds\right) \middle| \Psi_t\right) = \exp(\alpha(T-t) + \beta(T-t)\Psi_t),$$

where

$$\beta(\tau) = \frac{-2\rho^1(e^{\gamma\tau} - 1)}{\gamma - \kappa + e^{\gamma\tau}(\gamma + \kappa)},$$
$$\alpha(\tau) = -\rho^0\tau + 2\frac{\kappa\bar{\theta}}{\sigma^2} \ln\left(\frac{2\gamma e^{\tau(\gamma+\kappa)/2}}{\gamma - \kappa + e^{\gamma\tau}(\gamma + \kappa)}\right),$$

and  $\tau := T - t$ ,  $\gamma := \sqrt{\kappa^2 + 2\sigma^2\rho^1}$



## 10.5.5 Extensions


- It is possible to extend the above to CIR models with jumps,

$$d\Psi_t = \kappa(\bar{\theta} - \Psi_t) dt + \sigma\sqrt{\Psi_t} dW_t + dZ_t$$

where  $Z_t = \sum_{T_n \leq t} Z_n$  is a compound Poisson process with jump intensity  $\lambda_t = \lambda_0 + \lambda_1 \Psi_t$ ,  $\lambda_0$  and  $\lambda_1 > 0$  and the  $Z_n$  are iid positive rvs, for instance exponentially distributed.

- The computation of payment-at-default claims is also possible with “affine model technology”.

## Numerical example



Spreads of defaultable zero-coupon bonds in an affine model for various recovery assumptions. It holds  $\Psi_0 \approx 0.0533$ ,  $r = 6\%$  and  $\delta = 0.5$ . Note that under the RF recovery model (dashed line) the spread becomes negative for large times to maturity; this is not true under other recovery assumptions.

# 11 Portfolio credit risk management

11.1 Threshold models


11.2 Mixture models

11.3 Statistical inference for portfolio credit models

## Importance of default dependence

Dependence between defaults (and downgrades) is a key issue in credit risk management. There are two main sources of dependence between defaults:

- Dependence caused by **common factors** (for example, interest rates and changes in economic growth) affecting all obligors
- Default of company A may have direct impact on default probability of company B and vice versa because of **direct business relations**, a phenomenon known as **contagion**



Comparison of the loss distribution of a homogeneous portfolio of 1000 loans with a default probability of  $p_1 = \dots = p_{1000} = 1\%$  assuming (i) independent defaults and (ii) a default correlation of  $\rho(Y_i, Y_j) = 0.5\%$ . Case (ii) can be considered as roughly representative for BB-rated loans.

# 11.1 Threshold models

## 11.1.1 Notation for one-period portfolio models

- Consider portfolio of  $m$  firms and time horizon  $T = 1$  (say one year).
- For  $1 \leq i \leq m$ , let  $R_i$  be a state indicator for obligor  $i$  at time  $T$  taking values in the set  $\{0, 1, \dots, n\}$ ; we interpret the value 0 as default and non-zero values as states of increasing credit quality. At time  $t = 0$  obligors are assumed to be in some non-default state.
- Mostly we will concentrate on the binary outcomes of default and non-default. We write  $Y_i$  for the default indicator variables so that  $Y_i = 1 \iff R_i = 0$  and  $Y_i = 0 \iff R_i > 0$ .
- The random vector  $\mathbf{Y} = (Y_1, \dots, Y_m)'$  is a vector of default indicators for the portfolio and  $p(\mathbf{y}) = \mathbb{P}(Y_1 = y_1, \dots, Y_m = y_m)$ ,  $\mathbf{y} \in \{0, 1\}^m$ , is its joint probability function; the marginal default probabilities are denoted by  $p_i = \mathbb{P}(Y_i = 1)$ ,  $i = 1, \dots, m$ .

- **Default or event correlation.** Noting that

$$\text{var}(Y_i) = \mathbb{E}(Y_i^2) - p_i^2 = \mathbb{E}(Y_i) - p_i^2 = p_i - p_i^2,$$

we obtain, for firms  $i$  and  $j$  with  $i \neq j$ , the formula

$$\rho(Y_i, Y_j) = \frac{\mathbb{E}(Y_i Y_j) - p_i p_j}{\sqrt{(p_i - p_i^2)(p_j - p_j^2)}}. \quad (66)$$

- Let the rv  $M := \sum_{i=1}^m Y_i$  denote the **number of defaulted obligors** at  $T$ .
- The actual loss if company  $i$  defaults is modelled by the random quantity  $\delta_i e_i$ , where  $e_i$  represents the overall exposure to company  $i$  and  $0 \leq \delta_i \leq 1$  represents the LGD.
- We denote the **overall portfolio loss** by  $L := \sum_{i=1}^m \delta_i e_i Y_i$ .
- It is possible to set up different credit risk models leading to the same multivariate distribution for  $\mathbf{R}$  or  $\mathbf{Y}$ . We call two models with state vectors  $\mathbf{R}$  and  $\tilde{\mathbf{R}}$  (or  $\mathbf{Y}$  and  $\tilde{\mathbf{Y}}$ ) **equivalent** if  $\mathbf{R} \stackrel{d}{=} \tilde{\mathbf{R}}$  (or  $\mathbf{Y} \stackrel{d}{=} \tilde{\mathbf{Y}}$ ).

## 11.1.2 Threshold models and copulas

### Definition 11.1

Let  $\mathbf{X} = (X_1, \dots, X_m)'$  be an  $m$ -dimensional random vector and let  $D \in \mathbb{R}^{m \times n}$  be a deterministic matrix with elements  $d_{ij}$  such that, for every  $i$ , the elements of the  $i$ th row form a set of increasing thresholds satisfying  $d_{i1} < \dots < d_{in}$ . Augment these thresholds by setting  $d_{i0} = -\infty$  and  $d_{i(n+1)} = \infty$  for all obligors and then set

$$R_i = j \iff d_{ij} < X_i \leq d_{i(j+1)}, \quad j \in \{0, \dots, n\}, \quad i \in \{1, \dots, m\}.$$

Then  $(\mathbf{X}, D)$  is said to define a threshold model for  $\mathbf{R} = (R_1, \dots, R_m)'$ .

- $\mathbf{X}$  are the **critical variables** and the  $i$ th row of  $D$  contains the **critical thresholds** for firm  $i$ .
- Default occurs if  $X_i \leq d_{i1}$  so that the default probability of company  $i$  is given by  $p_i = F_{X_i}(d_{i1})$ .



- When working with a default-only model we simply write  $d_i = d_{i1}$  and denote the threshold model by  $(\mathbf{X}, \mathbf{d})$ .
- **Default correlation and asset correlation.** It is important to distinguish the default correlation  $\rho(Y_i, Y_j)$  of two firms  $i \neq j$  from the correlation of the critical variables  $X_i$  and  $X_j$ .
- Since the critical variables are often interpreted in terms of asset values, the latter correlation is often referred to as **asset correlation**.
- For given default probabilities,  $\rho(Y_i, Y_j)$  is determined by  $\mathbb{E}(Y_i Y_j)$  according to (66), and in a threshold model  $\mathbb{E}(Y_i Y_j) = \mathbb{P}(X_i \leq d_{i1}, X_j \leq d_{j1})$ , so default correlation depends on the joint df of  $X_i$  and  $X_j$ .
- If  $\mathbf{X}$  is multivariate normal, as in many models used in practice, the correlation of  $X_i$  and  $X_j$  determines the **copula** of their joint distribution and hence the default correlation.
- If two threshold models lead to the same state/default probabilities and if the critical variables have the same copula, they are **equivalent**.

## 11.1.3 Gaussian threshold models

### Multivariate Merton model:

- Assume that the multivariate asset-value process  $\mathbf{V}_t = (V_{t,1}, \dots, V_{t,m})'$  follows an  $m$ -dimensional GBM with drift vector  $\boldsymbol{\mu}_V = (\mu_1, \dots, \mu_m)'$ , vector of volatilities  $\boldsymbol{\sigma}_V = (\sigma_1, \dots, \sigma_m)'$  and correlation matrix  $P$ .
- This means that  $(\mathbf{V}_t)$  solves the stochastic differential equations

$$dV_{t,i} = \mu_i V_{t,i} dt + \sigma_i V_{t,i} dW_{t,i}, \quad i = 1, \dots, m,$$

for correlated BMs with correlation  $\rho(W_{t,i}, W_{t,j}) = \rho_{ij}$ ,  $t \geq 0$ .

- For all  $i$  the asset value  $V_{T,i}$  is of the form

$$V_{T,i} = V_{0,i} \exp\left((\mu_i - \frac{1}{2}\sigma_i^2)T + \sigma_i W_{T,i}\right),$$

where  $\mathbf{W}_T \sim N_m(\mathbf{0}, TP)$ .

- In its basic form the Merton model is a default-only model where the firm defaults if  $V_{T,i} \leq B_i$  and  $B_i$  is the liability of firm  $i$ .

- Writing  $\mathbf{B} = (B_1, \dots, B_m)'$  the threshold model representation is  $(\mathbf{V}_T, \mathbf{B})$ .
- The multivariate Merton model is equivalent to the model  $(\mathbf{X}, \mathbf{d})$  with

$$X_i := \frac{\ln V_{T,i} - \ln V_{0,i} - (\mu_i - \frac{1}{2}\sigma_i^2)T}{\sigma_i\sqrt{T}},$$

$$d_i := \frac{\ln B_i - \ln V_{0,i} - (\mu_i - \frac{1}{2}\sigma_i^2)T}{\sigma_i\sqrt{T}}.$$

- The transformed variables satisfy  $\mathbf{X} \sim N_m(\mathbf{0}, P)$  and their copula is the Gauss copula  $C_P^{\text{Ga}}$ .

## Gaussian threshold models in practice

- In practice it is usual to start directly with threshold models of the form  $(\mathbf{X}, \mathbf{d})$  with  $\mathbf{X} \sim N_m(\mathbf{0}, P)$ .

- There are two practical challenges:
  - 1) calibration of  $d$  (or, in the case of a multi-state model, the threshold matrix  $D$ ) in line with exogenously given default and transition probabilities;
  - 2) calibration of the correlation matrix  $P$  in a **parsimonious** way.
- The problem of embedding state transition probabilities in a threshold matrix  $D$  has already been discussed. In a default-only model we set  $d_i = \Phi^{-1}(p_i)$  for  $i = 1, \dots, m$ .

## Factor models

- In its most general form  $P$  has  $m(m - 1)/2$  distinct parameters.
- $m$  is typically large and it is important to use a more parsimonious parametrization of this matrix based on a factor model.
- Factor models also lend themselves to **economic interpretation** and the factors are commonly interpreted as **country and industry effects**.

- We assume that

$$X_i = \sqrt{\beta_i} \tilde{F}_i + \sqrt{1 - \beta_i} \varepsilon_i, \quad (67)$$

where  $\tilde{F}_i$  and  $\varepsilon_1, \dots, \varepsilon_m$  are independent standard normal variables, and where  $0 \leq \beta_i \leq 1$  for all  $i$ .

- In this formulation  $\tilde{F}_i$  are the **systematic** variables, which are correlated, and  $\varepsilon_i$  are **idiosyncratic** variables.
- It follows that  $\beta_i$  can be viewed as a measure of the **systematic risk** of  $X_i$ : that is, the part of the variance of  $X_i$  which is explained by the systematic variable.
- The systematic variables are assumed to be of the form  $\tilde{F}_i = \mathbf{a}_i' \mathbf{F}$  where  $\mathbf{F}$  is a vector of common factors satisfying  $\mathbf{F} \sim N_p(\mathbf{0}, \Omega)$  with  $p < m$ , and where  $\Omega$  is a correlation matrix.
- These factors typically represent country and industry effects.
- The assumption that  $\text{var}(\tilde{F}_i) = 1$  means that  $\mathbf{a}_i' \Omega \mathbf{a}_i = 1$  for all  $i$ .

- Since  $\text{var}(X_i) = 1$  and since  $\tilde{F}_i$  and  $\varepsilon_1, \dots, \varepsilon_m$  are independent and standard normal, the asset correlations in this model are given by

$$\rho(X_i, X_j) = \text{cov}(X_i, X_j) = \sqrt{\beta_i \beta_j} \text{cov}(\tilde{F}_i, \tilde{F}_j) = \sqrt{\beta_i \beta_j} \mathbf{a}_i' \Omega \mathbf{a}_j.$$

- In order to set up the model we have to determine  $\mathbf{a}_i$  and  $\beta_i$  for each obligor and  $\Omega$ , with the additional constraint that  $\mathbf{a}_i' \Omega \mathbf{a}_i = 1$  for all  $i$ .
- Since  $\Omega$  has  $p(p-1)/2$  parameters, the loading vectors  $\mathbf{a}_i$  and coefficients  $\beta_i$  have collectively  $mp + m$  parameters, and we are applying  $m$  constraints, this gives  $mp + p(p-1)/2$  parameters.

## 11.2 Mixture models

### 11.2.1 Bernoulli mixture models

- In a mixture model the default risk of an obligor is assumed to depend on a set of **common factors**, usually interpreted as macroeconomic variables, which are also modelled stochastically.
- Given a realization of the factors, defaults of individual firms are assumed to be independent.
- Dependence between defaults stems from the dependence of individual default probabilities on the set of common factors.
- Bernoulli mixture models provide a way of capturing the dependence between Bernoulli events (i.e. defaults/non-defaults).
- They can be extended to multinomial mixture models to capture dependent migrations in a rating system.

### Definition 11.2 (Bernoulli mixture model)

Given some  $p < m$  and a  $p$ -dimensional random vector  $\Psi = (\Psi_1, \dots, \Psi_p)'$ , the default indicator vector  $\mathbf{Y}$  follows a Bernoulli mixture model with factor vector  $\Psi$  if there are functions  $p_i : \mathbb{R}^p \rightarrow (0, 1)$ , such that conditional on  $\Psi$  the components of  $\mathbf{Y}$  are independent Bernoulli rvs with  $\mathbb{P}(Y_i = 1 \mid \Psi = \psi) = p_i(\psi)$ .

The **conditional independence given factors** makes these models relatively easy to analyse. For  $\mathbf{y} = (y_1, \dots, y_m)'$  in  $\{0, 1\}^m$  we get

$$\mathbb{P}(\mathbf{Y} = \mathbf{y} \mid \Psi = \psi) = \prod_{i=1}^m p_i(\psi)^{y_i} (1 - p_i(\psi))^{1-y_i}$$
$$\mathbb{P}(\mathbf{Y} = \mathbf{y}) = \int_{\mathbb{R}^p} \prod_{i=1}^m p_i(\psi)^{y_i} (1 - p_i(\psi))^{1-y_i} g(\psi) d\psi,$$

where  $g(\psi)$  is the probability density of the factors. The default probabilities are given by  $p_i = \mathbb{E}(Y_i = 1) = \mathbb{E}(p_i(\Psi))$ .



- Consider the portfolio loss  $L = \sum_{i=1}^m e_i \delta_i Y_i$  in the case where the exposures  $e_i$  and LGDs  $\delta_i$  are deterministic.
- It is difficult to compute the df  $F_L$  of  $L$ .
- However, it is easy to use the conditional independence of the defaults to show that the *Laplace–Stieltjes transform* of  $F_L$  is for  $t \in \mathbb{R}$  given by

$$\begin{aligned}\hat{F}_L(t) &= \mathbb{E}(e^{-tL}) = \mathbb{E}\left(E(e^{-t \sum_{i=1}^m e_i \delta_i Y_i} \mid \Psi)\right) \\ &= \mathbb{E}\left(\prod_{i=1}^m \mathbb{E}(e^{-te_i \delta_i Y_i} \mid \Psi)\right) \\ &= \mathbb{E}\left(\prod_{i=1}^m (p_i(\Psi)e^{-te_i \delta_i} + 1 - p_i(\Psi))\right)\end{aligned}$$

which can be obtained by integrating over distribution of factors  $\Psi$ .

- This is useful for: sampling losses from model with **importance sampling**; approximating probability mass function using **Fourier inversion**.

## 11.2.2 Threshold models as mixture models

- Although the mixture models of this section seem, at first glance, to be different in structure to the threshold models, it is important to realize that the majority of useful threshold models, including all the examples we have given, **can be represented as Bernoulli mixture models**.
- In a threshold model default occurs for counterparty  $i$  if a critical variable  $X_i$  lies below a critical threshold  $d_i$ .
- Moreover  $X_i$  follows a **linear factor model**

$$X_i = \sqrt{\beta_i} \mathbf{a}_i' \mathbf{F} + \sqrt{1 - \beta_i} \varepsilon_i, \quad \text{where}$$

- ▶  $\mathbf{F} \sim N_p(\mathbf{0}, \Omega)$  is a random vector of normally distributed common economic factors;
- ▶  $0 \leq \beta_i \leq 1$  and  $\text{var}(\mathbf{a}_i' \mathbf{F}) = 1$ ;
- ▶  $\varepsilon_1, \dots, \varepsilon_m$  are iid standard normal and are also independent of  $\mathbf{F}$ .

- We will write the Gaussian threshold model as a Bernoulli mixture model with factor vector  $\Psi = -F$ . (This makes the conditional default probabilities **increasing** in the factors for positive  $a_i$ .)
- Conditioning on  $\Psi = -F$ , the vector  $X$  is multivariate normally distributed with a diagonal covariance matrix and therefore the components of  $X$  are **conditionally independent**.
- The conditional default probabilities are

$$\begin{aligned}
 p_i(\psi) &= \mathbb{P}(Y_i = 1 \mid \Psi = \psi) = \mathbb{P}(X_i \leq d_i \mid \Psi = \psi) \\
 &= \mathbb{P}(X_i \leq d_i \mid F = -\psi) \\
 &= \mathbb{P}(\sqrt{1 - \beta_i} \varepsilon_i \leq d_i + \sqrt{\beta_i} \mathbf{a}_i' \psi) \\
 &= \Phi \left( \frac{d_i + \sqrt{\beta_i} \mathbf{a}_i' \psi}{\sqrt{1 - \beta_i}} \right) \\
 &= \Phi \left( \frac{\Phi^{-1}(p_i) + \sqrt{\beta_i} \mathbf{a}_i' \psi}{\sqrt{1 - \beta_i}} \right).
 \end{aligned}$$

### 11.2.3 Poisson mixture models and CreditRisk+

- Since default is typically a rare event, it is possible to approximate Bernoulli indicator rvs for default with Poisson rvs and Bernoulli mixture models with Poisson mixture models.
- By choosing independent gamma distributions for the economic factors  $\Psi$ , we obtain a tractable model known as CreditRisk+, proposed by Credit Suisse Financial Products in 1997.
- Assume that, given the factors  $\Psi$ , the default indicators  $Y_1, \dots, Y_m$  for a particular time horizon are conditionally independent Bernoulli variables satisfying  $\mathbb{P}(Y_i = 1 \mid \Psi = \psi) = p_i(\psi)$ .
- Moreover assume that the distribution of  $\Psi$  is such that the conditional default probabilities  $p_i(\psi)$  tend to be very small.
- The  $Y_i$  variables can be approximated by conditionally independent Poisson variables  $\tilde{Y}_i$  satisfying  $\tilde{Y}_i \mid \Psi = \psi \sim \text{Poi}(p_i(\psi))$ .

- This follows because

$$\mathbb{P}(\tilde{Y}_i = 0 \mid \Psi = \psi) = e^{-p_i(\psi)} \approx 1 - p_i(\psi),$$

$$\mathbb{P}(\tilde{Y}_i = 1 \mid \Psi = \psi) = p_i(\psi)e^{-p_i(\psi)} \approx p_i(\psi).$$

- The portfolio loss  $L = \sum_{i=1}^m e_i \delta_i Y_i$  can be approximated by  $\tilde{L} = \sum_{i=1}^m e_i \delta_i \tilde{Y}_i$ .
- It is possible for a company to “default more than once” in the approximating Poisson model, albeit with a very low probability.
- In **CreditRisk+** the parameter  $\lambda_i(\Psi)$  of the conditional Poisson distribution for firm  $i$  is assumed to take the form

$$\lambda_i(\Psi) = k_i \mathbf{w}_i' \Psi \quad (68)$$

for  $k_i > 0$ , non-negative weights  $\mathbf{w}_i = (w_{i1}, \dots, w_{ip})'$  satisfying  $\sum_j w_{ij} = 1$ , and  $p$  independent  $\text{Ga}(\alpha_j, \beta_j)$ -distributed factors  $\Psi_1, \dots, \Psi_p$ .

- The parameters are set to be  $\alpha_j = \beta_j = \sigma_j^{-2}$  for  $\sigma_j > 0$  and  $j = 1, \dots, p$ .

- This parametrization of the gamma variables ensures that we have  $\mathbb{E}(\Psi_j) = 1$  and  $\text{var}(\Psi_j) = \sigma_j^2$ .
- It is easy to verify that

$$\mathbb{E}(\tilde{Y}_i) = \mathbb{E}(\mathbb{E}(\tilde{Y}_i | \boldsymbol{\Psi})) = \mathbb{E}(\lambda_i(\boldsymbol{\Psi})) = k_i \mathbb{E}(\mathbf{w}'_i \boldsymbol{\Psi}) = k_i,$$

so  $k_i$  is the expected number of defaults for obligor  $i$  in the time period.

- The assumptions in **CreditRisk+** make it possible to compute the distribution of the number of defaults and the aggregate portfolio loss fairly explicitly using techniques for compound distributions and mixture distributions that are well known in actuarial mathematics.

## Distribution of the number of defaults

In **CreditRisk+** we have that given  $\boldsymbol{\Psi} = \boldsymbol{\psi}$ ,  $\tilde{Y}_i \sim \text{Poi}(k_i \mathbf{w}'_i \boldsymbol{\psi})$ , which implies that the distribution of the number of defaults  $\tilde{M} := \sum_{i=1}^m \tilde{Y}_i$  satisfies

$$\tilde{M} \mid \boldsymbol{\Psi} = \boldsymbol{\psi} \sim \text{Poi}\left(\sum_{i=1}^m k_i \mathbf{w}'_i \boldsymbol{\psi}\right). \quad (69)$$

- This uses the fact that the **sum of independent Poisson variables is also Poisson** with a rate parameter given by the sum of the rate parameters
- To compute the unconditional distribution of  $\tilde{M}$  we require a well-known result on **mixed Poisson distributions**.

### Proposition 11.3

If the rv  $N$  is conditionally Poisson with a gamma-distributed rate parameter  $\Lambda \sim \text{Ga}(\alpha, \beta)$ , then  $N$  has a **negative binomial distribution**,  $N \sim \text{NB}(\alpha, \beta/(\beta + 1))$ .

In the case when  $p = 1$  we may apply this result directly to (69) to deduce that  $\tilde{M}$  has a negative binomial distribution. The general result is:

### Proposition 11.4

$\tilde{M}$  is distributed as a **sum of  $p$  independent negative binomial rvs**.

## Distribution of the aggregate loss

- To obtain a tractable model, exposures are discretized in CreditRisk+ using the concept of exposure bands.
- The LGD is subsumed in the exposure by multiplying the actual exposure by a typical value for the LGD for an obligor with the same credit rating.
- The losses arising from the individual obligors are of the form  $\tilde{L}_i = e_i \tilde{Y}_i$  where the  $e_i$  are known (LGD-adjusted) exposures.
- For all  $i$ , the exposure  $e_i$  is discretized in units of an amount  $\epsilon$  so that  $e_i$  is replaced by a value  $\ell_i \epsilon \geq e_i$  where  $\ell_i$  is a positive integer multiplier.
- Exposure bands  $b = 1, \dots, n$  are defined corresponding to the distinct values  $\ell^{(1)}, \dots, \ell^{(n)}$  for the multipliers so that obligors are grouped in exposure bands according to the values of their discretized exposures.
- It is then possible to derive the distribution of the aggregate loss 
$$\tilde{L} = \sum_{i=1}^m \ell_i \epsilon \tilde{Y}_i.$$



## Theorem 11.5

Let  $\tilde{L}$  represent the aggregate loss in the general  $p$ -factor CreditRisk+ model with exposures discretized into exposure bands as described above. Then the following hold.

- i) The **Laplace–Stieltjes transform** of the df of  $\tilde{L}$  is given by

$$\hat{F}_{\tilde{L}}(s) = \prod_{j=1}^p \left( 1 + \sigma_j^2 \sum_{i=1}^m k_i w_{ij} \left( 1 - \sum_{b=1}^n e^{-s\ell^{(b)}} q_{jb} \right) \right)^{-\sigma_j^{-2}}, \quad (70)$$

where  $q_{jb} = \sum_{i \in s_b} k_i w_{ij} / \sum_{i=1}^m k_i w_{ij}$  for  $b = 1, \dots, n$ .

- ii) The distribution of  $\tilde{L}$  has the structure  $\tilde{L} \stackrel{d}{=} \sum_{j=1}^p Z_j$  where the  $Z_j$  are independent variables that follow a **compound negative binomial distribution**. More precisely, it holds that  $Z_j \sim \text{CNB}(\sigma_j^{-2}, \theta_j, G_{X_j})$  with  $\theta_j = (1 + \sigma_j^2 \sum_{i=1}^m k_i w_{ij})^{-1}$  and  $G_{X_j}$  the df of a **multinomial random variable**  $X_j$  taking the value  $\ell^{(b)}$  with probability  $q_{jb}$ .

## 11.3 Statistical inference for portfolio credit models

### 11.3.1 Industry factor models

- Recall that portfolio models in industry often take the form of a **Gaussian threshold model**  $(\mathbf{X}, \mathbf{d})$  with  $\mathbf{X} \sim N_m(\mathbf{0}, P)$ , where the random vector  $\mathbf{X}$  contains the critical variables, the deterministic vector  $\mathbf{d}$  contains the critical default thresholds and  $P$  is the so-called asset correlation matrix, which is estimated with the help of a factor model for  $\mathbf{X}$ .
- Industry models generally **separate the calibration of the vector  $\mathbf{d}$**  (or the threshold matrix  $D$  in a multi-state model) and the calibration of **the factor model for  $\mathbf{X}$** .
- In a default-only model the threshold  $d_i$  is usually set at  $d_i = \Phi^{-1}(p_i)$  where  $p_i$  is an estimate of the default probability for obligor  $i$  for the time period in question (generally one year).

- The default probability may be estimated in different ways: for larger corporates it may be estimated using credit ratings or using a firm-value approach, such as the Moody's public-firm EDF model; for retail obligors it may be estimated on the basis of credit scores.
- Recall that the factor model for  $\mathbf{X}$  takes the form

$$X_i = \sqrt{\beta_i} \tilde{F}_i + \sqrt{1 - \beta_i} \varepsilon_i, \quad i = 1, \dots, m, \quad (71)$$

where  $\tilde{F}_i$  and  $\varepsilon_1, \dots, \varepsilon_m$  are independent standard normal variables, and where  $0 \leq \beta_i \leq 1$  for all  $i$ .

- The systematic variables  $\tilde{F}_i$  are assumed to be of the form  $\tilde{F}_i = \mathbf{a}'_i \mathbf{F}$  where  $\mathbf{F}$  is a vector of common factors satisfying  $\mathbf{F} \sim N_p(\mathbf{0}, \Omega)$  with  $p < m$ , and where  $\Omega$  is a correlation matrix.
- The factors typically represent country and industry effects.
- The assumption that  $\text{var}(\tilde{F}_i) = 1$  implies that  $\mathbf{a}'_i \Omega \mathbf{a}_i = 1$  for all  $i$ .

- Different industry models use different **data for  $X$**  to calibrate the factor model (71).
- The **Moody's Analytics Global Correlation or GCorr model** has sub-models for many different kinds of obligor including public corporate firms, private firms, small and medium enterprises (SMEs), retail customers and sovereigns. Huang et al. (2012)
- The sub-model for public firms (GCorr Corporate) is calibrated using data on **weekly asset value returns**, where asset values are determined as part of the public-firm EDF methodology.
- In the **CreditMetrics** framework **weekly equity returns** are viewed as a **proxy for asset returns** and used to estimate the factor model.
- We sketch a generic procedure for estimating a factor model for corporates where the factors have country and industry-sector interpretations.

## Estimating a credit risk factor model

- We assume that we have a **high-dimensional multivariate time series**  $(\mathbf{X}_t)_{1 \leq t \leq n}$  of asset returns (or other proxy data for changing credit quality) over a period of time in which **stationarity** can be assumed.
  - We also assume that each component time series has been **scaled to have mean zero and variance one**.
- 1) We first fix the structure of the factor vector  $\mathbf{F}$  so that, for example, the first block of components might represent country factors and the second block of components might represent industry factors. We then assign vectors of factor weights  $\mathbf{a}_i$  to each obligor based on our knowledge of the companies. The elements of  $\mathbf{a}_i$  may simply consist of ones and zeros if the company can be clearly identified with a single country and industry, but may also consist of weights if the company has significant activity in more than one country or sector.

2) We then use **cross-sectional estimation techniques** to estimate the factor values  $\mathbf{F}_t$  at each time point  $t$ . Effectively the factor estimates  $\hat{\mathbf{F}}_t$  are constructed as weighted sums of the  $X_{t,i}$  data for obligors  $i$  that are exposed to each factor. One way of achieving this is to construct a matrix  $A$  with rows  $\mathbf{a}_i$  and then to estimate a **fundamental factor model** of the form  $\mathbf{X}_t = A\mathbf{F}_t + \boldsymbol{\varepsilon}_t$  at each time point  $t$ .

- We have a regression model

$$\mathbf{X}_t = A\mathbf{F}_t + \boldsymbol{\varepsilon}_t, \quad (72)$$

where  $\mathbf{X}_t \in \mathbb{R}^m$  are the return data,  $A \in \mathbb{R}^{m \times p}$  is a known matrix of factor loadings,  $\mathbf{F}_t \in \mathbb{R}^p$  are the factors to be estimated and  $\boldsymbol{\varepsilon}_t$  are errors with diagonal covariance matrix  $\Upsilon$ .

- Note that the components of the error vector  $\boldsymbol{\varepsilon}_t$  can not generally be assumed to have equal variance, so that (72) is a regression problem with so-called **heteroskedastic errors**.

- Unbiased estimators of the factors  $\mathbf{F}_t$  may be obtained by forming the **ordinary least squares (OLS)** estimates

$$\hat{\mathbf{F}}_t^{\text{OLS}} = (\mathbf{A}'\mathbf{A})^{-1}\mathbf{A}'\mathbf{X}_t.$$

- Since the errors are heteroskedastic, slightly more efficient estimators can be obtained by using the method of **generalized least squares (GLS)**.
- 3) The raw factor estimates form a multivariate time series of dimension  $p$ . We standardize each component series to have mean zero and variance one to obtain  $(\hat{\mathbf{F}}_t)_{1 \leq t \leq n}$  and calculate the sample covariance matrix of the standardized factor estimates, which serves as our estimate of  $\Omega$ .
  - 4) We then scale the vectors of factor weights  $\mathbf{a}_i$  so that the conditions  $\mathbf{a}_i'\hat{\Omega}\mathbf{a}_i = 1$  are met for each obligor.
  - 5) Time series of estimated systematic variables for each obligor are then constructed by calculating  $\hat{F}_{t,i} = \mathbf{a}_i'\hat{\mathbf{F}}_t$  for  $t = 1, \dots, n$ .

6) Finally we estimate the  $\beta_i$  parameters by performing a time series regression of  $X_{t,i}$  on  $\hat{F}_{t,i}$  for each obligor.

Note that the accurate estimation of the  $\beta_i$  in the last step is particularly important (as it effects tail behaviour). The estimate of  $\beta_i$  is the so-called R-squared of the time series regression model in Step 6 and will be largest for the firms whose credit-quality changes are best explained by systematic factors.



# 14 Multivariate time series

14.1 Fundamentals of multivariate time series

14.2 Multivariate GARCH Processes

# 14.1 Fundamentals of multivariate time series

## 14.1.1 Basic definitions

### Definition 14.1

The mean function  $\boldsymbol{\mu}(t)$  and the covariance matrix function  $\Gamma(t+h, t)$  of  $(\mathbf{X}_t)_{t \in \mathbb{Z}}$  are given by

$$\begin{aligned}\boldsymbol{\mu}(t) &= E(\mathbf{X}_t), & t \in \mathbb{Z}, \\ \Gamma(t+h, t) &= E((\mathbf{X}_{t+h} - \boldsymbol{\mu}(t+h))(\mathbf{X}_t - \boldsymbol{\mu}(t))'), & t, h \in \mathbb{Z}.\end{aligned}$$

- Analogously to the univariate case, we have  $\Gamma(t, t) = \text{cov}(\mathbf{X}_t)$ . By observing that the elements  $\gamma_{ij}(t+h, t)$  of  $\Gamma(t+h, t)$  satisfy

$$\gamma_{ij}(t+h, t) = \text{cov}(X_{t+h,i}, X_{t,j}) = \text{cov}(X_{t,j}, X_{t+h,i}) = \gamma_{ji}(t, t+h),$$

it is clear that  $\Gamma(t+h, t) = \Gamma(t, t+h)'$  for all  $t, h$ .

- However, the matrix  $\Gamma$  **need not be symmetric**, so in general  $\Gamma(t+h, t) \neq \Gamma(t, t+h)$ . **One series can lead other series.**

### Definition 14.2 (strict stationarity)

The multivariate time series  $(\mathbf{X}_t)_{t \in \mathbb{Z}}$  is **strictly stationary** if

$$(\mathbf{X}'_{t_1}, \dots, \mathbf{X}'_{t_n}) \stackrel{d}{=} (\mathbf{X}'_{t_1+k}, \dots, \mathbf{X}'_{t_n+k}),$$

for all  $t_1, \dots, t_n, k \in \mathbb{Z}$  and for all  $n \in \mathbb{N}$ .

### Definition 14.3 (covariance (weak, second-order) stationarity)

The multivariate time series  $(\mathbf{X}_t)_{t \in \mathbb{Z}}$  is **covariance stationary** if the first **two moments exist** and satisfy

$$\begin{aligned}\boldsymbol{\mu}(t) &= \boldsymbol{\mu}, & t \in \mathbb{Z}, \\ \Gamma(t+h, t) &= \Gamma(h, 0), & t, h \in \mathbb{Z}.\end{aligned}$$

- For a covariance-stationary process we write  $\Gamma(h) := \Gamma(h, 0)$ .
- Note that  $\Gamma(0) = \text{cov}(\mathbf{X}_t)$ , for all  $t$ .

- Write  $\Delta$  for the diagonal matrix whose entries are the square roots of the diagonal entries of  $\Gamma(0)$  (standard deviations of component series).

**Definition 14.4 (correlation matrix function)**

The correlation matrix function  $P(h)$  of a covariance-stationary multivariate time series is

$$P(h) = \Delta^{-1}\Gamma(h)\Delta^{-1}, \quad \forall h \in \mathbb{Z}. \quad (73)$$

- The diagonal entries  $\rho_{ii}(h)$  of this matrix-valued function give the autocorrelation function of the  $i$ th component series  $(X_{t,i})_{t \in \mathbb{Z}}$ .
- The off-diagonal entries give so-called cross-correlations between different component series at different times.

**Definition 14.5 (multivariate white noise)**

$(\mathbf{X}_t)_{t \in \mathbb{Z}}$  is multivariate white noise if it is covariance stationary with correlation matrix function given by

$$P(h) = \begin{cases} P, & h = 0, \\ 0, & h \neq 0, \end{cases}$$

for some positive-definite correlation matrix  $P$ .

Such a process has **no cross-correlation between component series**, except for contemporaneous cross-correlation at lag zero.

**Definition 14.6 (multivariate strict white noise)**

$(\mathbf{X}_t)_{t \in \mathbb{Z}}$  is multivariate strict white noise if it is a series of **iid random vectors**.

A strict white noise process with mean zero and covariance matrix  $\Sigma$  will be denoted  $\text{SWN}(\mathbf{0}, \Sigma)$ .

## 14.1.2 Analysis in the time domain

- Assume we have a random sample  $\mathbf{X}_1, \dots, \mathbf{X}_n$  from a covariance-stationary multivariate time series model  $(\mathbf{X}_t)_{t \in \mathbb{Z}}$ .
- In the time domain we construct empirical estimators of the covariance matrix function and the correlation matrix function.
- The *sample covariance matrix function* is calculated according to

$$\hat{\Gamma}(h) = \frac{1}{n} \sum_{t=1}^{n-h} (\mathbf{X}_{t+h} - \bar{\mathbf{X}})(\mathbf{X}_t - \bar{\mathbf{X}})', \quad 0 \leq h < n,$$


where  $\bar{\mathbf{X}} = \sum_{t=1}^n \mathbf{X}_t / n$  is the sample mean.

- Writing  $\hat{\Delta}$ , for the diagonal matrix of sample standard deviations (square root of the diagonal of  $\hat{\Gamma}(0)$ ) the *sample correlation matrix function* is

$$\hat{P}(h) = \hat{\Delta}^{-1} \hat{\Gamma}(h) \hat{\Delta}^{-1}, \quad 0 \leq h < n.$$

- The information contained in the elements  $\hat{\rho}_{ij}(h)$  of the sample correlation matrix function is generally displayed in the *cross-correlogram*.

# Cross-correlogram of index returns



The **US market** leads Europe and Japan.

### 14.1.3 Multivariate ARMA processes

- ARMA models extend to higher dimensions where they are called VARMA. They provide [models for the conditional mean vector](#).
- The VAR class is most widely used in practice.
- The first-order VAR process satisfies the set of equations

$$\mathbf{X}_t = \Phi \mathbf{X}_{t-1} + \boldsymbol{\varepsilon}_t, \quad \forall t. \quad (74)$$

where  $\Phi \in \mathbb{R}^{d \times d}$  is a matrix and  $(\boldsymbol{\varepsilon}_t)$  is a white noise process.

- The process is covariance stationary if and only if all eigenvalues of the matrix  $\Phi$  are less than one in absolute value.
- The covariance matrix function of this process is

$$\Gamma(h) = \Phi^h \Gamma(0), \quad h = 0, 1, 2, \dots$$



## 14.2 Multivariate GARCH Processes

Recall that the Cholesky factor  $A$  of a positive-definite matrix  $\Sigma$  is the lower-triangular matrix satisfying  $AA' = \Sigma$ .

### Definition 14.7

Let  $(\mathbf{Z}_t)_{t \in \mathbb{Z}}$  be  $\text{SWN}(\mathbf{0}, I_d)$ . The process  $(\mathbf{X}_t)_{t \in \mathbb{Z}}$  is said to be a multivariate GARCH process if it is strictly stationary and satisfies equations of the form

$$\mathbf{X}_t = A_t \mathbf{Z}_t, \quad t \in \mathbb{Z}, \quad (75)$$

where  $A_t \in \mathbb{R}^{d \times d}$  is the Cholesky factor of a positive-definite matrix  $\Sigma_t$  which is measurable with respect to  $\mathcal{F}_{t-1} = \sigma(\{\mathbf{X}_s : s \leq t-1\})$ , the history of the process up to time  $t-1$ .

### Conditional moments:

$$\blacksquare \quad \mathbb{E}(\mathbf{X}_t \mid \mathcal{F}_{t-1}) = \mathbf{0}$$

- $\text{cov}(\mathbf{X}_t \mid \mathcal{F}_{t-1}) = A_t A_t' = \Sigma_t$  is the **conditional covariance matrix**.
- Could add a non-zero conditional mean term  $\boldsymbol{\mu}_t$  so that  $\mathbf{X}_t = \boldsymbol{\mu}_t + A_t \mathbf{Z}_t$  where, for example,
  - ▶  $\boldsymbol{\mu}_t = \boldsymbol{\mu}$  for a constant conditional mean;
  - ▶ or  $\boldsymbol{\mu}_t$  could follow a VARMA specification, such as  $\boldsymbol{\mu}_t = \Phi \mathbf{X}_{t-1}$ .
- We can write  $\Sigma_t = \Delta_t P_t \Delta_t$ , where  $\Delta_t$  is the diagonal **volatility matrix** and  $P_t$  is the **conditional correlation matrix**.
- The art of building multivariate GARCH models is to specify the dependence of  $\Sigma_t$  (or of  $\Delta_t$  and  $P_t$ ) on the past in such a way that  $\Sigma_t$  always remains **symmetric and positive definite**.
- The innovations are generally taken to be from either a multivariate Gaussian distribution ( $\mathbf{Z}_t \sim N_d(\mathbf{0}, I_d)$ ) or an appropriately scaled **spherical** multivariate  $t$  distribution ( $\mathbf{Z}_t \sim t_d(\nu, \mathbf{0}, (\nu - 2)I_d/\nu)$ ). Any distribution with mean zero and covariance matrix  $I_d$  is permissible.

## 14.2.1 Models for conditional correlation

### Definition 14.8

The process  $(\mathbf{X}_t)_{t \in \mathbb{Z}}$  is a CCC-GARCH process if it is a multivariate GARCH process with conditional covariance matrix of the form  $\Sigma_t = \Delta_t P_c \Delta_t$ , where

- $P_c$  is a constant, positive-definite correlation matrix; and
- $\Delta_t$  is a diagonal volatility matrix with elements  $\sigma_{t,k}$  satisfying

$$\sigma_{t,k}^2 = \alpha_{k0} + \sum_{i=1}^{p_k} \alpha_{ki} X_{t-i,k}^2 + \sum_{j=1}^{q_k} \beta_{kj} \sigma_{t-j,k}^2, \quad k = 1, \dots, d, \quad (76)$$

where  $\alpha_{k0} > 0$ ,  $\alpha_{ki} \geq 0$ ,  $i = 1, \dots, p_k$ ,  $\beta_{kj} \geq 0$ ,  $j = 1, \dots, q_k$ .

- Alternatives to ordinary GARCH( $p_k, q_k$ ) model may of course be used.
- In the CCC GARCH model the process  $\mathbf{Y}_t = \Delta_t^{-1} \mathbf{X}_t$  (known as the **de-volatilized process**) satisfies  $(\mathbf{Y}_t)_{t \in \mathbb{Z}} \sim \text{SWN}(\mathbf{0}, P_c)$ .

- Estimation can be accomplished in **two stages**:
  - 1) Fit univariate GARCH models to each component series;
  - 2) Form residuals  $\hat{\mathbf{Y}}_t = \hat{\Delta}_t^{-1} \mathbf{X}_t$ , for  $t = 1, \dots, n$  and estimate  $P_c$  (either by using the standard correlation estimator or by fitting an appropriate distribution).
- Alternatively all parameters can be maximized in **one step**.
- The CCC model is often a **useful starting point** from which to proceed to more complex models.
- In some empirical settings it gives an adequate performance, but it is generally considered that the constancy of conditional correlation in this model is **an unrealistic feature** and that the impact of news on financial markets requires models that allow a **dynamic evolution of conditional correlation as well as a dynamic evolution of volatilities**.

## Definition 14.9

The process  $(\mathbf{X}_t)_{t \in \mathbb{Z}}$  is a DCC-GARCH process if it is a multivariate GARCH process where the volatilities comprising  $\Delta_t$  follow univariate GARCH specifications as in (76) and the conditional correlation matrices  $P_t$  satisfy, for  $t \in \mathbb{Z}$ , the equations

$$P_t = \wp \left( \left( 1 - \sum_{i=1}^p \alpha_i - \sum_{j=1}^q \beta_j \right) P_c + \sum_{i=1}^p \alpha_i \mathbf{Y}_{t-i} \mathbf{Y}_{t-i}' + \sum_{j=1}^q \beta_j P_{t-j} \right), \quad (77)$$

where

- $P_c$  is a positive-definite correlation matrix,
- $\wp$  is the operator that extracts correlation matrices from covariance matrices,
- $\mathbf{Y}_t = \Delta_t^{-1} \mathbf{X}_t$  denotes the devolatized process,
- and the coefficients satisfy  $\alpha_i \geq 0$ ,  $\beta_j \geq 0$  and  $\sum_{i=1}^p \alpha_i + \sum_{j=1}^q \beta_j < 1$ .

- If all the  $\alpha_i$  and  $\beta_j$  coefficients (77) are zero, model reduces to CCC.
- In a covariance-stationary **univariate GARCH model** with unconditional variance  $\sigma^2$ , the volatility equation can be written

$$\sigma_t^2 = \left(1 - \sum_{i=1}^p \alpha_i - \sum_{j=1}^q \beta_j\right) \sigma^2 + \sum_{i=1}^p \alpha_i X_{t-i} + \sum_{j=1}^q \beta_j \sigma_{t-j}^2.$$

- Thus, in DCC, the correlation matrix  $P_c$  in (77) can be thought of as representing the **long-run correlation structure**.
- The usual estimation method for the DCC model is as follows.
  - 1) Fit **univariate GARCH-type models** to the component series to estimate the volatility matrix  $\Delta_t$ . Form an estimated realization of the devolatilized process by taking  $\hat{\mathbf{Y}}_t = \hat{\Delta}_t^{-1} \mathbf{X}_t$ .
  - 2) Estimate  $P_c$  by estimating correlation matrix of the devolatilized data.
  - 3) Estimate the remaining parameters  $\alpha_i$  and  $\beta_j$  in equation (77) by fitting the implied dynamic model to the devolatilized data ( $\hat{\mathbf{Y}}_t$ ).

Consider a **first-order model** ( $p = q = 1$ ):

- Given  $\mathcal{F}_{t-1}$  (comprising  $\mathbf{Y}_{t-k}, P_{t-k}, k = 1, 2, \dots$ ) and using an estimate of  $P_c$  (known as **variance targeting**), we have

$$\mathbf{Y}_t = B_t \mathbf{Z}_t, \quad \text{where}$$

$$B_t B_t' = P_t, \quad (\text{Cholesky decomposition})$$

$$P_t = \wp(Q_t), \quad (\text{correlation from covariance})$$

$$Q_t = (1 - \alpha_1 - \beta_1)P_c + \alpha_1 \mathbf{Y}_{t-1} \mathbf{Y}_{t-1}' + \beta_1 P_{t-1}$$

- Usually estimated by **conditional maximum likelihood**.
- The likelihood is built up recursively from **starting values** (for example  $P_0 = \mathbf{Y}_0 \mathbf{Y}_0' = P_c$ ).
- There are two parameters to estimate for dynamics, in addition to parameters of innovation distribution (if non-Gaussian).

## Relationship to dynamic copula models:

- In terms of **copulas**, using a Gaussian innovation distribution means estimating a 2-parameter model where

$$\mathbf{Y}_t \mid \mathcal{F}_{t-1} \sim C_{P_t}^{\text{Ga}}(\Phi, \dots, \Phi).$$

- Using a Student innovation distribution means estimating a 3-parameter model where

$$\mathbf{Y}_t \mid \mathcal{F}_{t-1} \sim C_{\nu, P_t}^{\text{t}}(F_{\nu}, \dots, F_{\nu})$$

where  $F_{\nu}$  is a scaled Student  $t$  distribution.

## Copula-MGARCH models

- Note that models of the form

$$\mathbf{Y}_t \mid \mathcal{F}_{t-1} \sim C_{\nu, P_t^*}^{\text{t}}(F_{\nu_1}, \dots, F_{\nu_d})$$

with  $P_t^*$  updating as in (77) have also been considered.

This has  $d + 3$  parameters.



- Previous model **doesn't quite fit into the DCC class** as we have defined it because  $\text{cov}(\mathbf{Y}_t \mid \mathcal{F}_{t-1}) = P_t \neq P_t^*$ .
- It is **not the parameters of the conditional correlation matrix** but rather the **parameters of the copula that update according to (77)**.
- However it fits into a bigger class of **copula-MGARCH models** where

$$\mathbf{X}_t = \boldsymbol{\mu}_t + \Delta_t \mathbf{Y}_t, \quad \mathbf{Y}_t \mid \mathcal{F}_{t-1} \sim C_t(F_1, \dots, F_d)$$

and

- ▶ the volatility components of  $\Delta_t$  follow GARCH schemes;
  - ▶ the conditional mean terms  $\boldsymbol{\mu}_t$  follow VARMA schemes;
  - ▶ the conditional copula  $C_t$  evolves as function of information in  $\mathcal{F}_{t-1}$ ;
  - ▶  $F_1, \dots, F_d$  are zero-mean, unit-variance distributions.
- See Patton (2006), Patton (2012), and Fan and Patton (2014)

## 14.2.2 Dimension reduction in MGARCH

- While the multi-stage estimation procedure for DCC makes it possible to estimate in quite high dimensions, it is usual to first apply **dimension reduction through factor modelling** and then fit MGARCH models to the **most important factors**.
- Can easily fit MGARCH models to factors derived from **macroeconomic** and **fundamental** factor models. The factors are typically correlated.
- The use of so-called **PC-GARCH (principal components GARCH)** is quite popular and **avoids need for multivariate models**.
  - ▶ Here we assume that the principal components of  $X_t$  follow a CCC model with  $P = I_d$ .
  - ▶ To estimate such a model, we estimate the principal components of the data and fit **univariate GARCH models** to each principal components series.

# 17 Introduction to counterparty risk

## 17.1 Introduction

## 17.2 Credit value adjustments

## 17.1 Introduction

- A substantial part of all derivative transactions is carried out over the counter and there is no central clearing counterparty to guarantee fulfilment of the contractual obligations.
- These trades are subject to the risk that a contracting party defaults during the transaction, thus affecting the cash flows that are actually received by the other party. This is known as **counterparty credit risk**.
- Counterparty risk received a lot of attention during the financial crisis of 2007-2009 as some of the institutions heavily involved in derivative transactions experienced worsening credit quality or—in the case of Lehman Brothers—even a default event.
- **Counterparty risk management** is now a key issue for all financial institutions and the focus of many new regulatory developments.

## Example of Interest-Rate Swap

- Two parties  $A$  and  $B$  agree to exchange a series of interest payments on a given nominal amount of money for a given period.
- $A$  receives payments at a fixed interest rate and makes floating payments at a rate equal to the three-month LIBOR rate.
- Suppose that  $A$  defaults at time  $\tau_A$  before the maturity of the contract.
- If interest rates have risen relative to their value at inception of contract:
  - ▶ The fixed interest payments have decreased in value and the value of the contract has increased for  $B$ .
  - ▶ The default of  $A$  constitutes a loss for  $B$ ; the loss size depends on the term structure of interest rates at  $\tau_A$ .
- If interest rates have fallen relative to their value at  $t = 0$ :
  - ▶ The fixed payments have increased in value so that the swap has a negative value for  $B$ .

- ▶  $B$  will still has to pay the value of the contract into the bankruptcy pool, and there is **no upside for  $B$**  in  $A$ 's default.
- If  $B$  defaults first the situation is **reversed**: falling rates lead to a counterparty-risk-related loss for  $A$ .

## Management of counterparty risk

- Counterparty risk has to be taken into account in pricing and valuation. This has led to the notion of **credit value adjustments (CVA)**.
- Counterparty risk needs to be controlled using risk-mitigation techniques such as **netting** and **collateralization**.
- Under a **netting agreement** the value of all derivatives transactions between  $A$  and  $B$  is computed and only the aggregated value is subject to counterparty risk; since offsetting transactions cancel each other out, this has the potential to reduce counterparty risk substantially.
- Under a **collateralization agreement** the parties exchange collateral (cash and securities) that serves as a pledge for the receiver. The value of the collateral is adjusted dynamically to reflect changes in the value of the underlying transactions.

## 17.2 Credit value adjustments

**General definition.** The price (for the protection buyer ) satisfies

$$\begin{aligned}\text{True price} = & \text{(counterparty) risk-free price} \\ & - \text{adjustment for default of seller (CVA)} \\ & + \text{adjustment for default of buyer (DVA) ,}\end{aligned}$$

where CVA and DVA stand for Credit Value Adjustment and Debt Value Adjustment respectively.



## General adjustment formulas

Denote by  $V_t$  the market value of the CDS (assuming that  $B$  and  $S$  are default-free), by  $\tau = \min \tau_R, \tau_S, \tau_B$  the first default time and by  $\xi \in \{R, S, B\}$  the identity of first defaulting firm. Recall that  $x^+ = \max(x, 0)$  and  $x^- = -\min(x, 0)$  and denote by  $D(0, t)$  the discount factor over the period  $[0, t]$  (with constant interest rate,  $D(0, t) = e^{-rt}$ ).

It can be shown that

$$\begin{aligned}\text{CVA} &= \mathbb{E}^{\mathbb{Q}}(I_{\{\tau < T\}} I_{\{\xi=S\}} D(0, \tau) \delta^S V_{\tau}^+) \\ \text{DVA} &= \mathbb{E}^{\mathbb{Q}}(I_{\{\tau < T\}} I_{\{\xi=B\}} D(0, \tau) \delta^B V_{\tau}^-)\end{aligned}$$

## Comments.

- CVA gives loss of B due to premature default of S; DVA gives loss of S due to premature default of B.
- The value adjustments involve an option on the market value  $V = (V_t)_{t \leq T}$  of the swap with strike  $K = 0$  (a call for the CVA and a put for the DVA).
- Similar formula holds if  $V$  is the market value of another derivative such as an interest swap or even a reinsurance contract.
- DVA is a bit problematic: a worsening credit quality of B leads to an accounting profit for B.

## A simplified formula

In order to evaluate the CVA and DVA formulas one needs a model with stochastic credit spreads that takes **dependence** between the default of S, B and the **market value**  $V$  of the CDS into account (a **dynamic portfolio credit risk model**). Markets often work with a simpler formula that assumes that the default of S and B and  $V$  are **independent**:

$$\begin{aligned}\text{CVA}^{\text{indep}} &= \delta^S \int_0^T \bar{F}_B(t) D(0, t) E^Q(V_t^+) f_S(t) dt, \\ \text{DVA}^{\text{indep}} &= \delta^B \int_0^T \bar{F}_S(t) D(0, t) E^Q(V_t^-) f_B(t) dt.\end{aligned}$$

Here  $f_S$  is the density of  $\tau_S$  and  $\bar{F}_B$  resp  $\bar{F}_S$  is the survival function of  $\tau_B$  resp  $\tau_S$ .

## Comments.

- In order to evaluate the simplified formula one only needs to determine the marginal distribution of  $\tau_S$  and  $\tau_B$  and the so-called **expected exposures**  $\mathbb{E}^{\mathbb{Q}}(V_t^+)$  and  $\mathbb{E}^{\mathbb{Q}}(V_t^-)$ .
- The independence assumption underlying the simplified value adjustment formula between the price of the CDS on  $R$ , that is  $V_t$ , and the default event of  $S$  and  $B$  is often unrealistic; in practice this is known as **wrong way risk**.

Examples:

- ▶ CDS on a financial institution: given that  $S$  defaults it is quite likely that credit quality of  $R$  is low.
- ▶ Reinsurance.

For further reading on counterparty risk see Gregory (2012).

# References

Artzner, P., Delbaen, F., Eber, J. M., and Heath, D. (1999), Coherent measures of risk, *Mathematical Finance*, 9, 203–228.

Balkema, A. A. and de Haan, L. (1974), Residual life time at great age, *The Annals of Probability*, 2, 792–804.

CME SPAN: Standard Portfolio Analysis of Risk (2010), [www.cmegroup.com/cme](http://www.cmegroup.com/cme)  
Chicago Mercantile Exchange.

Cox, J. C., Ingersoll, J. E., and Ross, S. A. (1985), A Theory of the Term Structure of Interest Rates, *Econometrica. Journal of the Econometric Society*, 53(2), 385–407.

D'Agostino, R. B. and Stephens, M. A. (1986), Goodness-of-fit techniques, Dekker.

Demarta, S. and McNeil, A. J. (2005), The  $t$  Copula and Related Copulas, *International Statistical Review*, 73(1), 111–129.

- Embrechts, P., Klüppelberg, C., and Mikosch, T. (1997), Modelling Extremal Events for Insurance and Finance, Berlin: Springer.
- Fan, Y. and Patton, A. (2014), Copulas in econometrics, *Annual Review of Economics*, 6, 179–200.
- Föllmer, H. and Schied, A. (2002), Convex measures of risk and trading constraints, *Finance and Stochastics*, 6, 429–447.
- Gnedenko, B. V. (1943), Sur la distribution limite du terme maximum d'une série aléatoire, *Annals of Mathematics*, 44, 423–453.
- Gneiting, T. (2011), Making and Evaluating Point Forecasts, *Journal of the American Statistical Association*, 106(494), 746–762.
- Gregory, J. (2012), Counterparty Credit Risk and Credit Value Adjustment: A Continuing Challenge for Global Financial Markets, Chichester: John Wiley and Sons Ltd.
- Huang, J.-Z., Lanfranconi, M., Patel, N., and Pospisil, L. (2012), Modeling credit correlations: an overview of the Moody's Analytics GCorr Model, tech. rep., Moody's Analytics.

- Jarrow, R. A. and Turnbull, S. M. (1995), Pricing Derivatives on Financial Securities Subject to Credit Risk, *The Journal of Finance*, 50(1), 53–85.
- Joenssen, D. W. and Vogel, J. (2014), A power study of goodness-of-fit tests for multivariate normality implemented in R, *Journal of Statistical Computation and Simulation*, 84, 1055–1078.
- Kloman, H. F. (1990), Risk management agonists, *Risk Analysis*, 10, 201–205.
- Kolmogorov, A. N. (1933), Grundbegriffe der Wahrscheinlichkeitsrechnung, Berlin: Ergebnisse der Mathematik.
- Lando, D. and Skodeberg, T. (2002), Analyzing Rating Transitions and Rating Drift with Continuous Observations, *Journal of Banking and Finance*, 26, 423–444.
- McNeil, A. J. and Frey, R. (2000), Estimation of tail-related risk measures for heteroscedastic financial time series: An extreme value approach, *Journal of Empirical Finance*, 7, 271–300.

- McNeil, A. J., Frey, R., and Embrechts, P. (2005), Quantitative Risk Management: Concepts, Techniques and Tools, Princeton: Princeton University Press.
- McNeil, A. J., Frey, R., and Embrechts, P. (2015), Quantitative Risk Management: Concepts, Techniques and Tools, 2nd, Princeton: Princeton University Press.
- Nelson, C. R. and Siegel, A. F. (1987), Parsimonious modeling of yield curves, *Journal of Business*, 60, 473–489.
- Ou, S. (2013), Annual Default Study: Corporate Default and recovery Rates, 1920-2012, Technical document, Moody's Investor Service.
- Patton, A. J. (2006), Modelling asymmetric exchange rate dependence, *International Economic Review*, 47, 527–556.
- Patton, A. (2012), A review of copula models for economic time series, *Journal of Multivariate Analysis*, 110, 4–18.
- Pickands, J. (1975), Statistical inference using extreme order statistics, *The Annals of Statistics*, 3, 119–131.



- Press, W. H., Teukolsky, S. A., Vetterling, W. T., and Flannery, B. P. (1992), *Numerical Recipes in C*, Cambridge: Cambridge University Press.
- RiskMetrics (1996), RiskMetrics Technical Document, 3rd, J.P. Morgan, New York.
- Shreve, S. E. (2008), Don't blame the quants, Available at [www.forbes.com/2008/1008shreve.html](http://www.forbes.com/2008/1008shreve.html).
- Smith, R. L. (1987), Estimating Tails of Probability Distributions, *The Annals of Statistics*, 15, 1174–1207.