

<p><b>Statistics One</b></p> <p>Lecture 3 Variables, Distributions, &amp; Scales</p>
1

<p><b>Three segments</b></p> <ul style="list-style-type: none"><li>• Variables</li><li>• Distributions</li><li>• Scales</li></ul>
2

<p><b>Lecture 3 ~ Segment 1</b></p> <p>Types of variables</p>
3

<p><b>Variables</b></p> <ul style="list-style-type: none"><li>• Variables can take on multiple values</li><li>• In contrast, a constant has only one value</li></ul>
4

## Apples and gravity



5

## Variables

- The size, shape, weight, and type of apple are all variables
- Gravity, or gravitational force, is a constant on Earth

6

## Types of variables

- Nominal
- Ordinal
- Interval
- Ratio

7

## Stevens (1946)

### SCIENCE

Vol. 103, No. 2684

Friday, June 7, 1946

#### On the Theory of Scales of Measurement

S. S. Stevens  
*Director, Psycho-Acoustic Laboratory, Harvard University*

8

## Types of variables

- Nominal variables
  - Used to assign individual cases to categories
    - For example, Coursera students come from many different countries
    - *Country of Origin* is a nominal variable

9

## Types of variables

- Ordinal variables
  - Used to rank order cases
    - For example, countries may be ranked according to overall population
    - *Ranking* is an ordinal variable


10

## Types of variables

- Interval variables
  - Used to rank order cases and the distance, or interval, between each value is equal
    - For example, each country has a longitude and latitude
    - *Longitude* and *Latitude* are interval variables

11

## Types of variables



12

## Types of variables

- Ratio variables

- The same as interval variables but they have a “true zero”
  - For example, *Population* (Population = 0 = extinct)
  - For example, *Age* (Age = 0 literally means NO age)
  - For example, *Temperature K°* (the Kelvin scale)

13

## Stevens (1946)

TABLE 1			
Scale	Basic Empirical Operations	Mathematical Group Structure	Permissible Statistics (Invariance)
NOMINAL	Determination of equality	Permutation group $f(x) = f(y)$ $f(x)$ means any one-to-one function	Number of cases Mode Contingency correlation
ORDINAL	Determination of greater or less	Isotonic group $f(x) \leq f(y)$ $f(x)$ means any monotonic increasing function	Median Percentiles
INTERVAL	Determination of equality of intervals or differences	General linear group $x' = ax + b$	Mean Standard deviation Rank-order correlation Product-moment correlation
RATIO	Determination of equality of ratios	Similarity group $x'' = ax + b$	Coefficient of variation

14

## Types of variables

- Preview of variables in this course
  - Nominal variables
    - Independent variables in experimental research
      - For example, treatment to prevent polio (vaccine, placebo)
    - Quasi-independent variables in correlational research
      - For example, gender (female, male)

15

## Types of variables

- Preview of variables in this course
  - Interval and Ratio variables
    - Dependent variables in experimental research
      - For example, rate of polio in a community
    - Measured variables in correlational research
      - For example, intelligence test scores

16

## Types of variables

- Preview of variables in this course
  - Discrete vs. continuous variables
    - Nominal variables are discrete (categorical)
    - Interval and ratio variables are continuous
    - Ordinal variables are technically discrete but they are often treated as continuous in statistical analyses (more on this later)

17

## Segment summary

- Types of variables
  - Nominal
  - Ordinal
  - Interval
  - Ratio

18

**END SEGMENT**

19

## Lecture 3 ~ Segment 2

Distributions: Histograms

20

## Histograms

- A histogram is a type of graph used to display a distribution


21

## Histograms

- Why start with histograms?
  - To overcome the natural tendency to rely upon summary information, such as an average


22

## An example: Body temperature



23

## An example: Body temperature




24

## Histograms

- Histograms can reveal information not captured by summary statistics
  - Suppose a few children in a school are sick with influenza (flu) and have a high temperature
    - The distribution will be positively skewed


25

## An example: Body temperature



26

## An example: Body temperature



27


## Histograms

- Not all distributions are normal
  - Suppose one group of children had the flu a week prior to a second sick group of children
  - Assume the first group received antibiotics, which temporarily caused their body temperatures to be slightly below normal, while the second group was still above normal


28

## An example: Body temperature

Normal, below average




Normal, above average



29

## An example: Body temperature


Bimodal




30

## An example: Body temperature

Normal, below average




Normal, above average



31

## An example: Body temperature

Bimodal



32

## Histograms

- Not all distributions are normal
  - Simply viewing a histogram often reveals whether a distribution is normal or not normal
  - However, sometimes it is hard to determine
    - Summary statistics help in such cases

33

## Histograms

- Not all distributions are normal
  - As you view more and more distributions you will get a better sense of what is normal and what is not normal
  - So, let's look at more distributions

34

## Wine tasting!



35

## An example: Wine ratings

- Suppose that 100 wine experts rated the overall quality of 8 different wines on a scale of 1 to 100
  - Higher scores indicate higher quality

36

## An example: Wine ratings

- Suppose four countries submitted two wines each, one red and one white
  - Argentina
  - Australia
  - France
  - USA

37

## An example: Wine ratings

Malbec & Chardonnay      Shiraz & Pinot Grigio



38

## An example: Wine ratings

Bourdeaux & Sauvignon

Blanc



Cabernet & Reisling



39

## An example: Wine ratings


### • Preview

- The ratings of the red wines are normal
- The ratings of the whites are not normal


40

## An example: Wine ratings

Red, Argentina




Red, Australia




41

## Four histograms

Red, France




Red, USA




42

## An example: Wine ratings

White, Argentina




White, Australia




43

## An example: Wine ratings

White, France



White, USA



44

## Segment summary

- Histograms are used to display distributions
- Many distributions are normal


45

## Segment summary

- Some distributions are not normal, for example:
  - Bi-modal
  - Positively skewed
  - Negatively skewed
  - Uniform (platykurtic)
  - Leptokurtic

46

## Advanced graphs



47

## Advanced graphs



48

## Advanced graphs



49

## Advanced graphs



50

**END SEGMENT**

51

**Lecture 3 ~ Segment 3**

Scales of measurement

52

## Scales

- Scales of measurement
  - For example, in the last segment body temperature was presented in both Fahrenheit and in Celsius
    - Different scales but both measure temperature
    - F° can be converted to C° and vice-versa

53

## Scales

- In statistics, there is a standard scale
  - The Z scale
- Any score from any scale can be converted
  - To Z scores
- Allows for efficient communication

54

## Z scores

- $Z = (X - M) / SD$ 
  - X is a score on an original scale (raw score)
  - M is the mean
  - SD is the standard deviation


55

## Z scores

- $Z = (X - M) / SD$ 
  - The mean Z-score is  $Z = 0$
  - Positive Z scores are above average
  - Negative Z scores are below average


56

## Body temperature F°




57

## Body temperature C°



58

## Body temperature Z



59

## Z scores

- For example, assume  $M = 98.6$ ,  $SD = .5$
- Suppose an individual,  $X = 99.6$
- Convert  $X$  to  $Z$

60

## Z scores

- Convert X to Z

$$\begin{aligned} \bullet Z &= (X - M) / SD \\ \bullet Z &= (99.6 - 98.6) / .5 = 2 \\ \bullet Z &= 2 \end{aligned}$$


61

## Percentile rank

- Percentile rank
  - The percentage of scores that fall at or below a score in a distribution
  - Assume a normal distribution
  - If  $Z = 0$  then the percentile rank = 50<sup>th</sup>
  - 50 percent of the distribution falls below the mean

62

## Body temperature Z



63

## Segment summary

- The Z-scale is the standard scale in statistics
- Raw scores can be converted to Z-scores
- Z-scores can be used to find percentile rank
  - Raw score ~ Z-score ~ Percentile rank

64

**END SEGMENT**

65

**END LECTURE 3**

66