

Statistics One

Lecture 10
Confidence intervals

1

Two segments

- Confidence intervals for sample means (M)
- Confidence intervals for regression coefficients (B)

2

Lecture 10 ~ Segment 1

Confidence intervals for
sample means (M)


3

Confidence intervals

- All sample statistics, for example, a sample mean (M), are *point estimates*
- More specifically, a sample mean (M) represents a single point in a sampling distribution

4

The family of t distributions



5

Confidence intervals

- The logic of confidence intervals is to report a range of values, rather than a single value
- In other words, report an *interval estimate* rather than a *point estimate*

6

Confidence intervals

- *Confidence interval*: an interval estimate of a population parameter, based on a random sample
 - Degree of confidence, for example 95%, represents the probability that the interval captures the true population parameter

7

Confidence intervals

- The main argument for interval estimates is the reality of sampling error
- Sampling error implies that point estimates will vary from one study to the next
- A researcher will therefore be more confident about accuracy with an interval estimate

8

Confidence intervals for M

- Example, IMPACT
- Assume $N = 30$ and multiple samples...
 - Symptom Score (Baseline), $M = 0.05$
 - Symptom Score (Baseline), $M = 0.07$
 - Symptom Score (Baseline), $M = 0.03$
 - ...

9

Confidence intervals for M

- Example, IMPACT
- Assume $N = 10$ and multiple samples...
 - Symptom Score (Baseline), $M = 0.01$
 - Symptom Score (Baseline), $M = 0.20$
 - Symptom Score (Baseline), $M = 0.00$
 - ...

10

Confidence intervals for M

- Example, IMPACT
- Assume $N = 30$ and multiple samples...
 - Symptom Score (Post-injury), $M = 12.03$
 - Symptom Score (Post-injury), $M = 12.90$
 - Symptom Score (Post-injury), $M = 14.13$
 - ...

11

Confidence intervals for M

- Example, IMPACT
- Assume $N = 10$ and multiple samples...
 - Symptom Score (Post-injury), $M = 19.70$
 - Symptom Score (Post-injury), $M = 8.40$
 - Symptom Score (Post-injury), $M = 13.30$
 - ...

12

Confidence intervals for M

- The width of a confidence interval is influenced by
 - Sample size
 - Variance in the population (and sample)
- Standard error (SE) = SD / \sqrt{N}

13

Confidence intervals for M

- Example, IMPACT
- Assume $N = 30$
 - Symptom Score (Baseline)
 - 95% confidence interval
 - $-0.03 - 0.10$

14

Confidence intervals for M

- Example, IMPACT
- Assume $N = 10$
 - Symptom Score (Baseline)
 - 95% confidence interval
 - $-0.10 - 0.50$

15

Confidence intervals for M

- Example, IMPACT
- Assume $N = 30$
 - Symptom Score (Post-injury)
 - 95% confidence interval
 - $7.5 - 18.3$

16

Confidence intervals for M

- Example, IMPACT
- Assume $N = 10$
 - Symptom Score (Post-injury)
 - 95% confidence interval
 - 2.7 – 23.9


17

Confidence interval for M

- Upper bound = $M + t(SE)$
- Lower bound = $M - t(SE)$
- $SE = SD / \sqrt{N}$
- t depends on level of confidence desired and sample size

18

The family of t distributions



19

Segment summary

- All sample statistics, for example, a sample mean (M), are *point estimates*
- More specifically, a sample mean (M) represents a single point in a sampling distribution

20

Segment summary

- The logic of confidence intervals is to report a range of values, rather than a single value
- In other words, report an *interval estimate* rather than a *point estimate*

21

Segment summary

- The width of a confidence interval is influenced by
 - Sample size
 - Variance in the population (and sample)
 - Standard error (SE) = SD / \sqrt{N}

22

END SEGMENT

23

Lecture 10 ~ Segment 2

Confidence intervals for
regression coefficients (B)

24

Confidence intervals for B

- All sample statistics, for example, a regression coefficient (B), are *point estimates*
- More specifically, a regression coefficient (B) represents a single point in a sampling distribution


25

Confidence intervals for B

- In regression, $t = B / SE$

26

The family of t distributions



27

Confidence intervals for B

- The logic of confidence intervals is to report a range of values, rather than a single value
- In other words, report an *interval estimate* rather than a *point estimate*

28

Confidence intervals for B

- The main argument for interval estimates is the reality of sampling error
- Sampling error implies that point estimates will vary from one study to the next
- A researcher will therefore be more confident about accuracy with an interval estimate

29

Confidence intervals for B

- The width of a confidence interval is influenced by
 - Sample size
 - Variance in the population (and sample)
- Standard error (SE) = SD / \sqrt{N}


30

Confidence intervals for B

- Example, IMPACT
- Assume $N = 40$
 - Visual memory = $B_0 + (B)\text{Verbal memory}$

31

Confidence intervals for B




Confidence intervals for B

- Example, IMPACT
- Assume $N = 20$
 - Visual memory = $B_0 + (B)\text{Verbal memory}$

33

Confidence intervals for B



34

Segment summary

- All sample statistics are *point estimates*
- More specifically, a sample mean (M) or a regression coefficient (B) represents a single point in a sampling distribution

35

Segment summary

- The logic of confidence intervals is to report a range of values, rather than a single value
- In other words, report an *interval estimate* rather than a *point estimate*

36

Segment summary

- The width of a confidence interval is influenced by
 - Sample size
 - Variance in the population (and sample)
 - Standard error (SE) = SD / \sqrt{N}

37

END SEGMENT

38

END LECTURE 10

39