

# NCERT-9.4.6

EE24BTECH11035 - KOTHAPALLI AKHIL

**Question:** Find the solution of the differential equation:

$$\frac{dy}{dx} = (1 + x^2)(1 + y^2).$$

**Solution:** Rewriting the equation:

$$\frac{dy}{1 + y^2} = (1 + x^2)dx.$$

Integrating both sides:

$$\int \frac{dy}{1 + y^2} = \int (1 + x^2)dx.$$

After simplification:

$$\tan^{-1}(y) = x + \frac{x^3}{3} + k.$$

where  $k$  is the constant of integration(Here, it is assumed as 0).

## Numerical Approach:

I used a for loop for finding the  $y$  values as the loop proceeds with iterative formula given below. I took some initial value of  $x$  and as loop proceeds I assigned it the value as  $x + h$ , where  $h$  is the step size, representing the rate of change.

2. Assigned the values of  $y$  for different  $x$ -values using a for loop.

The iterative formula for updating  $y$ -values is:

$$y_n = y_{n-1} + \left(\frac{dy}{dx}\right)h, \quad (1)$$

The iterative formula for updating  $x$ -values is:

$$x_n = x_{n-1} + h \quad (2)$$

## Initial Conditions:

- $x = 0$
- $y = 0$
- $h = 0.0002$

Using Matplotlib, I plotted the computed points and the graph of the exact solution to verify that they approximately match.




Fig. 0.1: verifying through graph of sim and theory values