



MAT1320-Linear Algebra Lecture Notes

Linear Dependence and Independence of Vectors and Spanning Sets

Mehmet E. KÖROĞLU

Fall 2020

YILDIZ TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS

mkoroglu@yildiz.edu.tr

Table of contents

1. Linear Dependence and Independence of Vectors
2. Linear Combination of Vectors
3. Spanning Sets

Linear Dependence and Independence of Vectors

Linear Dependence and Independence of Vectors

Definition

Let V be a real vector space, $x_1, x_2, \dots, x_n \in \mathbb{R}$ and

$\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n \in V$.

Linear Dependence and Independence of Vectors

Definition

Let V be a real vector space, $x_1, x_2, \dots, x_n \in \mathbb{R}$ and

$\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n \in V$. We say vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are **linearly dependent** if there exists scalars $x_1, x_2, \dots, x_n \in \mathbb{R}$, not all of them zero, such that $x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n = \vec{0}$.

Linear Dependence and Independence of Vectors

Definition

Let V be a real vector space, $x_1, x_2, \dots, x_n \in \mathbb{R}$ and

$\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n \in V$. We say vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are **linearly dependent** if there exists scalars $x_1, x_2, \dots, x_n \in \mathbb{R}$, not all of them zero, such that $x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n = \vec{0}$. We say vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are **linearly independent** if there exists scalars $x_1, x_2, \dots, x_n \in \mathbb{R}$, all of them zero, such that $x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n = \vec{0}$.

Linear Dependence and Independence of Vectors

Definition

Let V be a real vector space, $x_1, x_2, \dots, x_n \in \mathbb{R}$ and $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n \in V$. We say vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are **linearly dependent** if there exists scalars $x_1, x_2, \dots, x_n \in \mathbb{R}$, not all of them zero, such that $x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n = \vec{0}$. We say vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are **linearly independent** if there exists scalars $x_1, x_2, \dots, x_n \in \mathbb{R}$, all of them zero, such that $x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n = \vec{0}$.

Note: If the only solution of the homogeneous system $x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n = \vec{0}$ is zero solution, then we say vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are linearly independent.

Linear Dependence and Independence of Vectors

Definition

Let V be a real vector space, $x_1, x_2, \dots, x_n \in \mathbb{R}$ and

$\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n \in V$. We say vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are **linearly dependent** if there exists scalars $x_1, x_2, \dots, x_n \in \mathbb{R}$, not all of them zero, such that $x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n = \vec{0}$. We say vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are **linearly independent** if there exists scalars $x_1, x_2, \dots, x_n \in \mathbb{R}$, all of them zero, such that $x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n = \vec{0}$.

Note: If the only solution of the homogeneous system

$x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n = \vec{0}$ is zero solution, then we say vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are linearly independent. If the

homogeneous system $x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n = \vec{0}$ has a nonzero solution, then we say vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ are linearly dependant.

Linear Dependence and Independence of Vectors

Example

The vectors

$\{\vec{e}_1 = (1, 0, 0), \vec{e}_2 = (0, 1, 0), \vec{e}_3 = (0, 0, 1)\} \subset \mathbb{R}^3$ are linearly independent.

Linear Dependence and Independence of Vectors

Example

The vectors

$\{\vec{e}_1 = (1, 0, 0), \vec{e}_2 = (0, 1, 0), \vec{e}_3 = (0, 0, 1)\} \subset \mathbb{R}^3$ are linearly independent. For $x_1, x_2, x_3 \in \mathbb{R}$ we have

$$\Rightarrow x_1 \vec{e}_1 + x_2 \vec{e}_2 + x_3 \vec{e}_3 = (0, 0, 0)$$

Linear Dependence and Independence of Vectors

Example

The vectors

$\{\vec{e}_1 = (1, 0, 0), \vec{e}_2 = (0, 1, 0), \vec{e}_3 = (0, 0, 1)\} \subset \mathbb{R}^3$ are linearly independent. For $x_1, x_2, x_3 \in \mathbb{R}$ we have

$$\Rightarrow x_1 \vec{e}_1 + x_2 \vec{e}_2 + x_3 \vec{e}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 0, 0) + x_2(0, 1, 0) + x_3(0, 0, 1) = (0, 0, 0)$$

Linear Dependence and Independence of Vectors

Example

The vectors

$\{\vec{e}_1 = (1, 0, 0), \vec{e}_2 = (0, 1, 0), \vec{e}_3 = (0, 0, 1)\} \subset \mathbb{R}^3$ are linearly independent. For $x_1, x_2, x_3 \in \mathbb{R}$ we have

$$\Rightarrow x_1 \vec{e}_1 + x_2 \vec{e}_2 + x_3 \vec{e}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 0, 0) + x_2(0, 1, 0) + x_3(0, 0, 1) = (0, 0, 0)$$

$$\Rightarrow (x_1, x_2, x_3) = (0, 0, 0)$$

Linear Dependence and Independence of Vectors

Example

The vectors

$\{\vec{e}_1 = (1, 0, 0), \vec{e}_2 = (0, 1, 0), \vec{e}_3 = (0, 0, 1)\} \subset \mathbb{R}^3$ are linearly independent. For $x_1, x_2, x_3 \in \mathbb{R}$ we have

$$\Rightarrow x_1 \vec{e}_1 + x_2 \vec{e}_2 + x_3 \vec{e}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 0, 0) + x_2(0, 1, 0) + x_3(0, 0, 1) = (0, 0, 0)$$

$$\Rightarrow (x_1, x_2, x_3) = (0, 0, 0)$$

$$\Rightarrow x_1 = 0, x_2 = 0, x_3 = 0.$$

Linear Dependence and Independence of Vectors

Example

The vectors

$\{\vec{e}_1 = (1, 0, 0), \vec{e}_2 = (0, 1, 0), \vec{e}_3 = (0, 0, 1)\} \subset \mathbb{R}^3$ are linearly independent. For $x_1, x_2, x_3 \in \mathbb{R}$ we have

$$\Rightarrow x_1 \vec{e}_1 + x_2 \vec{e}_2 + x_3 \vec{e}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 0, 0) + x_2(0, 1, 0) + x_3(0, 0, 1) = (0, 0, 0)$$

$$\Rightarrow (x_1, x_2, x_3) = (0, 0, 0)$$

$$\Rightarrow x_1 = 0, x_2 = 0, x_3 = 0.$$

Thus $\vec{e}_1, \vec{e}_2, \vec{e}_3$ are linearly independent.

Linear Dependence and Independence of Vectors

Example

Let's show that the vectors

$\{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (3, 2, 1)\} \subset \mathbb{R}^3$ are linearly dependent.

Linear Dependence and Independence of Vectors

Example

Let's show that the vectors

$\{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (3, 2, 1)\} \subset \mathbb{R}^3$ are linearly dependent. To do this, let $x_1, x_2, x_3 \in \mathbb{R}$. Then we have

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

Linear Dependence and Independence of Vectors

Example

Let's show that the vectors

$\{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (3, 2, 1)\} \subset \mathbb{R}^3$ are linearly dependent. To do this, let $x_1, x_2, x_3 \in \mathbb{R}$. Then we have

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(3, 2, 1) = (0, 0, 0)$$

Linear Dependence and Independence of Vectors

Example

Let's show that the vectors

$\{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (3, 2, 1)\} \subset \mathbb{R}^3$ are linearly dependent. To do this, let $x_1, x_2, x_3 \in \mathbb{R}$. Then we have

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(3, 2, 1) = (0, 0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 + 3x_3 = 0 \\ 2x_1 + 2x_3 = 0 \\ x_2 + x_3 = 0 \end{cases}$$

Linear Dependence and Independence of Vectors

Example

Let's show that the vectors

$\{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (3, 2, 1)\} \subset \mathbb{R}^3$ are linearly dependent. To do this, let $x_1, x_2, x_3 \in \mathbb{R}$. Then we have

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(3, 2, 1) = (0, 0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 + 3x_3 = 0 \\ 2x_1 + 2x_3 = 0 \\ x_2 + x_3 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 & 3 \\ 2 & 0 & 2 \\ 0 & 1 & 1 \end{pmatrix}$$

Linear Dependence and Independence of Vectors

Example

Let's show that the vectors

$\{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (3, 2, 1)\} \subset \mathbb{R}^3$ are linearly dependent. To do this, let $x_1, x_2, x_3 \in \mathbb{R}$. Then we have

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(3, 2, 1) = (0, 0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 + 3x_3 = 0 \\ 2x_1 + 2x_3 = 0 \\ x_2 + x_3 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 & 3 \\ 2 & 0 & 2 \\ 0 & 1 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

Linear Dependence and Independence of Vectors

Example

Let's show that the vectors

$\{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (3, 2, 1)\} \subset \mathbb{R}^3$ are linearly dependent. To do this, let $x_1, x_2, x_3 \in \mathbb{R}$. Then we have

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(3, 2, 1) = (0, 0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 + 3x_3 = 0 \\ 2x_1 + 2x_3 = 0 \\ x_2 + x_3 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 & 3 \\ 2 & 0 & 2 \\ 0 & 1 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\Rightarrow x_1 + x_3 = 0, x_2 + x_3 = 0$$

Linear Dependence and Independence of Vectors

Example

Let's show that the vectors

$\{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (3, 2, 1)\} \subset \mathbb{R}^3$ are linearly dependent. To do this, let $x_1, x_2, x_3 \in \mathbb{R}$. Then we have

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(3, 2, 1) = (0, 0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 + 3x_3 = 0 \\ 2x_1 + 2x_3 = 0 \\ x_2 + x_3 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 & 3 \\ 2 & 0 & 2 \\ 0 & 1 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\Rightarrow x_1 + x_3 = 0, x_2 + x_3 = 0 \Rightarrow \text{If } x_3 = r, \text{ then } x_1 = x_2 = -r.$$

Linear Dependence and Independence of Vectors

Example

Let's show that the vectors

$\{\vec{v}_1 = (1, 2, 0), \vec{v}_2 = (2, 0, 1), \vec{v}_3 = (3, 2, 1)\} \subset \mathbb{R}^3$ are linearly dependent. To do this, let $x_1, x_2, x_3 \in \mathbb{R}$. Then we have

$$\Rightarrow x_1 \vec{v}_1 + x_2 \vec{v}_2 + x_3 \vec{v}_3 = (0, 0, 0)$$

$$\Rightarrow x_1(1, 2, 0) + x_2(2, 0, 1) + x_3(3, 2, 1) = (0, 0, 0)$$

$$\Rightarrow \begin{cases} x_1 + 2x_2 + 3x_3 = 0 \\ 2x_1 + 2x_3 = 0 \\ x_2 + x_3 = 0 \end{cases} \Rightarrow \begin{pmatrix} 1 & 2 & 3 \\ 2 & 0 & 2 \\ 0 & 1 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\Rightarrow x_1 + x_3 = 0, x_2 + x_3 = 0 \Rightarrow \text{If } x_3 = r, \text{ then } x_1 = x_2 = -r.$$

This means that the system has infinitely many solution. So

$\vec{v}_1, \vec{v}_2, \vec{v}_3$ are linearly dependent.

Linear Combination of Vectors

Linear Combination of Vectors

Definition

Let V be a real vector space, $x_1, x_2, \dots, x_n \in \mathbb{R}$ and

$\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n \in V$.

Linear Combination of Vectors

Definition

Let V be a real vector space, $x_1, x_2, \dots, x_n \in \mathbb{R}$ and

$\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n \in V$. If there exists $x_1, x_2, \dots, x_n \in \mathbb{R}$ such that

$$\vec{w} = x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n,$$

Linear Combination of Vectors

Definition

Let V be a real vector space, $x_1, x_2, \dots, x_n \in \mathbb{R}$ and

$\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n \in V$. If there exists $x_1, x_2, \dots, x_n \in \mathbb{R}$ such that

$$\vec{w} = x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n,$$

then we say \vec{w} is a **linear combination** of the vectors

$\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$.

Linear Combination of Vectors

Definition

Let V be a real vector space, $x_1, x_2, \dots, x_n \in \mathbb{R}$ and

$\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n \in V$. If there exists $x_1, x_2, \dots, x_n \in \mathbb{R}$ such that

$$\vec{w} = x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n,$$

then we say \vec{w} is a **linear combination** of the vectors

$\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$. Here, again $\vec{w} = x_1 \vec{v}_1 + x_2 \vec{v}_2 + \dots + x_n \vec{v}_n$ is an element of V .

Linear Combination of Vectors

Example

Show that $\vec{\mathbf{w}} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{\mathbf{u}} = (1, 2, -1)$ and $\vec{\mathbf{v}} = (6, 4, 2)$, but $\vec{\mathbf{w}}' = (4, -1, 8)$ is not.

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\vec{w} = x_1 \vec{u} + x_2 \vec{v}$$

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\vec{w} = x_1 \vec{u} + x_2 \vec{v}$$

$$(9, 2, 7) = x_1 (1, 2, -1) + x_2 (6, 4, 2)$$

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\vec{w} = x_1 \vec{u} + x_2 \vec{v}$$

$$(9, 2, 7) = x_1 (1, 2, -1) + x_2 (6, 4, 2)$$

$$\Rightarrow \begin{cases} x_1 + 6x_2 = 9 \\ 2x_1 + 4x_2 = 2 \\ -x_1 + 2x_2 = 7 \end{cases}$$

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\vec{w} = x_1 \vec{u} + x_2 \vec{v}$$

$$(9, 2, 7) = x_1 (1, 2, -1) + x_2 (6, 4, 2)$$

$$\Rightarrow \begin{cases} x_1 + 6x_2 = 9 \\ 2x_1 + 4x_2 = 2 \\ -x_1 + 2x_2 = 7 \end{cases} \Rightarrow \begin{pmatrix} 1 & 6 & 9 \\ 2 & 4 & 2 \\ -1 & 2 & 7 \end{pmatrix}$$

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\vec{w} = x_1 \vec{u} + x_2 \vec{v}$$

$$(9, 2, 7) = x_1 (1, 2, -1) + x_2 (6, 4, 2)$$

$$\Rightarrow \begin{cases} x_1 + 6x_2 = 9 \\ 2x_1 + 4x_2 = 2 \\ -x_1 + 2x_2 = 7 \end{cases} \Rightarrow \begin{pmatrix} 1 & 6 & 9 \\ 2 & 4 & 2 \\ -1 & 2 & 7 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & -3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}$$

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\vec{w} = x_1 \vec{u} + x_2 \vec{v}$$

$$(9, 2, 7) = x_1 (1, 2, -1) + x_2 (6, 4, 2)$$

$$\Rightarrow \begin{cases} x_1 + 6x_2 = 9 \\ 2x_1 + 4x_2 = 2 \\ -x_1 + 2x_2 = 7 \end{cases} \Rightarrow \begin{pmatrix} 1 & 6 & 9 \\ 2 & 4 & 2 \\ -1 & 2 & 7 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & -3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\Rightarrow x_1 = -3, x_2 = 2.$$

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\begin{aligned}\vec{w} &= x_1 \vec{u} + x_2 \vec{v} \\ (9, 2, 7) &= x_1 (1, 2, -1) + x_2 (6, 4, 2) \\ \Rightarrow \begin{cases} x_1 + 6x_2 = 9 \\ 2x_1 + 4x_2 = 2 \\ -x_1 + 2x_2 = 7 \end{cases} &\Rightarrow \begin{pmatrix} 1 & 6 & 9 \\ 2 & 4 & 2 \\ -1 & 2 & 7 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & -3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{pmatrix} \\ \Rightarrow x_1 = -3, x_2 = 2. &\end{aligned}$$

This means that the system has a unique solution and we have

$$(9, 2, 7) = -3(1, 2, -1) + 2(6, 4, 2).$$

Linear Combination of Vectors

Example

Show that $\vec{\mathbf{w}} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{\mathbf{u}} = (1, 2, -1)$ and $\vec{\mathbf{v}} = (6, 4, 2)$, but $\vec{\mathbf{w}}' = (4, -1, 8)$ is not.

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\vec{w}' = x_1 \vec{u} + x_2 \vec{v}$$

Linear Combination of Vectors

Example

Show that $\vec{\mathbf{w}} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{\mathbf{u}} = (1, 2, -1)$ and $\vec{\mathbf{v}} = (6, 4, 2)$, but $\vec{\mathbf{w}}' = (4, -1, 8)$ is not.

$$\begin{aligned}\vec{\mathbf{w}}' &= x_1 \vec{\mathbf{u}} + x_2 \vec{\mathbf{v}} \\ (4, -1, 8) &= x_1 (1, 2, -1) + x_2 (6, 4, 2)\end{aligned}$$

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\vec{w}' = x_1 \vec{u} + x_2 \vec{v}$$

$$(4, -1, 8) = x_1 (1, 2, -1) + x_2 (6, 4, 2)$$

$$\begin{cases} x_1 + 6x_2 = 4 \\ 2x_1 + 4x_2 = -1 \\ -x_1 + 2x_2 = 8 \end{cases}$$

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\begin{aligned}\vec{w}' &= x_1 \vec{u} + x_2 \vec{v} \\ (4, -1, 8) &= x_1 (1, 2, -1) + x_2 (6, 4, 2) \\ \left\{ \begin{array}{l} x_1 + 6x_2 = 4 \\ 2x_1 + 4x_2 = -1 \\ -x_1 + 2x_2 = 8 \end{array} \right. &\Rightarrow \begin{pmatrix} 1 & 6 & 4 \\ 2 & 4 & -2 \\ -1 & 2 & 8 \end{pmatrix}\end{aligned}$$

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\begin{aligned}\vec{w}' &= x_1 \vec{u} + x_2 \vec{v} \\ (4, -1, 8) &= x_1 (1, 2, -1) + x_2 (6, 4, 2) \\ \left\{ \begin{array}{l} x_1 + 6x_2 = 4 \\ 2x_1 + 4x_2 = -1 \\ -x_1 + 2x_2 = 8 \end{array} \right. &\Rightarrow \begin{pmatrix} 1 & 6 & 4 \\ 2 & 4 & -2 \\ -1 & 2 & 8 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}\end{aligned}$$

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\begin{aligned}\vec{w}' &= x_1 \vec{u} + x_2 \vec{v} \\ (4, -1, 8) &= x_1 (1, 2, -1) + x_2 (6, 4, 2) \\ \left\{ \begin{array}{l} x_1 + 6x_2 = 4 \\ 2x_1 + 4x_2 = -1 \\ -x_1 + 2x_2 = 8 \end{array} \right. &\Rightarrow \begin{pmatrix} 1 & 6 & 4 \\ 2 & 4 & -2 \\ -1 & 2 & 8 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &\Rightarrow 0 = 1.\end{aligned}$$

Linear Combination of Vectors

Example

Show that $\vec{w} = (9, 2, 7) \in \mathbb{R}^3$ is a linear combination of the vectors $\vec{u} = (1, 2, -1)$ and $\vec{v} = (6, 4, 2)$, but $\vec{w}' = (4, -1, 8)$ is not.

$$\begin{aligned}\vec{w}' &= x_1 \vec{u} + x_2 \vec{v} \\ (4, -1, 8) &= x_1 (1, 2, -1) + x_2 (6, 4, 2) \\ \begin{cases} x_1 + 6x_2 = 4 \\ 2x_1 + 4x_2 = -1 \\ -x_1 + 2x_2 = 8 \end{cases} &\Rightarrow \begin{pmatrix} 1 & 6 & 4 \\ 2 & 4 & -2 \\ -1 & 2 & 8 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\ &\Rightarrow 0 = 1.\end{aligned}$$

This means that the system is inconsistent. Thus \vec{w}' can not be written as a combination of the vectors \vec{u} and \vec{v} .

Linear Combination of Vectors

Example

For which values of k , $\vec{w} = (1, -2, k) \in \mathbb{R}^3$ can be written as a linear combination of the vectors $\vec{u} = (3, 0, -2)$ and $\vec{v} = (2, -1, 5)$.

Linear Combination of Vectors

Example

For which values of k , $\vec{w} = (1, -2, k) \in \mathbb{R}^3$ can be written as a linear combination of the vectors $\vec{u} = (3, 0, -2)$ and $\vec{v} = (2, -1, 5)$.

$$\vec{w} = x_1 \vec{u} + x_2 \vec{v}$$

Linear Combination of Vectors

Example

For which values of k , $\vec{w} = (1, -2, k) \in \mathbb{R}^3$ can be written as a linear combination of the vectors $\vec{u} = (3, 0, -2)$ and $\vec{v} = (2, -1, 5)$.

$$\begin{aligned}\vec{w} &= x_1 \vec{u} + x_2 \vec{v} \\ (1, -2, k) &= x_1 (3, 0, -2) + x_2 (2, -1, 5)\end{aligned}$$

Linear Combination of Vectors

Example

For which values of k , $\vec{w} = (1, -2, k) \in \mathbb{R}^3$ can be written as a linear combination of the vectors $\vec{u} = (3, 0, -2)$ and $\vec{v} = (2, -1, 5)$.

$$\vec{w} = x_1 \vec{u} + x_2 \vec{v}$$

$$(1, -2, k) = x_1 (3, 0, -2) + x_2 (2, -1, 5)$$

$$\begin{cases} 3x_1 + 2x_2 = 1 \\ -x_2 = -2 \\ -2x_1 + 5x_2 = k \end{cases}$$

Linear Combination of Vectors

Example

For which values of k , $\vec{w} = (1, -2, k) \in \mathbb{R}^3$ can be written as a linear combination of the vectors $\vec{u} = (3, 0, -2)$ and $\vec{v} = (2, -1, 5)$.

$$\begin{aligned}\vec{w} &= x_1 \vec{u} + x_2 \vec{v} \\ (1, -2, k) &= x_1 (3, 0, -2) + x_2 (2, -1, 5) \\ \left\{ \begin{array}{l} 3x_1 + 2x_2 = 1 \\ -x_2 = -2 \\ -2x_1 + 5x_2 = k \end{array} \right. &\Rightarrow x_2 = 2, x_1 = -1\end{aligned}$$

Linear Combination of Vectors

Example

For which values of k , $\vec{w} = (1, -2, k) \in \mathbb{R}^3$ can be written as a linear combination of the vectors $\vec{u} = (3, 0, -2)$ and $\vec{v} = (2, -1, 5)$.

$$\begin{aligned}\vec{w} &= x_1 \vec{u} + x_2 \vec{v} \\ (1, -2, k) &= x_1 (3, 0, -2) + x_2 (2, -1, 5) \\ \left\{ \begin{array}{l} 3x_1 + 2x_2 = 1 \\ -x_2 = -2 \\ -2x_1 + 5x_2 = k \end{array} \right. &\Rightarrow x_2 = 2, x_1 = -1 \\ &\Rightarrow k = -2(-1) + 5 \cdot 2 = 12.\end{aligned}$$

Linear Dependence and Independence of Vectors

Theorem

Let $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m$ be m linearly independent vectors in V . If $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m, \vec{v}_{m+1}$ are linearly dependent, then \vec{v}_{m+1} can be written as a linear combination of the vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m$.

Linear Dependence and Independence of Vectors

Theorem

Let $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m$ be m linearly independent vectors in V . If $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m, \vec{v}_{m+1}$ are linearly dependent, then \vec{v}_{m+1} can be written as a linear combination of the vectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m$.

Theorem

For $r < m$, if r vectors among $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m \in V$ are linearly dependent, then $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m$ are also linearly dependent.

Linear Dependence and Independence of Vectors

Theorem

Let V a vector space, and for $m \leq n$, $\vec{v}_1 = (a_{11}, a_{12}, \dots, a_{1n})$,
 $\vec{v}_2 = (a_{21}, a_{22}, \dots, a_{2n})$, \dots , $\vec{v}_m = (a_{m1}, a_{m2}, \dots, a_{mn})$ such that

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}.$$

Linear Dependence and Independence of Vectors

Theorem

Let V a vector space, and for $m \leq n$, $\vec{v}_1 = (a_{11}, a_{12}, \dots, a_{1n})$,
 $\vec{v}_2 = (a_{21}, a_{22}, \dots, a_{2n})$, \dots , $\vec{v}_m = (a_{m1}, a_{m2}, \dots, a_{mn})$ such that

$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}$. If $\text{rank}(A) = r$, then r vectors
among m vectors are linearly independent.

Linear Dependence and Independence of Vectors

Theorem

Let V a vector space, and for $m \leq n$, $\vec{v}_1 = (a_{11}, a_{12}, \dots, a_{1n})$,
 $\vec{v}_2 = (a_{21}, a_{22}, \dots, a_{2n})$, \dots , $\vec{v}_m = (a_{m1}, a_{m2}, \dots, a_{mn})$ such that

$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}$. If $\text{rank}(A) = r$, then r vectors
among m vectors are linearly independent.

1. If $r < m$, then the remaining $m - r$ vectors can be written as a linear combination of these r vectors.

Linear Dependence and Independence of Vectors

Theorem

Let V a vector space, and for $m \leq n$, $\vec{v}_1 = (a_{11}, a_{12}, \dots, a_{1n})$,
 $\vec{v}_2 = (a_{21}, a_{22}, \dots, a_{2n})$, \dots , $\vec{v}_m = (a_{m1}, a_{m2}, \dots, a_{mn})$ such that

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}. \text{ If } \text{rank}(A) = r, \text{ then } r \text{ vectors}$$

among m vectors are linearly independent.

1. If $r < m$, then the remaining $m - r$ vectors can be written as a linear combination of these r vectors.
2. If $n = m$, then m vectors are linearly independent $\Leftrightarrow |A| \neq 0$.

Linear Dependence and Independence of Vectors

Theorem

Let V a vector space, and for $m \leq n$, $\vec{v}_1 = (a_{11}, a_{12}, \dots, a_{1n})$,
 $\vec{v}_2 = (a_{21}, a_{22}, \dots, a_{2n})$, \dots , $\vec{v}_m = (a_{m1}, a_{m2}, \dots, a_{mn})$ such that

$$A = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}. \text{ If } \text{rank}(A) = r, \text{ then } r \text{ vectors}$$

among m vectors are linearly independent.

1. If $r < m$, then the remaining $m - r$ vectors can be written as a linear combination of these r vectors.
2. If $n = m$, then m vectors are linearly independent $\Leftrightarrow |A| \neq 0$.
3. If $n < m$, then $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m$ are linearly dependent.

Linear Dependence and Independence of Vectors

Example

Let $\vec{\mathbf{a}} = (1, 0, 0, 1)$, $\vec{\mathbf{b}} = (0, -1, 2, 1)$, $\vec{\mathbf{c}} = (1, 2, 2, 1)$ and $\vec{\mathbf{d}} = (-2, 1, 0, 0) \in \mathbb{R}^4$ are given. Then

Linear Dependence and Independence of Vectors

Example

Let $\vec{a} = (1, 0, 0, 1)$, $\vec{b} = (0, -1, 2, 1)$, $\vec{c} = (1, 2, 2, 1)$ and $\vec{d} = (-2, 1, 0, 0) \in \mathbb{R}^4$ are given. Then

1. Determine, whether or not the vectors \vec{a} , \vec{b} , \vec{c} , \vec{d} are linearly independent or dependent?

$$\begin{array}{cccc} 1 & 0 & 0 & 1 \\ 0 & 1 & -2 & -1 \\ 1 & 2 & 2 & 1 \\ -2 & 1 & 0 & 0 \end{array}$$

$$\begin{array}{cccc} 1 & 0 & 0 & 1 \\ 0 & 1 & -2 & -1 \\ 0 & 2 & 2 & 0 \\ 0 & 1 & 0 & 2 \end{array}$$

$$\begin{array}{cccc} 1 & 0 & 0 & 1 \\ 0 & 1 & -2 & -1 \\ 0 & 0 & 1 & \frac{1}{3} \\ 0 & 0 & 0 & 1 \end{array}$$

Linear Dependence and Independence of Vectors

Example

Let $\vec{a} = (1, 0, 0, 1)$, $\vec{b} = (0, -1, 2, 1)$, $\vec{c} = (1, 2, 2, 1)$ and $\vec{d} = (-2, 1, 0, 0) \in \mathbb{R}^4$ are given. Then

1. Determine, whether or not the vectors \vec{a} , \vec{b} , \vec{c} , \vec{d} are linearly independent or dependent?
2. Express $\vec{u} = (1, -1, 2, 1)$ as a linear combination of \vec{a} , \vec{b} , \vec{c} , \vec{d} .

Linear Dependence and Independence of Vectors

Solution (1)

$$c_1 \vec{a} + c_2 \vec{b} + c_3 \vec{c} + c_4 \vec{d} = (0, 0, 0, 0)$$

Linear Dependence and Independence of Vectors

Solution (1)

$$c_1 \vec{a} + c_2 \vec{b} + c_3 \vec{c} + c_4 \vec{d} = (0, 0, 0, 0)$$

$$\begin{aligned} \Rightarrow \quad & c_1 \quad \quad \quad + c_3 \quad - 2c_4 = 0 \\ & \quad - c_2 \quad + 2c_3 \quad + c_4 = 0 \\ & \quad \quad 2c_2 \quad + 2c_3 \quad \quad = 0 \\ & c_1 \quad + c_2 \quad + c_3 \quad \quad = 0 \end{aligned}$$

Linear Dependence and Independence of Vectors

Solution (1)

$$c_1 \vec{a} + c_2 \vec{b} + c_3 \vec{c} + c_4 \vec{d} = (0, 0, 0, 0)$$

$$\Rightarrow \begin{array}{rrrr} c_1 & & +c_3 & -2c_4 & = 0 \\ & -c_2 & +2c_3 & +c_4 & = 0 \\ & 2c_2 & +2c_3 & & = 0 \\ c_1 & +c_2 & +c_3 & & = 0 \end{array} \Rightarrow A = \begin{pmatrix} 1 & 0 & 1 & -2 \\ 0 & -1 & 2 & 1 \\ 0 & 2 & 2 & 0 \\ 1 & 1 & 1 & 0 \end{pmatrix}$$

Linear Dependence and Independence of Vectors

Solution (1)

$$c_1 \vec{a} + c_2 \vec{b} + c_3 \vec{c} + c_4 \vec{d} = (0, 0, 0, 0)$$

$$\Rightarrow \begin{array}{rrcr} c_1 & & +c_3 & -2c_4 = 0 \\ & -c_2 & +2c_3 & +c_4 = 0 \\ & 2c_2 & +2c_3 & = 0 \\ c_1 & +c_2 & +c_3 & = 0 \end{array} \Rightarrow A = \begin{pmatrix} 1 & 0 & 1 & -2 \\ 0 & -1 & 2 & 1 \\ 0 & 2 & 2 & 0 \\ 1 & 1 & 1 & 0 \end{pmatrix}$$

$$\sim \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Linear Dependence and Independence of Vectors

Solution (1)

$$c_1 \vec{a} + c_2 \vec{b} + c_3 \vec{c} + c_4 \vec{d} = (0, 0, 0, 0)$$

$$\Rightarrow \begin{array}{rrcr} c_1 & & +c_3 & -2c_4 = 0 \\ & -c_2 & +2c_3 & +c_4 = 0 \\ & 2c_2 & +2c_3 & = 0 \\ c_1 & +c_2 & +c_3 & = 0 \end{array} \Rightarrow A = \begin{pmatrix} 1 & 0 & 1 & -2 \\ 0 & -1 & 2 & 1 \\ 0 & 2 & 2 & 0 \\ 1 & 1 & 1 & 0 \end{pmatrix}$$

$$\sim \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \Rightarrow c_1 = c_2 = c_3 = c_4 = 0$$

Linear Dependence and Independence of Vectors

Solution (1)

$$c_1 \vec{a} + c_2 \vec{b} + c_3 \vec{c} + c_4 \vec{d} = (0, 0, 0, 0)$$

$$\Rightarrow \begin{array}{rrcr} c_1 & & +c_3 & -2c_4 = 0 \\ & -c_2 & +2c_3 & +c_4 = 0 \\ & 2c_2 & +2c_3 & = 0 \\ c_1 & +c_2 & +c_3 & = 0 \end{array} \Rightarrow A = \begin{pmatrix} 1 & 0 & 1 & -2 \\ 0 & -1 & 2 & 1 \\ 0 & 2 & 2 & 0 \\ 1 & 1 & 1 & 0 \end{pmatrix}$$

$$\sim \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \Rightarrow c_1 = c_2 = c_3 = c_4 = 0$$

is the unique solution of the system. Hence \vec{a} , \vec{b} , \vec{c} , \vec{d} are linearly independent.

Linear Dependence and Independence of Vectors

Solution (2)

$$c_1 \vec{a} + c_2 \vec{b} + c_3 \vec{c} + c_4 \vec{d} = (1, -1, 2, 1)$$

Linear Dependence and Independence of Vectors

Solution (2)

$$c_1 \vec{a} + c_2 \vec{b} + c_3 \vec{c} + c_4 \vec{d} = (1, -1, 2, 1)$$

$$\Rightarrow \begin{array}{rcccc} c_1 & & +c_3 & -2c_4 & = 1 \\ & -c_2 & +2c_3 & +c_4 & = -1 \\ & 2c_2 & +2c_3 & & = 2 \\ c_1 & +c_2 & +c_3 & & = 1 \end{array}$$

Linear Dependence and Independence of Vectors

Solution (2)

$$c_1 \vec{a} + c_2 \vec{b} + c_3 \vec{c} + c_4 \vec{d} = (1, -1, 2, 1)$$

$$\Rightarrow \begin{array}{rrrrr} c_1 & & +c_3 & -2c_4 & = 1 \\ & -c_2 & +2c_3 & +c_4 & = -1 \\ & 2c_2 & +2c_3 & & = 2 \\ c_1 & +c_2 & +c_3 & & = 1 \end{array}$$

$$\Rightarrow [A|\mathbf{b}] = \begin{pmatrix} 1 & 0 & 1 & -2 & 1 \\ 0 & -1 & 2 & 1 & -1 \\ 0 & 2 & 2 & 0 & 2 \\ 1 & 1 & 1 & 0 & 1 \end{pmatrix}$$

Linear Dependence and Independence of Vectors

Solution (2)

$$c_1 \vec{a} + c_2 \vec{b} + c_3 \vec{c} + c_4 \vec{d} = (1, -1, 2, 1)$$

$$\Rightarrow \begin{array}{rrrr} c_1 & & +c_3 & -2c_4 & = 1 \\ & -c_2 & +2c_3 & +c_4 & = -1 \\ & 2c_2 & +2c_3 & & = 2 \\ c_1 & +c_2 & +c_3 & & = 1 \end{array}$$

$$\Rightarrow [A|\mathbf{b}] = \begin{pmatrix} 1 & 0 & 1 & -2 & 1 \\ 0 & -1 & 2 & 1 & -1 \\ 0 & 2 & 2 & 0 & 2 \\ 1 & 1 & 1 & 0 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & \frac{6}{7} \\ 0 & 0 & 1 & 0 & \frac{1}{7} \\ 0 & 0 & 0 & 1 & -\frac{3}{7} \end{pmatrix}$$

$$\Rightarrow c_1 = 0, c_2 = \frac{6}{7}, c_3 = \frac{1}{7}, c_4 = -\frac{3}{7}$$

Linear Dependence and Independence of Vectors

Solution (2)

$$c_1 \vec{a} + c_2 \vec{b} + c_3 \vec{c} + c_4 \vec{d} = (1, -1, 2, 1)$$

$$\Rightarrow \begin{array}{rrrr} c_1 & & +c_3 & -2c_4 & = 1 \\ & -c_2 & +2c_3 & +c_4 & = -1 \\ & 2c_2 & +2c_3 & & = 2 \\ c_1 & +c_2 & +c_3 & & = 1 \end{array}$$

$$\Rightarrow [A|\mathbf{b}] = \begin{pmatrix} 1 & 0 & 1 & -2 & 1 \\ 0 & -1 & 2 & 1 & -1 \\ 0 & 2 & 2 & 0 & 2 \\ 1 & 1 & 1 & 0 & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & \frac{6}{7} \\ 0 & 0 & 1 & 0 & \frac{1}{7} \\ 0 & 0 & 0 & 1 & -\frac{3}{7} \end{pmatrix}$$

$$\Rightarrow c_1 = 0, c_2 = \frac{6}{7}, c_3 = \frac{1}{7}, c_4 = -\frac{3}{7}$$

So we have $\frac{6}{7} \vec{b} + \frac{1}{7} \vec{c} - \frac{3}{7} \vec{d} = (1, -1, 2, 1)$.

Spanning Sets

Spanning Sets

Definition

Let $S = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_r\} \subset V$ be given. The set spanned by S is denoted by $\text{span}(S)$ or $\langle S \rangle$ and defined as the set of possible all linear combinations of S .

Spanning Sets

Definition

Let $S = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_r\} \subset V$ be given. The set spanned by S is denoted by $\text{span}(S)$ or $\langle S \rangle$ and defined as the set of possible all linear combinations of S . That is,

$$\text{span}(S) = \{k_1 \vec{v}_1 + k_2 \vec{v}_2 + \dots + k_r \vec{v}_r \mid k_1, k_2, \dots, k_r \in \mathbb{R}\}.$$

Spanning Sets

Definition

Let $S = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_r\} \subset V$ be given. The set spanned by S is denoted by $\text{span}(S)$ or $\langle S \rangle$ and defined as the set of possible all linear combinations of S . That is,

$$\text{span}(S) = \{k_1 \vec{v}_1 + k_2 \vec{v}_2 + \dots + k_r \vec{v}_r \mid k_1, k_2, \dots, k_r \in \mathbb{R}\}.$$

Note: $\text{span}(S)$ is a subspace of V .

Spanning Sets

Definition

Let $S = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_r\} \subset V$ be given. The set spanned by S is denoted by $\text{span}(S)$ or $\langle S \rangle$ and defined as the set of possible all linear combinations of S . That is,

$$\text{span}(S) = \{k_1 \vec{v}_1 + k_2 \vec{v}_2 + \dots + k_r \vec{v}_r \mid k_1, k_2, \dots, k_r \in \mathbb{R}\}.$$

Note: $\text{span}(S)$ is a subspace of V .

Example

The spanning set of the vector $\vec{u} = (1, -2, 1) \in \mathbb{R}^3$ is

Spanning Sets

Definition

Let $S = \{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_r\} \subset V$ be given. The set spanned by S is denoted by $\text{span}(S)$ or $\langle S \rangle$ and defined as the set of possible all linear combinations of S . That is,

$$\text{span}(S) = \{k_1 \vec{v}_1 + k_2 \vec{v}_2 + \dots + k_r \vec{v}_r \mid k_1, k_2, \dots, k_r \in \mathbb{R}\}.$$

Note: $\text{span}(S)$ is a subspace of V .

Example

The spanning set of the vector $\vec{u} = (1, -2, 1) \in \mathbb{R}^3$ is

$$\text{span}(\{(1, -2, 1)\}) = \{k(1, -2, 1) \mid k \in \mathbb{R}\}.$$

Example

The spanning set of the set

$$S = \{ \vec{u} = (1, -2, 1, 3), \vec{v} = (0, 2, -1, 0) \} \subset \mathbb{R}^4 \text{ is}$$

Example

The spanning set of the set

$$S = \{ \vec{u} = (1, -2, 1, 3), \vec{v} = (0, 2, -1, 0) \} \subset \mathbb{R}^4 \text{ is}$$

$$\text{span}(\{(1, -2, 1, 3), (0, 2, -1, 0)\})$$

$$= \{ a(1, -2, 1, 3) + b(0, 2, -1, 0) \mid a, b \in \mathbb{R} \}.$$

?