



MAT1320-Linear Algebra

Lecture Notes

Cramer's Rule, Inverse Matrix Method

Mehmet E. KÖROĞLU
Summer 2020

YILDIZ TECHNICAL UNIVERSITY, DEPARTMENT OF MATHEMATICS
mkoroglu@yildiz.edu.tr

Table of contents

1. Cramer's Rule

2. Inverse Matrix Method

Cramer's Rule

Cramer's Rule

Let

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \vdots \\ a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nn}x_n = b_n \end{cases} \quad (1)$$

be a linear system of n equations with n unknowns.

Cramer's Rule

Let

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n = b_2 \\ \vdots \\ a_{n1}x_1 + a_{n2}x_2 + \cdots + a_{nn}x_n = b_n \end{cases} \quad (1)$$

be a linear system of n equations with n unknowns. Let denote the coefficient matrix of the system by $A = [a_{ij}]$.

Cramer's Rule

Then for the vector of unknowns $\mathbf{X} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$ and the vector of

constants $\mathbf{b} = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$, the system given in (1) can be written as

$$A\mathbf{X} = \mathbf{b}.$$

Cramer's Rule

Then for the vector of unknowns $\mathbf{X} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$ and the vector of constants $\mathbf{b} = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$, the system given in (1) can be written as

$$A\mathbf{X} = \mathbf{b}.$$

The system $A\mathbf{X} = \mathbf{b}$ has a unique solution if and only if $\det(A) \neq 0$.

Cramer's Rule

Then for the vector of unknowns $\mathbf{X} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$ and the vector of constants $\mathbf{b} = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$, the system given in (1) can be written as

$$A\mathbf{X} = \mathbf{b}.$$

The system $A\mathbf{X} = \mathbf{b}$ has a unique solution if and only if $\det(A) \neq 0$. In this case the solution is given by

Cramer's Rule

Then for the vector of unknowns $\mathbf{X} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$ and the vector of constants $\mathbf{b} = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$, the system given in (1) can be written as

$$A\mathbf{X} = \mathbf{b}.$$

The system $A\mathbf{X} = \mathbf{b}$ has a unique solution if and only if $\det(A) \neq 0$. In this case the solution is given by

$$x_1 = \frac{\det(A_1)}{\det(A)},$$

Cramer's Rule

Then for the vector of unknowns $\mathbf{X} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$ and the vector of constants $\mathbf{b} = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$, the system given in (1) can be written as

$$A\mathbf{X} = \mathbf{b}.$$

The system $A\mathbf{X} = \mathbf{b}$ has a unique solution if and only if $\det(A) \neq 0$. In this case the solution is given by

$$x_1 = \frac{\det(A_1)}{\det(A)}, \quad x_2 = \frac{\det(A_2)}{\det(A)}$$

Cramer's Rule

Then for the vector of unknowns $\mathbf{X} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$ and the vector of constants $\mathbf{b} = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$, the system given in (1) can be written as

$$A\mathbf{X} = \mathbf{b}.$$

The system $A\mathbf{X} = \mathbf{b}$ has a unique solution if and only if $\det(A) \neq 0$. In this case the solution is given by

$$x_1 = \frac{\det(A_1)}{\det(A)}, \quad x_2 = \frac{\det(A_2)}{\det(A)}, \quad \dots, \quad x_n = \frac{\det(A_n)}{\det(A)},$$

Cramer's Rule

Then for the vector of unknowns $\mathbf{X} = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix}$ and the vector of

constants $\mathbf{b} = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix}$, the system given in (1) can be written as

$$A\mathbf{X} = \mathbf{b}.$$

The system $A\mathbf{X} = \mathbf{b}$ has a unique solution if and only if $\det(A) \neq 0$. In this case the solution is given by

$$x_1 = \frac{\det(A_1)}{\det(A)}, \quad x_2 = \frac{\det(A_2)}{\det(A)}, \quad \dots, \quad x_n = \frac{\det(A_n)}{\det(A)},$$

where A_i ($i = 1, 2, \dots, n$) is the matrix obtained from A by replacing i -th column with the vector of constants \mathbf{b} .

Example (Cramer's Rule)

By using Cramer's Rule solve the system

$$\begin{cases} 2x_1 + 3x_2 + x_3 = 1 \\ x_1 - x_2 + 2x_3 = -3 \\ -3x_1 + 4x_2 - x_3 = 4 \end{cases}$$

Example (Cramer's Rule)

By using Cramer's Rule solve the system

$$\begin{cases} 2x_1 + 3x_2 + x_3 = 1 \\ x_1 - x_2 + 2x_3 = -3 \\ -3x_1 + 4x_2 - x_3 = 4 \end{cases}$$

Solution

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 1 & -1 & 2 \\ -3 & 4 & -1 \end{bmatrix},$$

Example (Cramer's Rule)

By using Cramer's Rule solve the system

$$\begin{cases} 2x_1 + 3x_2 + x_3 = 1 \\ x_1 - x_2 + 2x_3 = -3 \\ -3x_1 + 4x_2 - x_3 = 4 \end{cases}$$

Solution

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 1 & -1 & 2 \\ -3 & 4 & -1 \end{bmatrix}, \quad A_1 = \begin{bmatrix} 1 & 3 & 1 \\ -3 & -1 & 2 \\ 4 & 4 & -1 \end{bmatrix},$$

Example (Cramer's Rule)

By using Cramer's Rule solve the system

$$\begin{cases} 2x_1 + 3x_2 + x_3 = 1 \\ x_1 - x_2 + 2x_3 = -3 \\ -3x_1 + 4x_2 - x_3 = 4 \end{cases}$$

Solution

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 1 & -1 & 2 \\ -3 & 4 & -1 \end{bmatrix}, \quad A_1 = \begin{bmatrix} 1 & 3 & 1 \\ -3 & -1 & 2 \\ 4 & 4 & -1 \end{bmatrix},$$

$$A_2 = \begin{bmatrix} 2 & 1 & 1 \\ 1 & -3 & 2 \\ -3 & 4 & -1 \end{bmatrix},$$

Example (Cramer's Rule)

By using Cramer's Rule solve the system

$$\begin{cases} 2x_1 + 3x_2 + x_3 = 1 \\ x_1 - x_2 + 2x_3 = -3 \\ -3x_1 + 4x_2 - x_3 = 4 \end{cases}$$

Solution

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 1 & -1 & 2 \\ -3 & 4 & -1 \end{bmatrix}, \quad A_1 = \begin{bmatrix} 1 & 3 & 1 \\ -3 & -1 & 2 \\ 4 & 4 & -1 \end{bmatrix},$$

$$A_2 = \begin{bmatrix} 2 & 1 & 1 \\ 1 & -3 & 2 \\ -3 & 4 & -1 \end{bmatrix}, \quad A_3 = \begin{bmatrix} 2 & 3 & 1 \\ 1 & -1 & -3 \\ -3 & 4 & 4 \end{bmatrix}$$

Solution (cont.)

$$\det(A) = -28,$$

Solution (cont.)

$$\det(A) = -28, \quad \det(A_1) = 0$$

Solution (cont.)

$$\begin{aligned}\det(A) &= -28, & \det(A_1) &= 0 \\ \det(A_2) &= -20,\end{aligned}$$

Solution (cont.)

$$\det(A) = -28, \quad \det(A_1) = 0$$

$$\det(A_2) = -20, \quad \det(A_3) = 32$$

Solution (cont.)

$$\begin{aligned}\det(A) &= -28, & \det(A_1) &= 0 \\ \det(A_2) &= -20, & \det(A_3) &= 32\end{aligned}$$

$$x_1 = \frac{\det(A_1)}{\det(A)} = 0$$

Solution (cont.)

$$\begin{aligned}\det(A) &= -28, & \det(A_1) &= 0 \\ \det(A_2) &= -20, & \det(A_3) &= 32\end{aligned}$$

$$\begin{aligned}x_1 &= \frac{\det(A_1)}{\det(A)} = 0 \\ x_2 &= \frac{\det(A_2)}{\det(A)} = \frac{5}{7}\end{aligned}$$

Solution (cont.)

$$\begin{aligned}\det(A) &= -28, & \det(A_1) &= 0 \\ \det(A_2) &= -20, & \det(A_3) &= 32\end{aligned}$$

$$\begin{aligned}x_1 &= \frac{\det(A_1)}{\det(A)} = 0 \\ x_2 &= \frac{\det(A_2)}{\det(A)} = \frac{5}{7} \\ x_3 &= \frac{\det(A_3)}{\det(A)} = \frac{-8}{7}\end{aligned}$$

Inverse Matrix Method

Let $A\mathbf{X} = \mathbf{b}$ be matrix representation of the linear system of equations given in (1).

Inverse Matrix Method

Let $A\mathbf{X} = \mathbf{b}$ be matrix representation of the linear system of equations given in (1). If $\det(A) \neq 0$, then A^{-1} exists and the system has a solution.

Inverse Matrix Method

Let $A\mathbf{X} = \mathbf{b}$ be matrix representation of the linear system of equations given in (1). If $\det(A) \neq 0$, then A^{-1} exists and the system has a solution. In this case the solution is given by

$$A^{-1}(A\mathbf{X}) = A^{-1}\mathbf{b}$$

Inverse Matrix Method

Let $A\mathbf{X} = \mathbf{b}$ be matrix representation of the linear system of equations given in (1). If $\det(A) \neq 0$, then A^{-1} exists and the system has a solution. In this case the solution is given by

$$\begin{aligned} A^{-1}(A\mathbf{X}) &= A^{-1}\mathbf{b} \\ \underbrace{(A^{-1}A)}_{I_n} \mathbf{X} &= A^{-1}\mathbf{b} \end{aligned}$$

Inverse Matrix Method

Let $A\mathbf{X} = \mathbf{b}$ be matrix representation of the linear system of equations given in (1). If $\det(A) \neq 0$, then A^{-1} exists and the system has a solution. In this case the solution is given by

$$\begin{aligned} A^{-1}(A\mathbf{X}) &= A^{-1}\mathbf{b} \\ \underbrace{(A^{-1}A)}_{I_n} \mathbf{X} &= A^{-1}\mathbf{b} \\ \Rightarrow \mathbf{X} &= A^{-1}\mathbf{b}. \end{aligned}$$

Example (Inverse Matrix Method)

By using Inverse Matrix Method solve the system

$$\begin{cases} 2x_1 + 3x_2 + x_3 = 1 \\ x_1 - x_2 + 2x_3 = -3 \\ -3x_1 + 4x_2 - x_3 = 4 \end{cases} .$$

Example (Inverse Matrix Method)

By using Inverse Matrix Method solve the system

$$\begin{cases} 2x_1 + 3x_2 + x_3 = 1 \\ x_1 - x_2 + 2x_3 = -3 \\ -3x_1 + 4x_2 - x_3 = 4 \end{cases} .$$

Solution

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 1 & -1 & 2 \\ -3 & 4 & -1 \end{bmatrix}$$

Example (Inverse Matrix Method)

By using Inverse Matrix Method solve the system

$$\begin{cases} 2x_1 + 3x_2 + x_3 = 1 \\ x_1 - x_2 + 2x_3 = -3 \\ -3x_1 + 4x_2 - x_3 = 4 \end{cases}.$$

Solution

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 1 & -1 & 2 \\ -3 & 4 & -1 \end{bmatrix} \Rightarrow A^{-1} = \begin{bmatrix} \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{5}{28} & -\frac{1}{28} & \frac{3}{28} \\ -\frac{1}{28} & \frac{17}{28} & \frac{5}{28} \end{bmatrix}$$

Example (Inverse Matrix Method)

By using Inverse Matrix Method solve the system

$$\begin{cases} 2x_1 + 3x_2 + x_3 = 1 \\ x_1 - x_2 + 2x_3 = -3 \\ -3x_1 + 4x_2 - x_3 = 4 \end{cases}.$$

Solution

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 1 & -1 & 2 \\ -3 & 4 & -1 \end{bmatrix} \Rightarrow A^{-1} = \begin{bmatrix} \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{5}{28} & -\frac{1}{28} & \frac{3}{28} \\ -\frac{1}{28} & \frac{17}{28} & \frac{5}{28} \end{bmatrix}$$

$$\mathbf{X} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$$

Example (Inverse Matrix Method)

By using Inverse Matrix Method solve the system

$$\begin{cases} 2x_1 + 3x_2 + x_3 = 1 \\ x_1 - x_2 + 2x_3 = -3 \\ -3x_1 + 4x_2 - x_3 = 4 \end{cases}.$$

Solution

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 1 & -1 & 2 \\ -3 & 4 & -1 \end{bmatrix} \Rightarrow A^{-1} = \begin{bmatrix} \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{5}{28} & -\frac{1}{28} & \frac{3}{28} \\ -\frac{1}{28} & \frac{17}{28} & \frac{5}{28} \end{bmatrix}$$

$$\mathbf{X} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = A^{-1}\mathbf{b}$$

Example (Inverse Matrix Method)

By using Inverse Matrix Method solve the system

$$\begin{cases} 2x_1 + 3x_2 + x_3 = 1 \\ x_1 - x_2 + 2x_3 = -3 \\ -3x_1 + 4x_2 - x_3 = 4 \end{cases}.$$

Solution

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 1 & -1 & 2 \\ -3 & 4 & -1 \end{bmatrix} \Rightarrow A^{-1} = \begin{bmatrix} \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{5}{28} & -\frac{1}{28} & \frac{3}{28} \\ -\frac{1}{28} & \frac{17}{28} & \frac{5}{28} \end{bmatrix}$$

$$\mathbf{X} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = A^{-1}\mathbf{b} = \begin{bmatrix} \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{5}{28} & -\frac{1}{28} & \frac{3}{28} \\ -\frac{1}{28} & \frac{17}{28} & \frac{5}{28} \end{bmatrix} \begin{bmatrix} 1 \\ -3 \\ 4 \end{bmatrix}$$

Example (Inverse Matrix Method)

By using Inverse Matrix Method solve the system

$$\begin{cases} 2x_1 + 3x_2 + x_3 = 1 \\ x_1 - x_2 + 2x_3 = -3 \\ -3x_1 + 4x_2 - x_3 = 4 \end{cases}.$$

Solution

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 1 & -1 & 2 \\ -3 & 4 & -1 \end{bmatrix} \Rightarrow A^{-1} = \begin{bmatrix} \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{5}{28} & -\frac{1}{28} & \frac{3}{28} \\ -\frac{1}{28} & \frac{17}{28} & \frac{5}{28} \end{bmatrix}$$

$$\mathbf{X} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = A^{-1}\mathbf{b} = \begin{bmatrix} \frac{1}{4} & -\frac{1}{4} & -\frac{1}{4} \\ \frac{5}{28} & -\frac{1}{28} & \frac{3}{28} \\ -\frac{1}{28} & \frac{17}{28} & \frac{5}{28} \end{bmatrix} \begin{bmatrix} 1 \\ -3 \\ 4 \end{bmatrix} = \begin{bmatrix} 0 \\ \frac{5}{7} \\ -\frac{8}{7} \end{bmatrix}$$

Example

Let the determinant of the matrix $\mathbf{F} = \begin{bmatrix} a & a^2 & a^3 & a^4 \\ b & b^2 & b^3 & b^4 \\ c & c^2 & c^3 & c^4 \\ d & d^2 & d^3 & d^4 \end{bmatrix}$ be $|\mathbf{F}|$ such that $|\mathbf{F}| \neq 0$.

Example

Let the determinant of the matrix $\mathbf{F} = \begin{bmatrix} a & a^2 & a^3 & a^4 \\ b & b^2 & b^3 & b^4 \\ c & c^2 & c^3 & c^4 \\ d & d^2 & d^3 & d^4 \end{bmatrix}$ be $|\mathbf{F}|$

such that $|\mathbf{F}| \neq 0$.

1. By using properties of determinants, find $|\mathbf{F}|$.

Example

Let the determinant of the matrix $\mathbf{F} = \begin{bmatrix} a & a^2 & a^3 & a^4 \\ b & b^2 & b^3 & b^4 \\ c & c^2 & c^3 & c^4 \\ d & d^2 & d^3 & d^4 \end{bmatrix}$ be $|\mathbf{F}|$

such that $|\mathbf{F}| \neq 0$.

1. By using properties of determinants, find $|\mathbf{F}|$.

2. For $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix}$ and $\mathbf{b} = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix}$ solve $\mathbf{F}\mathbf{x} = \mathbf{b}$ by Cramer's

Rule.

Solution (1)

$$abcd \begin{vmatrix} 1 & a & a^2 & a^3 \\ 1 & b & b^2 & b^3 \\ 1 & c & c^2 & c^3 \\ 1 & d & d^2 & d^3 \end{vmatrix} \begin{matrix} H_{21}(-1) \\ H_{31}(-1) \\ H_{41}(-1) \\ = \end{matrix}$$

Solution (1)

$$abcd \begin{vmatrix} 1 & a & a^2 & a^3 \\ 1 & b & b^2 & b^3 \\ 1 & c & c^2 & c^3 \\ 1 & d & d^2 & d^3 \end{vmatrix} \begin{matrix} H_{21}(-1) \\ H_{31}(-1) \\ H_{41}(-1) \\ = \end{matrix}$$

$$abcd \begin{vmatrix} 1 & a & a^2 & a^3 \\ 0 & b-a & b^2-a^2 & b^3-a^3 \\ 0 & c-a & c^2-a^2 & c^3-a^3 \\ 0 & d-a & d^2-a^2 & d^3-a^3 \end{vmatrix}$$

Solution (1)

$$\begin{aligned}
 & abcd \begin{vmatrix} 1 & a & a^2 & a^3 \\ 1 & b & b^2 & b^3 \\ 1 & c & c^2 & c^3 \\ 1 & d & d^2 & d^3 \end{vmatrix} \begin{matrix} H_{21}(-1) \\ H_{31}(-1) \\ H_{41}(-1) \\ = \end{matrix} \\
 & abcd \begin{vmatrix} 1 & a & a^2 & a^3 \\ 0 & b-a & b^2-a^2 & b^3-a^3 \\ 0 & c-a & c^2-a^2 & c^3-a^3 \\ 0 & d-a & d^2-a^2 & d^3-a^3 \end{vmatrix} \\
 & = abcd \begin{vmatrix} b-a & b^2-a^2 & b^3-a^3 \\ c-a & c^2-a^2 & c^3-a^3 \\ d-a & d^2-a^2 & d^3-a^3 \end{vmatrix}
 \end{aligned}$$

Solution (1 cont.)

$$= \underbrace{abcd(b-a)(c-a)(d-a)}_r$$
$$\left| \begin{array}{ccc} 1 & b+a & b^2+ab+a^2 \\ 1 & c+a & c^2+ac+a^2 \\ 1 & d+a & d^2+ad+a^2 \end{array} \right| \begin{array}{l} H_{21}(-1) \\ H_{31}(-1) \end{array}$$

Solution (1 cont.)

$$\begin{aligned}
 &= \underbrace{abcd(b-a)(c-a)(d-a)}_r \\
 &\quad \begin{vmatrix} 1 & b+a & b^2+ab+a^2 \\ 1 & c+a & c^2+ac+a^2 \\ 1 & d+a & d^2+ad+a^2 \end{vmatrix} \begin{matrix} H_{21}(-1) \\ H_{31}(-1) \end{matrix} \\
 &= r \begin{vmatrix} 1 & b+a & b^2+ab+a^2 \\ 0 & c-b & c^2-b^2+a(c-b) \\ 0 & d-b & d^2-b^2+a(d-b) \end{vmatrix}
 \end{aligned}$$

Solution (1 cont.)

$$\begin{aligned}
 &= \underbrace{abcd(b-a)(c-a)(d-a)}_r \\
 &\quad \begin{vmatrix} 1 & b+a & b^2+ab+a^2 \\ 1 & c+a & c^2+ac+a^2 \\ 1 & d+a & d^2+ad+a^2 \end{vmatrix} \begin{matrix} H_{21}(-1) \\ H_{31}(-1) \end{matrix} \\
 &= r \begin{vmatrix} 1 & b+a & b^2+ab+a^2 \\ 0 & c-b & c^2-b^2+a(c-b) \\ 0 & d-b & d^2-b^2+a(d-b) \end{vmatrix} \\
 &= \underbrace{r(c-b)(d-b)}_s \begin{vmatrix} 1 & b+a & b^2+ab+a^2 \\ 0 & 1 & c+b+a \\ 0 & 1 & d+b+a \end{vmatrix}
 \end{aligned}$$

Solution (1 cont.)

$$= rs \begin{vmatrix} 1 & c+b+a \\ 1 & d+b+a \end{vmatrix} H_{21}(-1) = rs \begin{vmatrix} 1 & c+b+a \\ 0 & d-c \end{vmatrix}$$

Solution (1 cont.)

$$\begin{aligned} &= rs \begin{vmatrix} 1 & c+b+a \\ 1 & d+b+a \end{vmatrix} H_{21}(-1) = rs \begin{vmatrix} 1 & c+b+a \\ 0 & d-c \end{vmatrix} \\ &= abcd (b-a) (c-a) (d-a) (c-b) (d-b) (d-c). \end{aligned}$$

Solution (1 cont.)

$$\begin{aligned} &= rs \begin{vmatrix} 1 & c+b+a \\ 1 & d+b+a \end{vmatrix} H_{21}(-1) = rs \begin{vmatrix} 1 & c+b+a \\ 0 & d-c \end{vmatrix} \\ &= abcd(b-a)(c-a)(d-a)(c-b)(d-b)(d-c). \end{aligned}$$

Solution (2)

We know that $\Delta = |\mathbf{F}| \neq 0$ and Δ_i ($i = 1, 2, 3, 4$) is the determinant of the matrix obtained from \mathbf{F} by replacing i -th column with \mathbf{b} .

Solution (1 cont.)

$$\begin{aligned} &= rs \begin{vmatrix} 1 & c+b+a \\ 1 & d+b+a \end{vmatrix} H_{21}(-1) = rs \begin{vmatrix} 1 & c+b+a \\ 0 & d-c \end{vmatrix} \\ &= abcd(b-a)(c-a)(d-a)(c-b)(d-b)(d-c). \end{aligned}$$

Solution (2)

We know that $\Delta = |\mathbf{F}| \neq 0$ and Δ_i ($i = 1, 2, 3, 4$) is the determinant of the matrix obtained from \mathbf{F} by replacing i -th column with \mathbf{b} . Then $\Delta_1 = |\mathbf{F}|$

Solution (1 cont.)

$$\begin{aligned} &= rs \begin{vmatrix} 1 & c+b+a \\ 1 & d+b+a \end{vmatrix} H_{21}(-1) = rs \begin{vmatrix} 1 & c+b+a \\ 0 & d-c \end{vmatrix} \\ &= abcd(b-a)(c-a)(d-a)(c-b)(d-b)(d-c). \end{aligned}$$

Solution (2)

We know that $\Delta = |\mathbf{F}| \neq 0$ and Δ_i ($i = 1, 2, 3, 4$) is the determinant of the matrix obtained from \mathbf{F} by replacing i -th column with \mathbf{b} . Then $\Delta_1 = |\mathbf{F}|$ and $\Delta_2 = \Delta_3 = \Delta_4 = 0$.

Solution (1 cont.)

$$\begin{aligned} &= rs \begin{vmatrix} 1 & c+b+a \\ 1 & d+b+a \end{vmatrix} H_{21}(-1) = rs \begin{vmatrix} 1 & c+b+a \\ 0 & d-c \end{vmatrix} \\ &= abcd(b-a)(c-a)(d-a)(c-b)(d-b)(d-c). \end{aligned}$$

Solution (2)

We know that $\Delta = |\mathbf{F}| \neq 0$ and Δ_i ($i = 1, 2, 3, 4$) is the determinant of the matrix obtained from \mathbf{F} by replacing i -th column with \mathbf{b} . Then $\Delta_1 = |\mathbf{F}|$ and $\Delta_2 = \Delta_3 = \Delta_4 = 0$. So, we have $x_1 = 1, x_2 = x_3 = x_4 = 0$.

?