

Optimization Techniques

Section 4

M. Fatih Amasyalı

Convergence

- We have tried to find some x^k that converge to the desired point x^* (most often a local minimizer). The fundamental question is how fast the convergence is.
- Let's define errors at each iteration step as $|e_k| = |x^k - x^*|$ and $|e_{k+1}| = |x^{k+1} - x^*|$

Convergence

- Linear Convergence :

There exist a constant $0 < c_1 < 1$

where $|e_{k+1}| \leq c_1 * |e_k|$

- If c_1 is very close to 0 then superlinear convergence.

- Quadratic Convergence :

There exist a constant $0 < c_2 < 1$

where $|e_{k+1}| \leq c_2 * |e_k|^2$

Convergence Example

- Two methods are given; one of them showing linear convergence and the other one showing quadratic convergence.
- After certain number of steps errors reach 3 digit precision for both of the methods ($|e_k| < 0,001$).
- How many more steps (iterations) will be necessary if we require 12 digits of precision ?
- $c_1 = c_2 = 1/2$

Convergence Example

- In case of Linear Convergence : $c_1=0.5$

- $|e_{k+1}| \leq 0.5 * |e_k|$
- $|e_{k+2}| \leq 0.5 * |e_{k+1}|$
- $|e_{k+2}| \leq 0.5 * 0.5 * |e_k|$
- ...
- $|e_{k+n}| \leq 0.5^n * |e_k|$

- $|e_k| \leq 0.001$ (It is given.)
- $|e_{k+n}| \leq 10^{-12}$ (It is required.)
- $|e_{k+n}| \leq 0.5^n * 10^{-3}$
- $0.5^n \approx 10^{-9}$
- $\log_{10} 2^{-n} = \log_{10} 10^{-9}$
- $-n * \log_{10} 2 = -9$ ($\log_{10} 2 \approx 0.301$)

Convergence Example

- In case of quadratic Convergence : $c_2=0.5$

- $|e_{k+1}| \leq 0.5 * |e_k|^2$
- $|e_{k+2}| \leq 0.5 * |e_{k+1}|^2$
- $|e_{k+2}| \leq 0.5 * (0.5 * |e_k|^2)^2$
- $|e_{k+3}| \leq 0.5 * (0.5 * (0.5 * |e_k|^2)^2)^2$

- $|e_k| \leq 10^{-3}$ (It is given.)
- $|e_{k+n}| \leq 10^{-12}$ (It is required.)
- $|e_k|^2 = (10^{-3})^2 = 10^{-6}$
- $(|e_k|^2)^2 = 10^{-12}$
- $n = 2$

with N dimensions

- $f(x) = (1/2) * x^T * q * x + b^T * x + c$
- $f: \mathbb{R}^n \rightarrow \mathbb{R}^1$
- $q \rightarrow n * n$
- $b \rightarrow n * 1$
- $c \rightarrow 1 * 1$
- $x \rightarrow n * 1$
-
- $df = q * x + b$
- $ddf = q$
- `opt_Ndim_v2.m`

To avoid local minimums

- Random restart
- Non Derivative Techniques

References

- <http://math.tutorvista.com/calculus/newton-raphson-method.html>
- <http://math.tutorvista.com/calculus/linear-approximation.html>
- http://en.wikipedia.org/wiki/Newton's_method
- http://en.wikipedia.org/wiki/Steepest_descent
- http://www.pitt.edu/~nak54/Unconstrained_Optimization_KN.pdf
- <http://mathworld.wolfram.com/MatrixInverse.html>
- <http://ipsa.swarthmore.edu/BackGround/RevMat/MatrixReview.html>
- <http://www.cut-the-knot.org/arithmetic/algebra/Determinant.shtml>
- Matematik Dünyası, MD 2014-II, Determinantlar
- http://www.sharetechnote.com/html/EngMath_Matrix_Main.html
- Advanced Engineering Mathematics , Erwin Kreyszig, 10th Edition, John Wiley & Sons, 2011
- http://en.wikipedia.org/wiki/Finite_difference
- http://ocw.usu.edu/Civil_and_Environmental_Engineering/Numerical_Methods_in_Civil_Engineering/NonLinearEquationsMatlab.pdf
- http://www-math.mit.edu/~djik/calculus_beginners/chapter09/section02.html
- <http://stanford.edu/class/ee364a/lectures/intro.pdf>