

ALF Tutorial

The *ALF* (Algorithms for Lattice Fermions) project release 2.0 tutorial

Florian Goth, Johannes S. Hofmann, Jonas Schwab,
Jefferson S. E. Portela, Fakher F. Assaad

August 17, 2020

The ALF package provides a general code for auxiliary-field Quantum Monte Carlo simulations and default analysis. In this tutorial we show how users from beginners to specialists can profit from ALF. This document is divided in two parts:

Part I. The first, introductory part of the tutorial is based on ALF's python interface – **pyALF** – which greatly simplifies using the code, making it ideal for: obtaining *benchmark* results for established models; *getting started* with QMC and ALF; or just *quickly* running a simulation.

Part II. The second part is independent of the first and aimed at more advanced users who want to simulate their own systems. It guides the user on how to modify the package's Fortran source code and presents the resources implemented to facilitate this task.

This document is intended to be self-contained, but the interested reader should check [ALF's documentation](#), which contains a thorough, systematic description of the package. [\[Update link when ALF 2 is open.\]](#)

Part I. Just run it

This a collection of Jupyter notebooks, each centered on a detailed example followed by a few simple exercises, while Part II consists in a set of guided advanced exercises.

can be found, together with its accompanying files, in the [pyAL repository](#).

ALF is compiled from source, which is downloaded from the [ALF repository](#) when not found locally.




Figure 1: $\Delta \tau t$ extrapolation for the 4-site Hubbard ring.

Part II. Getting your hands dirty - changing the code

A lot already comes implemented in ALF, but unavoidably, as one proceeds in their own investigations, a new model has to be implemented or a new observable defined – and for that one has to grapple with the package’s Fortran source code. However, in ALF this is made easy by means of predefined structures, templates, and the examples below.

Downloading the code and tutorial

To download the code, type `git clone git@git.physik.uni-wuerzburg.de:ALF/ALF_code.git` in a shell.

To download the tutorial including solutions type:

`git clone git@git.physik.uni-wuerzburg.de:ALF/ALF_Tutorial.git` again in a shell.

Exercise 1 – Dimensional crossover

[To be updated.]

1a) Modifying the hopping

Here we will modify the code so as to allow for different hopping matrix elements along the x and y directions of a square lattice. To do so, one merely has to do the following

- Add an extra variable, `Ham_Ty`, in the parameter file in the `VAR_Hubbard` name space
- Declare the variable `Ham_Ty` in the `Hamiltonian_Examples.f90` .
- Read in this variable in `Ham_set` subroutine of the `Hamiltonian_Examples.f90` file.
- Modify the hopping matrix in the subroutine `Ham_Hop` in the `Hamiltonian_Examples.f90` file.

```
DO I = 1, Latt%N
  I1 = Latt%nnlist(I,1,0)
  I2 = Latt%nnlist(I,0,1)
  Op_T(nc,n)%O(I,I1) = cmplx(-Ham_T, 0.d0, kind(0.D0))
  Op_T(nc,n)%O(I1,I) = cmplx(-Ham_T, 0.d0, kind(0.D0))
  !!!!! Modifications for Exercise 2 (a)
  !Op_T(nc,n)%O(I,I2) = cmplx(-Ham_T, 0.d0, kind(0.D0))
  !Op_T(nc,n)%O(I2,I) = cmplx(-Ham_T, 0.d0, kind(0.D0))
  Op_T(nc,n)%O(I,I2) = cmplx(-Ham_Ty, 0.d0, kind(0.D0))
  Op_T(nc,n)%O(I2,I) = cmplx(-Ham_Ty, 0.d0, kind(0.D0))
  !!!!!
  Op_T(nc,n)%O(I ,I) = cmplx(-Ham_chem, 0.d0, kind(0.D0))
ENDDO
```

In the directory `Solutions/Exercise_2` we have duplicated the ALF and commented the changes that have to be carried out to the file `Hamiltonian_Examples.f90` in the `Prog` directory.

As an application of this code, one can consider a ladder system, defined by the parameter file:

```
!=====
!  Variables for the Hubb program
!-----

&VAR_lattice
L1   = 14
L2   = 2
```

```

Lattice_type = "Square"
Model = "Hubbard_Mz" ! Hubbard_SU2, Hubbard_Mz
/

&VAR_Hubbard
ham_T = 1.0D0
!!!!!! Modifications for Exercise 2
ham_Ty = 2.0D0
!!!!!!
ham_chem= 0.0D0
ham_U = 4.0
Beta = 10.0
dtau = 0.1
/

&VAR_QMC
Nwrap = 10
NSweep = 100
NBin = 10
Ltau = 1
LOBS_ST = 1
LOBS_EN = 100
CPU_MAX = 0.1
/

&VAR_errors
n_skip = 2
N_rebin = 1
N_Cov = 0
/
! slash terminates namelist statement - DO NOT REMOVE

```

When running the code for the above ladder system and analyzing the spin correlation functions, `SpinZ_eqJR` and `SpinXY_eqJR` one will notice that it is hard to restore the $SU(2)$ spin symmetry and that very long runs are required to obtain the desired equality

$$\langle S_i^z S_j^z \rangle = \langle S_i^y S_j^y \rangle = \langle S_i^x S_j^x \rangle. \quad (1)$$

The structure of the output files `SpinZ_eqJR` and `SpinXY_eqJR` is described in the documentation. For the Mz Hubbard-Stratonovitch transformation it is hence better to consider the improved estimator

$$\langle \vec{S}_i \cdot \vec{S}_j \rangle \quad (2)$$

to compute the spin-spin correlations.

1b) Adding a new observable

Here the aim is to include the new observable equal time observable $\langle \vec{S}_i \cdot \vec{S}_j \rangle$ in the `Hubbard_Mz` code. To achieve this, you will have to carry out the following steps.

- In the subroutine `Alloc_obs` in the `Hamiltonian_example.f90` file you will have to add a new equal time observable with a call to `Call Obser_Latt_make(Observe(I),Ns,Nt,No,Filename)` with `Ns = Latt%N`; `No = Norb`; `Filename = "SpinT"`, `Nt=1`, `I=5`
- In the subroutine `Obser` you will have to add the Wick decomposition of this observable.

In the program `Hamiltonian_Examples.f90` to be found in the directory `Solutions/Exercise_2/Prog/` we have commented the changes that have to be carried out to add this observable. The new variable takes the name `SpinT` and the results you should obtain are summarized in Fig. 2.




Figure 2: Spin correlation functions along one leg for the Hubbard ladder. As t_y grows the spin gap becomes large enough so as to detect the exponential decay of the spin correlation function on this small lattice size. The underlying physics of odd-even ladder systems is introduced in the article: Elbio Dagotto and T. M. Rice, Surprises on the way from one- to two-dimensional quantum magnets: The ladder materials, Science 271 (1996), no. 5249, 618-623.

1c) The SU(2) Hubbard-Stratonovich transformation

The SU(2) Hubbard-Stratonovich decomposition, conserves spin rotational symmetry. Run the ladder code with the SU(2) flag in the parameter file switched on (i.e. `Model = Hubbard_SU2`) and compare results.

Exercise 2 – Defining a new model: The one-dimensional t-V model

[To be updated.]

2a) Define new model

In this section, one we will show what modifications have to be carried out for computing the physics of the one dimensional t-V model of spinless fermions.

$$H = -t \sum_i \left(c_i^\dagger c_{i+a} + c_{i+a}^\dagger c_i \right) - \frac{V}{2} \sum_i \left(c_i^\dagger c_{i+a} + c_{i+a}^\dagger c_i \right)^2 \quad (3)$$

The above form is readily included in the ALF since the interaction is written in terms of a perfect square. Expanding the square yields (up to a constant) the desired model:

$$H = -t \sum_i \left(c_i^\dagger c_{i+a} + c_{i+a}^\dagger c_i \right) + V \sum_i (n_i - 1/2) (n_{i+a} - 1/2) \quad (4)$$

In the directory `Solutions/Exercise_3` we have duplicated the ALF and commented the changes that have to be carried out to the file `Hamiltonian_Examples.f90` in the `Prog` directory so as to include the `t_V` model. Here are the steps to be carried out.

- Add the $t - V$ name space in the parameter file so as to read in the appropriate variables.
- Declare new variables in the `Hamiltonian_Examples.f90` file.
- In the `Ham_set` subroutine of the file `Hamiltonian_Examples.f90` set and read in the parameters for the new model, `t_V`. For this model `NF=1` and `N_SUN=1` since we are working with spinless fermions
- In the `Ham_V` subroutine you will have to add the new interaction. For a given bond at a given time-slice, we need to decouple the interaction:

$$e^{\Delta\tau \frac{V}{2} (c_i^\dagger c_{i+a} + c_{i+a}^\dagger c_i)^2} = \sum_{l=\pm 1, \pm 2} \gamma_l e^{\sqrt{\Delta\tau \frac{V}{2}} \eta_l (c_i^\dagger c_{i+a} + c_{i+a}^\dagger c_i)} = \sum_{l=\pm 1, \pm 2} \gamma_l e^{g \eta_l (c_i^\dagger c_{i+a} + c_{i+a}^\dagger c_i)} O(c_i, c_{i+a})^T \quad (5)$$

Here is how this translates in the code.

```
Allocate(Op_V(Latt%N,N_FL))
do nf = 1,N_FL
  do i = 1, N_coord*Ndim
    call Op_make(Op_V(i,nf),2)
  enddo
enddo
Do nc = 1, Latt%N ! Runs over bonds = # of lattice sites in one-dimension.
  I1 = nc
  I2 = Latt%nnlist(I1,1,0)
  Op_V(nc,1)%P(1) = I1
  Op_V(nc,1)%P(2) = I2
  Op_V(nc,1)%O(1,2) = cmplx(1.d0 ,0.d0, kind(0.D0))
  Op_V(nc,1)%O(2,1) = cmplx(1.d0 ,0.d0, kind(0.D0))
  Op_V(nc,1)%g      = SQRT(CMPLX( DTAU*Ham_Vint/2.d0, 0.D0, kind(0.D0)))
  Op_V(nc,1)%alpha  = cmplx(0d0,0.d0, kind(0.D0))
  Op_V(nc,1)%type =2
  Call Op_set( Op_V(nc,1) )
enddo
```

- Finally you will have to update the `Obser` and `ObserT` routines for the calculation of the equal and time displaced correlations. For the `t_V` model you can essentially use the same observables as for the Hubbard_SU(2) model.

You can now run the code for various values of V/t . A Jordan-Wigner transformation will map the t - V model onto the XXZ chain:

$$H = J_{xx} \sum_i S_i^x S_{i+a}^x + S_i^y S_{i+a}^y + J_{zz} \sum_i S_i^z S_{i+a}^z \quad (6)$$

with $J_{zz} = V$ and $J_{xx} = 2t$. Hence when $V/t = 2$ we reproduce the Heisenberg model. For $V/t > 2$ the model is in the Ising regime with long-range charge density wave order and is an insulator. In the regime $-2 < V/t < 2$ the model is metallic and corresponds to a Luttinger liquid. Finally, at $V/t < -2$ phase separation between hole rich and electron rich phases occur. Fig. 3 shows typical results.




Figure 3: Density-Density correlation functions of the t - V model. In the Luttinger liquid phase, $-2 < V/t < 2$ it is known that the density-density correlations decay as $\langle n(r)n(0) \rangle \propto \cos(\pi r) r^{-(1+K_\rho)}$ with $(1 + K_\rho)^{-1} = \frac{1}{2} + \frac{1}{\pi} \arcsin\left(\frac{V}{2|t|}\right)$ (A. Luther and I. Peschel, Calculation of critical exponents in two dimensions from quantum field theory in one dimension, Phys. Rev. B 12 (1975), 3908.) The interested reader can try to reproduce this result.

2b) Challenge

How would you use the code to carry out simulations at $V/t < 0$?