

# ALF Tutorial

## The *ALF* (Algorithms for Lattice Fermions) project release 2.0 tutorial

Florian Goth, Johannes S. Hofmann, Jonas Schwab,  
Jefferson S. E. Portela, Fakher F. Assaad

August 24, 2020

The ALF package provides a general code for auxiliary-field Quantum Monte Carlo simulations and default analysis. In this tutorial we show how users from beginners to specialists can profit from ALF. This document is divided in two parts:

**Part I.** The first, introductory part of the tutorial is based on ALF's python interface – **pyALF** – which greatly simplifies using the code, making it ideal for: *obtaining benchmark* results for established models; *getting started* with QMC and ALF; or just *quickly running* a simulation.

**Part II.** The second part is independent of the first and aimed at more advanced users who want to simulate their own systems. It guides the user on how to modify the package's Fortran source code and presents the resources implemented to facilitate this task.

This document is intended to be self-contained, but the interested reader should check [ALF's documentation](#), which contains a thorough, systematic description of the package.

[Update link when ALF 2 is open.]

## Part I. Just run it

What follows is a collection of self-explanatory Jupyter notebooks written in Python, each centered on a detailed example followed by a few simple exercises. The notebooks printed below can be found, together with the necessary files and an increasing number of additional notebooks exploring ALF's capabilities, in the [pyALF repository](#).

## Requirements

You can download pyALF from its repository linked above or, from the command line:

```
git clone git@git.physik.uni-wuerzburg.de:ALF/pyALF.git
```

To run the notebooks you need the following installed in your machine:

- Python
- Jupyter
- the libraries Lapack and Blas
- a Fortran compiler, such as **gfortran** or **ifort**,

where the last two are required by the main package [ALF](#). Also, add pyALF's path to your environment variable PYTHONPATH. In Linux, this can be achieved, e.g., by adding the following line to **.bashrc**:

```
export PYTHONPATH="/local/path/to/pyALF:$PYTHONPATH"
```

# Starting

Jupyter notebooks<sup>1</sup> are run through a Jupyter server started, e.g., from the command line:

```
jupyter notebook
```

(or, depending on the installation, `jupyter-notebook`) which opens the “notebook dashboard” in your default browser, where you can navigate through your file structure to the `pyALF` directory. There you will find the interface’s core module, `py_alf.py`, some auxiliary files, and notebooks such as the ones included bellow. Have fun.

## Notebooks

### 1. A minimal ALF run

In this bare-bones example we use the `pyALF` interface to run the canonical Hubbard model on a default configuration: a  $6 \times 6$  square grid, with interaction strength  $U = 4$  and inverse temperature  $\beta = 5$ .

Bellow we go through the steps for performing the simulation and outputting observables.

---

1. Import `Simulation` class from the `py_alf` python module, which provides the interface with ALF:

```
[1]: from py_alf import Simulation          # Interface with ALF
```

2. Create an instance of `Simulation`, setting parameters as desired:

```
[2]: sim = Simulation(
    "Hubbard",          # Hamiltonian
    {                   # Model and simulation parameters for each Simulation instance
        "Model": "Hubbard",    # Base model
        "Lattice_type": "Square"}, # Lattice type
    )
```

3. Compile ALF, downloading it first from the [ALF repository](#) if not found locally. This may take a few minutes:

```
[3]: sim.compile()          # Compilation needs to be performed only once
```

```
Repository /home/stafusa/ALF/pyALF/Notebooks/ALF does not exist, cloning from
git@git.physik.uni-wuerzburg.de:ALF/ALF.git
Compiling ALF... Done.
```

4. Perform the simulation as specified in `sim`:

```
[4]: sim.run()              # Perform the actual simulation in ALF
```

```
Prepare directory "/home/stafusa/ALF/pyALF/Notebooks/Hubbard_Square" for Monte
Carlo run.
Create new directory.
Run /home/stafusa/ALF/pyALF/Notebooks/ALF/Prog/Hubbard.out
```

5. Perform some simple analyses:

```
[5]: sim.analysis()         # Perform default analysis; list observables
```

---

<sup>1</sup>Note that `pyALF` can also be used to start a simulation from the command line, without starting a Jupyter server or using a notebook. For instance: `python3.7 Run.py -R -alfdir /home/debian/ALF-1.2/ -config "Intel" -executable_R Hubbard -mpi True` starts a parallel run of the Hubbard model, using ALF compiled with `ifort`. Notice that `Run.py` requires a configuration file `Sims`, which defines the simulation parameters. An entry of `Sims` might read as: `"Model": "Hubbard", "Lattice_type": "Square", "L1": 4, "L2": 4, "NBin": 5, "ham_T": 0.0, "Nsweep": 2000, "Beta": 1.0, "ham_chem": -1.0`

```
Analysing Ener_scal
Analysing Part_scal
Analysing Pot_scal
Analysing Kin_scal
Analysing Den_eq
Analysing SpinZ_eq
Analysing Green_eq
Analysing SpinXY_eq
Analysing SpinT_eq
Analysing SpinXY_tau
Analysing SpinZ_tau
Analysing Den_tau
Analysing Green_tau
Analysing SpinT_tau
```

6. Store computed observables list:

```
[6]: obs = sim.get_obs() # Dictionary for the observables
```

which are available for further analyses. For instance, the internal energy of the system (and its error) is accessed by:

```
[7]: obs['Ener_scalJ']['obs']
```

```
[7]: array([[ -29.983503,    0.232685]])
```

---

7. Running again: The simulation can be resumed to increase the precision of the results.

```
[8]: sim.run()
sim.analysis()
obs2 = sim.get_obs()
print(obs2['Ener_scalJ']['obs'])
print("\nRunning again reduced the error from ", obs['Ener_scalJ']['obs'][0][1], " to ",
↳obs2['Ener_scalJ']['obs'][0][1], ".")
```

Prepare directory "/home/stafusa/ALF/pyALF/Notebooks/Hubbard\_Square" for Monte Carlo run.

Resuming previous run.

Run /home/stafusa/ALF/pyALF/Notebooks/ALF/Prog/Hubbard.out

```
Analysing Ener_scal
Analysing Part_scal
Analysing Pot_scal
Analysing Kin_scal
Analysing Den_eq
Analysing SpinZ_eq
Analysing Green_eq
Analysing SpinXY_eq
Analysing SpinT_eq
Analysing SpinXY_tau
Analysing SpinZ_tau
Analysing Den_tau
Analysing Green_tau
Analysing SpinT_tau
[[-29.819654    0.135667]]
```

Running again reduced the error from 0.232685 to 0.135667 .

**Note:** To run a fresh simulation - instead of performing a refinement over previous run(s) - the Monte Carlo run directory should be deleted before rerunning.

---

## 1.1. Exercises

1. Rerun once again and check the new improvement in precision.
2. Look at a few other observables (`sim.analysis()` outputs the names of those available).

3. Change the lattice size by adding, e.g., "L1": 4, and "L2": 1, to the simulation parameters definitions of `sim` (step 2).

## 2. Trotter systematic error - Hubbard on the square lattice

In this example we use the [pyALF](#) interface to run ALF with the Mz choice of Hubbard-Stratonovich transformation (i.e., coupled to the  $z$ -component of the spin) on a  $6 \times 6$  site square lattice, at  $U/t = 4$  half-band filling, and inverse temperature  $\beta t = 5$ .

We carry out a systematic  $\Delta\tau t$  extrapolation keeping  $\Delta\tau t L_{\text{Trotter}} = 2$  constant. Recall that the formulation of the auxiliary field QMC approach is based on the Trotter decomposition

$$e^{-\Delta\tau(A+B)} = e^{-\Delta\tau A} e^{-\Delta\tau B} + \mathcal{O}(\Delta\tau^2)$$

The overall error produced by this approximation is of the order  $\Delta\tau^2$ .

Bellow we go through the steps for performing this extrapolation: setting the simulation parameters, running it and analysing the data. A reference plot for this analyses is found in [ALF documentation](#), Sec. 2.3.2 (Symmetric Trotter decomposition).

1. Import Simulation class from the `py_alf` python module, which provides the interface with ALF, as well as mathematics and plotting packages:

```
[1]: from py_alf import Simulation      # Interface with ALF
#
import numpy as np                    # Numerical library
from scipy.optimize import curve_fit  # Numerical library
import matplotlib.pyplot as plt       # Plotting library
```

2. Create instances of `Simulation`, specifying the necessary parameters, in particular the different  $\Delta\tau$  values:

```
[2]: sims = []                        # Vector of Simulation instances
print('dtau values used:')
for dtau in [0.05, 0.1, 0.15]:      # Values of dtau
    print(dtau)
    sim = Simulation(
        'Hubbard',                    # Hamiltonian
        {                             # Model and simulation parameters for each Simulation instance
            'Model': 'Hubbard',        # Base model
            'Lattice_type': 'Square',  # Lattice type
            'L1': 6,                   # Lattice length in the first unit vector direction
            'L2': 6,                   # Lattice length in the second unit vector direction
            'Checkerboard': False,     # Whether checkerboard decomposition is used or not
            'Symm': True,               # Whether symmetrization takes place
            'ham_T': 1.0,               # Hopping parameter
            'ham_U': 4.0,               # Hubbard interaction
            'ham_Tperp': 0.0,           # For bilayer systems
            'beta': 5.0,                # Inverse temperature
            'Ltau': 0,                  # '1' for time-displaced Green functions; '0' otherwise
            'NSweep': 200,              # Number of sweeps
            'NBin': 10,                 # Number of bins
            'Dtau': dtau,               # Only dtau varies between simulations, Ltrot=beta/Dtau
            'Mz': True,                 # If true, sets the M_z-Hubbard model: Nf=2, N_sum=1,
            },                          # HS field couples to z-component of magnetization
        alf_dir='~/Programs/ALF',      # Local ALF copy, if present
    )
    sims.append(sim)
```

```
dtau values used:
0.05
0.1
0.15
```

3. Compile ALF, downloading it first if not found locally. This may take a few minutes:

```
[3]: sims[0].compile() # Compilation needs to be performed only once
```

Compiling ALF... Done.

4. Perform the simulations, as specified in each element of `sim`:

```
[4]: for i, sim in enumerate(sims):
      sim.run() # Perform the actual simulation in ALF
```

Prepare directory "/home/stafusa/ALF/pyALF/Hubbard\_Square\_L1=6\_L2=6\_Checkerboard=False\_Symm=True\_T=1.0\_U=4.0\_Tperp=0.0\_beta=5.0\_Dtau=0.05\_Mz=True" for Monte Carlo run.

Resuming previous run.

Run /home/stafusa/Programs/ALF/Prog/Hubbard.out

Prepare directory "/home/stafusa/ALF/pyALF/Hubbard\_Square\_L1=6\_L2=6\_Checkerboard=False\_Symm=True\_T=1.0\_U=4.0\_Tperp=0.0\_beta=5.0\_Dtau=0.1\_Mz=True" for Monte Carlo run.

Resuming previous run.

Run /home/stafusa/Programs/ALF/Prog/Hubbard.out

Prepare directory "/home/stafusa/ALF/pyALF/Hubbard\_Square\_L1=6\_L2=6\_Checkerboard=False\_Symm=True\_T=1.0\_U=4.0\_Tperp=0.0\_beta=5.0\_Dtau=0.15\_Mz=True" for Monte Carlo run.

Resuming previous run.

Run /home/stafusa/Programs/ALF/Prog/Hubbard.out

5. Calculate the internal energies:

```
[5]: ener = np.empty((len(sims), 2)) # Matrix for storing energy values
      dtaus = np.empty((len(sims),)) # Matrix for Dtau values, for plotting
      for i, sim in enumerate(sims):
          print(sim.sim_dir) # Directory containing the simulation output
          sim.analysis() # Perform default analysis
          dtaus[i] = sim.sim_dict['Dtau'] # Store Dtau value
          ener[i] = sim.get_obs(['Ener_scalJ'])['Ener_scalJ']['obs'] # Store internal energy
```

/home/stafusa/ALF/pyALF/Hubbard\_Square\_L1=6\_L2=6\_Checkerboard=False\_Symm=True\_T=1.0\_U=4.0\_Tperp=0.0\_beta=5.0\_Dtau=0.05\_Mz=True

Analysing Ener\_scal

Analysing Part\_scal

Analysing Pot\_scal

Analysing Kin\_scal

Analysing Den\_eq

Analysing SpinZ\_eq

Analysing Green\_eq

Analysing SpinXY\_eq

Analysing SpinT\_eq

/home/stafusa/ALF/pyALF/Hubbard\_Square\_L1=6\_L2=6\_Checkerboard=False\_Symm=True\_T=1.0\_U=4.0\_Tperp=0.0\_beta=5.0\_Dtau=0.1\_Mz=True

Analysing Ener\_scal

Analysing Part\_scal

Analysing Pot\_scal

Analysing Kin\_scal

Analysing Den\_eq

Analysing SpinZ\_eq

Analysing Green\_eq

Analysing SpinXY\_eq

Analysing SpinT\_eq

/home/stafusa/ALF/pyALF/Hubbard\_Square\_L1=6\_L2=6\_Checkerboard=False\_Symm=True\_T=1.0\_U=4.0\_Tperp=0.0\_beta=5.0\_Dtau=0.15\_Mz=True

Analysing Ener\_scal

Analysing Part\_scal

Analysing Pot\_scal

Analysing Kin\_scal

Analysing Den\_eq

Analysing SpinZ\_eq

Analysing Green\_eq

Analysing SpinXY\_eq

Analysing SpinT\_eq

```
[6]: print('For Dtau values', dtaus, 'the measured energies are:\n', ener)
```

For Dtau values [0.05 0.1 0.15] the measured energies are:

```
[[ -29.714802  0.041044]]
```

```
[-29.784918  0.043263]
[-29.818716  0.029992]]
```


```
[7]: plt.errorbar(dtaus**2, ener[:, 0], ener[:, 1])

def func(x, y0, a):
    return y0 + a*x**2
popt1, pcov = curve_fit(func, dtaus, ener[:, 0], sigma=ener[:, 1], absolute_sigma=True)
perr1 = np.sqrt(np.diag(pcov))
print(popt1, perr1)
xs = np.linspace(0., dtaus.max())
plt.plot(xs**2, func(xs, *popt1))

plt.errorbar(0, popt1[0], perr1[0])
```

```
[-29.71520822 -4.77614903] [0.04068013 2.44507809]
```

```
[7]: <ErrorbarContainer object of 3 artists>
```



## 2.1. Exercises

1. Try out the four different combinations for **Checkerboard** and **Symm** settings in order to observe their effect on the output and run time. Reference: Sec. 2.3.2 - Symmetric Trotter decomposition - of the [ALF documentation](#), especially Fig. 2.

## 3. Testing against ED - Hubbard on a ring

In this example we use the [pyALF](#) interface to run ALF with the Mz choice of Hubbard Stratonovitch transformation on a four site ring, at  $U/t = 4$  and inverse temperature  $\beta t = 2$ . For this set of parameters, the exact internal energy reads:

$$\left\langle -t \sum_{\langle i,j \rangle, \sigma} c_{i,\sigma}^\dagger c_{j,\sigma} + U \sum_{i=1}^N n_{i,\uparrow} n_{i,\downarrow} \right\rangle = -1.47261997t$$

To reproduce this result we will have to carry out a systematic  $\Delta\tau t$  extrapolation keeping  $\Delta\tau t L_{\text{Trotter}} = 2$  constant.

Recall that the formulation of the auxiliary field QMC approach is based on the Trotter decomposition

$$e^{-\Delta\tau(A+B)} = e^{-\Delta\tau A} e^{-\Delta\tau B} + \mathcal{O}(\Delta\tau^2)$$

The overall error produced by this approximation is of the order  $\Delta\tau^2$ .

Bellow we go through the steps for performing this extrapolation: setting the simulation parameters, running it and analyzing the data.

1. Import Simulation class from the py\_alf python module, which provides the interface with ALF, as well as mathematics and plotting packages:

```
[1]: from py_alf import Simulation          # Interface with ALF
      #
      import numpy as np                  # Numerical library
      from scipy.optimize import curve_fit # Numerical library
      import matplotlib.pyplot as plt     # Plotting library
```

2. Create instances of Simulation, specifying the necessary parameters, in particular the different  $\Delta\tau$  values:

```
[2]: sims = []                          # Vector of Simulation instances
      print('dtau values used:')
      for dtau in [0.05, 0.1, 0.15]:    # Values of dtau
          print(dtau)
          sim = Simulation(
              'Hubbard',                  # Hamiltonian
              {                           # Model and simulation parameters for each Simulation instance
                  'Model': 'Hubbard',     # Base model
                  'Lattice_type': 'N_leg_ladder', # Lattice type
                  'L1': 4,                 # Lattice length in the first unit vector direction
                  'L2': 1,                 # Lattice length in the second unit vector direction
                  'Checkerboard': False,   # Whether checkerboard decomposition is used or not
                  'Symm': True,            # Whether symmetrization takes place
                  'ham_T': 1.0,            # Hopping parameter
                  'ham_U': 4.0,            # Hubbard interaction
                  'ham_Tperp': 0.0,        # For bilayer systems
                  'beta': 2.0,             # Inverse temperature
                  'Ltau': 0,               # '1' for time-displaced Green functions; '0' otherwise
                  'NSweep': 400,           # Number of sweeps
                  'NBin': 100,            # Number of bins
                  'Dtau': dtau,            # Only dtau varies between simulations, Ltrot=beta/Dtau
                  'Mz': True,              # If true, sets the M_z-Hubbard model: Nf=2, N_sum=1,
              },                          # HS field couples to z-component of magnetization
              alf_dir='~/Programs/ALF',   # Local ALF copy, if present
          )
          sims.append(sim)
```

```
dtau values used:
0.05
0.1
0.15
```

3. Compile ALF, downloading it first if not found locally. This may take a few minutes:

```
[3]: sims[0].compile()                  # Compilation needs to be performed only once
```

Compiling ALF... Done.

4. Perform the simulations, as specified in each element of `sims`:

```
[4]: for i, sim in enumerate(sims):
      sim.run()                          # Perform the actual simulation in ALF
```

```
Prepare directory "/home/stafusa/ALF/pyALF/Hubbard_N_leg_ladder_L1=4_L2=1_Checke
rboard=False_Symm=True_T=1.0_U=4.0_Tperp=0.0_beta=2.0_Dtau=0.05_Mz=True" for
Monte Carlo run.
Resuming previous run.
Run /home/stafusa/Programs/ALF/Prog/Hubbard.out
Prepare directory "/home/stafusa/ALF/pyALF/Hubbard_N_leg_ladder_L1=4_L2=1_Checke
rboard=False_Symm=True_T=1.0_U=4.0_Tperp=0.0_beta=2.0_Dtau=0.1_Mz=True" for
Monte Carlo run.
Resuming previous run.
Run /home/stafusa/Programs/ALF/Prog/Hubbard.out
```

```

Prepare directory "/home/stafusa/ALF/pyALF/Hubbard_N_leg_ladder_L1=4_L2=1_Checke
rboard=False_Symm=True_T=1.0_U=4.0_Tperp=0.0_beta=2.0_Dtau=0.15_Mz=True" for
Monte Carlo run.
Resuming previous run.
Run /home/stafusa/Programs/ALF/Prog/Hubbard.out

```

## 5. Calculate the internal energies:

```

[5]: ener = np.empty((len(sims), 2))          # Matrix for storing energy values
      dtaus = np.empty((len(sims),))          # Matrix for Dtau values, for plotting
      for i, sim in enumerate(sims):
          print(sim.sim_dir)                  # Directory containing the simulation output
          sim.analysis()                      # Perform default analysis
          dtaus[i] = sim.sim_dict['Dtau']      # Store Dtau value
          ener[i] = sim.get_obs(['Ener_scalJ'])['Ener_scalJ']['obs'] # Store internal energy

```

```

/home/stafusa/ALF/pyALF/Hubbard_N_leg_ladder_L1=4_L2=1_Checkerboard=False_Symm=T
rue_T=1.0_U=4.0_Tperp=0.0_beta=2.0_Dtau=0.05_Mz=True
Analysing Ener_scal
Analysing Part_scal
Analysing Pot_scal
Analysing Kin_scal
Analysing Den_eq
Analysing SpinZ_eq
Analysing Green_eq
Analysing SpinXY_eq
Analysing SpinT_eq
/home/stafusa/ALF/pyALF/Hubbard_N_leg_ladder_L1=4_L2=1_Checkerboard=False_Symm=T
rue_T=1.0_U=4.0_Tperp=0.0_beta=2.0_Dtau=0.1_Mz=True
Analysing Ener_scal
Analysing Part_scal
Analysing Pot_scal
Analysing Kin_scal
Analysing Den_eq
Analysing SpinZ_eq
Analysing Green_eq
Analysing SpinXY_eq
Analysing SpinT_eq
/home/stafusa/ALF/pyALF/Hubbard_N_leg_ladder_L1=4_L2=1_Checkerboard=False_Symm=T
rue_T=1.0_U=4.0_Tperp=0.0_beta=2.0_Dtau=0.15_Mz=True
Analysing Ener_scal
Analysing Part_scal
Analysing Pot_scal
Analysing Kin_scal
Analysing Den_eq
Analysing SpinZ_eq
Analysing Green_eq
Analysing SpinXY_eq
Analysing SpinT_eq

```

```

[6]: print('For Dtau values', dtaus, 'the measured energies are:\n', ener)

```

```

For Dtau values [0.05 0.1 0.15] the measured energies are:
[[-1.473383  0.002569]
 [-1.478316  0.002525]
 [-1.46798   0.002359]]

```

```

[7]: plt.errorbar(dtaus**2, ener[:, 0], ener[:, 1])

def func(x, y0, a):
    return y0 + a*x**2
popt1, pcov = curve_fit(func, dtaus, ener[:, 0], sigma=ener[:, 1], absolute_sigma=True)
perr1 = np.sqrt(np.diag(pcov))
print(popt1, perr1)
xs = np.linspace(0., dtaus.max())
plt.plot(xs**2, func(xs, *popt1))

plt.errorbar(0, popt1[0], perr1[0])

```

```


[-1.47718877  0.3429858 ] [0.00254357 0.17130454]

```

```

[7]: <ErrorbarContainer object of 3 artists>

```



### 3.1. Exercises

1. Redo the extrapolation for different values of  $\beta t$  (e.g., for  $\beta t = 1$ , the internal energy is -0.62186692t, and for  $\beta t = 4$ , it is -1.90837196t).
2. Experiment with different settings for **Checkerboard** and **Symm**.

## 4. Projective algorithm

In this example we use the [pyALF](#) interface to run ALF's projective algorithm with the Mz choice of Hubbard Stratonovich transformation on a 4-site ring.

The projective approach is the method of choice if one is interested in ground-state properties. The starting point is a pair of trial wave functions,  $|\Psi_{T,L/R}\rangle$ , that are not orthogonal to the ground state  $|\Psi_0\rangle$ :

$$\langle \Psi_{T,L/R} | \Psi_0 \rangle \neq 0.$$

The ground-state expectation value of any observable  $\hat{O}$  can then be computed by propagation along the imaginary time axis:

$$\frac{\langle \Psi_0 | \hat{O} | \Psi_0 \rangle}{\langle \Psi_0 | \Psi_0 \rangle} = \lim_{\theta \rightarrow \infty} \frac{\langle \Psi_{T,L} | e^{-\theta \hat{H}} e^{-(\beta-\tau) \hat{H}} \hat{O} e^{-\tau \hat{H}} e^{-\theta \hat{H}} | \Psi_{T,R} \rangle}{\langle \Psi_{T,L} | e^{-(2\theta+\beta) \hat{H}} | \Psi_{T,R} \rangle},$$

where  $\beta$  defines the imaginary time range where observables (time displaced and equal time) are measured and  $\tau$  varies from 0 to  $\beta$  in the calculation of time-displace observables. For further details, see Sec. 3 of [ALF documentation](#).

1. Import **Simulation** class from the `py_alf` python module, which provides the interface with ALF, as well as numerical and plotting packages:

```
[1]: from py_alf import Simulation          # Interface with ALF
#
import numpy as np                        # Numerical library
from scipy.optimize import curve_fit      # Numerical library
import matplotlib.pyplot as plt          # Plotting library
```

2. Create instances of **Simulation**, specifying the necessary parameters, in particular the **Projector** to **True**:

```
[2]: sims = []                                # Vector of Simulation instances
print('Theta values used:')
for theta in [5, 10, 20]:                    # Values of Theta
    print(theta)
    sim = Simulation(
        'Hubbard',                           # Hamiltonian
        {                                     # Model and simulation parameters for each Simulation instance
            'Model': 'Hubbard',               # Base model
            'Lattice_type': 'N_leg_ladder',   # Lattice type
            'L1': 4,                          # Lattice length in the first unit vector direction
            'L2': 1,                          # Lattice length in the second unit vector direction
            'Checkerboard': False,            # Whether checkerboard decomposition is used or not
            'Symm': True,                     # Whether symmetrization takes place
            'Projector': True,                # Whether to use the projective algorithm
            'Theta': theta,                   # Projector parameter
            'ham_T': 1.0,                     # Hopping parameter
            'ham_U': 4.0,                     # Hubbard interaction
            'ham_Tperp': 0.0,                 # For bilayer systems
            'beta': 1.0,                      # Inverse temperature
            'Ltau': 0,                        # '1' for time-displaced Green functions; '0' otherwise
            'NSweep': 400,                    # Number of sweeps
            'NBin': 10,                       # Number of bins
            'Dtau': 0.05,                     # Only dtau varies between simulations, Ltrot=beta/Dtau
            'Mz': True,                       # If true, sets the M_z-Hubbard model: Nf=2, N_sum=1,
                                                # HS field couples to z-component of magnetization
        },
        alf_dir='~/Programs/ALF',            # Local ALF copy, if present
    )
    sims.append(sim)
```

Theta values used:

5  
10  
20

3. Compile ALF, downloading it first if not found locally. This may take a few minutes:

```
[3]: sims[0].compile()                      # Compilation needs to be performed only once
```

Compiling ALF... Done.

4. Perform the simulations, as specified in each element of `sim`:

```
[4]: for i, sim in enumerate(sims):
    sim.run()                                # Perform the actual simulation in ALF
```

Prepare directory "/home/stafusa/ALF/pyALF/Hubbard\_N\_leg\_ladder\_L1=4\_L2=1\_Checke  
rboard=False\_Symm=True\_Projector=True\_Theta=5\_T=1.0\_U=4.0\_Tperp=0.0\_beta=1.0\_Dt  
au=0.05\_Mz=True" for Monte Carlo run.

Create new directory.

Run /home/stafusa/Programs/ALF/Prog/Hubbard.out

Prepare directory "/home/stafusa/ALF/pyALF/Hubbard\_N\_leg\_ladder\_L1=4\_L2=1\_Checke  
rboard=False\_Symm=True\_Projector=True\_Theta=10\_T=1.0\_U=4.0\_Tperp=0.0\_beta=1.0\_Dt  
au=0.05\_Mz=True" for Monte Carlo run.

Create new directory.

Run /home/stafusa/Programs/ALF/Prog/Hubbard.out

Prepare directory "/home/stafusa/ALF/pyALF/Hubbard\_N\_leg\_ladder\_L1=4\_L2=1\_Checke  
rboard=False\_Symm=True\_Projector=True\_Theta=20\_T=1.0\_U=4.0\_Tperp=0.0\_beta=1.0\_Dt  
au=0.05\_Mz=True" for Monte Carlo run.

Create new directory.

Run /home/stafusa/Programs/ALF/Prog/Hubbard.out

5. Calculate the internal energies:

```
[6]: %capture
ener = np.empty((len(sims), 2))              # Matrix for storing energy values
thetas = np.empty((len(sims),))             # Matrix for Thetas values, for plotting
for i, sim in enumerate(sims):
    print(sim.sim_dir)                       # Directory containing the simulation output
    sim.analysis()                           # Perform default analysis
    thetas[i] = sim.sim_dict['Theta']         # Store Theta value
    ener[i] = sim.get_obs(['Ener_scalJ'])['Ener_scalJ']['obs'] # Store internal energy
```


Where the *cell magic* `%%capture` suppresses the output of `sim.analysis()`, which lists the data directories and observables.

```
[7]: print('For Theta values', thetas, 'the measured energies are:\n', ener)
```

```
For Theta values [ 5. 10. 20.] the measured energies are:
[[-2.137697  0.025552]
 [-2.108235  0.023964]
 [-2.089294  0.025426]]
```

```
[8]: plt.errorbar(1/thetas, ener[:, 0], ener[:, 1])
```

```
[8]: <ErrorbarContainer object of 3 artists>
```



## 4.1. Exercises

1. [ **TO BE FLESHED OUT** ] Kondo phase transition. References:  
<https://journals.aps.org/prb/abstract/10.1103/PhysRevB.63.155114>  
<https://journals.aps.org/prl/abstract/10.1103/PhysRevLett.83.796>

# Part II. Getting your hands dirty - writing new code

Part II consists in a set of guided advanced exercises.

A lot already comes implemented in ALF, but unavoidably, as one proceeds in their own investigations, a new model has to be implemented or a new observable defined – and for that one has to grapple with the package’s Fortran source code. However, in ALF this is made easy by means of predefined structures, templates, and the examples below.

## Downloading the code and tutorial

To download the code, type `git clone git@git.physik.uni-wuerzburg.de:ALF/ALF_code.git` in a shell.

To download the tutorial including solutions type:

git clone git@git.physik.uni-wuerzburg.de:ALF/ALF\_Tutorial.git again in a shell.

## Exercise 1 – Dimensional crossover

[To be updated.]

### 1a) Modifying the hopping

Here we will modify the code so as to allow for different hopping matrix elements along the x and y directions of a square lattice. To do so, one merely has to do the following

- Add an extra variable, `Ham_Ty`, in the parameter file in the `VAR_Hubbard` name space
- Declare the variable `Ham_Ty` in the `Hamiltonian_Examples.f90` .
- Read in this variable in `Ham_set` subroutine of the `Hamiltonian_Examples.f90` file.
- Modify the hopping matrix in the subroutine `Ham_Hop` in the `Hamiltonian_Examples.f90` file.

```
DO I = 1, Latt%N
  I1 = Latt%nnlist(I,1,0)
  I2 = Latt%nnlist(I,0,1)
  Op_T(nc,n)%0(I,I1) = cmplx(-Ham_T, 0.d0, kind(0.D0))
  Op_T(nc,n)%0(I1,I) = cmplx(-Ham_T, 0.d0, kind(0.D0))
  !!!!! Modifications for Exercise 2 (a)
  !Op_T(nc,n)%0(I,I2) = cmplx(-Ham_T, 0.d0, kind(0.D0))
  !Op_T(nc,n)%0(I2,I) = cmplx(-Ham_T, 0.d0, kind(0.D0))
  Op_T(nc,n)%0(I,I2) = cmplx(-Ham_Ty, 0.d0, kind(0.D0))
  Op_T(nc,n)%0(I2,I) = cmplx(-Ham_Ty, 0.d0, kind(0.D0))
  !!!!!
  Op_T(nc,n)%0(I,I) = cmplx(-Ham_chem, 0.d0, kind(0.D0))
ENDDO
```

In the directory `Solutions/Exercise_2` we have duplicated the ALF and commented the changes that have to be carried out to the file `Hamiltonian_Examples.f90` in the `Prog` directory.

As an application of this code, one can consider a ladder system, defined by the parameter file:

```
!=====
! Variables for the Hubb program
!-----

&VAR_lattice
L1    = 14
L2    = 2
Lattice_type = "Square"
Model = "Hubbard_Mz" ! Hubbard_SU2, Hubbard_Mz
/

&VAR_Hubbard
ham_T    = 1.0D0
!!!!!! Modifications for Exercise 2
ham_Ty   = 2.0D0
!!!!!!
ham_chem= 0.0D0
ham_U    = 4.0
Beta     = 10.0
dtau     = 0.1
/

&VAR_QMC
```

```

Nwrap   = 10
NSweep  = 100
NBin    = 10
Ltau    = 1
LOBS_ST = 1
LOBS_EN = 100
CPU_MAX = 0.1
/

&VAR_errors
n_skip  = 2
N_rebin = 1
N_Cov   = 0
/
! slash terminates namelist statement - DO NOT REMOVE

```

When running the code for the above ladder system and analyzing the spin correlation functions, `SpinZ_eqJR` and `SpinXY_eqJR` one will notice that it is hard to restore the SU(2) spin symmetry and that very long runs are required to obtain the desired equality

$$\langle S_i^z S_j^z \rangle = \langle S_i^y S_j^y \rangle = \langle S_i^x S_j^x \rangle. \quad (1)$$

The structure of the output files `SpinZ_eqJR` and `SpinXY_eqJR` is described in the documentation. For the Mz Hubbard-Stratonovitch transformation it is hence better to consider the improved estimator

$$\langle \vec{S}_i \cdot \vec{S}_j \rangle \quad (2)$$


to compute the spin-spin correlations.

## 1b) Adding a new observable

Here the aim is to include the new observable equal time observable  $\langle \vec{S}_i \cdot \vec{S}_j \rangle$  in the `Hubbard_Mz` code. To achieve this, you will have to carry out the following steps.

- In the subroutine `Alloc_obs` in the `Hamiltonian_example.f90` file you will have to add a new equal time observable with a call to `Call Obser_Latt_make(Observeq(I),Ns,Nt,No,Filename)` with `Ns = Latt%N`; `No = Norb`; `Filename = "SpinT"`, `Nt=1`, `I=5`
- In the subroutine `Obser` you will have to add the Wick decomposition of this observable.

In the program `Hamiltonian_Examples.f90` to be found in the directory `Solutions/Exercise_2/Prog/` we have commented the changes that have to be carried out to add this observable. The new variable takes the name `SpinT` and the results you should obtain are summarized in Fig. ??.



### 1c) The $SU(2)$ Hubbard-Stratonovich transformation

The  $SU(2)$  Hubbard-Stratonovich decomposition, conserves spin rotational symmetry. Run the ladder code with the  $SU(2)$  flag in the parameter file switched on (i.e. `Model = Hubbard_SU2`) and compare results.

## Exercise 2 – Defining a new model: The one-dimensional t-V model

[To be updated.]

### 2a) Define new model

In this section, one we will show what modifications have to be carried out for computing the physics of the one dimensional t-V model of spinless fermions.

$$H = -t \sum_i \left( c_i^\dagger c_{i+a} + c_{i+a}^\dagger c_i \right) - \frac{V}{2} \sum_i \left( c_i^\dagger c_{i+a} + c_{i+a}^\dagger c_i \right)^2 \quad (3)$$

The above form is readily included in the ALF since the interaction is written in terms of a perfect square. Expanding the square yields (up to a constant) the desired model:

$$H = -t \sum_i \left( c_i^\dagger c_{i+a} + c_{i+a}^\dagger c_i \right) + V \sum_i (n_i - 1/2) (n_{i+a} - 1/2) \quad (4)$$

In the directory `Solutions/Exercise_3` we have duplicated the ALF and commented the changes that have to be carried out to the file `Hamiltonian_Examples.f90` in the `Prog` directory so as to include the `t_V` model. Here are the steps to be carried out.

- Add the  $t - V$  name space in the parameter file so as to read in the appropriate variables.
- Declare new variables in the `Hamiltonian_Examples.f90` file.
- In the `Ham_set` subroutine of the file `Hamiltonian_Examples.f90` set and read in the parameters for the new model, `t_V`. For this model `NF=1` and `N_SUN=1` since we are working with spinless fermions
- In the `Ham_V` subroutine you will have to add the new interaction. For a given bond at a given time-slice, we need to decouple the interaction:

$$e^{\Delta\tau \frac{V}{2} (c_i^\dagger c_{i+a} + c_{i+a}^\dagger c_i)^2} = \sum_{l=\pm 1, \pm 2} \gamma_l e^{\sqrt{\Delta\tau \frac{V}{2}} \eta_l (c_i^\dagger c_{i+a} + c_{i+a}^\dagger c_i)} = \sum_{l=\pm 1, \pm 2} \gamma_l e^{g \eta_l (c_i^\dagger, c_{i+a}^\dagger) O(c_i, c_{i+a})^T} \quad (5)$$

Here is how this translates in the code.


```
Allocate(Op_V(Latt%N,N_FL))
do nf = 1,N_FL
  do i = 1, N_coord*Ndim
    call Op_make(Op_V(i,nf),2)
  enddo
enddo
Do nc = 1, Latt%N ! Runs over bonds = # of lattice sites in one-dimension.
  I1 = nc
  I2 = Latt%nnlist(I1,1,0)
  Op_V(nc,1)%P(1) = I1
  Op_V(nc,1)%P(2) = I2
  Op_V(nc,1)%O(1,2) = cmplx(1.d0 ,0.d0, kind(0.D0))
  Op_V(nc,1)%O(2,1) = cmplx(1.d0 ,0.d0, kind(0.D0))
  Op_V(nc,1)%g = SQRT(CMPLX( DTAU*Ham_Vint/2.d0, 0.D0, kind(0.D0)))
  Op_V(nc,1)%alpha = cmplx(0d0,0.d0, kind(0.D0))
  Op_V(nc,1)%type =2
  Call Op_set( Op_V(nc,1) )
enddo
```

- Finally you will have to update the `Obser` and `ObserT` routines for the calculation of the equal and time displaced correlations. For the `t_V` model you can essentially use the same observables as for the `Hubbard_SU(2)` model.

You can now run the code for various values of  $V/t$ . A Jordan-Wigner transformation will map the  $\mathfrak{t}_V$  model onto the XXZ chain:

$$H = J_{xx} \sum_i S_i^x S_{i+a}^x + S_i^y S_{i+a}^y + J_{zz} \sum_i S_i^z S_{i+a}^z \quad (6)$$

with  $J_{zz} = V$  and  $J_{xx} = 2t$ . Hence when  $V/t = 2$  we reproduce the Heisenberg model. For  $V/t > 2$  the model is in the Ising regime with long-range charge density wave order and is an insulator. In the regime  $-2 < V/t < 2$  the model is metallic and corresponds to a Luttinger liquid. Finally, at  $V/t < -2$  phase separation between hole rich and electron rich phases occur. Fig. ?? shows typical results.



## 2b) Challenge

How would you use the code to carry out simulations at  $V/t < 0$ ?