

Antimicrobial Resistance (AMR)

Surveillance report

Hospital name: Hypothetical Hospital

Country name: Hypothetical Country

Data from:

02 Jan 2016 to 31 Dec 2016

Contact person: xxx_Can be changed in the dictionary_of_variable_data.csv_xxx

Contact address: xxx_Can be changed in the dictionary_of_variable_data.csv_xxx

Contact email: xxx@xxx.xxx

Generated on: 16 Nov 2023 13:37

Software version: 3.0 (BETA) Build 3019 on 16 Nov 2023

Generated by

AutoMated tool for Antimicrobial resistance Surveillance System (AMASS) version 3.0
(released on 16 Nov 2023)

The AMASS application is available under the Creative Commons Attribution 4.0 International Public License (CC BY 4.0). The application can be downloaded at :
<https://www.amass.website>

The AMASS application used microbiology_data and hospital_admission_data files that are stored in the same folder as the application (AMASS.bat) to generate this report.

The goal of the AMASS application is to enable hospitals with microbiology data available in electronic formats to analyze their own data and generate AMR surveillance reports promptly. If hospital admission data are available, the reports will additionally be stratified by infection origin (community–origin or hospital–origin). If mortality data (such as patient discharge outcome data) are available, a report on mortality involving AMR infection will be added.

This automatically generated report has limitations, and requires users to understand those limitations and use the summary data in the report with careful interpretation.

A valid report could have local implications and much wider benefits if shared with national and international organizations.

This automatically generated report is under the jurisdiction of the hospital to copy, redistribute, and share with any individual or organization.

This automatically generated report contains no patient identifier, similar to standard reports on cumulative antimicrobial susceptibility.

For any query on AMASS, please contact:
Chalida Rangsiwutisak (chalida@tropmedres.ac),
Cherry Lim (cherry@tropmedres.ac), and
Direk Limmathurotsakul (direk@tropmedres.ac)

Suggested title for citation:

Antimicrobial resistance surveillance report, Hypothetical Hospital,
Hypothetical Country, 02 Jan 2016 to 31 Dec 2016.

Content

Introduction	1
Section [1]: Data overview	3
Section [2]: AMR proportion report	5
Section [3]: AMR proportion report with stratification by infection origin	12
Section [4]: AMR frequency report	24
Section [5]: AMR frequency report with stratification by infection origin	27
Section [6]: Mortality involving AMR and antimicrobial–susceptible infections	32
Annex A: Supplementary report on notifiable bacterial infections	38
Annex B: Supplementary report on data indicators	41
Annex C: Cluster signals	43
Methods	58
Acknowledgements	62

Introduction

Antimicrobial resistance (AMR) is a global health crisis [1]. The report by Lord Jim O'Neill estimated that 700,000 global deaths could be attributable to AMR in 2015, and projected that the annual death toll could reach 10 million by 2050 [1]. However, data of AMR surveillance from low and middle-income countries (LMICs) are scarce [1,2], and data of mortality associated with AMR infections are rarely available. A recent study estimated that 19,000 deaths are attributable to AMR infections in Thailand annually, using routinely available microbiological and hospital databases [3]. The study also proposed that hospitals in LMICs should utilize routinely available microbiological and hospital admission databases to generate reports on AMR surveillance systematically [3].

Reports on AMR surveillance can have a wide range of benefits [2]; including

- characterization of the frequency of resistance and organisms in different facilities and regions;
- prospective and retrospective information on emerging public health threats;
- evaluation and optimization of local and national standard treatment guidelines;
- evaluation of the impact of interventions beyond antimicrobial guidelines that aim to reduce AMR; and
- data sharing with national and international organizations to support decisions on resource allocation for interventions against AMR and to inform the implementation of action plans at national and global levels.

When reporting AMR surveillance results, it is generally recommended that (a) duplicate results of bacterial isolates are removed, and (b) reports are stratified by infection origin (community-origin or hospital-origin), if possible [2]. Many hospitals in LMICs lack time and resources needed to analyze the data (particularly to deduplicate data and to generate tables and figures), write the reports, and to release the data or reports [4].

AutoMated tool for Antimicrobial resistance Surveillance System (AMASS) was developed as an offline, open-access and easy-to-use application that allows a hospital to perform data analysis independently and generate AMR proportion and AMR frequency reports stratified by infection origin from routinely collected electronic databases. The application was built in R, which is a free software environment. The application has been placed within a user-friendly interface that only requires the user to double-click on the application icon. The AMASS application can be downloaded at: <https://www.amass.website>

The AMASS version 3.0 additionally generates reports on notifiable bacterial diseases in Annex A and on data indicators (including proportion of contaminants and discordant AST results) in Annex B for the "microbiology_data" file that is used to generate this report. A careful review of the Annex B could help readers and data owners to identify potential errors in the microbiology data used to generate the report.

The AMASS version 3.0 also separately generates Supplementary data indicators report (in PDF and Excel formats) in a new folder "Report_with_patient_identifiers" to support users to check and validate records with notifiable bacteria, notifiable antibiotic-pathogen combinations, infrequent phenotypes or potential errors in the AST results at the local level. The identifiers listed include hospital number and specimen collection date. The files are generated in a separate folder "Report_with_patient_identifiers" so that it is clear that users should not share or transfer the Supplementary Data Indicators report (in PDF and Excel format) to any party outside of the hospital without data security management and confidential agreement.

References:

- [1] O'Neill J. (2014) Antimicrobial resistance: tackling a crisis for the health and wealth of nations. Review on antimicrobial resistance. <http://amr-review.org>. (accessed on 3 Dec 2018).
- [2] World Health Organization (2018) Global Antimicrobial Resistance Surveillance System (GLASS) Report. Early implantation 2016–2017. <http://apps.who.int/iris/bitstream/handle/10665/259744/9789241513449-eng.pdf>. (accessed on 3 Dec 2018)
- [3] Lim C., et al. (2016) Epidemiology and burden of multidrug-resistant bacterial infection in a developing country. *Elife* 5: e18082.
- [4] Ashley EA, Shetty N, Patel J, et al. Harnessing alternative sources of antimicrobial resistance data to support surveillance in low-resource settings. *J Antimicrob Chemother*. 2019; 74(3):541–546.
- [5] Clinical and Laboratory Standards Institute (CLSI). Analysis and Presentation of Cumulative Antimicrobial Susceptibility Test Data, 4th Edition. 2014. (accessed on 21 Jan 2020)
- [6] European Antimicrobial Resistance Surveillance Network (EARS-Net). Antimicrobial resistance (AMR) reporting protocol 2018. (accessed on 21 Jan 2020)
- [7] European Committee on Antimicrobial Susceptibility Testing (EUCAST). www.eucast.org (accessed on 21 Jan 2020)

Section [1]: Data overview

Introduction

An overview of the data detected by the AMASS application is generated by default. The summary is based on the raw data files saved within the same folder as the application file (AMASS.bat).

Please review and validate this section carefully before proceeds to the next section.

Results

The microbiology_data file (stored in the same folder as the application file) had:

50414 specimen data records with collection dates ranging from
02 Jan 2016 to **10 Jan 2017**

The hospital_admission_data file (stored in the same folder as the application file) had:

247260 admission data records with hospital admission dates ranging from
01 Jan 2016 to **31 Dec 2016**

The total number of patient-days was **3393075**.

The total number of patient-days at risk of BSI of hospital-origin was **2898895**.

Note:

[1] If the periods of the data in microbiology_data and hospital_admission_data files are not similar, the automatically-generated report should be interpreted with caution. The AMASS generates the reports based on the available data.

[2] A patient is defined as at risk of BSI of hospital-origin when the patient is admitted to the hospital for more than two calendar days with calendar day one equal to the day of admission.

Reporting period by months:

Data was stratified by month to assist detection of missing data, and verification of whether the month distribution of data records in microbiology_data file and hospital_admission_data file reflected the microbiology culture frequency and admission rate of the hospital, respectively. For example if the number of specimens in the microbiology_data file reported below is lower than what is expected, please check the raw data file and data dictionary files.

Month	Number of specimen data records in microbiology_data file	Number of admission data records in hospital_admission_data file
January	4197	20760
February	4059	19900
March	4337	21400
April	4269	21170
May	4317	21105
June	4022	19800
July	4301	21115
August	4296	20840
September	3975	19660
October	4302	20965
November	4136	20150
December	4203	20395
Total	50414	247260

Note:

[1] Additional general demographic data will be made available in the next version of the AMASS application.

Section [2]: AMR proportion report

Introduction

An AMR proportion report is generated by default, even if the hospital_admission_data file is unavailable. This is to enable hospitals with only microbiology data available to utilize the de-duplication and report generation functions of AMASS. This report is without stratification by origin of infection.

The report generated by the AMASS application version 3.0 includes only blood samples. The next version of AMASS will include other specimen types, including cerebrospinal fluid (CSF), urine, stool, and other specimens.

Organisms under this survey:

- *Staphylococcus aureus*
- *Enterococcus faecalis*
- *Enterococcus faecium*
- *Streptococcus pneumoniae*
- *Salmonella* spp.
- *Escherichia coli*
- *Klebsiella pneumoniae*
- *Pseudomonas aeruginosa*
- *Acinetobacter baumannii*

Results

The microbiology_data file had:

*Sample collection dates ranged from **02 Jan 2016** to **10 Jan 2017***

Number of records of blood specimens collected within the above date range:

15878 blood specimens records

*Number of records of blood specimens with *negative culture (no growth):*

13315 blood specimens records

Number of records of blood specimens with culture positive for a microorganism:

2563 blood specimens records

Number of records of blood specimens with culture positive for organism under this survey:

857 blood specimens records

The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period as described in the method. The number of patients with positive samples is as follows:

Organism	Number of records of blood specimens culture positive for the organism	**Number of patients with blood culture positive for the organism (de-duplicated)
<i>Staphylococcus aureus</i>	113	96
<i>Enterococcus faecalis</i>	0	0
<i>Enterococcus faecium</i>	0	0
<i>Streptococcus pneumoniae</i>	25	20
<i>Salmonella</i> spp.	43	35
<i>Escherichia coli</i>	384	339
<i>Klebsiella pneumoniae</i>	135	120
<i>Pseudomonas aeruginosa</i>	56	48
<i>Acinetobacter baumannii</i>	101	90
Total:	857	748

*The negative culture included data values specified as 'no growth' in the dictionary_for_microbiology_data file (details on data dictionary files are in the method section) to represent specimens with negative culture for any microorganism.


**Only the first isolate for each patient per specimen type, per pathogen, and per evaluation period was included in the analysis.

The following figures and tables show the proportion of patients with blood culture positive for antimicrobial resistant isolates.

Section [2]: AMR proportion report

Blood: *Staphylococcus aureus*


(No. of patients = 96)



Antibiotic agent	Proportion of R	95% CI
Methicillin	16% (15/96)	10% - 24%
Cefoxitin	16% (15/96)	10% - 24%
Oxacillin by MIC	NA	-
Vancomycin	3% (3/96)	1% - 9%
Clindamycin	19% (18/96)	12% - 28%
Chloramphenicol	NA	-

Blood: *Enterococcus faecalis*

(No. of patients = 0)



Antibiotic agent	Proportion of R	95% CI
Penicillin G	NA	-
Ampicillin	NA	-
Vancomycin	NA	-
Teicoplanin	NA	-
Linezolid	NA	-
Daptomycin	NA	-

*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; Methicillin: cefoxitin or oxacillin by MIC


Section [2]: AMR proportion report

Blood: *Enterococcus faecium*

(No. of patients = 0)


Penicillin G
Ampicillin
Vancomycin
Teicoplanin
Linezolid
Daptomycin

Antibiotic agent	Proportion of R	95% CI
Penicillin G	NA	-
Ampicillin	NA	-
Vancomycin	NA	-
Teicoplanin	NA	-
Linezolid	NA	-
Daptomycin	NA	-



Blood: *Streptococcus pneumoniae*

(No. of patients = 20)




Antibiotic agent	Proportion of R	95% CI
Penicillin G	0% (0/16)	0% - 19%
Oxacillin	NA	-
Co-trimoxazole	58% (11/19)	36% - 77%
3GC	0% (0/19)	0% - 17%
Ceftriaxone	NA	-
Cefotaxime	NA	-
Erythromycin	20% (4/20)	8% - 42%
Clindamycin	20% (4/20)	8% - 42%
Levofloxacin	NA	-

*Proportion of R represents represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; 3GC=3rd-generation cephalosporin

Section [2]: AMR proportion report

Blood: *Salmonella* spp.

(No. of patients = 35)




Antibiotic agent	Proportion of R	95% CI
FLUOROQUINOLONES	40% (14/35)	26% - 56%
Ciprofloxacin	40% (14/35)	26% - 56%
Levofloxacin	NA	-
3GC	14% (5/35)	6% - 29%
Ceftriaxone	14% (5/35)	6% - 29%
Ceftazidime	NA	-
Cefotaxime	14% (5/35)	6% - 29%
CARBAPENEMS	NA	-
Imipenem	NA	-
Meropenem	NA	-
Doripenem	NA	-
Ertapenem	NA	-

*Proportion of R represents represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; FLUOROQUINOLONES: ciprofloxacin or levofloxacin; 3GC=3rd-generation cephalosporin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [2]: AMR proportion report

Blood: *Escherichia coli*

(No. of patients = 339)




Antibiotic agent	Proportion of R	95% CI
Ampicillin	50% (118/236)	44% - 56%
Gentamicin	22% (74/339)	18% - 26%
Amikacin	100% (100/100)	96% - 100%
Co-trimoxazole	60% (144/239)	54% - 66%
FLUOROQUINOLONES	50% (118/236)	44% - 56%
Ciprofloxacin	50% (118/236)	44% - 56%
Levofloxacin	NA	-
3GC	47% (113/239)	41% - 54%
Cefpodoxime	NA	-
Ceftriaxone	47% (113/239)	41% - 54%
Ceftazidime	45% (79/174)	38% - 53%
Cefotaxime	100% (113/113)	97% - 100%
Cefepime	33% (9/27)	19% - 52%
CARBAPENEMS	2% (4/239)	0.7% - 4%
Imipenem	2% (4/239)	0.7% - 4%
Meropenem	2% (4/199)	0.8% - 5%
Ertapenem	2% (4/239)	0.7% - 4%
Doripenem	NA	-
Colistin	NA	-
Piperacillin/tazobactam	NA	-
Cefoperazone/sulbactam	NA	-

*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; FLUOROQUINOLONES: ciprofloxacin or levofloxacin; 3GC=3rd-generation cephalosporin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [2]: AMR proportion report

Blood: *Klebsiella pneumoniae*

(No. of patients = 120)




Antibiotic agent	Proportion of R	95% CI
Ampicillin	NA	-
Gentamicin	19% (23/120)	13% - 27%
Amikacin	NA	-
Co-trimoxazole	48% (58/120)	40% - 57%
FLUOROQUINOLONES	46% (52/112)	38% - 56%
Ciprofloxacin	46% (52/112)	38% - 56%
Levofloxacin	NA	-
3GC	48% (58/120)	40% - 57%
Cefpodoxime	NA	-
Ceftriaxone	48% (57/120)	39% - 56%
Ceftazidime	46% (54/117)	37% - 55%
Cefotaxime	48% (58/120)	40% - 57%
Cefepime	90% (36/40)	77% - 96%
CARBAPENEMS	3% (4/120)	1% - 8%
Imipenem	0% (0/120)	0% - 3%
Meropenem	0% (0/31)	0% - 11%
Ertapenem	3% (4/120)	1% - 8%
Doripenem	NA	-
Colistin	NA	-
Piperacillin/tazobactam	NA	-
Cefoperazone/sulbactam	NA	-

*Proportion of R represents represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; FLUOROQUINOLONES: ciprofloxacin or levofloxacin; 3GC=3rd-generation cephalosporin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [2]: AMR proportion report

Blood: *Pseudomonas aeruginosa*


(No. of patients = 48)



Antibiotic agent	Proportion of R	95% CI
Ceftazidime	10% (5/48)	4% - 22%
Ciprofloxacin	6% (3/48)	2% - 17%
AMINOGLYCOSIDES	10% (5/48)	4% - 22%
Gentamicin	10% (5/48)	4% - 22%
Amikacin	NA	-
CARBAPENEMS	10% (5/48)	4% - 22%
Imipenem	10% (5/48)	4% - 22%
Meropenem	12% (5/42)	5% - 25%
Doripenem	NA	-
Colistin	NA	-
Piperacillin/tazobactam	20% (2/10)	6% - 51%
Cefoperazone/sulbactam	NA	-

Blood: *Acinetobacter baumannii*

(No. of patients = 90)



Antibiotic agent	Proportion of R	95% CI
Tigecycline	0% (0/90)	0% - 4%
Minocycline	NA	-
AMINOGLYCOSIDES	52% (47/90)	42% - 62%
Gentamicin	52% (47/90)	42% - 62%
Amikacin	NA	-
CARBAPENEMS	59% (53/90)	49% - 68%
Imipenem	59% (53/90)	49% - 68%
Meropenem	52% (47/90)	42% - 62%
Doripenem	NA	-
Colistin	NA	-
Piperacillin/tazobactam	NA	-
Cefoperazone/sulbactam	NA	-

*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem; AMINOGLYCOSIDES: either gentamicin or amikacin

Section [3]: AMR proportion report with stratification by infection origin

Introduction

An AMR proportion report with stratification by origin of infection is generated only if admission date data are available in the raw data file(s) with the appropriate specification in the data dictionaries.

Stratification by origin of infection is used as a proxy to define where the bloodstream infection (BSI) was contracted (hospital versus community).

The definitions of infection origin proposed by the WHO GLASS are used. In brief, community–origin BSI is defined as patients in the hospital for less than or equal to two calendar days when the first specimen culture positive for the pathogen was taken. Hospital–origin BSI is defined as patients admitted for more than two calendar days when the first specimen culture positive for the pathogen was taken.

Results

The data included in the analysis to generate the report had:

*Sample collection dates ranged from **02 Jan 2016** to **10 Jan 2017***

**Number of patients with blood culture positive for pathogen under the survey:*

748 patients

***Number of patients with community–origin BSI:*

132 patients

***Number of patients with hospital–origin BSI:*

515 patients

****Number of patients with unknown infection of origin status:*

101 patients

Organism	Number of patients with blood culture positive for the organism	Community -origin**	Hospital -origin**	Unknown -origin***
<i>Staphylococcus aureus</i>	96	18	78	0
<i>Enterococcus faecalis</i>	0	0	0	0
<i>Enterococcus faecium</i>	0	0	0	0
<i>Streptococcus pneumoniae</i>	20	20	0	0
<i>Salmonella</i> spp.	35	8	27	0
<i>Escherichia coli</i>	339	35	203	101
<i>Klebsiella pneumoniae</i>	120	26	94	0
<i>Pseudomonas aeruginosa</i>	48	9	39	0
<i>Acinetobacter baumannii</i>	90	16	74	0
Total:	748	132	515	101

Note

NA=not applicable (hospital admission date or infection origin data are not available)

*Only the first isolate for each patient per specimen type per pathogen under the reporting period is included in the analysis. Please refer to Section [2] for details on how this number was calculated from the raw microbiology_data file.

**The definitions of infection origin proposed by the WHO GLASS is used. In brief, community–origin BSI was defined as patients in the hospital for less than or equal to two calendar days when the first blood culture positive for the pathogen was taken.

Hospital–origin BSI was defined as patients admitted for more than two calendar days when the first specimen culture positive for the pathogen was taken.

Please refer to the 'Methods' section for more details on the definitions used.


***Unknown origin could be because admission date data are not available or the patient was not hospitalised.

The following figures and tables below show the proportion of patients with blood culture positive for antimicrobial resistant isolates stratified by infection of origin.

Section [3]: AMR proportion report with stratification by infection origin

Blood: *Staphylococcus aureus*


Community-origin (*No. of patients = 18*)



Antibiotic agent	Proportion of R	95% CI
Methicillin	0% (0/18)	0% - 18%
Cefoxitin	0% (0/18)	0% - 18%
Oxacillin by MIC	NA	-
Vancomycin	0% (0/18)	0% - 18%
Clindamycin	6% (1/18)	1% - 26%
Chloramphenicol	NA	-

Blood: *Staphylococcus aureus*

Hospital-origin (*No. of patients = 78*)



Antibiotic agent	Proportion of R	95% CI
Methicillin	19% (15/78)	12% - 29%
Cefoxitin	19% (15/78)	12% - 29%
Oxacillin by MIC	NA	-
Vancomycin	4% (3/78)	1% - 11%
Clindamycin	22% (17/78)	14% - 32%
Chloramphenicol	NA	-

*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; Methicillin: cefoxitin or oxacillin by MIC


Section [3]: AMR proportion report with stratification by infection origin

Blood: *Enterococcus faecalis*

Community-origin (*No. of patients = 0*)

Penicillin G
Ampicillin
Vancomycin
Teicoplanin
Linezolid
Daptomycin

Antibiotic agent	Proportion of R	95% CI
Penicillin G	NA	-
Ampicillin	NA	-
Vancomycin	NA	-
Teicoplanin	NA	-
Linezolid	NA	-
Daptomycin	NA	-




Blood: *Enterococcus faecalis*

Hospital-origin (*No. of patients = 0*)

Penicillin G
Ampicillin
Vancomycin
Teicoplanin
Linezolid
Daptomycin

Antibiotic agent	Proportion of R	95% CI
Penicillin G	NA	-
Ampicillin	NA	-
Vancomycin	NA	-
Teicoplanin	NA	-
Linezolid	NA	-
Daptomycin	NA	-



*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested


Section [3]: AMR proportion report with stratification by infection origin

Blood: *Enterococcus faecium*

Community-origin (No. of patients = 0)

Penicillin G
Ampicillin
Vancomycin
Teicoplanin
Linezolid
Daptomycin

Antibiotic agent	Proportion of R	95% CI
Penicillin G	NA	-
Ampicillin	NA	-
Vancomycin	NA	-
Teicoplanin	NA	-
Linezolid	NA	-
Daptomycin	NA	-




Blood: *Enterococcus faecium*

Hospital-origin (No. of patients = 0)

Penicillin G
Ampicillin
Vancomycin
Teicoplanin
Linezolid
Daptomycin

Antibiotic agent	Proportion of R	95% CI
Penicillin G	NA	-
Ampicillin	NA	-
Vancomycin	NA	-
Teicoplanin	NA	-
Linezolid	NA	-
Daptomycin	NA	-




*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested

Section [3]: AMR proportion report with stratification by infection origin

Blood: *Streptococcus pneumoniae*

Community-origin (No. of patients = 20)




Antibiotic agent	Proportion of R	95% CI
Penicillin G	0% (0/16)	0% - 19%
Oxacillin	NA	-
Co-trimoxazole	58% (11/19)	36% - 77%
3GC	0% (0/19)	0% - 17%
Ceftriaxone	NA	-
Cefotaxime	NA	-
Erythromycin	20% (4/20)	8% - 42%
Clindamycin	20% (4/20)	8% - 42%
Levofloxacin	NA	-

Blood: *Streptococcus pneumoniae*

Hospital-origin

(No. of patients = 0)




Antibiotic agent	Proportion of R	95% CI
Penicillin G	NA	-
Oxacillin	NA	-
Co-trimoxazole	NA	-
3GC	NA	-
Ceftriaxone	NA	-
Cefotaxime	NA	-
Erythromycin	NA	-
Clindamycin	NA	-
Levofloxacin	NA	-

*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; 3GC=3rd-generation cephalosporin

Section [3]: AMR proportion report with stratification by infection origin

Blood: *Salmonella* spp.


Community-origin (No. of patients = 8)



Antibiotic agent	Proportion of R	95% CI
FLUOROQUINOLONES	62% (5/8)	31% - 86%
Ciprofloxacin	62% (5/8)	31% - 86%
Levofloxacin	NA	-
3GC	38% (3/8)	14% - 69%
Ceftriaxone	38% (3/8)	14% - 69%
Ceftazidime	NA	-
Cefotaxime	38% (3/8)	14% - 69%
CARBAPENEMS	NA	-
Imipenem	NA	-
Meropenem	NA	-
Doripenem	NA	-
Ertapenem	NA	-

Blood: *Salmonella* spp.

Hospital-origin (No. of patients = 27)




Antibiotic agent	Proportion of R	95% CI
FLUOROQUINOLONES	33% (9/27)	19% - 52%
Ciprofloxacin	33% (9/27)	19% - 52%
Levofloxacin	NA	-
3GC	7% (2/27)	2% - 23%
Ceftriaxone	7% (2/27)	2% - 23%
Ceftazidime	NA	-
Cefotaxime	7% (2/27)	2% - 23%
CARBAPENEMS	NA	-
Imipenem	NA	-
Meropenem	NA	-
Doripenem	NA	-
Ertapenem	NA	-

*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; FLUOROQUINOLONES: ciprofloxacin or levofloxacin; 3GC=3rd-generation cephalosporin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [3]: AMR proportion report with stratification by infection origin

Blood: *Escherichia coli*



Community-origin (*No. of patients = 35*)

Antibiotic agent	Proportion of R	95% CI
Ampicillin	63% (22/35)	46% - 77%
Gentamicin	37% (13/35)	23% - 54%
Amikacin	NA	-
Co-trimoxazole	69% (24/35)	52% - 81%
FLUOROQUINOLONES	63% (22/35)	46% - 77%
Ciprofloxacin	63% (22/35)	46% - 77%
Levofloxacin	NA	-
3GC	60% (21/35)	44% - 74%
Cefpodoxime	NA	-
Ceftriaxone	60% (21/35)	44% - 74%
Ceftazidime	54% (13/24)	35% - 72%
Cefotaxime	100% (21/21)	84% - 100%
Cefepime	50% (2/4)	15% - 85%
CARBAPENEMS	0% (0/35)	0% - 10%
Imipenem	0% (0/35)	0% - 10%
Meropenem	0% (0/26)	0% - 13%
Ertapenem	0% (0/35)	0% - 10%
Doripenem	NA	-
Colistin	NA	-
Piperacillin/tazobactam	NA	-
Cefoperazone/sulbactam	NA	-


*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; FLUOROQUINOLONES: ciprofloxacin or levofloxacin; 3GC=3rd-generation cephalosporin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [3]: AMR proportion report with stratification by infection origin

Blood: *Escherichia coli*

Hospital-origin

(No. of patients = 203)




Antibiotic agent	Proportion of R	95% CI
Ampicillin	48% (96/200)	41% - 55%
Gentamicin	30% (61/203)	24% - 37%
Amikacin	NA	-
Co-trimoxazole	59% (119/203)	52% - 65%
FLUOROQUINOLONES	48% (96/200)	41% - 55%
Ciprofloxacin	48% (96/200)	41% - 55%
Levofloxacin	NA	-
3GC	45% (92/203)	39% - 52%
Cefpodoxime	NA	-
Ceftriaxone	45% (92/203)	39% - 52%
Ceftazidime	44% (66/150)	36% - 52%
Cefotaxime	100% (92/92)	96% - 100%
Cefepime	30% (7/23)	16% - 51%
CARBAPENEMS	2% (4/203)	0.8% - 5%
Imipenem	2% (4/203)	0.8% - 5%
Meropenem	2% (4/172)	0.9% - 6%
Ertapenem	2% (4/203)	0.8% - 5%
Doripenem	NA	-
Colistin	NA	-
Piperacillin/tazobactam	NA	-
Cefoperazone/sulbactam	NA	-

*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; FLUOROQUINOLONES: ciprofloxacin or levofloxacin; 3GC=3rd-generation cephalosporin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [3]: AMR proportion report with stratification by infection origin

Blood: *Klebsiella pneumoniae*

Community-origin (*No. of patients = 26*)



Antibiotic agent	Proportion of R	95% CI
Ampicillin	NA	-
Gentamicin	27% (7/26)	14% - 46%
Amikacin	NA	-
Co-trimoxazole	54% (14/26)	36% - 71%
FLUOROQUINOLONES	52% (12/23)	33% - 71%
Ciprofloxacin	52% (12/23)	33% - 71%
Levofloxacin	NA	-
3GC	54% (14/26)	36% - 71%
Cefpodoxime	NA	-
Ceftriaxone	54% (14/26)	36% - 71%
Ceftazidime	52% (13/25)	34% - 70%
Cefotaxime	54% (14/26)	36% - 71%
Cefepime	100% (10/10)	72% - 100%
CARBAPENEMS	8% (2/26)	2% - 24%
Imipenem	0% (0/26)	0% - 13%
Meropenem	0% (0/9)	0% - 30%
Ertapenem	8% (2/26)	2% - 24%
Doripenem	NA	-
Colistin	NA	-
Piperacillin/tazobactam	NA	-
Cefoperazone/sulbactam	NA	-


*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; FLUOROQUINOLONES: ciprofloxacin or levofloxacin; 3GC=3rd-generation cephalosporin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [3]: AMR proportion report with stratification by infection origin

Blood: *Klebsiella pneumoniae*

Hospital-origin

(No. of patients = 94)




Antibiotic agent	Proportion of R	95% CI
Ampicillin	NA	-
Gentamicin	17% (16/94)	11% - 26%
Amikacin	NA	-
Co-trimoxazole	47% (44/94)	37% - 57%
FLUOROQUINOLONES	45% (40/89)	35% - 55%
Ciprofloxacin	45% (40/89)	35% - 55%
Levofloxacin	NA	-
3GC	47% (44/94)	37% - 57%
Cefpodoxime	NA	-
Ceftriaxone	46% (43/94)	36% - 56%
Ceftazidime	45% (41/92)	35% - 55%
Cefotaxime	47% (44/94)	37% - 57%
Cefepime	87% (26/30)	70% - 95%
CARBAPENEMS	2% (2/94)	0.6% - 7%
Imipenem	0% (0/94)	0% - 4%
Meropenem	0% (0/22)	0% - 15%
Ertapenem	2% (2/94)	0.6% - 7%
Doripenem	NA	-
Colistin	NA	-
Piperacillin/tazobactam	NA	-
Cefoperazone/sulbactam	NA	-

*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; FLUOROQUINOLONES: ciprofloxacin or levofloxacin; 3GC=3rd-generation cephalosporin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [3]: AMR proportion report with stratification by infection origin

Blood: *Pseudomonas aeruginosa*


Community-origin (No. of patients = 9)



Antibiotic agent	Proportion of R	95% CI
Ceftazidime	11% (1/9)	2% - 44%
Ciprofloxacin	11% (1/9)	2% - 44%
AMINOGLYCOSIDES	11% (1/9)	2% - 44%
Gentamicin	11% (1/9)	2% - 44%
Amikacin	NA	-
CARBAPENEMS	11% (1/9)	2% - 44%
Imipenem	11% (1/9)	2% - 44%
Meropenem	17% (1/6)	3% - 56%
Doripenem	NA	-
Colistin	NA	-
Piperacillin/tazobactam	50% (1/2)	10% - 90%
Cefoperazone/sulbactam	NA	-

Blood: *Pseudomonas aeruginosa*

Hospital-origin (No. of patients = 39)




Antibiotic agent	Proportion of R	95% CI
Ceftazidime	10% (4/39)	4% - 24%
Ciprofloxacin	5% (2/39)	1% - 17%
AMINOGLYCOSIDES	10% (4/39)	4% - 24%
Gentamicin	10% (4/39)	4% - 24%
Amikacin	NA	-
CARBAPENEMS	10% (4/39)	4% - 24%
Imipenem	10% (4/39)	4% - 24%
Meropenem	11% (4/36)	4% - 25%
Doripenem	NA	-
Colistin	NA	-
Piperacillin/tazobactam	12% (1/8)	2% - 47%
Cefoperazone/sulbactam	NA	-

*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; AMINOGLYCOSIDES: either gentamicin or amikacin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [3]: AMR proportion report with stratification by infection origin

Blood: *Acinetobacter baumannii*


Community-origin (No. of patients = 16)



Antibiotic agent	Proportion of R	95% CI
Tigecycline	0% (0/16)	0% - 19%
Minocycline	NA	-
AMINOGLYCOSIDES	62% (10/16)	39% - 82%
Gentamicin	62% (10/16)	39% - 82%
Amikacin	NA	-
CARBAPENEMS	62% (10/16)	39% - 82%
Imipenem	62% (10/16)	39% - 82%
Meropenem	56% (9/16)	33% - 77%
Doripenem	NA	-
Colistin	NA	-
Piperacillin/tazobactam	NA	-
Cefoperazone/sulbactam	NA	-

Blood: *Acinetobacter baumannii*

Hospital-origin (No. of patients = 74)



Antibiotic agent	Proportion of R	95% CI
Tigecycline	0% (0/74)	0% - 5%
Minocycline	NA	-
AMINOGLYCOSIDES	50% (37/74)	39% - 61%
Gentamicin	50% (37/74)	39% - 61%
Amikacin	NA	-
CARBAPENEMS	58% (43/74)	47% - 69%
Imipenem	58% (43/74)	47% - 69%
Meropenem	51% (38/74)	40% - 62%
Doripenem	NA	-
Colistin	NA	-
Piperacillin/tazobactam	NA	-
Cefoperazone/sulbactam	NA	-

*Proportion of R represents the number of patients with blood culture positive for resistant isolates (numerator) over the total number of patients with blood culture positive for the organism and the organism was tested for susceptibility against the antibiotic (denominator). The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period. Grey bars indicate that AST unknown results are more than 30% of the total number of patients with blood culture positive for the organism. CI=confidence interval; NA=not available/reported/tested; AMINOGLYCOSIDES: either gentamicin or amikacin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [4]: AMR frequency report

Introduction

A AMR frequency report is generated if data of culture negative is available.

The AMR frequency approach involves the collection of data on all blood samples taken for microbiological testing and includes information on the number of positive blood samples for a specific specimen type (both pathogens under the survey and other bacteria) as well as number of negative (no microbial growth) samples. After removal of duplicate results and assuming that routine blood culture testing is applied systematically, we can use the number of tested patients as a proxy for a number of patients with new cases of bloodstream infection (BSI).

Results

The microbiology_data file had:

*Specimen collection dates ranged from **02 Jan 2016** to **10 Jan 2017***

Number of records on blood specimen collected within the above date range:

15878 blood specimen records

**Number of patients sampled for blood culture within the above date range:*

15638 patients sampled for blood culture

Note

***Number of patients sampled for blood culture is used as denominator to estimate the frequency of infections per 100,000 tested patients**

The following figures show the frequency of infections for patients with blood culture tested.

Section [4]: AMR frequency report

Blood: Pathogens

(No. of patients = 15638)




Pathogens	*Frequency of infection (per 100,000 tested patients; 95% CI)
<i>S. aureus</i>	614 (504-750)
<i>E. faecalis</i>	0 (0-25)
<i>E. faecium</i>	0 (0-25)
<i>S. pneumoniae</i>	128 (83-198)
<i>Salmonella</i> spp.	224 (161-312)
<i>E. coli</i>	2168 (1952-2409)
<i>K. pneumoniae</i>	768 (643-917)
<i>P. aeruginosa</i>	307 (232-407)
<i>A. baumannii</i>	576 (469-707)

*Frequency of infection per 100,000 tested patients represents the number of patients with blood culture positive for a pathogen (numerator) over the total number of tested patients (denominator). The AMASS application de-duplicates the data by included only the first isolate of each patient per specimen type per reporting period. CI=confidence interval; R=resistant; NA=not available/reported/tested

Section [4]: AMR frequency report

Blood: Resistant pathogens

(No. of patients = 15638)



Resistant (NS) pathogens	*Frequency of infection (per 100,000 tested patients; 95% CI)
MRSA	96 (59-159)
Vancomycin-R <i>E. faecalis</i>	0 (0-25)
Vancomycin-R <i>E. faecium</i>	0 (0-25)
Penicillin-R <i>S. pneumoniae</i>	0 (0-25)
Fluoroquinolone-R <i>Salmonella</i> spp.	90 (54-151)
3GC-R <i>E. coli</i>	723 (602-868)
Carbapenem-R <i>E. coli</i>	26 (10-66)
3GC-R <i>K. pneumoniae</i>	371 (288-480)
Carbapenem-R <i>K. pneumoniae</i>	26 (10-66)
Carbapenem-R <i>P. aeruginosa</i>	32 (14-75)
Carbapenem-R <i>A. baumannii</i>	339 (260-444)

*Frequency of infection per 100,000 tested patients represents the number of patients with blood culture positive for a pathogen (numerator) over the total number of tested patients (denominator). The AMASS application de-duplicates the data by included only the first isolate of each patient per specimen type per reporting period. CI=confidence interval; R=resistant; NA=not available/reported/tested ; FLUOROQUINOLONES: ciprofloxacin or levofloxacin; 3GC=3rd-generation cephalosporin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [5]: AMR frequency report with stratification by infection origin

Introduction

A AMR frequency report with stratification by origin of infection is generated only if data of culture negative is available and admission date or a variable containing the classification is available in the raw data file with the appropriate specification in the data dictionaries.

Results

The data included in the analysis had:

*Specimen collection dates ranged from **02 Jan 2016** to **10 Jan 2017***

Number of records on blood specimen collected within the above date range:

15878 blood specimen records

Number of patients sampled for blood culture within the above date range:

15638 patients sampled for blood culture

2919 patients had at least one admission having the first blood culture drawn within first 2 calendar days of hospital admission.

This parameter is used as a denominators for frequency of community–origin bacteraemia (per 100,000 patients tested for blood culture on admission).

11765 patients had at least one admission having the first blood culture drawn after 2 calendar days of hospital admission.

This parameter is used as a denominators for frequency of hospital–origin bacteraemia (per 100,000 patients tested for blood culture for HAI).

1041 patients had a blood drawn for culture and with unknown origin of infection.

Validation of this statistics is highly recommended.

Note:


87 patients had more than one admissions, of which at least one admission had the first blood culture drawn within the first 2 calendar days of hospital admission AND at least one admission had the first blood culture drawn after 2 calendar days of hospital admission.

The following figures show the frequency of infections for patients with blood culture tested and stratified by infection origin, under this surveillance.

Section [5]: AMR frequency report with stratification by infection origin

Blood: Pathogens

Community-origin (*No. of patients = 2919*)



Pathogens	*Frequency of infection (per 100,000 tested patients; 95% CI)
<i>S. aureus</i>	617 (391-973)
<i>E. faecalis</i>	0 (0-132)
<i>E. faecium</i>	0 (0-132)
<i>S. pneumoniae</i>	686 (444-1056)
<i>Salmonella</i> spp.	275 (139-540)
<i>E. coli</i>	1200 (864-1663)
<i>K. pneumoniae</i>	891 (609-1302)
<i>P. aeruginosa</i>	309 (163-585)
<i>A. baumannii</i>	549 (338-889)


*Frequency of infection per 100,000 tested patients on admission represents the number of patients with blood culture positive for a pathogen (numerator) over the total number of tested population on admission (denominator). The AMASS application de-duplicates the data by included only the first isolate of each patient per specimen type per reporting period. CI=confidence interval; NA=not available/reported/tested

Section [5]: AMR frequency report with stratification by infection origin

Blood: Pathogens

Hospital-origin

(No. of patients = 11765)




Pathogens	*Frequency of infection (per 100,000 tested patients; 95% CI)
<i>S. aureus</i>	663 (532-827)
<i>E. faecalis</i>	0 (0-33)
<i>E. faecium</i>	0 (0-33)
<i>S. pneumoniae</i>	0 (0-33)
<i>Salmonella</i> spp.	230 (158-334)
<i>E. coli</i>	1726 (1506-1978)
<i>K. pneumoniae</i>	799 (654-977)
<i>P. aeruginosa</i>	332 (243-453)
<i>A. baumannii</i>	629 (502-789)

*Frequency of infection per 100,000 tested population at risk of HAI represents the number of patients with blood culture positive for a pathogen (numerator) over the total number of tested population at risk of HAI (denominator). The AMASS application de-duplicates the data by included only the first isolate of each patient per specimen type per reporting period. CI=confidence interval; NA=not available/reported/tested

Section [5]: AMR frequency report with stratification by infection origin

Blood: Resistant pathogens

Community-origin (*No. of patients = 2919*)



Pathogens	*Frequency of infection (per 100,000 tested patients; 95% CI)
MRSA	0 (0-132)
Vancomycin-R <i>E. faecalis</i>	0 (0-132)
Vancomycin-R <i>E. faecium</i>	0 (0-132)
Penicillin-R <i>S. pneumoniae</i>	0 (0-132)
Fluoroquinolone-R <i>Salmonella</i> spp.	172 (74-401)
3GC-R <i>E. coli</i>	720 (472-1098)
Carbapenem-R <i>E. coli</i>	0 (0-132)
3GC-R <i>K. pneumoniae</i>	480 (286-804)
Carbapenem-R <i>K. pneumoniae</i>	69 (19-250)
Carbapenem-R <i>P. aeruginosa</i>	35 (7-194)
Carbapenem-R <i>A. baumannii</i>	343 (187-630)


*Frequency of infection per 100,000 tested patients on admission represents the number of patients with blood culture positive for a pathogen (numerator) over the total number of tested population on admission (denominator). The AMASS application de-duplicates the data by included only the first isolate of each patient per specimen type per reporting period. CI=confidence interval; NA=not available/reported/tested; FLUOROQUINOLONES: ciprofloxacin or levofloxacin; 3GC=3rd-generation cephalosporin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [5]: AMR frequency report with stratification by infection origin

Blood: Resistant pathogens

Hospital-origin

(No. of patients = 11765)



Pathogens	*Frequency of infection (per 100,000 tested patients; 95% CI)
MRSA	128 (78-211)
Vancomycin-R <i>E. faecalis</i>	0 (0-33)
Vancomycin-R <i>E. faecium</i>	0 (0-33)
Penicillin-R <i>S. pneumoniae</i>	0 (0-33)
Fluoroquinolone-R <i>Salmonella</i> spp.	77 (41-146)
3GC-R <i>E. coli</i>	782 (639-958)
Carbapenem-R <i>E. coli</i>	34 (14-88)
3GC-R <i>K. pneumoniae</i>	374 (279-502)
Carbapenem-R <i>K. pneumoniae</i>	17 (5-62)
Carbapenem-R <i>P. aeruginosa</i>	34 (14-88)
Carbapenem-R <i>A. baumannii</i>	366 (272-492)

*Frequency of infection per 100,000 tested patients represents the number of patients with blood culture positive for a pathogen (numerator) over the total number of tested patients (denominator). The AMASS application de-duplicates the data by included only the first isolate of each patient per specimen type per reporting period. CI=confidence interval; NA=not available/reported/tested; FLUOROQUINOLONES: ciprofloxacin or levofloxacin; 3GC=3rd-generation cephalosporin; CARBAPENEMS: imipenem, meropenem, ertapenem or doripenem

Section [6] Mortality involving AMR and antimicrobial–susceptible infections

Introduction

A surveillance report on mortality involving AMR infections and antimicrobial–susceptible infections with stratification by origin of infection is generated only if data on patient outcomes (i.e. discharge status) are available. Antimicrobial–resistant infection is a threat to modern health care, and the impact of the infection on patient outcomes is largely unknown. Performing analyses and generating reports on mortality often takes time and resources.

The term 'mortality involving AMR and antimicrobial–susceptible infections' was used because the mortality reported was all–cause mortality. This measure of mortality included deaths caused by or related to other underlying and intermediate causes.

Here, AMASS summarized the overall mortality of patients with antimicrobial–resistant and antimicrobial–susceptible bacteria bloodstream infections (BSI).

Results

The data included in the analysis had:

*Sample collection dates ranged from **02 Jan 2016** to **10 Jan 2017***

Number of patients with blood culture positive for the organism under the survey:

748 patients

Number of patients with community–origin BSI:

132 patients

Number of patients with hospital–origin BSI:

515 patients

The hospital admission data file had:

*Hospital admission dates ranging from **01 Jan 2016** to **31 Dec 2016***

Number of records in the raw hospital admission data:

247260 records

Number of patients included in the analysis (de–duplicated):

242659 patients

Number of patients having death as an outcome in any admission data records:

30850 patients

Overall mortality:

13% (30850/242659)

The AMASS application merged the microbiology data file and hospital admission data file. The merged dataset was then de-duplicated so that only the first isolate per patient per specimen per reporting period was included in the analysis. The de-duplicated data was stratified by infection origin (community-origin infection or hospital-origin infection).


Organism	Mortality in patients with Community-origin BSI	Mortality in patients with Hospital-origin BSI
<i>Staphylococcus aureus</i>	11% (2/18)	6% (5/78)
<i>Enterococcus faecalis</i>	NA	NA
<i>Enterococcus faecium</i>	NA	NA
<i>Streptococcus pneumoniae</i>	0% (0/20)	NA
<i>Salmonella</i> spp.	12% (1/8)	0% (0/27)
<i>Escherichia coli</i>	9% (3/35)	9% (19/203)
<i>Klebsiella pneumoniae</i>	19% (5/26)	16% (15/94)
<i>Pseudomonas aeruginosa</i>	22% (2/9)	8% (3/39)
<i>Acinetobacter baumannii</i>	19% (3/16)	22% (16/74)
Total:	12% (16/132)	11% (58/515)

The following figures and tables show the mortality of patients who were blood culture positive for antimicrobial resistant and susceptible isolates.

Section [6] Mortality involving AMR and antimicrobial-susceptible infections

Blood: *Staphylococcus aureus*

Community-origin



Type of pathogen	Mortality (n)	95% CI
MRSA	NA	-
MSSA	11% (2/18)	3% - 33%

Blood: *Staphylococcus aureus*


Hospital-origin



Type of pathogen	Mortality (n)	95% CI
MRSA	7% (1/15)	1% - 30%
MSSA	6% (4/63)	2% - 15%

Blood: *Enterococcus faecalis*


Community-origin



Type of pathogen	Mortality (n)	95% CI
Vancomycin-R	NA	-
Vancomycin-S	NA	-

Blood: *Enterococcus faecalis*

Hospital-origin



Type of pathogen	Mortality (n)	95% CI
Vancomycin-R	NA	-
Vancomycin-S	NA	-

*Mortality is the proportion (%) of in-hospital deaths (all-cause deaths). This represents the number of in-hospital deaths (numerator) over the total number of patients with blood culture positive for the organism and the type of pathogen (denominator). The AMASS application de-duplicates the data by included only the first isolate per patient per specimen type per evaluation period. R=resistant; S=susceptible (including sensitive and intermediate categories); CI=confidence interval


Section [6] Mortality involving AMR and antimicrobial–susceptible infections

Blood: *Enterococcus faecium*

Community-origin

Vancomycin-R

Vancomycin-S




Type of pathogen	Mortality (n)	95% CI
Vancomycin-R	NA	-
Vancomycin-S	NA	-

Blood: *Enterococcus faecium*

Hospital-origin

Vancomycin-R

Vancomycin-S




Type of pathogen	Mortality (n)	95% CI
Vancomycin-R	NA	-
Vancomycin-S	NA	-

Blood: *Streptococcus pneumoniae*

Community-origin

Penicillin-R

Penicillin-S




Type of pathogen	Mortality (n)	95% CI
Penicillin-R	NA	-
Penicillin-S	0% (0/16)	0% - 19%

Blood: *Streptococcus pneumoniae*

Hospital-origin

Penicillin-R

Penicillin-S




Type of pathogen	Mortality (n)	95% CI
Penicillin-R	NA	-
Penicillin-S	NA	-

*Mortality is the proportion (%) of in-hospital deaths (all-cause deaths). This represents the number of in-hospital deaths (numerator) over the total number of patients with blood culture positive for the organism and the type of pathogen (denominator). The AMASS application de-duplicates the data by included only the first isolate per patient per specimen type per evaluation period. R=resistant; S=susceptible (including sensitive and intermediate categories); CI=confidence interval

Section [6] Mortality involving AMR and antimicrobial-susceptible infections

Blood: *Salmonella* spp.


Community-origin



Type of pathogen	Mortality (n)	95% CI
Fluoroquinolone-R	20% (1/5)	4% - 62%
Fluoroquinolone-S	0% (0/3)	0% - 56%

Blood: *Salmonella* spp.

Hospital-origin



Type of pathogen	Mortality (n)	95% CI
Fluoroquinolone-R	0% (0/9)	0% - 30%
Fluoroquinolone-S	0% (0/18)	0% - 18%

Blood: *Escherichia coli*


Community-origin



Type of pathogen	Mortality (n)	95% CI
Carbapenem-R	NA	-
3GC-R**	10% (2/21)	3% - 29%
3GC-S***	7% (1/14)	1% - 32%

Blood: *Escherichia coli*

Hospital-origin




Type of pathogen	Mortality (n)	95% CI
Carbapenem-R	25% (1/4)	5% - 70%
3GC-R**	8% (7/88)	4% - 16%
3GC-S***	10% (11/111)	6% - 17%

*Mortality is the proportion (%) of in-hospital deaths (all-cause deaths). This represents the number of in-hospital deaths (numerator) over the total number of patients with blood culture positive for the organism and the type of pathogen (denominator). The AMASS application de-duplicates the data by included only the first isolate per patient per specimen type per evaluation period. R=resistant; S=susceptible (including sensitive and intermediate categories); CI=confidence interval; Fluoroquinolone-R=R to any fluoroquinolone tested; Carbapenem-R=R to any Carbapenem tested; **3GC-R [for this section]: R to any 3rd-generation cephalosporin excluding isolates which are resistant to carbapenem; ***3GC-S [for this section]: S to all 3rd-generation cephalosporin tested excluding isolates which are resistant to carbapenem

Section [6] Mortality involving AMR and antimicrobial-susceptible infections

Blood: *Klebsiella pneumoniae*


Community-origin



Type of pathogen	Mortality (n)	95% CI
Carbapenem-R	0% (0/2)	0% - 66%
3GC-R**	33% (4/12)	14% - 61%
3GC-S***	8% (1/12)	2% - 35%

Blood: *Klebsiella pneumoniae*


Hospital-origin



Type of pathogen	Mortality (n)	95% CI
Carbapenem-R	100% (2/2)	34% - 100%
3GC-R**	29% (12/42)	17% - 44%
3GC-S***	2% (1/50)	0.4% - 10%

Blood: *Pseudomonas aeruginosa*


Community-origin



Type of pathogen	Mortality (n)	95% CI
Carbapenem-R	100% (1/1)	21% - 100%
Carbapenem-S	12% (1/8)	2% - 47%

Blood: *Pseudomonas aeruginosa*

Hospital-origin




Type of pathogen	Mortality (n)	95% CI
Carbapenem-R	50% (2/4)	15% - 85%
Carbapenem-S	3% (1/35)	0.5% - 14%

*Mortality is the proportion (%) of in-hospital deaths (all-cause deaths). This represents the number of in-hospital deaths (numerator) over the total number of patients with blood culture positive for the organism and the type of pathogen (denominator). The AMASS application de-duplicates the data by included only the first isolate per patient per specimen type per evaluation period. R=resistant; S=susceptible (including sensitive and intermediate categories); CI=confidence interval; Carbapenem-R=R to any Carbapenem tested; **3GC-R [for this section]: R to any 3rd-generation cephalosporin excluding isolates which are resistant to carbapenem; ***3GC-S [for this section]: S to all 3rd-generation cephalosporin tested excluding isolates which are resistant to carbapenem

Section [6] Mortality involving AMR and antimicrobial-susceptible infections

Blood: *Acinetobacter baumannii*


Community-origin



Type of pathogen	Mortality (n)	95% CI
Carbapenem-R	30% (3/10)	11% - 60%
Carbapenem-S	0% (0/6)	0% - 39%

Blood: *Acinetobacter baumannii*

Hospital-origin



Type of pathogen	Mortality (n)	95% CI
Carbapenem-R	33% (14/43)	20% - 48%
Carbapenem-S	6% (2/31)	2% - 21%

*Mortality is the proportion (%) of in-hospital deaths (all-cause deaths). This represents the number of in-hospital deaths (numerator) over the total number of patients with blood culture positive for the organism and the type of pathogen (denominator). The AMASS application de-duplicates the data by included only the first isolate per patient per specimen type per evaluation period. R=resistant; S=susceptible (including sensitive and intermediate categories); CI=confidence interval; Carbapenem-R=R to any Carbapenem tested

Annex A: Supplementary report on notifiable bacterial infections

Introduction

This supplementary report has two parts; including (A1) notifiable bacterial infections and (A2) mortality involving notifiable bacterial infections. The AMR proportion notifiable bacterial infections supplementary report is generated by default, even if the hospital_admission_data file is unavailable. This is to enable hospitals with only microbiology data available to utilize the de-duplication and report generation functions of AMASS.

Please note that the completion of this supplementary report is strongly associated with the availability of data (particularly, all bacterial pathogens and all types of specimens) and the completion of the data dictionary files to make sure that the AMASS application understands the notifiable bacteria and each type of specimens.

Annex A includes various type of specimens including blood, cerebrospinal fluid (CSF), respiratory tract specimens, urine, genital swab, stool and other or unknown sample types. The microorganisms in this report were initially selected from common notifiable bacterial diseases in Thailand.

Notifiable bacteria under the survey

- | | |
|----------------------------------------|------------------------|
| - <i>B. pseudomallei</i> | - <i>S. Paratyphi</i> |
| - <i>Brucella</i> spp. | - <i>S. Typhi</i> |
| - <i>C. diphtheriae</i> | - <i>Shigella</i> spp. |
| - <i>N. gonorrhoeae</i> | - <i>S. suis</i> |
| - <i>N. meningitidis</i> | - <i>Vibrio</i> spp. |
| - Non-typhoidal <i>Salmonella</i> spp. | |

Note: The list of notifiable bacteria included in the AMASS application version 3.0 was generated based on the literature review and the collaboration with Department of Disease Control, Ministry of Public Health, Thailand. The list could be expanded or modified in the next version of AMASS.

Annex A1: Notifiable bacterial infections

Results

The microbiology_data file had:

Sample collection dates ranged from 02 Jan 2016 to 10 Jan 2017

Number of records of clinical specimens collected with culture positive for a notifiable bacteria under this survey:

615 specimen records (176 , 3 , 3 , 99 , 125 , 175 , 34 were blood, CSF, genital swab, respiratory tract specimens, stool, urine, and other or unknown sample types, respectively)

The AMASS application de-duplicated the data by including only the first isolate per patient per specimen type per evaluation period as described in the method. The number of patients with positive samples is as follows:


Pathogens	Total number of patients*	Blood	CSF	Genital swab	RTS	Stool	Urine	Others
<i>B. pseudomallei</i>	331	109	3	3	92	0	155	34
<i>Brucella</i> spp.	0	0	0	0	0	0	0	0
<i>C. diphtheriae</i>	0	0	0	0	0	0	0	0
<i>N. gonorrhoeae</i>	0	0	0	0	0	0	0	0
<i>N. meningitidis</i>	0	0	0	0	0	0	0	0
Non-typhoidal <i>Salmonella</i> spp.	60	35	0	0	0	54	0	0
<i>S. Paratyphi</i>	0	0	0	0	0	0	0	0
<i>S. Typhi</i>	0	0	0	0	0	0	0	0
<i>Shigella</i> spp.	0	0	0	0	0	0	0	0
<i>S. suis</i>	0	0	0	0	0	0	0	0
<i>Vibrio</i> spp.	16	12	0	0	0	16	0	0
Total	407	156	3	3	92	70	155	34

*Some patients may have more than one type of clinical specimen culture positive for the notifiable bacteria under the survey, and some may have more than one notifiable organism per evaluation period.

CSF = Cerebrospinal fluid; RTS = Respiratory tract specimens; Others = Other or unknown sample types; NA = Not applicable (i.e. the specimen type is not available or identified in the microbiology_data file)

Annex A2: Mortality involving notifiable bacterial infections

A report on mortality involving notifiable bacterial infections is generated only if data on patient outcomes (i.e. discharge status) are available. The term "mortality involving notifiable bacterial infections" was used because the mortality reported was all-cause mortality. This measure of mortality included deaths caused by or related to other underlying and intermediate causes. The AMASS application merged the microbiology data file and hospital admission data file. The merged dataset was then de-duplicated so that only the first isolate per patient per specimen per reporting period was included in the analysis.



Pathogens	Mortality (n)	95% CI
<i>B. pseudomallei</i>	24% (79/330)	20% - 29%
<i>Brucella</i> spp.	NA	-
<i>C. diphtheriae</i>	NA	-
<i>N. gonorrhoeae</i>	NA	-
<i>N. meningitidis</i>	NA	-
Non-typhoidal <i>Salmonella</i> spp.	5% (3/60)	2% - 14%
<i>S. Paratyphi</i>	NA	-
<i>S. Typhi</i>	NA	-
<i>Shigella</i> spp.	NA	-
<i>S. suis</i>	NA	-
<i>Vibrio</i> spp.	6% (1/16)	1% - 28%

*Mortality is the proportion (%) of in-hospital deaths (all-cause deaths). This represents the number of in-hospital deaths (numerator) over the total number of patients with culture positive for each type of pathogen (denominator). Some patients may have the data of a clinical specimen culture positive for the notifiable bacteria under the survey in the microbiology data file, but do not have the data in the hospital admission data file. That is the most common cause of the discrepancy between total number of patients with notifiable bacterial infections presented in the Annex A1 and the Annex A2 (followed by typos in patient identifiers in either data file).

CI = confidence interval

Annex B: Supplementary report on data indicators

Introduction

This supplementary report is generated by default, even if the hospital_admission_data file is unavailable. The management of clinical and laboratory practice can be supported by some data indicators such as blood culture contamination rate, proportion of notifiable antibiotic-pathogen combinations, and proportion of isolates with infrequent phenotypes or potential errors in AST results. Isolates with infrequent phenotypes or potential errors in AST results include (a) reports of organisms which are intrinsically resistant to an antibiotic but are reported as susceptible and (b) reports of organisms with discordant AST results.

This supplementary report could support the clinicians, policy makers and the laboratory staff to understand their summary data quickly. The laboratory staff could also use "Supplementary_data_indicators_report.pdf" generated in the folder "Report_with_patient_identifiers" to check and validate individual data records further.

This supplementary report was estimated from data of blood specimens only. Please note that the data indicators do not represent quality of the clinical or laboratory practice.

Results

Indicators	Number of observations			
	Total (n)	Critical priority (n)	High priority (n)	Medium priority (n)
Blood culture contamination rate*	5% (742/15878)	NA	NA	5% (742/15878)
Proportion of notifiable antibiotic-pathogen combinations**	30% (308/1017)	27% (275/1017)	3% (33/1017)	0% (0/1017)
Proportion of isolates with infrequent phenotypes or potential errors in AST results***	10% (100/1017)	NA	NA	10% (100/1017)

*Blood culture contamination rate is defined as the number of raw contaminated cultures per number of blood cultures received by the laboratory per reporting period. Blood culture contamination rate will not be estimated in case that the data of negative culture (specified as 'no growth' in the dictionary_for_microbiology_data file) is not available. **Notifiable antibiotic-pathogen combinations and their classifications are defined as WHO list of AMR priority pathogen published in 2017. **, ***The proportion is estimated per number of blood specimens culture positive for any organisms with AST result in the raw microbiology data. *, **, ***Details of the criteria are available in Table 3 and Table 4 of "Supplementary_data_indicators_report.pdf", and "list_of_indicators.xlsx" in the folder "Configuration". NA = Not applicable

Annex B: Supplementary report on data indicators

Reporting period by months

Data was stratified by month to assist detection of missing data and understand the change of indicators by months.

Month	Blood culture contamination rate (n)*	Proportion of notifiable antibiotic-pathogen combinations (n)**	Proportion of isolates with infrequent phenotypes or potential errors in AST results (n)***
January	4% (59/1316)	32% (29/92)	13% (12/92)
February	6% (69/1256)	33% (28/85)	7% (6/85)
March	4% (53/1331)	28% (23/81)	6% (5/81)
April	4% (53/1382)	29% (21/73)	8% (6/73)
May	4% (57/1345)	32% (24/76)	5% (4/76)
June	4% (56/1269)	23% (15/66)	11% (7/66)
July	4% (58/1361)	31% (32/103)	10% (10/103)
August	5% (70/1344)	32% (30/94)	10% (9/94)
September	5% (58/1261)	43% (34/79)	9% (7/79)
October	4% (60/1365)	26% (28/108)	12% (13/108)
November	6% (78/1301)	28% (22/78)	15% (12/78)
December	5% (71/1347)	27% (22/82)	11% (9/82)

*Blood culture contamination rate is defined as the number of raw contaminated cultures per number of blood cultures received by the laboratory per reporting period. Blood culture contamination rate will not be estimated in case that the data of negative culture (specified as 'no growth' in the dictionary_for_microbiology_data file) is not available. **Notifiable antibiotic-pathogen combinations and their classifications are defined as WHO list of AMR priority pathogen published in 2017. **, ***The proportion is estimated per number of blood specimens culture positive for any organisms with AST result in the raw microbiology data. *, **, ***Details of the criteria are available in Table 3 and Table 4 of "Supplementary_data_indicators_report.pdf", and "list_of_indicators.xlsx" in the folder "Configuration". NA = Not applicable

Annex C: Cluster signals

Introduction

This supplementary report shows the information of potential clusters which are identified using the SatScan (www.satscan.org). An outbreak of hospital-acquired infection (HAI) is defined as an increase in the occurrence of hospital-origin bloodstream infection caused by AMR pathogens (BSI; i.e. blood specimen model) and of any specimen culture positive for the AMR pathogen (i.e. all specimen model) compared to the recorded baseline rates. This report is generated by default, even if hospital_admission_data is unavailable. This is to enable hospitals with only microbiology data to utilize the de-duplication and automation of AMASS-SatScan and report generation functions of AMASS.

The AMASS-SatScan considers each ward as an independent ward within the hospital. In case that there are no ward data in the microbiology data file or that the dictionary for ward are not available, the model will consider all wards in the hospital as a single ward.

Pathogens under the survey

Methicillin-resistant *S. aureus* (MRSA)
Vancomycin-resistant *E. faecalis* (VREfs)
Vancomycin-resistant *E. faecium* (VREfm)
Carbapenem-resistant *E. coli* (CREC)
Carbapenem-resistant *K. pneumoniae* (CRKP)
Carbapenem-resistant *P. aeruginosa* (CRPA)
Carbapenem-resistant *A. baumannii* (CRAB)

Results

Annex C: Cluster signals


Blood specimen: MRSA

Hospital-origin

No. of patients = 19 (3 [16%] were included in cluster signals)

No. of wards = 2 (1 [50%] were included in cluster signals)

No. of AMR profiles = 2 (1 [50%] were included in cluster signals)



Wards with cluster signals with p value < 0.05.

Ward ID	Total no. of observed cases in any cluster by ward	AMR Profile	Start date	End date	No. of observed cases in each cluster	P-value
ward_001	3	profile_MRSA_1	2016-10-18	2016-10-25	3	0.0066

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with blood culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with blood culture positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals


All specimens: MRSA

Hospital-origin

No. of patients = 56 (3 [6%] were included in cluster signals)

No. of wards = 2 (1 [50%] were included in cluster signals)

No. of AMR profiles = 2 (1 [50%] were included in cluster signals)



Wards with cluster signals with p value < 0.05.

Ward ID	Total no. of observed cases in any cluster by ward	AMR Profile	Start date	End date	No. of observed cases in each cluster	P-value
ward_001	3	profile_MRSA_1	2016-10-18	2016-10-25	3	0.005

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with a clinical specimen culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with a clinical specimen positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals

Blood specimen: VREfs

Hospital-origin

No. of patients = 0 (0 [0%] were included in cluster signals)

No. of wards = 0 (0 [0%] were included in cluster signals)

No. of AMR profiles = 0 (0 [0%] were included in cluster signals)

There is no cluster signal with p-value < 0.05.

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with blood culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with blood culture positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals

All specimens: VREfs

Hospital-origin

No. of patients = 0 (0 [0%] were included in cluster signals)

No. of wards = 0 (0 [0%] were included in cluster signals)

No. of AMR profiles = 0 (0 [0%] were included in cluster signals)

There is no cluster signal with p-value < 0.05.

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with a clinical specimen culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with a clinical specimen positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals

Blood specimen: VREfm

Hospital-origin

No. of patients = 0 (0 [0%] were included in cluster signals)

No. of wards = 0 (0 [0%] were included in cluster signals)

No. of AMR profiles = 0 (0 [0%] were included in cluster signals)

There is no cluster signal with p-value < 0.05.

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with blood culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with blood culture positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals

All specimens: VREfm

Hospital-origin

No. of patients = 0 (0 [0%] were included in cluster signals)

No. of wards = 0 (0 [0%] were included in cluster signals)

No. of AMR profiles = 0 (0 [0%] were included in cluster signals)

There is no cluster signal with p-value < 0.05.

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with a clinical specimen culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with a clinical specimen positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals


Blood specimen: CREC

Hospital-origin

No. of patients = 4 (0 [0%] were included in cluster signals)

No. of wards = 1 (0 [0%] were included in cluster signals)

No. of AMR profiles = 1 (0 [0%] were included in cluster signals)



There is no cluster signal with p-value < 0.05.

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with blood culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with blood culture positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals


All specimens: CREC

Hospital-origin

No. of patients = 12 (0 [0%] were included in cluster signals)

No. of wards = 1 (0 [0%] were included in cluster signals)

No. of AMR profiles = 1 (0 [0%] were included in cluster signals)



There is no cluster signal with p-value < 0.05.

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with a clinical specimen culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with a clinical specimen positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals


Blood specimen: CRKP

Hospital-origin

No. of patients = 2 (0 [0%] were included in cluster signals)

No. of wards = 1 (0 [0%] were included in cluster signals)

No. of AMR profiles = 1 (0 [0%] were included in cluster signals)



There is no cluster signal with p-value < 0.05.

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with blood culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with blood culture positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals


All specimens: CRKP

Hospital-origin

No. of patients = 15 (0 [0%] were included in cluster signals)

No. of wards = 1 (0 [0%] were included in cluster signals)

No. of AMR profiles = 2 (0 [0%] were included in cluster signals)



There is no cluster signal with p-value < 0.05.

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with a clinical specimen culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with a clinical specimen positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals


Blood specimen: CRPA

Hospital-origin

No. of patients = 4 (0 [0%] were included in cluster signals)

No. of wards = 1 (0 [0%] were included in cluster signals)

No. of AMR profiles = 2 (0 [0%] were included in cluster signals)



There is no cluster signal with p-value < 0.05.

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with blood culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with blood culture positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals


All specimens: CRPA

Hospital-origin

No. of patients = 28 (0 [0%] were included in cluster signals)

No. of wards = 1 (0 [0%] were included in cluster signals)

No. of AMR profiles = 2 (0 [0%] were included in cluster signals)



There is no cluster signal with p-value < 0.05.

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with a clinical specimen culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with a clinical specimen positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals


Blood specimen: CRAB

Hospital-origin

No. of patients = 44 (9 [21%] were included in cluster signals)

No. of wards = 2 (1 [50%] were included in cluster signals)

No. of AMR profiles = 4 (2 [50%] were included in cluster signals)



Wards with cluster signals with p value < 0.05.

Ward ID	Total no. of observed cases in any cluster by ward	AMR Profile	Start date	End date	No. of observed cases in each cluster	P-value
ward_004	9	profile_CRAB_1	2016-02-04	2016-02-14	4	0.0004
ward_004		profile_CRAB_3	2016-07-28	2016-08-12	5	<0.0001

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with blood culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with blood culture positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Annex C: Cluster signals


All specimens: CRAB

Hospital-origin

No. of patients = 154 (8 [6%] were included in cluster signals)

No. of wards = 2 (1 [50%] were included in cluster signals)

No. of AMR profiles = 4 (2 [50%] were included in cluster signals)



Wards with cluster signals with p value < 0.05.

Ward ID	Total no. of observed cases in any cluster by ward	AMR Profile	Start date	End date	No. of observed cases in each cluster	P-value
ward_004	8	profile_CRAB_1	2016-02-04	2016-02-14	4	0.0003
ward_004		profile_CRAB_3	2016-07-30	2016-08-12	4	0.009

* The AMASS-SatScan (Annex C) de-duplicated by including only the first resistant isolate per patient per specimen type per evaluation period. Bar graphs show patients with a clinical specimen culture positive with organism profiles which were identified in at least one cluster signal. Gray bars (Other profiles) represents patients with a clinical specimen positive for organisms with profiles that were not included in any cluster signals. Details of AMR profiles are available in "Supplementary_data_Annex_C.pdf" and files in the folder "Report_with_patient_identifiers"

Methods used by the AMASS application

Data source:

For each run (double-click on AMASS.bat file), the AMASS application used the microbiology data file (microbiology_data) and the hospital admission data file (hospital_admission_data) that were stored in the same folder as the application file. Hence, if the user would like to update, correct, revise or change the data, the data files in the folder should be updated before the AMASS.bat file is double-clicked again. A new report based on the updated data would then be generated.

Requirements:

– Computer with Microsoft Windows 7 or 10

AMASS may work in other versions of Microsoft Windows and other operating systems. However, thorough testing and adjustment have not been performed.

– AMASSv3.0.zip package file

The AMASS application is to be downloaded from <https://www.amass.website>, and unzipped to generate an AMASS folder that could be stored under any folder in the computer. The AMASS folder contains 3 files (AMASS.bat, dictionary_for_microbiology_data.xlsx, and dictionary_for_hospital_admission_data.xlsx), and 5 folders (Configuration, Example_Dataset_1_WHONET, Example_Dataset_2, Example_Dataset_3_longformat, Programs).

– Microbiology data file (microbiology_data in .csv or .xlsx file format)

The user needs to obtain microbiology data, and then copy & paste this data file into the same folder as the AMASS.bat file.

– [Optional] Hospital admission data file (hospital_admission_data)

If available, the user could obtain hospital admission data, and then copy & paste this data file into the same folder as the AMASS.bat file.

Not required:

– Internet to run AMASS application

The AMASS application will run offline. No data are transferred while the application is running and reports are being generated; the reports are in PDF format (do not contain any patient identifier) and can be shared under the user's jurisdiction.

– R and Python

The download package (AMASSv3.0.zip) included R portable, Python portable and their libraries that the AMASS application requires. The user does not need to install any programme before using the AMASS. The user also does not have to uninstall R or Python if the computer already has the programme installed. The user does not need to know how to use R and Python.

Note:

[1] Please ensure that the file names of microbiology data file (microbiology_data) and the hospital admission data file (hospital_admission_data) are identical to what is written here. Please make sure that all are lower-cases with an underscore '_' at each space.

[2] Please ensure that both microbiology and hospital admission data files have no empty rows before the row of the variable names (i.e. the variable names are the first row in both files).

[3] For the first run, an user may need to fill the data dictionary files to make sure that the AMASS application understands your variable names and values.

AMASS uses a tier-based approach. In cases when only the microbiology data file with the results of culture positive samples is available, only section one and two would be generated for users. Section three would be generated only when data on admission date are available. This is because these data are required for the stratification by origin of infection. Section four would be generated only when data of specimens with culture negative (no microbial growth) are available in the microbiology data. This is because these data are required for the AMR frequency approach. Section five would be generated only when both data of specimens with culture negative and admission date are available. Section six would be generated only when mortality data are available.

Mortality was calculated from the number of in-hospital deaths (numerator) over the total number of patients with blood culture positive for the organism (denominator). Please note that this is the all-cause mortality calculated using the outcome data in the data file, and may not necessarily represent the mortality directly due to the infections.

How to use data dictionary files

In cases when variable names in the microbiology and hospital admission data files were not the same as the one that AMASS used, the data dictionary files could be edited. The raw microbiology and hospital admission data files were to be left unchanged. The data dictionary files provided could be edited and re-used automatically when the microbiology and hospital admission data files were updated and the AMASS.bat were to be double-clicked again (i.e. the data dictionary files would allow the user to re-analyze data files without the need to adjust variable names and data value again every time).

For example:

If variable name for 'hospital number' is written as 'hn' in the raw data file, the user would need to add 'hn' in the cell next to 'hospital_number'. If data value for blood specimens is defined by 'Blood–Hemoculture' in the raw data file, then the user would need to add 'Blood–Hemoculture' in the cell next to 'blood_specimen'.

Dictionary file (dictionary_for_microbiology_data.xlsx) may show up as in the table below:

Variable names used in AMASS	Variable names used in your microbiology data file	Requirements
Don't change values in this column, but you can add rows with similar values if you need	Change values in this column to represent how variable names are written in your raw microbiology data file	
hospital_number		Required
Values described in AMASS	Values used in your microbiology data file	Requirements
blood_specimen		Required

Please fill in your variable names as follows:

Variable names used in AMASS	Variable names used in your microbiology data file	Requirements
Don't change values in this column, but you can add rows with similar values if you need	Change values in this column to represent how variable names are written in your raw microbiology data file	
hospital_number	hn	Required
Values described in AMASS	Values used in your microbiology data file	Requirements
blood_specimen	Blood–Hemoculture	Required

Then, save the file. For every time the user double-clicked AMASS.bat, the application would know that the variable named 'hn' is similar to 'hospital_number' and represents the patient identifier in the analysis.

Organisms included for the AMR Surveillance Report:

- *Staphylococcus aureus*
- *Enterococcus faecalis*
- *Enterococcus faecium*
- *Streptococcus pneumoniae*
- *Salmonella* spp.
- *Escherichia coli*
- *Klebsiella pneumoniae*
- *Pseudomonas aeruginosa*
- *Acinetobacter baumannii*

Definitions:

The definitions of infection origin proposed by the WHO GLASS was used [1]. In brief, community–origin bloodstream infection (BSI) was defined for patients in the hospital within the first two calendar days of admission when the first blood culture positive specimens were taken. Hospital–origin BSI was defined for patients in the hospital longer than the first two calendar days of admission when the first blood culture positive specimens were taken. In cases when the user had additional data on infection origin defined by infection control team or based on referral data, the user could edit the data dictionary file (variable name 'infection_origin') and the AMASS application would use the data of that variable to stratify the data by origin of infection instead of the above definition. However, in cases when data on infection origin were not available (as in many hospitals in LMICs), the above definition would be calculated based on admission date and specimen collection date (with cutoff of 2 calendar days) and used to classify infections as community–origin or hospital–origin.

De–duplication:

When more than one blood culture was collected during patient management, duplicated findings of the same patient were excluded (de–duplicated). Only one result was reported for each patient per sample type (blood) and surveyed organisms (listed above). For example, if two blood cultures from the same patient had *E. coli*, only the first would be included in the report. If there was growth of *E. coli* in one blood culture and of *K. pneumoniae* in the other blood culture, then both results would be reported. One would be for the report on *E. coli* and the other one would be for the report on *K. pneumoniae*.

References:

- [1] World Health Organization (2018) Global Antimicrobial Resistance Surveillance System (GLASS) Report. Early implantation 2016–2017. <http://apps.who.int/iris/bitstream/handle/10665/259744/9789241513449-eng.pdf>. (accessed on 3 Dec 2018)
- [2] World Health Organization (2017) Global priority list of antibiotic–resistant bacteria to guide research, discovery, and development of new antibiotics. https://www.who.int/medicines/publications/WHO-PPL-Short_Summary_25Feb-ET_NM_WHO.pdf. (accessed on 3 Dec 2018)

Investigator team

The AMASS application is being developed by Cherry Lim, Clare Ling, Elizabeth Ashley, Paul Turner, Rahul Batra, Rogier van Doorn, Soawapak Hinjoy, Sopon Iamsirithaworn, Susanna Dunachie, Tri Wangrangsimakul, Viriya Hantrakun, William Schilling, John Stelling, Jonathan Edgeworth, Guy Thwaites, Nicholas PJ Day, Ben Cooper and Direk Limmathurotskul.

The AMASS application version 2.0 and 3.0 was developed by Chalida Rangsiwutisak, Preeyarach Klaytong, Prapass Wannapinij, Paul Tuner, John Stelling, Cherry Lim and Direk Limmathurotsakul.

Funding

The AMASS application version 1.0 was funded by the Wellcome Trust (grant no. 206736 and 101103). C.L. was funded by a Research Training Fellowship (grant no. 206736) and D.L. was funded by an Intermediate Training Fellowship (grant no. 101103) from the Wellcome Trust.

The AMASS application version 2.0 and 3.0 was funded by the Wellcome Trust (grant no. 224681/Z/21/Z and Institutional Translational Partnership Award-MORU)

If you have any queries about AMASS, please contact:

For technical information:

Chalida Rangsiwutisak (chalida@tropmedes.ac),
Cherry Lim (cherry@tropmedres.ac), and
Direk Limmathurotsakul (direk@tropmedres.ac)

For implementation of AMASS at your hospitals in Thailand:

Preeyarach Klaytong (preeyarach@tropmedres.ac)