



DaTSCAN™ SPECT Visual Interpretation (VI) Assessment

Justin Albani, John Seibyl, Ken Marek, Institute for Neurodegenerative Disorders (IND)

Summary

DaTSCAN™ SPECT imaging was acquired at PPMI imaging centers per the PPMI imaging protocol and sent to the imaging core lab at the Institute for Neurodegenerative Disorders (IND) for visual interpretation. All DaTSCAN™ images read for visual assessment of dopamine transporter deficit were reviewed per the procedure detailed in the methods section.

Method

Visual assessment of screening and visit 6 (SWEDDs only) DaTSCAN™ SPECT imaging for determination of dopamine transporter deficit was performed at IND by two expert readers in the field of nuclear medicine. Scans were read as either showing evidence of dopamine transporter deficit (i.e. abnormal) or not showing evidence of dopamine transporter deficit (i.e. normal.)

Normal images are characterized by two relatively symmetric comma or crescent-shaped focal regions of radiotracer uptake mirrored about the midline in transaxial images. Striatal uptake, comprising both the caudate and putamen, is distinctly evident compared to surrounding brain tissue.

Abnormal images typically fall into at least one of the following three general categories:

Activity is asymmetric, e.g. uptake in the region of the putamen of one hemisphere is absent or greatly reduced with respect to the other. Uptake is still visible in the caudate nuclei of both hemispheres resulting in a comma or crescent shape in one and a circular or oval focus in the other. There may be reduced uptake between at least one striatum and surrounding tissues.

Ioflupane uptake is absent in the putamen of both hemispheres and confined to the caudate nuclei. The signal is relatively symmetric and forms two roughly circular or oval foci. Uptake of one or both is generally reduced.

Uptake is absent in the putamen of both hemispheres and greatly reduced in one or both caudate nuclei. Uptake of the striata with respect to the background is reduced (1).





PARKINSON'S
PROGRESSION
MARKERS
INITIATIVE

Play a Part in Parkinson's Research

References

1. DaTSCAN™ [package insert]. Arlington Heights, IL: GE Healthcare; 2011.

About the Authors

This document was prepared by Justin Albani, Imaging Technical Quality Control & Processing Specialist and John Seibyl, Executive Director and Senior Scientist, Institute for Neurodegenerative Disorders, New Haven, CT.

For more information please contact Justin Albani by email at jalbani@mnimaging.com

Notice: This document is presented by the author(s) as a service to PPMI data users. However, users should be aware that no formal review process has vetted this document and that PPMI cannot guarantee the accuracy or utility of this document.

Appendix 1: IND SPECT Visual Interpretation Report: Annotated

(**EXAMPLE Not for use or distribution**)





SPECT Visual Interpretation Report Reader 1

Study Name:

Protocol Number:

Study Subject ID

scan_date_E3_C10

Site Number:

Subject Number:

Scan Date:

Imaging Center Name:

Clinical Site Name:

Scan Number:

☐

N/A

Radiopharmaceutical:

☐

N/A

The DAT scan for the above subject was reviewed. Visual interpretation results are as follows:

overall_vi_score_E2_C9

☐ DAT SPECT scan **shows** evidence of Dopamine transporter deficit

☐ DAT SPECT scan **does not** show evidence of Dopamine transporter deficit



* M N I I D 1 0 4 2 *

CONFIDENTIAL

Molecular NeuroImaging, L.L.C.

© Copyright 2015. All rights reserved.

MNIID1042

V5_IND SPECT VI Report Reader 1_06/Nov/2015