



EVALUATION OF PD-LINKED TRANSCRIPTS IN PPMI

Principal Investigator: Clemens R. Scherzer

Institution: Harvard Medical School, Brigham & Women's Hospital

Summary

Neuronal and peripheral transcriptional changes are linked to Parkinson's disease (PD) --- including the earliest stages of Lewy body neuropathology¹⁻⁵. These transcriptional changes may provide leads for biomarkers, therapeutic targets, and elucidate the disease pathobiology.

The objective of this study was to evaluate associations between the expression of three susceptibility genes and their splice variants in peripheral blood and early-stage, *de novo* Parkinson's disease. A cross-sectional, case-control study of *de novo* cases with Parkinson's disease and age- and sex-matched controls enrolled in the Parkinson Progression Marker Initiative (PPMI) was performed. Transcript counts of the three PD-linked loci *PARK1* (*SNCA*), *PARK15* (*FBXO7*), and *PARK7* (*DJ-1*) were analyzed. Five *SNCA* probes targeting distinct transcript isoforms, four probes targeting distinct *FBXO7* transcript isoforms, and one probe targeting *DJ-1* transcripts were employed. In addition, probes targeting the genes *DHPR* and *ZNF746* were included. Transcript counts were assayed in human blood in a high-precision NanoString gene expression assay that directly measures RNA counts in total RNA without bias from reverse transcription into cDNA. This technology uses molecular "barcodes" and single molecule imaging to detect and digitally count transcripts.

The distinct *SNCA* probes used target the boundaries of exon 3 and exon 4 (termed *E3E4-SNCA*), transcripts specifically with a long 3'UTR region (termed *3UTR-1-SNCA* and *3UTR2-SNCA*), transcripts that skip exon 5 (termed *E4E6-SNCA*), or the rare short *SNCA-007* transcript isoform (Ensembl ID ENST00000506691) that comprises exons 1-4. The distinct *FBXO7* probes used were designed to target the Ensembl transcripts *FBXO7-001*, *-005*, *-007*, *-008*, *-010* (www.ensembl.org).

Method

Sample collection. *PPMI* was designed to minimize bias from sample processing by collecting, handling and analyzing specimens of cases and controls in a standardized and blinded manner according to rules of evidence (Ref.^{6,7} and "Eleven good practices for RNA biomarker studies", in Scherzer, *Biomarkers in Medicine*, 2008⁸). Phlebotomy and biospecimens processing was performed in a standardized manner. Cases and controls were processed in parallel to avoid bias due to "run order" of samples. Venous whole blood was collected in PAXgene tubes to preserve RNA quality.

RNA isolation and quality control was performed at Coriell according to the PPMI Biospecimens Manual. Briefly, from each subject venous blood was collected in





PAXgene (Qiagen, Valencia, CA) tubes, incubated at room temperature for 24 hours, and frozen and shipped on dry ice to Coriell. RNA was extracted following the PAXgene procedure. RNA quality was determined by using the RNA Integrity Number package⁹. Only RNAs meeting three stringent Q/C parameters will be included in the analysis. These were (1) RIN ≥ 7.3 by Agilent Bioanalyzer (high mRNA integrity), (2) ratio of absorbances at 260 nm/280 nm ≥ 1.8 (no protein contamination). Controlling assay performance. Probes were designed according to the manufacturer's design principles¹⁰, including screening for inter- and intra-reporter and capture probe interactions, and selection for probes with optimal melting temperatures¹⁰. Direct counts of the target RNAs were measured by digital expression analysis based on NanoString technology (without reverse transcription into cDNA). Probes for the target and control RNAs were multiplexed and assayed according to the manufacturer's protocol on the nCounter Digital Analyzer. Our previous low-density array qPCR analysis of 246 human blood samples highlighted five RNAs with low variation in abundance (*GUSB*, *MON1B*, *SRCAP*, *GLT25D1*, *RPL13*) and these were included together with UBC to control for input RNA (as reference genes). The technician was blinded to the diagnosis. No-template (negative) controls containing water substituted for template were included (NTC). To control for plate-to-plate variation and drift equal amounts of RNA derived from Human Universal Reference RNA (UR) were spotted at the beginning, end, and throughout the entire experiment.

About the Authors

This document was prepared by Clemens R. Scherzer, The Neurogenomics Laboratory, Harvard Medical School and Brigham & Women's Hospital. For more information please contact Dr. Scherzer by email at cscherzer at rics.bwh.harvard.edu.

Publications: Publications using these data should include a reference for the primary paper reporting this work (Locascio *et al.*, 2014, in preparation).

Notice: This document is presented by the author(s) as a service to PPMI data users. However, users should be aware that no formal review process has vetted this document and that the author and PPMI cannot guarantee the accuracy or utility of this document. Moreover, as this study is still active it is subject to change.





References

1. Scherzer CR. Chipping away at diagnostics for neurodegenerative diseases. *Neurobiol Dis*. Mar 10 2009.
2. Scherzer CR, Eklund AC, Morse LJ, et al. Molecular markers of early Parkinson's disease based on gene expression in blood. *Proc Natl Acad Sci U S A*. Jan 16 2007;104(3):955-960.
3. Scherzer CR, Grass JA, Liao Z, et al. GATA transcription factors directly regulate the Parkinson's disease-linked gene alpha-synuclein. *Proc Natl Acad Sci U S A*. Aug 5 2008;105(31):10907-10912.
4. Scherzer CR, Jensen RV, Gullans SR, Feany MB. Gene expression changes presage neurodegeneration in a *Drosophila* model of Parkinson's disease. *Hum Mol Genet*. Oct 1 2003;12(19):2457-2466.
5. Zheng B, Liao Z, Locascio JJ, et al. PGC-1alpha, a potential therapeutic target for early intervention in Parkinson's disease. *Sci Transl Med*. Oct 6 2010;2(52):52ra73.
6. Ransohoff DF. Rules of evidence for cancer molecular-marker discovery and validation. *Nat Rev Cancer*. Apr 2004;4(4):309-314.
7. Ransohoff DF. Bias as a threat to the validity of cancer molecular-marker research. *Nat Rev Cancer*. Feb 2005;5(2):142-149.
8. Hennecke G, Scherzer CR. RNA biomarkers of Parkinson's disease: developing tools for novel therapies. *Biomarkers in Medicine*. 2008;2:41-53.
9. Imbeaud S, Graudens E, Boulanger V, et al. Towards standardization of RNA quality assessment using user-independent classifiers of microcapillary electrophoresis traces. *Nucleic Acids Res*. 2005;33(6):e56.
10. Geiss GK, Bumgarner RE, Birditt B, et al. Direct multiplexed measurement of gene expression with color-coded probe pairs. *Nat Biotechnol*. Mar 2008;26(3):317-325.

