



Truncated octahedron

In geometry, the **truncated octahedron** is the Archimedean solid that arises from a regular octahedron by removing six pyramids, one at each of the octahedron's vertices. The truncated octahedron has 14 faces (8 regular hexagons and 6 squares), 36 edges, and 24 vertices. Since each of its faces has point symmetry the truncated octahedron is a **6-zonohedron**. It is also the Goldberg polyhedron $G_{IV}(1,1)$, containing square and hexagonal faces. Like the cube, it can tessellate (or "pack") 3-dimensional space, as a permutohedron.

The truncated octahedron was called the "mecon" by Buckminster Fuller.^[1]


Its dual polyhedron is the tetrakis hexahedron. If the original truncated octahedron has unit edge length, its dual tetrakis hexahedron has edge lengths $\frac{9}{8}\sqrt{2}$ and $\frac{3}{2}\sqrt{2}$.

Classifications

As an Archimedean solid


A truncated octahedron is constructed from a regular octahedron by cutting off all vertices. This resulting polyhedron has six squares and eight hexagons, leaving out six square pyramids. Setting the edge length of the regular octahedron equal to **3a**, it follows that the length of each edge of a square pyramid (to be removed) is **a** (the square pyramid has four equilateral triangles as faces, the first Johnson solid). From the equilateral square pyramid's property, its volume is $\frac{\sqrt{2}}{6}a^3$. Because six equilateral square pyramids are removed by truncation, the volume of a truncated octahedron **V**

Truncated octahedron




Type	<u>Archimedean solid</u> , <u>Parallelohedron</u> , <u>Permutohedron</u> , <u>Plesiohedron</u> , <u>Zonohedron</u>
Faces	14
Edges	36
Vertices	24
Symmetry group	<u>octahedral symmetry</u> O_h
Dual polyhedron	<u>tetrakis hexahedron</u>

Vertex figure



Net


is obtained by subtracting the volume of those six from that of a regular octahedron:^[2]



$$V = \frac{\sqrt{2}}{3}(3a)^3 - 6 \cdot \frac{\sqrt{2}}{6}a^3 = 8\sqrt{2}a^3 \approx 11.3137a^3.$$

The surface area of a truncated octahedron A can be obtained by summing all polygons' area, six squares and eight hexagons. Considering the edge length a , this is:^[2]

$$A = (6 + 12\sqrt{3})a^2 \approx 26.7846a^2.$$



3D model of a truncated octahedron

The truncated octahedron is one of the thirteen Archimedean solids. In other words, it has a highly symmetric and semi-regular polyhedron with two or more different regular polygonal faces that meet in a vertex.^[3] The dual polyhedron of a truncated octahedron is the tetrakis hexahedron. They both have the same three-dimensional symmetry group as the regular octahedron does, the octahedral symmetry O_h .^[4] A square and two hexagons surround each of its vertex, denoting its vertex figure as $4 \cdot 6^2$.^[5]

The dihedral angle of a truncated octahedron between square-to-hexagon is $\arccos(-1/\sqrt{3}) \approx 125.26^\circ$, and that between adjacent hexagonal faces is $\arccos(-1/3) \approx 109.47^\circ$.^[6]


The Cartesian coordinates of the vertices of a truncated octahedron with edge length 1 are all permutations of^[7]

$$(\pm \sqrt{2}, \pm \frac{\sqrt{2}}{2}, 0).$$

As a space-filling polyhedron


The truncated octahedron can be described as a permutohedron of order 4 or **4-permutohedron**, meaning it can be represented with even more symmetric coordinates in four dimensions: all permutations of $(1, 2, 3, 4)$ form the vertices of a truncated octahedron in the three-dimensional subspace $x + y + z + w = 10$.^[8] Therefore, each vertex corresponds to a permutation of $(1, 2, 3, 4)$ and each edge represents a single pairwise swap of two elements.^[9] With this labeling, the swaps are of elements

whose values differ by one. If, instead, the truncated octahedron is labeled by the inverse permutations, the edges correspond to swaps of elements whose positions differ by one. With this alternative labeling, the edges and vertices of the truncated octahedron form the Cayley graph of the symmetric group S_4 , the group of four-element permutations, as generated by swaps of consecutive positions.^[10]



The truncated octahedron can tile space. It is classified as plesiohedron, meaning it can be defined as the Voronoi cell of a symmetric Delone set.^[11] Plesiohedra, translated without rotating, can be repeated to fill space. There are five three-dimensional primary parallelhedrons, one of which is the truncated octahedron. This polyhedron is generated from six line segments with four triples of coplanar segments, with the most symmetric form being generated from six line segments parallel to the face diagonals of a cube;^[12] an example of the honeycomb is the bitruncated cubic honeycomb.^[13] More generally, every permutohedron and parallelhedron is a zonohedron, a polyhedron that is centrally symmetric and can be defined by a Minkowski sum.^[14]

Applications



In chemistry, the truncated octahedron is the sodalite cage structure in the framework of a faujasite-type of zeolite crystals.^[15]

In solid-state physics, the first Brillouin zone of the face-centered cubic lattice is a truncated octahedron.^[16]

The truncated octahedron (in fact, the generalized truncated octahedron) appears in the error analysis of quantization index modulation (QIM) in conjunction with repetition coding.^[17]


Dissection

The truncated octahedron can be dissected into a central octahedron, surrounded by 8 triangular cupolae on each face, and 6 square pyramids above the vertices.^[18]

Removing the central octahedron and 2 or 4 triangular cupolae creates two Stewart toroids, with dihedral and tetrahedral symmetry:

It is possible to slice a tesseract by a hyperplane so that its sliced cross-section is a truncated octahedron.^[19]

The cell-transitive bitruncated cubic honeycomb can also be seen as the Voronoi tessellation of the body-centered cubic lattice. The truncated octahedron is one of five three-dimensional primary parallelohedra.




Objects



ancient Chinese die



sculpture in Bonn




Rubik's Cube variant



model made with
Polydron construction
set



Pyrite crystal




Boleite crystal

Truncated octahedral graph

In the mathematical field of graph theory, a **truncated octahedral graph** is the graph of vertices and edges of the truncated octahedron. It has 24 vertices and 36 edges, and is a cubic Archimedean graph.^[20] It has book thickness 3 and queue number 2.^[21]

As a Hamiltonian cubic graph, it can be represented by LCF notation in multiple ways: $[3, -7, 7, -3]^6$, $[5, -11, 11, 7, 5, -5, -7, -11, 11, -5, -7, 7]^2$, and $[-11, 5, -3, -7, -9, 3, -5, 5, -3, 9, 7, 3, -5, 11, -3, 7, 5, -7, -9, 9, 7, -5, -7, 3]$.^[22]


Truncated octahedral graph



3-fold symmetric Schlegel diagram

<u>Vertices</u>	24
<u>Edges</u>	36
<u>Automorphisms</u>	48
<u>Chromatic number</u>	2
<u>Book thickness</u>	3
<u>Queue number</u>	2
<u>Properties</u>	<u>Cubic</u> , <u>Hamiltonian</u> , <u>regular</u> , <u>zero-symmetric</u>

Table of graphs and parameters



Three different Hamiltonian cycles described by the three different LCF notations for the truncated octahedral graph

References

1. Weisstein, Eric W. "Truncated Octahedron" (<https://mathworld.wolfram.com/TruncatedOctahedron.html>). *MathWorld*.
2. Berman, Martin (1971). "Regular-faced convex polyhedra". *Journal of the Franklin Institute*. **291** (5): 329–352. doi:10.1016/0016-0032(71)90071-8 (<https://doi.org/10.1016%2F0016-0032%2871%2990071-8>). MR 0290245 (<https://mathscinet.ams.org/mathscinet-getitem?mr=0290245>).
3. Diudea, M. V. (2018). *Multi-shell Polyhedral Clusters* (https://books.google.com/books?id=p_06DwAAQBAJ&pg=PA39). Carbon Materials: Chemistry and Physics. Vol. 10. Springer. p. 39. doi:10.1007/978-3-319-64123-2 (<https://doi.org/10.1007%2F978-3-319-64123-2>). ISBN 978-3-319-64123-2.
4. Koca, M.; Koca, N. O. (2013). "Coxeter groups, quaternions, symmetries of polyhedra and 4D polytopes" (<https://books.google.com/books?id=ILnBkuSxXGEC&pg=PA48>). *Mathematical Physics: Proceedings of the 13th Regional Conference, Antalya, Turkey, 27–31 October 2010*. World Scientific. p. 48.
5. Williams, Robert (1979). *The Geometrical Foundation of Natural Structure: A Source Book of Design* (<https://archive.org/details/geometricalfound00will/page/78>). Dover Publications, Inc. p. 78. ISBN 978-0-486-23729-9.
6. Johnson, Norman W. (1966). "Convex polyhedra with regular faces" (<https://doi.org/10.4153%2Fcjm-1966-021-8>). *Canadian Journal of Mathematics*. **18**: 169–200. doi:10.4153/cjm-1966-021-8 (<https://doi.org/10.4153%2Fcjm-1966-021-8>). MR 0185507 (<https://mathscinet.ams.org/mathscinet-getitem?mr=0185507>). S2CID 122006114 (<https://api.semanticscholar.org/CorpusID:122006114>). Zbl 0132.14603 (<https://zbmath.org/?format=complete&q=an:0132.14603>).
7. Chieh, C. (November 1979). "The Archimedean truncated octahedron, and packing of geometric units in cubic crystal structures" (<https://scholar.archive.org/work/lfzy6gofjvc55nygiq36od3cwi>). *Acta Crystallographica Section A*. **35** (6). International Union of Crystallography (IUCr): 946–952. Bibcode:1979AcCrA..35..946C (<https://ui.adsabs.harvard.edu/abs/1979AcCrA..35..946C>). doi:10.1107/s0567739479002114 (<https://doi.org/10.1107%2Fs0567739479002114>).
8. Johnson, Tom; Jedrzejewski, Franck (2014). *Looking at Numbers* (<https://books.google.com/books?id=xtE-AgAAQBAJ&pg=PA15>). Springer. p. 15. doi:10.1007/978-3-0348-0554-4 (<https://doi.org/10.1007%2F978-3-0348-0554-4>). ISBN 978-3-0348-0554-4.

9. Crisman, Karl-Dieter (2011). "The Symmetry Group of the Permutahedron". *The College Mathematics Journal*. **42** (2): 135–139. doi:10.4169/college.math.j.42.2.135 (<https://doi.org/10.4169%2Fcollege.math.j.42.2.135>). JSTOR college.math.j.42.2.135 (<https://www.jstor.org/stable/college.math.j.42.2.135>).
10. Budden, Frank (December 1985). "Cayley graphs for some well-known groups". *The Mathematical Gazette*. **69** (450). JSTOR: 271–278. doi:10.2307/3617571 (<https://doi.org/10.2307%2F3617571>). JSTOR 3617571 (<https://www.jstor.org/stable/3617571>).
11. Erdahl, R. M. (1999). "Zonotopes, dicings, and Voronoi's conjecture on parallelhedra" (<https://doi.org/10.1006%2Feujc.1999.0294>). *European Journal of Combinatorics*. **20** (6): 527–549. doi:10.1006/eujc.1999.0294 (<https://doi.org/10.1006%2Feujc.1999.0294>). MR 1703597 (<https://mathscinet.ams.org/mathscinet-getitem?mr=1703597>). Voronoi conjectured that all tilings of higher dimensional spaces by translates of a single convex polytope are combinatorially equivalent to Voronoi tilings, and Erdahl proves this in the special case of zonotopes. But as he writes (p. 429), Voronoi's conjecture for dimensions at most four was already proven by Delaunay. For the classification of three-dimensional parallelhedra into these five types, see Grünbaum, Branko; Shephard, G. C. (1980). "Tilings with congruent tiles" (<https://doi.org/10.1090%2FS0273-0979-1980-14827-2>). *Bulletin of the American Mathematical Society*. New Series. **3** (3): 951–973. doi:10.1090/S0273-0979-1980-14827-2 (<https://doi.org/10.1090%2FS0273-0979-1980-14827-2>). MR 0585178 (<https://mathscinet.ams.org/mathscinet-getitem?mr=0585178>).
12. Alexandrov, A. D. (2005). "8.1 Parallelhedra" (<https://books.google.com/books?id=R9vPatr5aqYC&pg=PA349>). *Convex Polyhedra*. Springer. pp. 349–359.
13. Thuswaldner, Jörg; Zhang, Shu-qin (2020). "On self-affine tiles whose boundary is a sphere". *Transactions of the American Mathematical Society*. **373** (1): 491–527. arXiv:1811.06718 (<https://arxiv.org/abs/1811.06718>). doi:10.1090/tran/7930 (<https://doi.org/10.1090%2Ftran%2F7930>). MR 4042883 (<https://mathscinet.ams.org/mathscinet-getitem?mr=4042883>).
14. Jensen, Patrick M.; Trinderup, Camilia H.; Dahl, Anders B.; Dahl, Vedrana A. (2019). "Zonohedral Approximation of Spherical Structuring Element for Volumetric Morphology" (<https://books.google.com/books?id=uLWZDwAAQBAJ&pg=PA131>). In Felsberg, Michael; Forssén, Per-Erik; Sintorn, Ida-Maria; Unger, Jonas (eds.). *Image Analysis: 21st Scandinavian Conference, SCIA 2019, Norrköping, Sweden, June 11–13, 2019, Proceedings*. Springer. p. 131–132. doi:10.1007/978-3-030-20205-7 (<https://doi.org/10.1007%2F978-3-030-20205-7>). ISBN 978-3-030-20205-7.
15. Yen, Teh F. (2007). *Chemical Processes for Environmental Engineering* (<https://books.google.com/books?id=KHGelcDf1qQC&pg=PA388>). Imperial College Press. p. 338. ISBN 978-1-86094-759-9.
16. Mizutani, Uichiro (2001). *Introduction to the Electron Theory of Metals* (https://books.google.com/books?id=zY5z_UGqAcwC&pg=PA112). Cambridge University Press. p. 112. ISBN 978-0-521-58709-9.
17. Perez-Gonzalez, F.; Balado, F.; Martin, J.R.H. (2003). "Performance analysis of existing and new methods for data hiding with known-host information in additive channels". *IEEE Transactions on Signal Processing*. **51** (4): 960–980. Bibcode:2003ITSP...51..960P (<https://ui.adsabs.harvard.edu/abs/2003ITSP...51..960P>). doi:10.1109/TSP.2003.809368 (<https://doi.org/10.1109%2FTSP.2003.809368>).
18. Doskey, Alex. "Adventures Among the Toroids – Chapter 5 – Simplest (R)(A)(Q)(T) Toroids of genus p=1" (<http://www.doskey.com/polyhedra/Stewart05.html>). *www.doskey.com*.
19. Borovik, Alexandre V.; Borovik, Anna (2010). "Exercise 14.4" (<https://books.google.com/books?id=bVdzDL8KW3wC&pg=PA109>). *Mirrors and Reflections*. Universitext. New York: Springer. p. 109. doi:10.1007/978-0-387-79066-4 (<https://doi.org/10.1007%2F978-0-387-79066-4>). ISBN 978-0-387-79065-7. MR 2561378 (<https://mathscinet.ams.org/mathscinet-getitem?mr=2561378>).
20. Read, R. C.; Wilson, R. J. (1998). *An Atlas of Graphs*. Oxford University Press. p. 269.

21. Wolz, Jessica; *Engineering Linear Layouts with SAT*. Master Thesis, University of Tübingen, 2018
22. Weisstein, Eric W. "Truncated octahedral graph" (<https://mathworld.wolfram.com/TruncatedOctahedralGraph.html>). *MathWorld*.

Further reading

- Freitas, Robert A. Jr (1999). "Figure 5.5: Uniform space-filling using only truncated octahedra" (<http://www.nanomedicine.com/NMI/Figures/5.5.jpg>). *Nanomedicine, Volume I: Basic Capabilities* (<http://www.nanomedicine.com/NMI.htm>). Georgetown, Texas: Landes Bioscience. Retrieved 2006-09-08.
- Gaiha, P. & Guha, S.K. (1977). "Adjacent vertices on a permutohedron". *SIAM Journal on Applied Mathematics*. **32** (2): 323–327. doi:10.1137/0132025 (<https://doi.org/10.1137%2F0132025>).
- Hart, George W. "VRML model of truncated octahedron" (http://www.georgehart.com/virtual-polyhedra/vrml/truncated_octahedron.wrl). *Virtual Polyhedra: The Encyclopedia of Polyhedra* (<http://www.georgehart.com/virtual-polyhedra/vp.html>). Retrieved 2006-09-08.
- Cromwell, P. (1997). *Polyhedra*. United Kingdom: Cambridge. pp. 79–86 *Archimedean solids*. ISBN 0-521-55432-2.

External links

- Klitzing, Richard. "3D convex uniform polyhedra x3x4o - toe" (<https://bendwavy.org/klitzing/dimensions/polyhedra.htm>).
- Mäder, Roman. "The Uniform Polyhedra: Truncated Octahedron" (<http://www.mathconsult.ch/showroom/unipoly/08.html>). Retrieved 2006-09-08.
- Editable printable net of a truncated octahedron with interactive 3D view (<http://www.dr-mike-s-math-games-for-kids.com/polyhedral-nets.html?net=3UtM7vifCnAn5PXmSbX99Eu3LJZAs0nAWn3JyT7et98rnPxTGYml4FXjuQ2tE4viYN0KMgAstBRd0otTWLThQWI9BNNC4uigRoZQUQOQibYqtCuLQw9Ui3OofXtQPEsqQ7#applet>)

Retrieved from "https://en.wikipedia.org/w/index.php?title=Truncated_octahedron&oldid=1300968822"