



# Electromagnetic spectrum

The **electromagnetic spectrum** is the full range of electromagnetic radiation, organized by frequency or wavelength. The spectrum is divided into separate bands, with different names for the electromagnetic waves within each band. From low to high frequency these are: radio waves, microwaves, infrared, visible light, ultraviolet, X-rays, and gamma rays. The electromagnetic waves in each of these bands have different characteristics, such as how they are produced, how they interact with matter, and their practical applications.


Radio waves, at the low-frequency end of the spectrum, have the lowest photon energy and the longest wavelengths—thousands of kilometers, or more. They can be emitted and received by antennas, and pass through the atmosphere, foliage, and most building materials.

Gamma rays, at the high-frequency end of the spectrum, have the highest photon energies and the shortest wavelengths—much smaller than an atomic nucleus. Gamma rays, X-rays, and extreme ultraviolet rays are called ionizing radiation because their high photon energy is able to ionize atoms, causing chemical reactions. Longer-wavelength radiation such as visible light is nonionizing; the photons do not have sufficient energy to ionize atoms.

Throughout most of the electromagnetic spectrum, spectroscopy can be used to separate waves of different frequencies, so that the intensity of the radiation can be measured as a function of frequency or wavelength. Spectroscopy is used to study the interactions of electromagnetic waves with matter.<sup>[1]</sup>

## History and discovery

Humans have always been aware of visible light and radiant heat but for most of history it was not known that these phenomena were connected or were representatives of a more extensive principle. The ancient Greeks recognized that light traveled in straight lines and studied some of its properties, including reflection and refraction. Light was intensively studied from the beginning of the 17th century leading to the invention of important instruments like the telescope and microscope. Isaac Newton was the first to use the term spectrum for the range of colours that white light could be split into with a prism. Starting in 1666, Newton showed that these colours were intrinsic to light and could be recombined into white light. A debate arose over whether light had a wave nature or a particle nature with René Descartes, Robert Hooke and Christiaan Huygens favouring a wave description and Newton favouring a particle



A diagram of the electromagnetic spectrum, showing various properties across the range of frequencies and wavelengths

description. Huygens in particular had a well developed theory from which he was able to derive the laws of reflection and refraction. Around 1801, Thomas Young measured the wavelength of a light beam with his two-slit experiment thus conclusively demonstrating that light was a wave.

In 1800, William Herschel discovered infrared radiation.<sup>[2]</sup> He was studying the temperature of different colours by moving a thermometer through light split by a prism. He noticed that the highest temperature was beyond red. He theorized that this temperature change was due to "calorific rays", a type of light ray that could not be seen. The next year, Johann Ritter, working at the other end of the spectrum, noticed what he called "chemical rays" (invisible light rays that induced certain chemical reactions). These behaved similarly to visible violet light rays, but were beyond them in the spectrum.<sup>[3]</sup> They were later renamed ultraviolet radiation.

The study of electromagnetism began in 1820 when Hans Christian Ørsted discovered that electric currents produce magnetic fields (Oersted's law). Light was first linked to electromagnetism in 1845, when Michael Faraday noticed that the polarization of light traveling through a transparent material responded to a magnetic field (see Faraday effect). During the 1860s, James Clerk Maxwell developed four partial differential equations (Maxwell's equations) for the electromagnetic field. Two of these equations predicted the possibility and behavior of waves in the field. Analyzing the speed of these theoretical waves, Maxwell realized that they must travel at a speed that was about the known speed of light. This startling coincidence in value led Maxwell to make the inference that light itself is a type of electromagnetic wave. Maxwell's equations predicted an infinite range of frequencies of electromagnetic waves, all traveling at the speed of light. This was the first indication of the existence of the entire electromagnetic spectrum.

Maxwell's predicted waves included waves at very low frequencies compared to infrared, which in theory might be created by oscillating charges in an ordinary electrical circuit of a certain type. Attempting to prove Maxwell's equations and detect such low frequency electromagnetic radiation, in 1886, the physicist Heinrich Hertz built an apparatus to generate and detect what are now called radio waves. Hertz found the waves and was able to infer (by measuring their wavelength and multiplying it by their frequency) that they traveled at the speed of light. Hertz also demonstrated that the new radiation could be both reflected and refracted by various dielectric media, in the same manner as light. For example, Hertz was able to focus the waves using a lens made of tree resin. In a later experiment, Hertz similarly produced and measured the properties of microwaves. These new types of waves paved the way for inventions such as the wireless telegraph and the radio.

In 1895, Wilhelm Röntgen noticed a new type of radiation emitted during an experiment with an evacuated tube subjected to a high voltage. He called this radiation "x-rays" and found that they were able to travel through parts of the human body but were reflected or stopped by denser matter such as bones. Before long, many uses were found for this radiography.

The last portion of the electromagnetic spectrum was filled in with the discovery of gamma rays. In 1900, Paul Villard was studying the radioactive emissions of radium when he identified a new type of radiation that he at first thought consisted of particles similar to known alpha and beta particles, but with the power of being far more penetrating than either. However, in 1910, British physicist William Henry Bragg demonstrated that gamma rays are electromagnetic radiation, not particles, and in 1914, Ernest Rutherford (who had named them gamma rays in 1903 when he realized that they were fundamentally different from charged alpha and beta particles) and Edward Andrade measured their wavelengths, and found that gamma rays were similar to X-rays, but with shorter wavelengths.

The wave-particle debate was rekindled in 1901 when Max Planck discovered that light is absorbed only in discrete "quanta", now called photons, implying that light has a particle nature. This idea was made explicit by Albert Einstein in 1905, but never accepted by Planck and many other contemporaries. The modern position of science is that electromagnetic radiation has both a wave and a particle nature, the wave-particle duality. The contradictions arising from this position are still being debated by scientists and philosophers.

## Range

---

Electromagnetic waves are typically described by any of the following three physical properties: the frequency  $f$ , wavelength  $\lambda$ , or photon energy  $E$ . Frequencies observed in astronomy range from  $2.4 \times 10^{23}$  Hz (1 GeV gamma rays) down to the local plasma frequency of the ionized interstellar medium (~1 kHz). Wavelength is inversely proportional to the wave frequency,<sup>[1]</sup> so gamma rays have very short wavelengths that are fractions of the size of atoms, whereas wavelengths on the opposite end of the spectrum can be indefinitely long. Photon energy is directly proportional to the wave frequency, so gamma ray photons have the highest energy (around a billion electron volts), while radio wave photons have very low energy (around a femtoelectronvolt). These relations are illustrated by the following equations:

$$f = \frac{c}{\lambda}, \quad f = \frac{E}{h}, \quad E = \frac{hc}{\lambda},$$

where:

- $c$  is the speed of light in vacuum
- $h$  is the Planck constant.

Whenever electromagnetic waves travel in a medium with matter, their wavelength is decreased. Wavelengths of electromagnetic radiation, whatever medium they are traveling through, are usually quoted in terms of the *vacuum wavelength*, although this is not always explicitly stated.

Generally, electromagnetic radiation is classified by wavelength into radio wave, microwave, infrared, visible light, ultraviolet, X-rays and gamma rays. The behavior of EM radiation depends on its wavelength. When EM radiation interacts with single atoms and molecules, its behavior also depends on the amount of energy per quantum (photon) it carries.

Spectroscopy can detect a much wider region of the EM spectrum than the visible wavelength range of 400 nm to 700 nm in a vacuum. A common laboratory spectroscope can detect wavelengths from 2 nm to 2500 nm.<sup>[1]</sup> Detailed information about the physical properties of objects, gases, or even stars can be obtained from this type of device. Spectroscopes are widely used in astrophysics. For example, many hydrogen atoms emit a radio wave photon that has a wavelength of 21.12 cm. Also, frequencies of 30 Hz and below can be produced by and are important in the study of certain stellar nebulae<sup>[4]</sup> and frequencies as high as  $2.9 \times 10^{27}$  Hz have been detected from astrophysical sources.<sup>[5]</sup>

# Regions

The types of electromagnetic radiation are broadly classified into the following classes (regions, bands or types):<sup>[1]</sup>


1. Gamma radiation
2. X-ray radiation
3. Ultraviolet radiation
4. Visible light (light that humans can see)
5. Infrared radiation
6. Microwave radiation
7. Radio waves

This classification goes in the increasing order of wavelength, which is characteristic of the type of radiation.<sup>[1]</sup>


There are no precisely defined boundaries between the bands of the electromagnetic spectrum; rather they fade into each other like the bands in a rainbow. Radiation of each frequency and wavelength (or in each band) has a mix of properties of the two regions of the spectrum that bound it. For example, red light resembles infrared radiation, in that it can excite and add energy to some chemical bonds and indeed must do so to power the chemical mechanisms responsible for photosynthesis and the working of the visual system.

In atomic and nuclear physics, the distinction between X-rays and gamma rays is based on sources: the photons generated from nuclear decay or other nuclear and subnuclear/particle process are termed gamma rays, whereas X-rays are generated by electronic transitions involving energetically deep inner atomic electrons.<sup>[6][7]</sup> Electronic transitions in muonic atoms transitions are also said to produce X-rays.<sup>[8]</sup> In astrophysics, energies below 100keV are called X-rays and higher energies are gamma rays.<sup>[9]</sup>

The region of the spectrum where electromagnetic radiation is observed may differ from the region it was emitted in due to relative velocity of the source and observer, (the Doppler shift), relative gravitational potential (gravitational redshift), or expansion of the universe (cosmological redshift).<sup>[9]:543</sup> For example,



The electromagnetic spectrum



A visualization of the electromagnetic spectrum.

the cosmic microwave background, relic blackbody radiation from the era of recombination, started out at energies around 1eV, but as has undergone enough cosmological red shift to put it into the microwave region of the spectrum for observers on Earth.<sup>[10]</sup>

Class			Wave-length $\lambda$	Freq- uency $f$	Energy per photon $E$
Ionizing radiation	$\gamma$	Gamma rays	10 pm	30 EHz	124 keV
	HX	Hard X-rays	100 pm	3 EHz	12.4 keV
	SX	Soft X-rays	10 nm	30 PHz	124 eV
	EUV	Extreme ultraviolet	121 nm	3 PHz	10.2 eV
	NUV	Near ultraviolet	400 nm	750 THz	3.1 eV
		Visible spectrum	700 nm	480 THz	1.77 eV
Infrared	NIR	Near infrared	1 $\mu\text{m}$	300 THz	1.24 eV
	MIR	Mid infrared	10 $\mu\text{m}$	30 THz	124 meV
	FIR	Far infrared	100 $\mu\text{m}$	3 THz	12.4 meV
Micro- waves <sup>[11]</sup>	EHF	Extremely high frequency	1 mm	300 GHz	1.24 meV
			1 cm	30 GHz	124 $\mu\text{eV}$
	SHF	Super high frequency	1 dm	3 GHz	12.4 $\mu\text{eV}$
	UHF	Ultra high frequency	1 m	300 MHz	1.24 $\mu\text{eV}$
Radio waves <sup>[11]</sup>	VHF	Very high frequency	10 m	30 MHz	124 neV
	HF	High frequency	100 m	3 MHz	12.4 neV
	MF	Medium frequency	1 km	300 kHz	1.24 neV
	LF	Low frequency	10 km	30 kHz	124 peV
	VLF	Very low frequency	100 km	3 kHz	12.4 peV
	3	Band 3	1 Mm	300 Hz	1.24 peV
	2	Band 2	10 Mm	30 Hz	124 feV
	1	Band 1	100 Mm	3 Hz	12.4 feV

Sources<sup>[12][13][14]</sup> Table shows the lower frequency limits (and higher wavelength limits) for the specified class

Explanation of units and prefixes.

Unit	Abbreviation	Name	Scale
Wavelength	pm	picometer	$1 \times 10^{-12}$ meters
Wavelength	nm	nanometer	$1 \times 10^{-9}$ meters
Wavelength	$\mu\text{m}$	micrometer	$1 \times 10^{-6}$ meters
Wavelength	mm	millimeter	$1 \times 10^{-3}$ meters
Wavelength	cm	centimeter	$1 \times 10^{-2}$ meters
Wavelength	dm	decimeter	$1 \times 10^{-1}$ meters
Wavelength	m	meter	1 meter
Wavelength	km	kilometer	$1 \times 10^3$ meters
Wavelength	Mm	megameter	$1 \times 10^6$ meters
Frequency	EHZ	exaHertz	$1 \times 10^{18}$ hertz
Frequency	PHz	petaHertz	$1 \times 10^{15}$ hertz
Frequency	THz	teraHertz	$1 \times 10^{12}$ hertz
Frequency	GHz	gigaHertz	$1 \times 10^9$ hertz
Frequency	MHz	megaHertz	$1 \times 10^6$ hertz
Frequency	KHz	kiloHertz	$1 \times 10^3$ hertz
Frequency	Hz	Hertz	1 Hertz
Energy Per Photon	keV	kilo-electronvolt	$1 \times 10^3$ electronvolts
Energy Per Photon	eV	electronvolt	1 electronvolt
Energy Per Photon	meV	milli-electronvolt	$1 \times 10^{-3}$ electronvolts
Energy Per Photon	$\mu\text{eV}$	micro-electronvolt	$1 \times 10^{-6}$ electronvolts
Energy Per Photon	neV	nano-electronvolt	$1 \times 10^{-9}$ electronvolts
Energy Per Photon	peV	pico-electronvolt	$1 \times 10^{-12}$ electronvolts
Energy Per Photon	feV	femto-electronvolt	$1 \times 10^{-15}$ electronvolts

## Rationale for names

Electromagnetic radiation interacts with matter in different ways across the spectrum. These types of interaction are so different that historically different names have been applied to different parts of the spectrum, as though these were different types of radiation. Thus, although these "different kinds" of electromagnetic radiation form a quantitatively continuous spectrum of frequencies and wavelengths, the spectrum remains divided for practical reasons arising from these qualitative interaction differences.

## Electromagnetic radiation interaction with matter

Region of the spectrum	Main interactions with matter
<u>Radio</u>	Collective oscillation of charge carriers in bulk material ( <u>plasma oscillation</u> ). An example would be the oscillatory travels of the electrons in an <u>antenna</u> .
<u>Microwave</u> through far <u>infrared</u>	Plasma oscillation, molecular rotation
Near <u>infrared</u>	Molecular vibration, plasma oscillation (in metals only)
<u>Visible</u>	Molecular electron excitation (including pigment molecules found in the human retina), plasma oscillations (in metals only)
<u>Ultraviolet</u>	Excitation of molecular and atomic valence electrons, including ejection of the electrons ( <u>photoelectric effect</u> )
<u>X-rays</u>	Excitation and ejection of core atomic electrons, <u>Compton scattering</u> (for low atomic numbers)
<u>Gamma rays</u>	Energetic ejection of core electrons in heavy elements, Compton scattering (for all atomic numbers), excitation of atomic nuclei, including dissociation of nuclei
High-energy <u>gamma rays</u>	Creation of <u>particle-antiparticle pairs</u> . At very high energies a single photon can create a shower of high-energy particles and antiparticles upon interaction with matter.

## Types of radiation

---

### Radio waves

Radio waves are emitted and received by antennas, which consist of conductors such as metal rod resonators. In artificial generation of radio waves, an electronic device called a transmitter generates an alternating electric current which is applied to an antenna. The oscillating electrons in the antenna generate oscillating electric and magnetic fields that radiate away from the antenna as radio waves. In reception of radio waves, the oscillating electric and magnetic fields of a radio wave couple to the electrons in an antenna, pushing them back and forth, creating oscillating currents which are applied to a radio receiver. Earth's atmosphere is mainly transparent to radio waves, except for layers of charged particles in the ionosphere which can reflect certain frequencies.

Radio waves are extremely widely used to transmit information across distances in radio communication systems such as radio broadcasting, television, two way radios, mobile phones, communication satellites, and wireless networking. In a radio communication system, a radio frequency current is modulated with an information-bearing signal in a transmitter by varying either the amplitude, frequency or phase, and applied to an antenna. The radio waves carry the information across space to a receiver, where they are received by an antenna and the information extracted by demodulation in the receiver. Radio waves are also used for navigation in systems like Global Positioning System (GPS) and navigational beacons, and locating distant objects in radiolocation and radar. They are also used for remote control, and for industrial heating.


The use of the radio spectrum is strictly regulated by governments, coordinated by the International Telecommunication Union (ITU) which allocates frequencies to different users for different uses.




## Microwaves

Microwaves are radio waves of short wavelength, from about 10 centimeters to one millimeter, in the SHF and EHF frequency bands. Microwave energy is produced with klystron and magnetron tubes, and with solid state devices such as Gunn and IMPATT diodes. Although they are emitted and absorbed by short antennas, they are also absorbed by polar molecules, coupling to vibrational and rotational modes, resulting in bulk heating. Unlike higher frequency waves such as infrared and visible light which are absorbed mainly at surfaces, microwaves can penetrate into materials and deposit their energy below the surface. This effect is used to heat food in microwave ovens, and for industrial heating and medical diathermy. Microwaves are the main wavelengths used in radar, and are used for satellite communication, and wireless networking technologies such as Wi-Fi. The copper cables (transmission lines) which are used to carry lower-frequency radio waves to antennas have excessive power losses at microwave frequencies, and metal pipes called waveguides are used to carry them. Although at the low end of the band the atmosphere is mainly transparent, at the upper end of the band absorption of microwaves by atmospheric gases limits practical propagation distances to a few kilometers.

Terahertz radiation or sub-millimeter radiation is a region of the spectrum from about 100 GHz to 30 terahertz (THz) between microwaves and far infrared which can be regarded as belonging to either band. Until recently, the range was rarely studied and few sources existed for microwave energy in the so-called *terahertz gap*, but applications such as imaging and communications are now appearing. Scientists are also looking to apply terahertz technology in the armed forces, where high-frequency waves might be directed at enemy troops to incapacitate their electronic equipment.<sup>[15]</sup> Terahertz radiation is strongly absorbed by atmospheric gases, making this frequency range useless for long-distance communication.



Plot of Earth's atmospheric opacity to various wavelengths of electromagnetic radiation. This is the surface-to-space opacity, the atmosphere is transparent to longwave radio transmissions within the troposphere but opaque to space due to the ionosphere.



Plot of atmospheric opacity for terrestrial to terrestrial transmission showing the molecules responsible for some of the resonances

## Infrared radiation

The infrared part of the electromagnetic spectrum covers the range from roughly 300 GHz to 400 THz (1 mm – 750 nm). It can be divided into three parts:<sup>[1]</sup>









- **Far-infrared**, from 300 GHz to 30 THz (1 mm – 10  $\mu\text{m}$ ). The lower part of this range may also be called microwaves or terahertz waves. This radiation is typically absorbed by so-called rotational modes in gas-phase molecules, by molecular motions in liquids, and by phonons in solids. The water in Earth's atmosphere absorbs so strongly in this range that it renders the atmosphere in effect opaque. However, there are certain wavelength ranges ("windows") within the opaque range that allow partial transmission, and can be used for astronomy. The wavelength range from approximately 200  $\mu\text{m}$  up to a few mm is often referred to as Submillimetre astronomy, reserving far infrared for wavelengths below 200  $\mu\text{m}$ .
- **Mid-infrared**, from 30 THz to 120 THz (10–2.5  $\mu\text{m}$ ). Hot objects (black-body radiators) can radiate strongly in this range, and human skin at normal body temperature radiates strongly at the lower end of this region. This radiation is absorbed by molecular vibrations, where the different atoms in a molecule vibrate around their equilibrium positions. This range is sometimes called the *fingerprint region*, since the mid-infrared absorption spectrum of a compound is very specific for that compound.
- **Near-infrared**, from 120 THz to 400 THz (2,500–750 nm). Physical processes that are relevant for this range are similar to those for visible light. The highest frequencies in this region can be detected directly by some types of photographic film, and by many types of solid state image sensors for infrared photography and videography.

## Visible light

Above infrared in frequency comes visible light. The Sun emits its peak power in the visible region, although integrating the entire emission power spectrum through all wavelengths shows that the Sun emits slightly more infrared than visible light.<sup>[16]</sup> By definition, visible light is the part of the EM spectrum the human eye is the most sensitive to. Visible light (and near-infrared light) is typically absorbed and emitted by electrons in molecules and atoms that move from one energy level to another. This action allows the chemical mechanisms that underlie human vision and plant photosynthesis. The light that excites the human visual system is a very small portion of the electromagnetic spectrum.

A rainbow shows the optical (visible) part of the electromagnetic spectrum; infrared (if it could be seen) would be located just beyond the red side of the rainbow whilst ultraviolet would appear just beyond the opposite violet end.

Electromagnetic radiation with a wavelength between 380 nm and 760 nm (400–790 terahertz) is detected by the human eye and perceived as visible light. Other wavelengths, especially near infrared (longer than 760 nm) and ultraviolet (shorter than 380 nm) are also sometimes referred to as light, especially when the

			
<u>Colour</u>	<u>Wavelength (nm)</u>	<u>Frequency (THz)</u>	<u>Photon energy (eV)</u>
 <u>violet</u>	380–450	670–790	2.75–3.26
 <u>blue</u>	450–485	620–670	2.56–2.75
 <u>cyan</u>	485–500	600–620	2.48–2.56
 <u>green</u>	500–565	530–600	2.19–2.48
 <u>yellow</u>	565–590	510–530	2.10–2.19
 <u>orange</u>	590–625	480–510	1.98–2.10
 <u>red</u>	625–750	400–480	1.65–1.98

visibility to humans is not relevant. White light is a combination of lights of different wavelengths in the visible spectrum. Passing white light through a prism splits it up into the several colours of light observed in the visible spectrum between 400 nm and 780 nm.

If radiation having a frequency in the visible region of the EM spectrum reflects off an object, say, a bowl of fruit, and then strikes the eyes, this results in visual perception of the scene. The brain's visual system processes the multitude of reflected frequencies into different shades and hues, and through this insufficiently understood psychophysical phenomenon, most people perceive a bowl of fruit.

At most wavelengths, however, the information carried by electromagnetic radiation is not directly detected by human senses. Natural sources produce EM radiation across the spectrum, and technology can also manipulate a broad range of wavelengths. Optical fiber transmits light that, although not necessarily in the visible part of the spectrum (it is usually infrared), can carry information. The modulation is similar to that used with radio waves.


## Ultraviolet radiation

Next in frequency comes ultraviolet (UV). In frequency (and thus energy), UV rays sit between the violet end of the visible spectrum and the X-ray range. The UV wavelength spectrum ranges from 399 nm to 10 nm and is divided into 3 sections: UVA, UVB, and UVC.

UV is the lowest energy range energetic enough to ionize atoms, separating electrons from them, and thus causing chemical reactions. UV, X-rays, and gamma rays are thus collectively called ionizing radiation; exposure to them can damage living tissue. UV can also cause substances to glow with visible light; this is called fluorescence. UV fluorescence is used by forensics to detect any evidence like blood and urine, that is produced by a crime scene. Also UV fluorescence is used to detect counterfeit money and IDs, as they are laced with material that can glow under UV.

At the middle range of UV, UV rays cannot ionize but can break chemical bonds, making molecules unusually reactive. Sunburn, for example, is caused by the disruptive effects of middle range UV radiation on skin cells, which is the main cause of skin cancer. UV rays in the middle range can irreparably damage the complex DNA molecules in the cells producing thymine dimers making it a very potent mutagen. Due to skin cancer caused by UV, the sunscreen industry was invented to combat UV damage. Mid UV wavelengths are called UVB and UVB lights such as germicidal lamps are used to kill germs and also to sterilize water.

The Sun emits UV radiation (about 10% of its total power), including extremely short wavelength UV that could potentially destroy most life on land (ocean water would provide some protection for life there). However, most of the Sun's damaging UV wavelengths are absorbed by the atmosphere before they reach the surface. The higher energy (shortest wavelength) ranges of UV (called "vacuum UV") are absorbed by nitrogen and, at longer wavelengths, by simple diatomic oxygen in the air. Most of the UV in the mid-range of energy is blocked by the ozone layer, which absorbs strongly in the important 200–315 nm range, the lower energy part of which is too long for ordinary dioxygen in air to absorb. This



The amount of penetration of UV relative to altitude in Earth's ozone

leaves less than 3% of sunlight at sea level in UV, with all of this remainder at the lower energies. The remainder is UV-A, along with some UV-B. The very lowest energy range of UV between 315 nm and visible light (called UV-A) is not blocked well by the atmosphere, but does not cause sunburn and does less biological damage. However, it is not harmless and does create oxygen radicals, mutations and skin damage.

## X-rays

After UV come X-rays, which, like the upper ranges of UV are also ionizing. However, due to their higher energies, X-rays can also interact with matter by means of the Compton effect. Hard X-rays have shorter wavelengths than soft X-rays and as they can pass through many substances with little absorption, they can be used to 'see through' objects with 'thicknesses' less than that equivalent to a few meters of water. One notable use is diagnostic X-ray imaging in medicine (a process known as radiography). X-rays are useful as probes in high-energy physics. In astronomy, the accretion disks around neutron stars and black holes emit X-rays, enabling studies of these phenomena. X-rays are also emitted by stellar corona and are strongly emitted by some types of nebulae. However, X-ray telescopes must be placed outside the Earth's atmosphere to see astronomical X-rays, since the great depth of the atmosphere of Earth is opaque to X-rays (with areal density of 1000 g/cm<sup>2</sup>), equivalent to 10 meters thickness of water.<sup>[17]</sup> This is an amount sufficient to block almost all astronomical X-rays (and also astronomical gamma rays—see below).

## Gamma rays

After hard X-rays come gamma rays, which were discovered by Paul Ulrich Villard in 1900. These are the most energetic photons, having no defined lower limit to their wavelength. In astronomy they are valuable for studying high-energy objects or regions, however as with X-rays this can only be done with telescopes outside the Earth's atmosphere. Gamma rays are used experimentally by physicists for their penetrating ability and are produced by a number of radioisotopes. They are used for irradiation of foods and seeds for sterilization, and in medicine they are occasionally used in radiation cancer therapy.<sup>[18]</sup> More commonly, gamma rays are used for diagnostic imaging in nuclear medicine, an example being PET scans. The wavelength of gamma rays can be measured with high accuracy through the effects of Compton scattering.

## See also

---

- |                                 |                           |
|---------------------------------|---------------------------|
| ▪ <u>Bandplan</u>               | ▪ <u>Radiant energy</u>   |
| ▪ <u>Cosmic ray</u>             | ▪ <u>Radiation</u>        |
| ▪ <u>Electroencephalography</u> | ▪ <u>Radio window</u>     |
| ▪ <u>Infrared window</u>        | ▪ <u>Spectral imaging</u> |
| ▪ <u>Ionizing radiation</u>     | ▪ <u>Spectroscopy</u>     |
| ▪ <u>Optical window</u>         | ▪ <u>V band</u>           |
| ▪ <u>Ozone layer</u>            | ▪ <u>W band</u>           |

## Notes and references

---

1. Mehta, Akul (25 August 2011). "Introduction to the Electromagnetic Spectrum and Spectroscopy" (<http://pharmaxchange.info/press/2011/08/introduction-to-the-electromagnetic-spectrum-and-spectroscopy/>). Pharmaxchange.info. Retrieved 2011-11-08.
2. "Herschel Discovers Infrared Light" ([https://web.archive.org/web/20120225094516/http://coolcosmos.ipac.caltech.edu/cosmic\\_classroom/classroom\\_activities/herschel\\_bio.html](https://web.archive.org/web/20120225094516/http://coolcosmos.ipac.caltech.edu/cosmic_classroom/classroom_activities/herschel_bio.html)). *Cool Cosmos Classroom activities*. Archived from the original ([http://coolcosmos.ipac.caltech.edu/cosmic\\_classroom/classroom\\_activities/herschel\\_bio.html](http://coolcosmos.ipac.caltech.edu/cosmic_classroom/classroom_activities/herschel_bio.html)) on 2012-02-25. Retrieved 4 March 2013. "He directed sunlight through a glass prism to create a spectrum [...] and then measured the temperature of each colour. [...] He found that the temperatures of the colours increased from the violet to the red part of the spectrum. [...] Herschel decided to measure the temperature just *beyond* the red of the spectrum in a region where no sunlight was visible. To his surprise, he found that this region had the highest temperature of all."
3. Davidson, Michael W. "Johann Wilhelm Ritter (1776–1810)" (<http://micro.magnet.fsu.edu/optics/timeline/people/ritter.html>). The Florida State University. Retrieved 5 March 2013. "Ritter [...] hypothesized that there must also be invisible radiation beyond the violet end of the spectrum and commenced experiments to confirm his speculation. He began working with silver chloride, a substance decomposed by light, measuring the speed at which different colours of light broke it down. [...] Ritter [...] demonstrated that the fastest rate of decomposition occurred with radiation that could not be seen, but that existed in a region beyond the violet. Ritter initially referred to the new type of radiation as chemical rays, but the title of ultraviolet radiation eventually became the preferred term."
4. Condon, J. J.; Ransom, S. M. "Essential Radio Astronomy: Pulsar Properties" (<https://web.archive.org/web/20110504064425/http://www.cv.nrao.edu/course/ast534/Pulsars.html>). National Radio Astronomy Observatory. Archived from the original (<http://www.cv.nrao.edu/course/ast534/Pulsars.html>) on 2011-05-04. Retrieved 2008-01-05.
5. Abdo, A. A.; Allen, B.; Berley, D.; Blaufuss, E.; Casanova, S.; Chen, C.; Coyne, D. G.; Delay, R. S.; Dingus, B. L.; Ellsworth, R. W.; Fleysher, L.; Fleysher, R.; Gebauer, I.; Gonzalez, M. M.; Goodman, J. A.; Hays, E.; Hoffman, C. M.; Kolterman, B. E.; Kelley, L. A.; Lansdell, C. P.; Linnemann, J. T.; McEnery, J. E.; Mincer, A. I.; Moskalenko, I. V.; Nemethy, P.; Noyes, D.; Ryan, J. M.; Samuelson, F. W.; Saz Parkinson, P. M.; et al. (2007). "Discovery of TeV Gamma-Ray Emission from the Cygnus Region of the Galaxy". *The Astrophysical Journal*. **658** (1): L33 – L36. arXiv:[astro-ph/0611691](https://arxiv.org/abs/astro-ph/0611691) (<https://arxiv.org/abs/astro-ph/0611691>). Bibcode:2007ApJ...658L..33A (<https://ui.adsabs.harvard.edu/abs/2007ApJ...658L..33A>). doi:10.1086/513696 (<https://doi.org/10.1086/513696>). S2CID 17886934 (<https://api.semanticscholar.org/CorpusID:17886934>).
6. Feynman, Richard; Leighton, Robert; Sands, Matthew (1963). *The Feynman Lectures on Physics, Vol.1* ([https://www.feynmanlectures.caltech.edu/I\\_toc.html](https://www.feynmanlectures.caltech.edu/I_toc.html)). US: Addison-Wesley. pp. 2–5 ([https://www.feynmanlectures.caltech.edu/I\\_02.html#Ch2-S2-p10](https://www.feynmanlectures.caltech.edu/I_02.html#Ch2-S2-p10)). ISBN 978-0-201-02116-5.
7. L'Annunziata, Michael; Baradei, Mohammad (2003). *Handbook of Radioactivity Analysis* (<https://books.google.com/books?id=b519e10OPT0C&q=gamma+x-ray&pg=PA58>). Academic Press. p. 58. ISBN 978-0-12-436603-9.
8. Corrections to muonic X-rays and a possible proton halo (<http://www.slac.stanford.edu/cgi-wrap/getdoc/slac-pub-0335.pdf>) Archived (<https://web.archive.org/web/20170313134112/http://www.slac.stanford.edu/cgi-wrap/getdoc/slac-pub-0335.pdf>) 2017-03-13 at the Wayback Machine slac-pub-0335 (1967)
9. Grupen, Claus; Cowan, G.; Eidelman, S. D.; Stroth, T. (2005). *Astroparticle Physics* (<https://archive.org/details/astroparticlephy00grup>). Springer. p. 109 (<https://archive.org/details/astroparticlephy00grup/page/n123>). ISBN 978-3-540-25312-9.

10. Samtleben, Dorothea; Staggs, Suzanne; Winstein, Bruce (2007-11-01). "The Cosmic Microwave Background for Pedestrians: A Review for Particle and Nuclear Physicists" (<http://www.annualreviews.org/doi/10.1146/annurev.nucl.54.070103.181232>). *Annual Review of Nuclear and Particle Science*. **57** (1): 245–283. arXiv:0803.0834 (<https://arxiv.org/abs/0803.0834>). Bibcode:2007ARNPS..57..245S (<https://ui.adsabs.harvard.edu/abs/2007ARNPS..57..245S>). doi:10.1146/annurev.nucl.54.070103.181232 (<https://doi.org/10.1146%2Fannurev.nucl.54.070103.181232>). ISSN 0163-8998 (<https://search.worldcat.org/issn/0163-8998>).
11. Article 2 "Final Acts WRC-15: World Radiocommunication Conference" ([https://www.itu.int/dms\\_pub/itu-r/opb/act/R-ACT-WRC.12-2015-PDF-E.pdf#page=6.34](https://www.itu.int/dms_pub/itu-r/opb/act/R-ACT-WRC.12-2015-PDF-E.pdf#page=6.34)), International Telecommunication Union Geneva, 2015
12. What is Light? ([http://cbst.ucdavis.edu/education/courses/winter-2006-IST8A/ist8a\\_2006\\_01\\_09light.pdf](http://cbst.ucdavis.edu/education/courses/winter-2006-IST8A/ist8a_2006_01_09light.pdf)) Archived ([https://web.archive.org/web/20131205005843/http://cbst.ucdavis.edu/education/courses/winter-2006-IST8A/ist8a\\_2006\\_01\\_09light.pdf](https://web.archive.org/web/20131205005843/http://cbst.ucdavis.edu/education/courses/winter-2006-IST8A/ist8a_2006_01_09light.pdf)) December 5, 2013, at the [Wayback Machine](#) – UC Davis lecture slides
13. Elert, Glenn. "The Electromagnetic Spectrum" (<https://physics.info/em-spectrum/>). *The Physics Hypertextbook*. Archived (<https://web.archive.org/web/20220121220019/https://physics.info/em-spectrum/>) from the original on 2022-01-21. Retrieved 2022-01-21.
14. Stimac, Tomislav. "Definition of frequency bands (VLF, ELF... etc.)" (<http://www.vlf.it/frequency/bands.html>). vlf.it. Archived (<https://web.archive.org/web/20100430012219/http://www.vlf.it/frequency/bands.html>) from the original on 2010-04-30. Retrieved 2022-01-21.
15. "Advanced weapon systems using lethal Short-pulse terahertz radiation from high-intensity-laser-produced plasmas" (<https://web.archive.org/web/20100106223741/http://indiadaily.com/editorial/1803.asp>). *India Daily*. March 6, 2005. Archived from the original (<http://www.indiadaily.com/editorial/1803.asp>) on 6 January 2010. Retrieved 2010-09-27.
16. "Reference Solar Spectral Irradiance: Air Mass 1.5" (<http://rredc.nrel.gov/solar/spectra/am1.5/>). Archived (<https://web.archive.org/web/20190512190812/https://rredc.nrel.gov/solar/spectra/am1.5/>) from the original on 2019-05-12. Retrieved 2009-11-12.
17. Koontz, Steve (26 June 2012) Designing Spacecraft and Mission Operations Plans to Meet Flight Crew Radiation Dose (<https://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/20120012405.pdf>) Archived (<https://web.archive.org/web/20170502131715/https://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/20120012405.pdf>) 2017-05-02 at the [Wayback Machine](#). NASA/MIT Workshop. See pages I-7 (atmosphere) and I-23 (for water).
18. Uses of Electromagnetic Waves | gcse-revision, physics, waves, uses-electromagnetic-waves | Revision World (<http://www.revisionworld.com/gcse-revision/physics/waves/uses-electromagnetic-waves>)

## External links

---

- Australian Radiofrequency Spectrum Allocations Chart ([https://web.archive.org/web/20170114181318/http://www.acma.gov.au/webwr/radcomm/frequency\\_planning/spectrum\\_plan/arsp-wc.pdf](https://web.archive.org/web/20170114181318/http://www.acma.gov.au/webwr/radcomm/frequency_planning/spectrum_plan/arsp-wc.pdf)) (from Australian Communications and Media Authority)
- Canadian Table of Frequency Allocations ([http://www.ic.gc.ca/epic/site/smt-gst.nsf/vwapj/spectrallocation-08.pdf/\\$FILE/spectrallocation-08.pdf](http://www.ic.gc.ca/epic/site/smt-gst.nsf/vwapj/spectrallocation-08.pdf/$FILE/spectrallocation-08.pdf)) Archived ([https://web.archive.org/web/20081209114827/http://www.ic.gc.ca/epic/site/smt-gst.nsf/vwapj/spectrallocation-08.pdf/\\$FILE/spectrallocation-08.pdf](https://web.archive.org/web/20081209114827/http://www.ic.gc.ca/epic/site/smt-gst.nsf/vwapj/spectrallocation-08.pdf/$FILE/spectrallocation-08.pdf)) 2008-12-09 at the [Wayback Machine](#) (from Industry Canada)
- U.S. Frequency Allocation Chart (<http://www.ntia.doc.gov/osmhome/allochrt.html>) – Covering the range 3 kHz to 300 GHz (from Department of Commerce)
- UK frequency allocation table ([http://www.ofcom.org.uk/static/archive/ra/topics/spectrum-strat/future/strat02/strategy02app\\_b.pdf](http://www.ofcom.org.uk/static/archive/ra/topics/spectrum-strat/future/strat02/strategy02app_b.pdf)) (from Ofcom, which inherited the Radiocommunications Agency's duties, pdf format)



- [Flash EM Spectrum Presentation / Tool \(https://web.archive.org/web/20190205001750/http://attic.e-motiv.net/em-spectrum\)](https://web.archive.org/web/20190205001750/http://attic.e-motiv.net/em-spectrum) – Very complete and customizable.
- [Poster "Electromagnetic Radiation Spectrum" \(http://unihedron.com/projects/spectrum/downloads/spectrum\\_20090210.pdf\)](http://unihedron.com/projects/spectrum/downloads/spectrum_20090210.pdf) (992 kB)

---

Retrieved from "[https://en.wikipedia.org/w/index.php?title=Electromagnetic\\_spectrum&oldid=1305968550](https://en.wikipedia.org/w/index.php?title=Electromagnetic_spectrum&oldid=1305968550)"