

# Independent Component Analysis


Group6 : Armandine, Samuael, Verlon

African Master's in Machine Intelligence (AMMI)

Supervised by Pr. MOUSTAPHA CISSE

April 12, 2023

# Overview

- 
- 1 Introduction
    - Motivation
    - Problem formalization
    - ICA Assumptions
  - 2 ICA Ambiguities
  - 3 Densities and linear transformations
  - 4 ICA Algorithm

# Overview

- 
- 1 Introduction
    - Motivation
      - Problem formalization
      - ICA Assumptions
  - 2 ICA Ambiguities
  - 3 Densities and linear transformations
  - 4 ICA Algorithm



- Two people speaking simultaneously in a room.
- Speeches are recorded by two microphones in separate locations.




Image ref



**AIMS**

African Institute for  
Mathematical Sciences  
SENEGAL


# Overview

- 
- 1 **Introduction**
    - Motivation
    - **Problem formalization**
    - ICA Assumptions
  - 2 ICA Ambiguities
  - 3 Densities and linear transformations
  - 4 ICA Algorithm



# Problem formalization

- $d$ -speakers are speaking simultaneously at a party.



# Problem formalization

- $d$ -**speakers** are speaking simultaneously at a party.
- $d$  different **microphones** place in the same room records an overlapping combination of the  $d$  speakers.

# Problem formalization

- $d$ -**speakers** are speaking simultaneously at a party.
- $d$  different **microphones** place in the same room records an overlapping combination of the  $d$  speakers.
- Using  $d$ -**microphones**, can you separate out the  $d$ - **speakers** speech signals?




# Problem formalization

- $d$ -**speakers** are speaking simultaneously at a party.
- $d$  different **microphones** place in the same room records an overlapping combination of the  $d$  speakers.
- Using  $d$ -**microphones**, can you separate out the  $d$ - **speakers** speech signals?
- To solve this problem, we use Independent Component Analysis (ICA) which is a statistical method for separating a multivariate signal into independent components.

# Problem formalization

Let us consider some data  $S^{(i)} \in \mathbb{R}^d$  that is generated via  $d$  independent sources at time  $i$  as :



# Problem formalization

Let us consider some data  $S^{(i)} \in \mathbb{R}^d$  that is generated via  $d$  independent sources at time  $i$  as :

$$X^{(i)} = AS^{(i)} \quad , X^{(i)} \in \mathbb{R}^n \quad (1.1)$$

Where :

- $A$  is an unknown square matrix called **mixing matrix**

# Problem formalization

Let us consider some data  $S^{(i)} \in \mathbb{R}^d$  that is generated via  $d$  independent sources at time  $i$  as :

$$X^{(i)} = AS^{(i)} \quad , X^{(i)} \in \mathbb{R}^n \quad (1.1)$$

Where :

- $A$  is an unknown square matrix called **mixing matrix**
- $X^{(i)}$  is the signal of each microphone at time  $i$



# Problem formalization

Let us consider some data  $S^{(i)} \in \mathbb{R}^d$  that is generated via  $d$  independent sources at time  $i$  as :

$$X^{(i)} = AS^{(i)} \quad , X^{(i)} \in \mathbb{R}^n \quad (1.1)$$

Where :

- $A$  is an unknown square matrix called **mixing matrix**
- $X^{(i)}$  is the signal of each microphone at time  $i$
- $S^{(i)}$  is the signal of each speaker records at time  $i$

# Problem formalization

Let us consider some data  $S^{(i)} \in \mathbb{R}^d$  that is generated via  $d$  independent sources at time  $i$  as :

$$X^{(i)} = AS^{(i)} \quad , X^{(i)} \in \mathbb{R}^n \quad (1.1)$$

Where :

- $A$  is an unknown square matrix called **mixing matrix**
- $X^{(i)}$  is the signal of each microphone at time  $i$
- $S^{(i)}$  is the signal of each speaker records at time  $i$
- $S_j^{(i)}$  is the signal from speaker  $j$  at time  $i$



# Problem formalization

Let us consider some data  $S^{(i)} \in \mathbb{R}^d$  that is generated via  $d$  independent sources at time  $i$  as :

$$X^{(i)} = AS^{(i)} \quad , X^{(i)} \in \mathbb{R}^n \quad (1.1)$$

Where :

- $A$  is an unknown square matrix called **mixing matrix**
- $X^{(i)}$  is the signal of each microphone at time  $i$
- $S^{(i)}$  is the signal of each speaker records at time  $i$
- $S_j^{(i)}$  is the signal from speaker  $j$  at time  $i$
- $X_j^{(i)}$  is the signal of one microphone  $j$  records at time  $i$

Let  $W = A^{-1}$  be the **unmixing matrix**. Our goal is to find  $W$ . for notation convenience, also let  $w_i^T$  denote the  $i$  –  $th$  row of  $W$ , so that

$$W = \begin{bmatrix} -w_1^T - \\ -w_2^T - \\ \vdots \\ -w_d^T - \end{bmatrix}$$

Thus,  $w_i \in \mathbb{R}^d$ , and the  $j$  –  $th$  source can be recovered as  $s_j^{(i)} = w_j^T x^{(i)}$




# Overview

- 
- 1 Introduction
    - Motivation
    - Problem formalization
    - ICA Assumptions
  - 2 ICA Ambiguities
  - 3 Densities and linear transformations
  - 4 ICA Algorithm



- The number of sources is equal to the number of recorders.



- The number of sources is equal to the number of recorders.
- The Sources are statistically independent.




- The number of sources is equal to the number of recorders.
- The Sources are statistically independent.
- The independent component have a non gaussian distribution.

- The number of sources is equal to the number of recorders.
- The Sources are statistically independent.
- The independent component have a non gaussian distribution.
- The mixing matrix  $A$  is invertible.

# ICA Ambiguities

- The perturbation of the original sources doesn't matter for most applications.



# ICA Ambiguities

- The perturbation of the original sources doesn't matter for most applications.
- If the  $w_i$  found by the algorithm is scaled by any non zero real number, the corresponding recovered source  $S_i$  will be scaled by the same factor. This usually doesn't matter.

# ICA Ambiguities

- The perturbation of the original sources doesn't matter for most applications.
- If the  $w_i$  found by the algorithm is scaled by any non zero real number, the corresponding recovered source  $S_i$  will be scaled by the same factor. This usually doesn't matter.
- Since ICA separates sources by maximizing their non-Gaussianity, perfect Gaussian sources can't be separated.



## Densities and linear transformations

Let's consider a random variable  $S$  draw according to some density  $P_S(S)$ .

Let  $x$  defined according to  $x = AS = W^{-1}S$ , ( $S = Wx$  with  $W = A^{-1}$ ).

So,

$$P_x(x) = P_S(S) = P_S(Wx) \quad \text{"is not correct"} \quad (3.1)$$

## Densities and linear transformations

Let's consider a random variable  $S$  draw according to some density  $P_S(S)$ .

Let  $x$  defined according to  $x = AS = W^{-1}S$ , ( $S = Wx$  with  $W = A^{-1}$ ).

So,

$$P_x(x) = P_s(S) = P_s(Wx) \quad \text{"is not correct"} \quad (3.1)$$

and in particular the right formula is :

$$P_x(x) = P_s(Wx) \cdot |W| \quad (3.2)$$

with  $|W|$  the determinant of matrix  $W$ .

# Densities and linear transformations

## Choice of $P_s(S_j)$ :

We can choose density or as equivalent, we can choose a CDF. For the historical reason and for convenience we need to choose some function that increase from 0 to 1. And we can't choose a Gaussian. For convenience we choose the CDF to be a sigmoïd function.

# ICA Algorithm

Two principles of ICA estimations :

- **Minimization of the mutual information (ICA gradient ascend and Fast ICA) and**
- **Maximum Likelihood Estimation (MLE)**

# MLE

We suppose that the distribution of each source  $S_j$  is given by a density  $P_s$  and that the joint distribution of the sources  $S$  is given by

$$P(S) = \prod_{j=1}^d P_s(S_j)$$

$$P_x(x) = \left( \prod_{j=1}^d P_s(W_j^T x) \right) |W|$$

$$L(W) = \log \left( \prod_{i=1}^n \left( \prod_{j=1}^d P_s(W_j^T x^{(i)}) \right) |W| \right)$$



$$L(W) = \sum_{i=1}^n \left( \left( \sum_{j=1}^d \log(P_s(W_j^T x^{(i)})) \right) + \log(|W|) \right) \quad (4.1)$$

Let's consider

$$g(S_j) = \frac{1}{1 + \exp(-S_j)} \quad \text{with} \quad S_j = W_j^T x$$

So

$$P_s(S_j) = g'(S_j) = g(S_j)(1 - g(S_j))$$

Then,

$$L(W) = \sum_{i=1}^n \left( \left( \sum_{j=1}^d \log(g'(S_j)) \right) + \log(|W|) \right) \quad (4.2)$$



**AIMS**

African Institute for  
Mathematical Sciences  
SENEGAL

# ICA Algorithm

Let us compute the gradient of  $L(W)$  given in (4.2) w.r.t  $W$  So,

$$\nabla_W L(W) = \sum_{i=1}^n \left( \left( \sum_{j=1}^d \nabla_W \log(g'(S_j)) \right) + \nabla_W \log(|W|) \right)$$

•

$$\nabla_W \log(|W|) = \frac{\nabla_W |W|}{|W|} = (W^{-1})^T = (W^T)^{-1},$$

because

$$\nabla_W |W| = |W|(W^{-1})^T$$

• Set  $f(g'(S_j)) = \log(g'(S_j))$ . So,

$$\nabla_W f = \frac{\partial f}{\partial g'(S_j)} \frac{\partial g'(S_j)}{\partial S_j} \frac{\partial S_j}{\partial W} \quad (\text{Chain rule's})$$



**AIMS**

African Institute for  
Mathematical Sciences  
SENEGAL

# ICA Algorithm

$$\frac{\partial S_j}{\partial w} = x, \frac{\partial g'(S_j)}{\partial S_j} = g'(S_j)(1 - 2g(S_j)) \quad \text{and} \quad \frac{\partial f}{\partial g'(S_j)} = \frac{1}{g'(S_j)}$$

Then,

$$\nabla_W f = (1 - 2g(S_j))$$

Therefore,

$$\nabla_W L(W) = \sum_{i=1}^n \left( (1 - 2g(w_j^T x^{(i)})) x^{(i)T} + (W^T)^{-1} \right) \quad \text{where } S_j^{(i)} = w_j^T x^{(i)} \quad (4.3)$$





# ICA Algorithm




Then, for the training example  $x^{(i)}$ , the update rule using the stochastic gradient ascent is :

$$W := W + \alpha \left( \begin{bmatrix} 1 - 2g(w_1^T x^{(i)}) \\ 1 - 2g(w_2^T x^{(i)}) \\ \vdots \\ 1 - 2g(w_d^T x^{(i)}) \end{bmatrix} x^{(i)T} + (W^T)^{-1} \right)$$

Where  $\alpha$  is the learning rate.

We then compute  $S^{(i)} = Wx^{(i)}$  to recover the original sources.  
After the algorithm converges.

## References

-  Andrew Ng, Tengyu.  
CS229 Lecture Notes, 2022.
-  Hyvarinen, Aapo.  
"Independent component analysis, a tutorial."  
[http://www.cs.helsinki.fi/u/ahyvarin/papers/NN00new](http://www.cs.helsinki.fi/u/ahyvarin/papers/NN00new.pdf), pdf  
(1999).
-  Andrew Ng.  
Lecture 15| Machine Learning  
[https://www.youtube.com/watch?v=QGd06MTRMHslist=PLFyHXC1T3SSje86s3ukLqTP8\\_3eS\\_xc-j](https://www.youtube.com/watch?v=QGd06MTRMHslist=PLFyHXC1T3SSje86s3ukLqTP8_3eS_xc-j)

# Acknowledgements



THANK YOU