

# On Shelling a Family of Symmetric Spaces

Fernando Díaz Morera  
(joint with Aram Bingham)



AMS Special Session: Geometry, Combinatorics, and flag varieties  
Saint Louis University, Saint Louis, MO  
October 18-19, 2025

# Outline

- 1 Poset
  - Weyl group
- 2 EL-labelling
- 3 Intermezzo: Stanley-Reisner correspondence
- 4 Symmetric spaces
  - Classification
  - Type AIII
  - Sects
- 5 Coverings and labelling on  $\mathcal{C}_{p,q}^\lambda$
- 6 Questions

## (Strong) Bruhat order: $(\mathfrak{S}_n, <_{\text{BC}})$

### Permutation group

$$\mathfrak{S}_n := \overbrace{\{v : [n] \rightarrow [n] \mid v \text{ is 1-1}\}}.$$

- $v = v_1 v_2 \cdots v_n \rightsquigarrow v(i) = v_i$ .
  - ▶  $v = 312 \rightsquigarrow v(2) = 1 = v_2$ .
- $\underbrace{|\{(i, j) : i \leq i < j \leq n, v_i > v_j\}|}_{\ell(v)}$ .
  - ▶  $\ell(312) = |(1, 2), (1, 3)| = 2$ .

## (Strong) Bruhat order: $(\mathfrak{S}_n, <_{\text{BC}})$

### Permutation group

$$\mathfrak{S}_n := \{v : [n] \rightarrow [n] \mid v \text{ is 1-1}\}.$$

- $v = v_1 v_2 \cdots v_n \rightsquigarrow v(i) = v_i$ .
  - ▶  $v = 312 \rightsquigarrow v(2) = 1 = v_2$ .
- $|\underbrace{\{(i, j) : i \leq i < j \leq n, v_i > v_j\}}_{\ell(v)}|$ .
  - ▶  $\ell(312) = |(1, 2), (1, 3)| = 2$ .
- $u <_{\text{BC}} v \iff v$  is attainable from  $u$  by a sequence of transposition  $(ij)$  such that  $i < j$  and  $u_i < u_j$ .

$$1324 \xrightarrow{(13)} 3124 \xrightarrow{(24)} 3142 \rightsquigarrow 1324 <_{\text{BC}} 3142.$$

# (Strong) Bruhat order: $(\mathfrak{S}_n, <_{\text{BC}})$

## Permutation group


$$\mathfrak{S}_n := \{v : [n] \rightarrow [n] \mid v \text{ is 1-1}\}.$$

- $v = v_1 v_2 \cdots v_n \rightsquigarrow v(i) = v_i$ .
  - ▶  $v = 312 \rightsquigarrow v(2) = 1 = v_2$ .
- $|\{(i, j) : i \leq i < j \leq n, v_i > v_j\}|$ .

$\ell(v)$

  - ▶  $\ell(312) = |(1, 2), (1, 3)| = 2$ .
- $u <_{\text{BC}} v \iff v$  is attainable from  $u$  by a sequence of transposition  $(ij)$  such that  $i < j$  and  $u_i < u_j$ .

$$1324 \xrightarrow{(13)} 3124 \xrightarrow{(24)} 3142 \rightsquigarrow 1324 <_{\text{BC}} 3142.$$




# (Strong) Bruhat order: $(\mathfrak{S}_n, <_{\text{BC}})$

## Permutation group

$$\mathfrak{S}_n := \{v : [n] \rightarrow [n] \mid v \text{ is 1-1}\}.$$

- $v = v_1 v_2 \cdots v_n \rightsquigarrow v(i) = v_i$ .  
 ▶  $v = 312 \rightsquigarrow v(2) = 1 = v_2$ .
- $|\{(i, j) : i \leq i < j \leq n, v_i > v_j\}|$ .  
 $\ell(v)$   
 ▶  $\ell(312) = |(1, 2), (1, 3)| = 2$ .
- $u <_{\text{BC}} v \iff v$  is attainable from  $u$  by a sequence of transposition  $(ij)$  such that  $i < j$  and  $u_i < u_j$ .

$$1324 \xrightarrow{(13)} 3124 \xrightarrow{(24)} 3142 \rightsquigarrow 1324 <_{\text{BC}} 3142.$$



$$\mathbf{T}_n \subset \mathbf{B}_n \subset \text{GL}_n(\mathbb{C}) \rightsquigarrow \text{GL}_n / \mathbf{B} = \bigsqcup \mathbf{B} v \mathbf{B} / \mathbf{B} \rightsquigarrow X_v \mathbf{B} := \overline{\mathbf{B} v \mathbf{B}} / \mathbf{B}.$$

$$\mathfrak{S}_n \cong \text{N}_{\text{GL}_n}(\mathbf{T}) / \mathbf{T}, \quad u <_{\text{BC}} v \iff X_u \mathbf{B} \subseteq X_v \mathbf{B}, \quad \ell(v) = \dim X_v \mathbf{B}$$

# Posets

- A poset  $\mathcal{P}$  with order relation  $<$  is bounded if there are elements  $\hat{0}$  and  $\hat{1}$  such that  $\hat{0} < v < \hat{1}$  for all  $v$  in  $\mathcal{P}$ .

# Posets

- A poset  $\mathcal{P}$  with order relation  $<$  is bounded if there are elements  $\hat{0}$  and  $\hat{1}$  such that  $\hat{0} < v < \hat{1}$  for all  $v$  in  $\mathcal{P}$ .
- If  $u < v$  and there is not  $u_1$  such that  $u < u_1 < v$ , we say  $v$  covers  $u$ , denoted as  $u \triangleleft v$ .
- A chain is saturated if it is of the form  $u = u_0 \triangleleft u_1 \triangleleft \cdots \triangleleft u_m = v$ .



# Posets

- A poset  $\mathcal{P}$  with order relation  $<$  is bounded if there are elements  $\hat{0}$  and  $\hat{1}$  such that  $\hat{0} < v < \hat{1}$  for all  $v$  in  $\mathcal{P}$ .
- If  $u < v$  and there is not  $u_1$  such that  $u < u_1 < v$ , we say  $v$  covers  $u$ , denoted as  $u \triangleleft v$ .
- A chain is saturated if it is of the form  $u = u_0 \triangleleft u_1 \triangleleft \cdots \triangleleft u_m = v$ .
- $\mathcal{P}$  is graded if all maximal chains have the same length.
- If  $\mathcal{P}$  is graded, then the length function  $\ell : \mathcal{P} \rightarrow \mathbb{N}$  assigns to  $v$  in  $\mathcal{P}$  the value  $m$  where  $u_0 \triangleleft u_1 \triangleleft \cdots \triangleleft u_m = v$  for any saturated chain to  $v$ .

# Posets

- A poset  $\mathcal{P}$  with order relation  $<$  is bounded if there are elements  $\hat{0}$  and  $\hat{1}$  such that  $\hat{0} < v < \hat{1}$  for all  $v$  in  $\mathcal{P}$ .
- If  $u < v$  and there is not  $u_1$  such that  $u < u_1 < v$ , we say  $v$  covers  $u$ , denoted as  $u \triangleleft v$ .
- A chain is saturated if it is of the form  $u = u_0 \triangleleft u_1 \triangleleft \cdots \triangleleft u_m = v$ .
- $\mathcal{P}$  is graded if all maximal chains have the same length.
- If  $\mathcal{P}$  is graded, then the length function  $\ell : \mathcal{P} \rightarrow \mathbb{N}$  assigns to  $v$  in  $\mathcal{P}$  the value  $m$  where  $u_0 \triangleleft u_1 \triangleleft \cdots \triangleleft u_m = v$  for any saturated chain to  $v$ .

Finding the number of partially order sets  
with  $n$  elements is still unknown...

## EL-labeling on $\mathfrak{S}_n$

Let  $u = u_1 u_2 \cdots u_n$  and  $v = v_1 v_2 \cdots v_n$  be in  $\mathfrak{S}_n$ . We say  $u \prec_{\text{BC}} v$  whether  $\ell(v) = \ell(u) + 1$ , and

(i)  $u_k = v_k$  for  $k$  in  $\{1, \dots, \hat{i}, \dots, \hat{j}, \dots, n\}$

(ii)  $u_i = v_j, u_j = v_i$ , and  $u_i < u_j$ .

►  $213 \prec_{\text{BC}} 231 \rightsquigarrow u_1 = v_1, u_2 = v_3, u_3 = v_2$  and  $u_2 < u_3$ .

## EL-labeling on $\mathfrak{S}_n$

Let  $u = u_1 u_2 \cdots u_n$  and  $v = v_1 v_2 \cdots v_n$  be in  $\mathfrak{S}_n$ . We say  $u \leq_{\text{BC}} v$  whether  $\ell(v) = \ell(u) + 1$ , and

(i)  $u_k = v_k$  for  $k$  in  $\{1, \dots, \hat{i}, \dots, \hat{j}, \dots, n\}$

(ii)  $u_i = v_j, u_j = v_i$ , and  $u_i < u_j$ .

►  $213 \leq_{\text{BC}} 231 \rightsquigarrow u_1 = v_1, u_2 = v_3, u_3 = v_2$  and  $u_2 < u_3$ .

Let  $\Lambda = [n] \times [n]$  denote poset of pair such that  $(i, j) \leq (r, s)$  if  $i < r$ , or  $i = r$  and  $j < s$ .

$$C(\mathfrak{S}_n) \xrightarrow{\eta} \Lambda$$

$$(u, v) \longrightarrow \eta(u, v) := (u_i, u_j)$$

## EL-labeling on $\mathfrak{S}_n$

Let  $u = u_1 u_2 \cdots u_n$  and  $v = v_1 v_2 \cdots v_n$  be in  $\mathfrak{S}_n$ . We say  $u \leq_{\text{BC}} v$  whether  $\ell(v) = \ell(u) + 1$ , and

(i)  $u_k = v_k$  for  $k$  in  $\{1, \dots, \hat{i}, \dots, \hat{j}, \dots, n\}$

(ii)  $u_i = v_j, u_j = v_i$ , and  $u_i < u_j$ .

►  $213 \leq_{\text{BC}} 231 \rightsquigarrow u_1 = v_1, u_2 = v_3, u_3 = v_2$  and  $u_2 < u_3$ .

Let  $\Lambda = [n] \times [n]$  denote poset of pair such that  $(i, j) \leq (r, s)$  if  $i < r$ , or  $i = r$  and  $j < s$ .

$$C(\mathfrak{S}_n) \xrightarrow{\eta} \Lambda$$

$$(u, v) \longrightarrow \eta(u, v) := (u_i, u_j)$$

E.g. Let  $v = 312$  and  $e = 123$  be in  $\mathfrak{S}_3$ . The max chain  $e \leq_{\text{BC}} 213 \leq_{\text{BC}} v$  is the **LEX** smallest. Moreover,  $\eta(e, 213) \leq_{\text{LEX}} \eta(213, v)$  is non-decreasing.

# Lexicographic shellability

Let  $C(\mathcal{P}) := \{(u, v) \in \mathcal{P} \times \mathcal{P} \mid u \lessdot v\}$  denote the set of covering relations in a poset  $\mathcal{P}$ . An **EL-labelling** on  $(\mathcal{P}, <)$  is a map  $\eta : C(\mathcal{P}) \rightarrow (\Lambda, \leq_{\text{LEX}})$  holding the following:

## Lexicographic shellability

Let  $C(\mathcal{P}) := \{(u, v) \in \mathcal{P} \times \mathcal{P} \mid u \triangleleft v\}$  denote the set of covering relations in a poset  $\mathcal{P}$ . An **EL-labelling** on  $(\mathcal{P}, <)$  is a map  $\eta : C(\mathcal{P}) \rightarrow (\Lambda, \leq_{\text{LEX}})$  holding the following:

- (i) For each  $u < v$ , there is a *unique non-decreasing sequence* from  $u$  to  $v$ . That is, there is a unique saturated chain  $u \triangleleft u_1 \triangleleft \cdots \triangleleft u_k \triangleleft v$  with  $\eta(u, u_1) \leq \eta(u_1, u_2) \leq \cdots \leq \eta(u_k, v)$ .

# Lexicographic shellability

Let  $C(\mathcal{P}) := \{(u, v) \in \mathcal{P} \times \mathcal{P} \mid u \triangleleft v\}$  denote the set of covering relations in a poset  $\mathcal{P}$ . An **EL-labelling** on  $(\mathcal{P}, <)$  is a map  $\eta : C(\mathcal{P}) \rightarrow (\Lambda, \leq_{\text{LEX}})$  holding the following:

- (i) For each  $u < v$ , there is a *unique non-decreasing sequence* from  $u$  to  $v$ . That is, there is a unique saturated chain  $u \triangleleft u_1 \triangleleft \cdots \triangleleft u_k \triangleleft v$  with  $\eta(u, u_1) \leq \eta(u_1, u_2) \leq \cdots \leq \eta(u_k, v)$ .
- (ii) The above label sequence is lexicographically *smaller* than the label sequence for every other saturated chain from  $u$  to  $v$ . That is, if  $u \triangleleft w < v$ , with  $w \neq u_1$  as defined above, then  $\eta(u, u_1) \leq \eta(u, w)$ .




## Lexicographic shellability

Let  $C(\mathcal{P}) := \{(u, v) \in \mathcal{P} \times \mathcal{P} \mid u \lessdot v\}$  denote the set of covering relations in a poset  $\mathcal{P}$ . An **EL-labelling** on  $(\mathcal{P}, <)$  is a map  $\eta : C(\mathcal{P}) \rightarrow (\Lambda, \leq_{\text{LEX}})$  holding the following:


- (i) For each  $u < v$ , there is a *unique non-decreasing sequence* from  $u$  to  $v$ . That is, there is a unique saturated chain  $u \lessdot u_1 \lessdot \cdots \lessdot u_k \lessdot v$  with  $\eta(u, u_1) \leq \eta(u_1, u_2) \leq \cdots \leq \eta(u_k, v)$ .
- (ii) The above label sequence is lexicographically *smaller* than the label sequence for every other saturated chain from  $u$  to  $v$ . That is, if  $u \lessdot w < v$ , with  $w \neq u_1$  as defined above, then  $\eta(u, u_1) \leq \eta(u, w)$ .

A poset  $\mathcal{P}$  is **EL-shellable or lexicographically shellable** if it admits an *EL*-labeling.


# EL-labeling of $\mathfrak{S}_3$




$\Delta(P) \rightarrow$



## EL-labeling of $\mathfrak{S}_3$



$\Delta(P) \rightarrow$




The **order complex** of a poset  $\mathcal{P}$  is the abstract *simplicial complex*, denoted  $\Delta(\mathcal{P})$ , whose  $k$ -dimensional faces are the chains  $u_0 < u_1 < \cdots < u_k$  of  $k + 1$  comparable poset elements.

## Intermezzo: commutative algebra

A *simplicial complex*  $\Delta$  is **shellable** if there is an ordering of the maximal faces  $F_1, F_2, \dots, F_m$  so that for all  $i, j$  with  $i < j$ , there exists  $k < j$  such that  $F_i \cap F_j \subseteq F_k \cap F_j = F_j \setminus \{p\}$  for some  $p$  in  $F_j$ .

## Intermezzo: commutative algebra


A simplicial complex  $\Delta$  is **shellable** if there is an ordering of the maximal faces  $F_1, F_2, \dots, F_m$  so that for all  $i, j$  with  $i < j$ , there exists  $k < j$  such that  $F_i \cap F_j \subseteq F_k \cap F_j = F_j \setminus \{p\}$  for some  $p$  in  $F_j$ .



$$\begin{aligned}
 I_\Delta &= (x_{i_1} \cdots x_{i_r} : \{i_1, \dots, i_r\} \notin \Delta) \\
 &\quad \downarrow \\
 S &:= \mathbf{k}[x_1, x_2, \dots, x_n] \\
 &\quad \downarrow \\
 S/I_\Delta &:= \mathbf{k}[\Delta] \text{ Stanley-Reisner}
 \end{aligned}$$

## Intermezzo: commutative algebra

A *simplicial complex*  $\Delta$  is **shellable** if there is an ordering of the maximal faces  $F_1, F_2, \dots, F_m$  so that for all  $i, j$  with  $i < j$ , there exists  $k < j$  such that  $F_i \cap F_j \subseteq F_k \cap F_j = F_j \setminus \{p\}$  for some  $p$  in  $F_j$ .



### Theorem (Björner '80)

If  $\mathcal{P}$  is a graded poset with an *EL*-labeling, then  $\Delta(\mathcal{P})$  is shellable.

### Theorem (Kind-Kleinschmidt '79)

The Stanley-Reisner ring of a shellable  $\Delta$  is *Cohen–Macaulay*.

## Results on shellability and Bruhat order

- $\mathfrak{S}_n$  Bruhat order is EL-shellable (Edelman, '81).

## Results on shellability and Bruhat order

- $\mathfrak{S}_n$  Bruhat order is EL-shellable (Edelman, '81).
- *Classical* Coxeter groups are EL-shellable (Proctor, '82)
  - ▶  $S/I_\lambda$  Cohen-Macaulay  $\rightsquigarrow X_\lambda \subset \mathbf{G}/\mathbf{P}$  normal.
- Bruhat order on  $\mathbf{W}/\mathbf{W}_J$  is CL-shellable for any Coxeter group  $\mathbf{W}$  and any parabolic subgroup  $\mathbf{W}_J \subset \mathbf{W}$  (Björner-Wachs, '82).



## Results on shellability and Bruhat order

- $\mathfrak{S}_n$  Bruhat order is EL-shellable (Edelman, '81).
- *Classical* Coxeter groups are EL-shellable (Proctor, '82)
  - ▶  $S/I_\lambda$  Cohen-Macaulay  $\rightsquigarrow X_\lambda \subset \mathbf{G}/\mathbf{P}$  normal.
- Bruhat order on  $\mathbf{W}/\mathbf{W}_J$  is CL-shellable for any Coxeter group  $\mathbf{W}$  and any parabolic subgroup  $\mathbf{W}_J \subset \mathbf{W}$  (Björner-Wachs, '82).
- Bruhat order on any  $\mathbf{W}/\mathbf{W}_J$  is EL-shellable (Dyer, '93).


# Results on shellability and Bruhat order

- $\mathfrak{S}_n$  Bruhat order is EL-shellable (Edelman, '81).
- *Classical* Coxeter groups are EL-shellable (Proctor, '82)
  - ▶  $S/I_\lambda$  Cohen-Macaulay  $\rightsquigarrow X_\lambda \subset \mathbf{G}/\mathbf{P}$  normal.
- Bruhat order on  $\mathbf{W}/\mathbf{W}_J$  is CL-shellable for any Coxeter group  $\mathbf{W}$  and any parabolic subgroup  $\mathbf{W}_J \subset \mathbf{W}$  (Björner-Wachs, '82).
- Bruhat order on any  $\mathbf{W}/\mathbf{W}_J$  is EL-shellable (Dyer, '93).
- Bruhat order on  $\text{Invol}(\mathbf{W})$  is EL-shellable where  $\mathbf{W}$  is classical (Incitti, '04/'06).

# Results on shellability and Bruhat order


- $\mathfrak{S}_n$  Bruhat order is EL-shellable (Edelman, '81).
- *Classical* Coxeter groups are EL-shellable (Proctor, '82)
  - ▶  $S/I_\lambda$  Cohen-Macaulay  $\rightsquigarrow X_\lambda \subset \mathbf{G}/\mathbf{P}$  normal.
- Bruhat order on  $\mathbf{W}/\mathbf{W}_J$  is CL-shellable for any Coxeter group  $\mathbf{W}$  and any parabolic subgroup  $\mathbf{W}_J \subset \mathbf{W}$  (Björner-Wachs, '82).
- Bruhat order on any  $\mathbf{W}/\mathbf{W}_J$  is EL-shellable (Dyer, '93).
- Bruhat order on  $\text{Invol}(\mathbf{W})$  is EL-shellable where  $\mathbf{W}$  is classical (Incitti, '04/'06).
- Fixed-point free involutions in  $\mathfrak{S}_n$  are EL-shellable (Can-Cherniavsky-Twelbeck, '15).
- The *rook monoid* is EL-shellable (Can, '19).

# Symmetric spaces



Let  $\theta : \mathbf{G} \rightarrow \mathbf{G}$  such that  $\theta^2 = \text{id}$ . We say  $\mathbf{G}/\mathbf{K}$  is a symmetric space if  $\mathbf{K} = \mathbf{G}^\theta$  (fixed-point subgroup).

# Symmetric spaces




Let  $\theta : \mathbf{G} \rightarrow \mathbf{G}$  such that  $\theta^2 = \text{id}$ . We say  $\mathbf{G}/\mathbf{K}$  is a symmetric space if  $\mathbf{K} = \mathbf{G}^\theta$  (fixed-point subgroup).

## Theorem (Matsuki '79)

$\mathbf{G}/\mathbf{K}$  is a spherical variety, i.e.  $\mathbf{B}$  acts on  $\mathbf{G}/\mathbf{K}$  with finitely many orbits.

# Symmetric spaces



Let  $\theta : \mathbf{G} \rightarrow \mathbf{G}$  such that  $\theta^2 = \text{id}$ . We say  $\mathbf{G}/\mathbf{K}$  is a symmetric space if  $\mathbf{K} = \mathbf{G}^\theta$  (fixed-point subgroup).

## Theorem (Matsuki '79)

$\mathbf{G}/\mathbf{K}$  is a spherical variety, i.e.  $\mathbf{B}$  acts on  $\mathbf{G}/\mathbf{K}$  with finitely many orbits.

$\mathbf{G} = \text{GL}_n \times \text{GL}_n$ ,  $\theta(g, h) = (h, g)$ ,  $\mathbf{K} = \{(g, g) : g \in \text{GL}_n\}$ . So  $\mathbf{G}/\mathbf{K} \cong \text{GL}_n$  acted on by  $\mathbf{B} = \mathbf{B}_n \times \mathbf{B}_n \subset \mathbf{G}$ . In particular,  $\mathbf{B}$ -orbits are the same as  $\mathbf{B}_n \nu \mathbf{B}_n$  double cosets,  $\nu$  in  $\mathfrak{S}_n \dots$

# Classification (à la Cartan)

Type	<b>G</b>	<b>K</b>	<b>B \ G / K</b>	Shellable
AI	$GL_n(\text{or } SL_n)$	$O_n(\text{or } SO_n)$	$\text{Invol}(n)$	✓(Incitti, '04)
AII	$SL_{2n}$	$Sp_{2n}$	$\text{Invol}^{FPF}(2n)$	✓(Can-C-T, '15)
AIII	$GL_{p+q}$	$GL_p \times GL_q$	$(p, q)\text{-clans}$	☕✍...
CI	$Sp_{2n}$	$GL_n$	$(n, n)\text{-clans}$	🦇
CII	$Sp_{p+q}$	$Sp_{2p} \times Sp_{2q}$	CII-clans	🦇
BDI	$SO_n$	$SO_k \times SO_{n-k}$	$(k, n - k)\text{-clans}$	🦇
DIII	$SO_{2n}$	$GL_n$	DIII-clans	🦇

# Clans

## Definition (Yamamoto '97 - Wyser '06)

Let  $p$  and  $q$  be two positive integers such that  $p + q = n$ . A  $(p, q)$ -**clan** is an ordered set of  $n$  symbols  $c_1 \dots c_n$  such that:

- (i) Each symbol  $c_i$  is either “+”, “−” or a  $\mathbb{N}_{>0}$ .
- (ii) If  $c_i \in \mathbb{N}$ , then there is a unique index  $j \neq i$  such that  $c_i = c_j$ .
- (iii) The difference between the numbers of “+” and “−” symbols in the clan is equal to  $p - q$ . If  $q > p$ , then we have  $q - p$  more “−” signs than “+” signs.

We denote the set of all  $(p, q)$ -clans by  $\mathcal{C}_{p,q}$ .



# Clans

## Definition (Yamamoto '97 - Wyser '06)

Let  $p$  and  $q$  be two positive integers such that  $p + q = n$ . A  $(p, q)$ -**clan** is an ordered set of  $n$  symbols  $c_1 \dots c_n$  such that:

- (i) Each symbol  $c_i$  is either “+”, “−” or a  $\mathbb{N}_{>0}$ .
- (ii) If  $c_i \in \mathbb{N}$ , then there is a unique index  $j \neq i$  such that  $c_i = c_j$ .
- (iii) The difference between the numbers of “+” and “−” symbols in the clan is equal to  $p - q$ . If  $q > p$ , then we have  $q - p$  more “−” signs than “+” signs.

We denote the set of all  $(p, q)$ -clans by  $\mathcal{C}_{p,q}$ .

## Example

$\gamma_1 = +1212-$  and  $\gamma_2 = +1717-$  are equivalent  $(3, 3)$ -clans. Likewise,  $\gamma_3 = 12+21$  is a  $(3, 2)$ -clan and  $\gamma_4 = +1+1$  is a  $(3, 1)$ -clan.

## Stemming from...

### Theorem (Matsuki-Oshima '90)

Let  $\mathcal{C}_{p,q}$  denote the set of  $(p, q)$ -clans. Then

$$\mathcal{C}_{p,q} \xleftrightarrow[1:1]{} \left\{ \begin{array}{c} \mathbf{B}\text{-orbits in } \mathrm{GL}_{p+q} / \mathrm{GL}_p \times \mathrm{GL}_q \\ \text{or} \\ \mathrm{GL}_p \times \mathrm{GL}_q\text{-orbits in } \mathrm{GL}_n / \mathbf{B} \end{array} \right\}.$$

Moreover, a clan is **matchless** if it consists only of  $+/-$ 's. Hence,

$$\{\text{matchless clans}\} \xleftrightarrow[1:1]{} \{\text{closed orbits/min. dim.}\}.$$

# Stemming from...

## Theorem (Matsuki-Oshima '90)

Let  $\mathcal{C}_{p,q}$  denote the set of  $(p, q)$ -clans. Then

$$\mathcal{C}_{p,q} \xleftrightarrow[1:1]{} \left\{ \begin{array}{c} \mathbf{B}\text{-orbits in } \mathrm{GL}_{p+q} / \mathrm{GL}_p \times \mathrm{GL}_q \\ \text{or} \\ \mathrm{GL}_p \times \mathrm{GL}_q\text{-orbits in } \mathrm{GL}_n / \mathbf{B} \end{array} \right\}.$$

Moreover, a clan is **matchless** if it consists only of  $+/-$ 's. Hence,

$$\{\text{matchless clans}\} \xleftrightarrow[1:1]{} \{\text{closed orbits/min. dim.}\}.$$

In other words, for clans  $\gamma_1, \gamma_2$  in  $\mathcal{C}_{p,q}$ , we have corresponding  $\mathbf{B}$ -orbits  $\mathcal{O}_{\gamma_1}$  and  $\mathcal{O}_{\gamma_2}$ . Then the Bruhat poset  $(\mathcal{C}_{p,q}, \leq)$  is defined by the relation

$$\gamma_1 \leq \gamma_2 \iff \mathcal{O}_{\gamma_1} \subseteq \overline{\mathcal{O}_{\gamma_2}}.$$

## Stemming from...

### Theorem (Matsuki-Oshima '90)

Let  $\mathcal{C}_{p,q}$  denote the set of  $(p, q)$ -clans. Then

$$\mathcal{C}_{p,q} \xleftrightarrow[1:1]{} \left\{ \begin{array}{c} \mathbf{B}\text{-orbits in } \mathrm{GL}_{p+q} / \mathrm{GL}_p \times \mathrm{GL}_q \\ \text{or} \\ \mathrm{GL}_p \times \mathrm{GL}_q\text{-orbits in } \mathrm{GL}_n / \mathbf{B} \end{array} \right\}.$$

Moreover, a clan is **matchless** if it consists only of  $+/-$ 's. Hence,


$$\{\text{matchless clans}\} \xleftrightarrow[1:1]{} \{\text{closed orbits/min. dim.}\}.$$

In other words, for clans  $\gamma_1, \gamma_2$  in  $\mathcal{C}_{p,q}$ , we have corresponding  $\mathbf{B}$ -orbits  $\mathcal{O}_{\gamma_1}$  and  $\mathcal{O}_{\gamma_2}$ . Then the Bruhat poset  $(\mathcal{C}_{p,q}, \leq)$  is defined by the relation

$$\gamma_1 \leq \gamma_2 \iff \mathcal{O}_{\gamma_1} \subseteq \overline{\mathcal{O}_{\gamma_2}}.$$

**Shortcoming:** There is no longer a unique minimal element...

# Bruhat poset $\mathcal{C}_{2,2}$ (à la Wyser '16)



# Sects

## Def/Prop (Bingham-Can '20)

Let  $C_\lambda$  denote the Schubert cell of  $\mathrm{GL}_{p+q} / \mathbf{P}$  associated to the partition  $\lambda \in \binom{[p+q]}{p}$ . Then the **sect**  $\mathcal{C}_{p,q}^\lambda$  is the collection of clans  $\gamma$  whose corresponding orbits satisfy  $\pi(\mathcal{O}_\gamma) = C_\lambda$  where  $\pi : \mathbf{G} / \mathbf{L} \rightarrow \mathbf{G} / \mathbf{P}$  is the natural projection map.

# Sects

## Def/Prop (Bingham-Can '20)

Let  $C_\lambda$  denote the Schubert cell of  $\mathrm{GL}_{p+q}/\mathbf{P}$  associated to the partition  $\lambda \in ([p+q]_p)$ . Then the **sect**  $\mathcal{C}_{p,q}^\lambda$  is the collection of clans  $\gamma$  whose corresponding orbits satisfy  $\pi(\mathcal{O}_\gamma) = C_\lambda$  where  $\pi : \mathbf{G}/\mathbf{L} \rightarrow \mathbf{G}/\mathbf{P}$  is the natural projection map.

## Combinatorially

(i)  $\{\text{matchless/base clans } \tau\} \xleftrightarrow{1:1} \{\text{sects/Schubert cells/partitions } \lambda\}$

$$\tau_\gamma = -+--++ \longleftrightarrow \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline & & \\ \hline & & \\ \hline \end{array}$$

(ii) Label the symbols of  $\gamma$  as  $\gamma = c_1 \cdots c_{p+q}$ . For each pair  $c_i = c_j \in \mathbb{N}$  with  $i < j$ , replace  $c_i$  by a  $-$  symbol and  $c_j$  by a  $+$  symbol.

$$\gamma = 1+-221 \mapsto -+--++ = \tau_\gamma$$

## From sects to rooks

Let  $R(\lambda)$  denote the set of **rook placements** of partition  $\lambda$ . For  $\rho, \pi$  in  $R(\lambda)$ , we say  $\rho \leq \pi \iff rt_\rho \leq rt_\pi$  where  $R(\lambda) \xrightarrow{rt} \mathbb{N}$  is a labeling of the boxes of  $[\lambda]$  by the number of rooks weakly NW of each box.



## From sects to rooks

Let  $R(\lambda)$  denote the set of **rook placements** of partition  $\lambda$ . For  $\rho, \pi$  in  $R(\lambda)$ , we say  $\rho \leq \pi \iff rt_\rho \leq rt_\pi$  where  $R(\lambda) \xrightarrow{rt} \mathbb{N}$  is a labeling of the boxes of  $[\lambda]$  by the number of rooks weakly NW of each box.

### Bingham '21

There is an isomorphism between  $R(\lambda)$  and  $\mathcal{C}_{p,q}^\lambda$  as follows.

- (i) Let the positions of the  $-$  symbols in  $\tau_\gamma$  be  $i_1, \dots, i_q$  and the positions of the  $+$  symbols be  $j_1, \dots, j_p$  from left to right,
- (ii) For each pair  $c_{i_k} = c_{j_l} \in \mathbb{N}$  in  $\gamma$ , we place a rook in the square with northeast corner  $(k, l)$ .


## From sects to rooks

Let  $R(\lambda)$  denote the set of **rook placements** of partition  $\lambda$ . For  $\rho, \pi$  in  $R(\lambda)$ , we say  $\rho \leq \pi \iff rt_\rho \leq rt_\pi$  where  $R(\lambda) \xrightarrow{rt} \mathbb{N}$  is a labeling of the boxes of  $[\lambda]$  by the number of rooks weakly NW of each box.

### Bingham '21

There is an isomorphism between  $R(\lambda)$  and  $\mathcal{C}_{p,q}^\lambda$  as follows.

- (i) Let the positions of the  $-$  symbols in  $\tau_\gamma$  be  $i_1, \dots, i_q$  and the positions of the  $+$  symbols be  $j_1, \dots, j_p$  from left to right,
- (ii) For each pair  $c_{i_k} = c_{j_l} \in \mathbb{N}$  in  $\gamma$ , we place a rook in the square with northeast corner  $(k, l)$ .

$\gamma = 1+-221 \rightsquigarrow \tau_\gamma = -+---++$ . The labels of  $\tau_\gamma$  are  $c_{i_1} c_{j_1} c_{i_2} c_{i_3} c_{j_2} c_{j_3}$ . As  $c_{i_1} = c_{j_3}$ , we have a  in the box with NE corner  $(1, 3)$  and so on.

## Partial permutations

The **partial permutation** associated to a clan  $\gamma \in \mathcal{C}_{p,q}$  is the function  $\phi_\gamma : [q] \rightarrow [p] \cup \{0\}$  defined by labeling the positions of the  $-$  symbols in  $\tau_\gamma$  by  $i_1, \dots, i_q$  and the positions of the  $+$  symbols as  $j_1, \dots, j_p$  in *ascending* order.

## Partial permutations

The **partial permutation** associated to a clan  $\gamma \in \mathcal{C}_{p,q}$  is the function  $\phi_\gamma : [q] \rightarrow [p] \cup \{0\}$  defined by labeling the positions of the  $-$  symbols in  $\tau_\gamma$  by  $i_1, \dots, i_q$  and the positions of the  $+$  symbols as  $j_1, \dots, j_p$  in *ascending* order. Then we read the symbols  $c_1 \cdots c_{p+q}$  left to right and construct  $\phi_\gamma$  as

## Partial permutations

The **partial permutation** associated to a clan  $\gamma \in \mathcal{C}_{p,q}$  is the function  $\phi_\gamma : [q] \rightarrow [p] \cup \{0\}$  defined by labeling the positions of the  $-$  symbols in  $\tau_\gamma$  by  $i_1, \dots, i_q$  and the positions of the  $+$  symbols as  $j_1, \dots, j_p$  in *ascending* order. Then we read the symbols  $c_1 \cdots c_{p+q}$  left to right and construct  $\phi_\gamma$  as

- (1) If  $c_{i_s} = c_{j_t} \in \mathbb{N}$  in  $\gamma$ , then  $\phi_\gamma(s) = t$ . These are the rooks that are placed within the associated partition  $\lambda$  under (B.'21)-iso.

## Partial permutations

The **partial permutation** associated to a clan  $\gamma \in \mathcal{C}_{p,q}$  is the function  $\phi_\gamma : [q] \rightarrow [p] \cup \{0\}$  defined by labeling the positions of the  $-$  symbols in  $\tau_\gamma$  by  $i_1, \dots, i_q$  and the positions of the  $+$  symbols as  $j_1, \dots, j_p$  in *ascending* order. Then we read the symbols  $c_1 \cdots c_{p+q}$  left to right and construct  $\phi_\gamma$  as

- (1) If  $c_{i_s} = c_{j_t} \in \mathbb{N}$  in  $\gamma$ , then  $\phi_\gamma(s) = t$ . These are the rooks that are placed within the associated partition  $\lambda$  under (B.'21)-iso.
- (2) Modify  $\gamma$  by iteratively replacing all 1212 patterns by 1221 patterns to obtain a clan which we call  $\hat{\gamma}_0 \in \mathcal{C}_{p,q}^\lambda$  and which has symbols  $\hat{c}_1 \cdots \hat{c}_{p+q}$ .

## Partial permutations

The **partial permutation** associated to a clan  $\gamma \in \mathcal{C}_{p,q}$  is the function  $\phi_\gamma : [q] \rightarrow [p] \cup \{0\}$  defined by labeling the positions of the  $-$  symbols in  $\tau_\gamma$  by  $i_1, \dots, i_q$  and the positions of the  $+$  symbols as  $j_1, \dots, j_p$  in *ascending* order. Then we read the symbols  $c_1 \cdots c_{p+q}$  left to right and construct  $\phi_\gamma$  as

- (1) If  $c_{i_s} = c_{j_t} \in \mathbb{N}$  in  $\gamma$ , then  $\phi_\gamma(s) = t$ . These are the rooks that are placed within the associated partition  $\lambda$  under (B.'21)-iso.
- (2) Modify  $\gamma$  by iteratively replacing all 1212 patterns by 1221 patterns to obtain a clan which we call  $\hat{\gamma}_0 \in \mathcal{C}_{p,q}^\lambda$  and which has symbols  $\hat{c}_1 \cdots \hat{c}_{p+q}$ .
- (3) For each  $1+-1$  pattern in  $\hat{\gamma}_0$  of the form  $\hat{c}_a \hat{c}_{j_l} \hat{c}_{i_k} \hat{c}_b$ , set  $\phi_\gamma(k) = l$ . Delete all of the symbols involved in any  $1+-1$  pattern to obtain a new clan  $\hat{\gamma}_1$  which inherits position labels from  $\hat{\gamma}_0$ .


## Partial permutations

The **partial permutation** associated to a clan  $\gamma \in \mathcal{C}_{p,q}$  is the function  $\phi_\gamma : [q] \rightarrow [p] \cup \{0\}$  defined by labeling the positions of the  $-$  symbols in  $\tau_\gamma$  by  $i_1, \dots, i_q$  and the positions of the  $+$  symbols as  $j_1, \dots, j_p$  in *ascending* order. Then we read the symbols  $c_1 \cdots c_{p+q}$  left to right and construct  $\phi_\gamma$  as

- (1) If  $c_{i_s} = c_{j_t} \in \mathbb{N}$  in  $\gamma$ , then  $\phi_\gamma(s) = t$ . These are the rooks that are placed within the associated partition  $\lambda$  under (B.'21)-iso.
- (2) Modify  $\gamma$  by iteratively replacing all 1212 patterns by 1221 patterns to obtain a clan which we call  $\hat{\gamma}_0 \in \mathcal{C}_{p,q}^\lambda$  and which has symbols  $\hat{c}_1 \cdots \hat{c}_{p+q}$ .
- (3) For each  $1+-1$  pattern in  $\hat{\gamma}_0$  of the form  $\hat{c}_a \hat{c}_{j_l} \hat{c}_{i_k} \hat{c}_b$ , set  $\phi_\gamma(k) = l$ . Delete all of the symbols involved in any  $1+-1$  pattern to obtain a new clan  $\hat{\gamma}_1$  which inherits position labels from  $\hat{\gamma}_0$ .
- (4) Repeat the procedure of the previous step on  $\hat{\gamma}_1$ , and so on until we obtain a clan  $\hat{\gamma}_s$  which is free of  $1+-1$  patterns.



## Example: *hidden* rooks



The hidden rook associated to the pattern  $2+-2 = \hat{c}_{i_2} \hat{c}_{j_2} \hat{c}_{i_5} \hat{c}_{j_5}$  is obtained from  $\hat{\gamma}_1$ , after changing the 1212 pattern to 1221 and deleting the symbols  $3+-3 = \hat{c}_{i_3} \hat{c}_{j_3} \hat{c}_{i_4} \hat{c}_{j_4}$ .

$$\gamma = 12+3++-3-1-2 \mapsto \phi_\gamma = (5, 6, 4, 3, 2, 1).$$

## Coverings on $\mathcal{C}_{p,q}^\lambda$ (à la Wyser '16)

Lemma (Bingham-D, '25)

Let  $\gamma = c_1 \cdots c_{p,q} \in \mathcal{C}_{p,q}^\lambda$  with partial permutation  $\phi_\gamma = (a_1, \dots, a_q)$ .

# Coverings on $\mathcal{C}_{p,q}^\lambda$ (à la Wyser '16)

## Lemma (Bingham-D, '25)

Let  $\gamma = c_1 \cdots c_{p,q} \in \mathcal{C}_{p,q}^\lambda$  with partial permutation  $\phi_\gamma = (a_1, \dots, a_q)$ .

- (1) An **ff-rise** move applied to  $(c_{i_k}, c_{j_l})$  changes the symbol  $a_k$  from 0 to  $l$ .

# Coverings on $\mathcal{C}_{p,q}^\lambda$ (à la Wyser '16)

## Lemma (Bingham-D, '25)

Let  $\gamma = c_1 \cdots c_{p,q} \in \mathcal{C}_{p,q}^\lambda$  with partial permutation  $\phi_\gamma = (a_1, \dots, a_q)$ .

- (1) An **ff-rise** move applied to  $(c_{i_k}, c_{j_l})$  changes the symbol  $a_k$  from 0 to  $l$ .
- (2) An **fe-rise** move applied to  $(c_{i_k}, c_{i_m})$  where  $c_{i_m} = c_{j_l} \in \mathbb{N}$  swaps the symbol pair  $(a_k, a_m)$  from  $(0, l)$  to  $(l, 0)$ .

# Coverings on $\mathcal{C}_{p,q}^\lambda$ (à la Wyser '16)

## Lemma (Bingham-D, '25)

Let  $\gamma = c_1 \cdots c_{p,q} \in \mathcal{C}_{p,q}^\lambda$  with partial permutation  $\phi_\gamma = (a_1, \dots, a_q)$ .

- (1) An **ff-rise** move applied to  $(c_{i_k}, c_{j_l})$  changes the symbol  $a_k$  from 0 to  $l$ .
- (2) An **fe-rise** move applied to  $(c_{i_k}, c_{i_m})$  where  $c_{i_m} = c_{j_l} \in \mathbb{N}$  swaps the symbol pair  $(a_k, a_m)$  from  $(0, l)$  to  $(l, 0)$ .
- (3) An **ef-rise** move applied to  $(c_{i_k}, c_{j_m})$  where  $c_{i_k} = c_{j_l} \in \mathbb{N}$  changes the symbol  $a_k$  from  $l$  to  $m$ .

# Coverings on $\mathcal{C}_{p,q}^\lambda$ (à la Wyser '16)

## Lemma (Bingham-D, '25)

Let  $\gamma = c_1 \cdots c_{p,q} \in \mathcal{C}_{p,q}^\lambda$  with partial permutation  $\phi_\gamma = (a_1, \dots, a_q)$ .

- (1) An **ff-rise** move applied to  $(c_{i_k}, c_{j_l})$  changes the symbol  $a_k$  from 0 to  $l$ .
- (2) An **fe-rise** move applied to  $(c_{i_k}, c_{i_m})$  where  $c_{i_m} = c_{j_l} \in \mathbb{N}$  swaps the symbol pair  $(a_k, a_m)$  from  $(0, l)$  to  $(l, 0)$ .
- (3) An **ef-rise** move applied to  $(c_{i_k}, c_{j_m})$  where  $c_{i_k} = c_{j_l} \in \mathbb{N}$  changes the symbol  $a_k$  from  $l$  to  $m$ .
- (4) A **non-crossing ee-rise** move applied to  $(c_{i_k}, c_{i_m})$  where  $c_{i_k} = c_{j_l} \in \mathbb{N}$  and  $c_{i_m} = c_{j_n} \in \mathbb{N}$  swaps the symbol pair  $(a_k, a_m)$  from  $(l, n)$  to  $(n, l)$ .

# Coverings on $\mathcal{C}_{p,q}^\lambda$ (à la Wyser '16)

## Lemma (Bingham-D, '25)

Let  $\gamma = c_1 \cdots c_{p,q} \in \mathcal{C}_{p,q}^\lambda$  with partial permutation  $\phi_\gamma = (a_1, \dots, a_q)$ .

- (1) An **ff-rise** move applied to  $(c_{i_k}, c_{j_l})$  changes the symbol  $a_k$  from 0 to  $l$ .
- (2) An **fe-rise** move applied to  $(c_{i_k}, c_{i_m})$  where  $c_{i_m} = c_{j_l} \in \mathbb{N}$  swaps the symbol pair  $(a_k, a_m)$  from  $(0, l)$  to  $(l, 0)$ .
- (3) An **ef-rise** move applied to  $(c_{i_k}, c_{j_m})$  where  $c_{i_k} = c_{j_l} \in \mathbb{N}$  changes the symbol  $a_k$  from  $l$  to  $m$ .
- (4) A **non-crossing ee-rise** move applied to  $(c_{i_k}, c_{i_m})$  where  $c_{i_k} = c_{j_l} \in \mathbb{N}$  and  $c_{i_m} = c_{j_n} \in \mathbb{N}$  swaps the symbol pair  $(a_k, a_m)$  from  $(l, n)$  to  $(n, l)$ .
- (5) A **crossing ee-rise** move applied to  $(c_{i_k}, c_{i_m})$  where  $c_{i_k} = c_{j_l} \in \mathbb{N}$  and  $c_{i_m} = c_{j_n} \in \mathbb{N}$  swaps the symbol pair  $(a_k, a_m)$  from  $(l, n)$  to  $(n, l)$ .

## Labelling on $\mathcal{C}_{p,q}^\lambda$

### Definition (Bingham-D, '25)

Suppose that  $\gamma \triangleleft \tau$  is a covering relation where  $\gamma$  and the covering moves are as before. Then we label the covering relation  $(\gamma, \tau)$  in  $\mathcal{C}(\mathcal{C}_{p,q}^\lambda)$  with an element of  $\mathbb{N}^2$  as follows.

- (1) If  $\tau$  is obtained from an  $ff$ -rise move, then apply the label  $(0, l)$ .
- (2) If  $\tau$  is obtained from an  $fe$ -rise move, then apply the label  $(0, l)$ .
- (3) If  $\tau$  is obtained from an  $ef$ -rise move, then apply the label  $(l, m)$ .
- (4) If  $\tau$  is obtained from a non-crossing  $ee$ -rise move, then apply the label  $(l, n)$ .
- (5) If  $\tau$  is obtained from a crossing  $ee$ -rise move, then apply the label  $(l, n)$ .



## Labelling on $\mathcal{C}_{p,q}^\lambda$

### Definition (Bingham-D, '25)


Suppose that  $\gamma \triangleleft \tau$  is a covering relation where  $\gamma$  and the covering moves are as before. Then we label the covering relation  $(\gamma, \tau)$  in  $\mathcal{C}(\mathcal{C}_{p,q}^\lambda)$  with an element of  $\mathbb{N}^2$  as follows.

- (1) If  $\tau$  is obtained from an *ff*-rise move, then apply the label  $(0, l)$ .
- (2) If  $\tau$  is obtained from an *fe*-rise move, then apply the label  $(0, l)$ .
- (3) If  $\tau$  is obtained from an *ef*-rise move, then apply the label  $(l, m)$ .
- (4) If  $\tau$  is obtained from a non-crossing *ee*-rise move, then apply the label  $(l, n)$ .
- (5) If  $\tau$  is obtained from a crossing *ee*-rise move, then apply the label  $(l, n)$ .

### Theorem (Bingham-D' 25)

The Bruhat order on  $GL_{p+q}/GL_p \times GL_q$  restricted to any sect  $\mathcal{C}_{p,q}^\lambda$  is an *EL*-shellable poset.

$C_{3,3}^{(3,2,1)}$



## Open questions...

- (1) How could we conclude that  $\mathcal{C}_{p,q}$  is shellable?
- (2) What about the Möbius function on  $\mathcal{C}_{p,q}^\lambda$ ?
- (2) Other (sects of) symmetric spaces (of Hermitian type)?

# Open questions...

- (1) How could we conclude that  $\mathcal{C}_{p,q}$  is shellable?
- (2) What about the Möbius function function on  $\mathcal{C}_{p,q}^\lambda$ ?
- (2) Other (sects of) symmetric spaces (of Hermitian type)?

*Thank You/Gracias/Obrigado 😊*



*"If you spend your time chasing butterflies, they will just fly away.  
But if you build a beautiful garden, the butterflies will come" Mario  
Quintana.*