

Staircase diagrams, pattern avoidance, and smooth Schubert varieties

Faqrudin Ali Azam and Edward Richmond*

Lewis-Clark State College and Oklahoma State University*

October 18, 2025

Let $n \in \mathbb{Z}_+$ and $\mathbb{C}^n = \text{Span}_{\mathbb{C}}\{e_1, \dots, e_n\}$.

Flag variety:

$$\text{Fl}(n) := \{V_{\bullet} = (V_1 \subset \dots \subset V_{n-1} \subset \mathbb{C}^n) \mid \dim(V_i) = i\}$$

For any permutation $w \in S_n$, define the coordinate flag $E_{\bullet}^w \in \text{Fl}(n)$ by

$$E_i^w := \text{Span}\{e_{w(1)}, \dots, e_{w(i)}\}.$$

Let $n \in \mathbb{Z}_+$ and $\mathbb{C}^n = \text{Span}_{\mathbb{C}}\{e_1, \dots, e_n\}$.

Flag variety:

$$\text{Fl}(n) := \{V_{\bullet} = (V_1 \subset \dots \subset V_{n-1} \subset \mathbb{C}^n) \mid \dim(V_i) = i\}$$

For any permutation $w \in S_n$, define the coordinate flag $E_{\bullet}^w \in \text{Fl}(n)$ by

$$E_i^w := \text{Span}\{e_{w(1)}, \dots, e_{w(i)}\}.$$

Bruhat decomposition:

Let B denote $n \times n$ invertible upper triangular matrices. Then

$$\text{Fl}(n) = \bigsqcup_{w \in S_n} B \cdot E_{\bullet}^w$$

Schubert variety:

$$X(w) := \overline{B \cdot E_{\bullet}^w}.$$


Remark: The geometry of Schubert varieties plays an important role in combinatorics and representation theory.

Question: For which $w \in S_n$ is $X(w)$ smooth?

Question: For which $w \in S_n$ is $X(w)$ smooth?

Pattern avoidance: Let $m \leq n$. We say a permutation $w = w(1) \cdots w(n)$ *contains* the pattern $u = u(1) \cdots u(m)$ if there is a subsequence of w with the same relative order as u . Otherwise, w *avoids* the pattern u .

Example: $w = 416253 =$



contains the pattern 3412, but avoids the pattern 1234.

Conventions:

- The matrix entries of w mark the points $(i, w(i))$ (col \rightarrow , row \downarrow).
- $(1, 1)$ represents the NW corner of the matrix.

Theorem (Lakshmibai-Sandhya 1990, Carrell 1994, Ryan 1987):

Let $w \in S_n$. The following are equivalent:

- 1 $X(w)$ is smooth.
- 2 w avoids 3412 and 4231.
- 3 The Bruhat interval $[e, w]$ is rank symmetric.
- 4 $X(w)$ is an iterated fiber-bundle of Grassmannian varieties.

Theorem (Lakshmibai-Sandhya 1990, Carrell 1994, Ryan 1987):

Let $w \in S_n$. The following are equivalent:

- ① $X(w)$ is smooth.
- ② w avoids 3412 and 4231.
- ③ The Bruhat interval $[e, w]$ is rank symmetric.
- ④ $X(w)$ is an iterated fiber-bundle of Grassmannian varieties.

Enumeration (Haiman preprint, Bousquet-Mélou-Butler 2007):

Let

$$a_n := \#\{w \in S_n \mid X(w) \text{ is smooth}\}.$$

Then

$$\sum_{n \geq 0} a_n x^n = \frac{1 - 5x + 3x^2 + x^2 \sqrt{1 - 4x}}{1 - 6x + 8x^2 - 4x^3}.$$

Alternate encoding of smooth permutations:

Let $[n-1] := \{1, 2, \dots, n-1\}$ and

$$\mathcal{D} = \{B_1, \dots, B_k\}$$

be a set of intervals in $[n-1]$ where if $B_i = [l_i, r_i]$, then

$$l_i < l_{i+1} \text{ and } r_i < r_{i+1} \text{ for all } i.$$

Alternate encoding of smooth permutations:

Let $[n-1] := \{1, 2, \dots, n-1\}$ and

$$\mathcal{D} = \{B_1, \dots, B_k\}$$


be a set of intervals in $[n-1]$ where if $B_i = [l_i, r_i]$, then

$$l_i < l_{i+1} \text{ and } r_i < r_{i+1} \text{ for all } i.$$



Definition: We say a partial order \preceq on \mathcal{D} is a *staircase diagram* on $[n-1]$ if the following conditions hold:

- 1 B_i covers B_j if and only if $j = i \pm 1$ and $B_i \cup B_j$ is connected.
- 2 If $B_i \preceq B_{i-1}, B_{i+1}$ or $B_i \succeq B_{i-1}, B_{i+1}$, then $B_{i-1} \cup B_{i+1}$ is disconnected.


Example: Let $n = 12$ and $\mathcal{D} = \{B_1, \dots, B_6\}$:



Other staircase diagrams on $\mathcal{D} = \{B_1, \dots, B_6\}$:



Other staircase diagrams on $\mathcal{D} = \{B_1, \dots, B_6\}$:




Permutation of a staircase diagram: Note that

$$S_n = \langle s_1, \dots, s_{n-1} \rangle \text{ where } s_i := (i, i+1).$$

For any $J \subseteq [n-1]$, let u_J denote the *longest permutation* generated by $\{s_i \mid i \in J\}$. For $B \in \mathcal{D}$ define

$$J(B) := B \cap \left(\bigcup_{B' \prec B} B' \right).$$

Other staircase diagrams on $\mathcal{D} = \{B_1, \dots, B_6\}$:



Permutation of a staircase diagram: Note that

$$S_n = \langle s_1, \dots, s_{n-1} \rangle \text{ where } s_i := (i, i+1).$$

For any $J \subseteq [n-1]$, let u_J denote the *longest permutation* generated by $\{s_i \mid i \in J\}$. For $B \in \mathcal{D}$ define

$$J(B) := B \cap \left(\bigcup_{B' \prec B} B' \right).$$

Define $\Lambda(\emptyset) := \text{id}$. Otherwise, let \mathcal{D} be a staircase diagram and choose a maximal element $B \in \mathcal{D}$. Define

$$\Lambda(\mathcal{D}) := u_B \cdot u_{J(B)} \cdot \Lambda(\mathcal{D} \setminus \{B\}).$$

Example:

$$\Lambda \left(\begin{array}{ccccccc} & & & 4 & 5 & 6 & 7 \\ & 2 & 3 & 4 & & & \\ 1 & 2 & 3 & & & 7 & 8 \end{array} \right) = u_{[4,7]} \cdot s_4 s_7 \cdot \Lambda \left(\begin{array}{cccc} & 2 & 3 & 4 \\ 1 & 2 & 3 & \\ & 7 & 8 & \end{array} \right)$$

Example:

$$\Lambda \left(\begin{array}{ccccccc} & & & 4 & 5 & 6 & 7 \\ & 2 & 3 & 4 & & & \\ 1 & 2 & 3 & & & 7 & 8 \end{array} \right) = u_{[4,7]} \cdot s_4 s_7 \cdot \Lambda \left(\begin{array}{cccc} & 2 & 3 & 4 \\ 1 & 2 & 3 & \\ & & 7 & 8 \end{array} \right)$$

Facts:

- $\Lambda(\mathcal{D})$ is well-defined.
- $u_B \cdot u_{J(B)}$ is maximal in $W_B \cap W^{J(B)}$.
- $\Lambda(\mathcal{D})^{-1} = \Lambda(\mathcal{D}^*)$ where \mathcal{D}^* denotes the dual poset.
- The projection map

$$X(\Lambda(\mathcal{D})) \twoheadrightarrow X^{J(B)}(u_B)$$

is a $X(\Lambda(\mathcal{D} \setminus \{B\}))$ -fiber bundle.

Example:

$$\Lambda \left(\begin{array}{ccccccc} & & & 4 & 5 & 6 & 7 \\ & 2 & 3 & 4 & & & \\ 1 & 2 & 3 & & & & \\ & & & 7 & 8 & & \end{array} \right) = u_{[4,7]} \cdot s_4 s_7 \cdot \Lambda \left(\begin{array}{cccc} & 2 & 3 & 4 \\ 1 & 2 & 3 & \\ & & & 7 & 8 \end{array} \right)$$

Facts:

- $\Lambda(\mathcal{D})$ is well-defined.
- $u_B \cdot u_{J(B)}$ is maximal in $W_B \cap W^{J(B)}$.
- $\Lambda(\mathcal{D})^{-1} = \Lambda(\mathcal{D}^*)$ where \mathcal{D}^* denotes the dual poset.
- The projection map

$$X(\Lambda(\mathcal{D})) \twoheadrightarrow X^{J(B)}(u_B)$$

is a $X(\Lambda(\mathcal{D} \setminus \{B\}))$ -fiber bundle.

Theorem (R-Slofstra 2017): The map Λ is a bijection:

$$\{\text{Staircase diagrams on } [n-1]\} \xrightarrow{\Lambda} \{w \in S_n \mid X(w) \text{ is smooth}\}$$

Remarks (R-Slofstra 2016, 2017, 2018):

- Versions of staircase diagrams exist for flag varieties in all finite types and affine type A.
- Haiman's enumeration (g.f.) of (rationally) smooth Schubert varieties generalizes to other types.

Remarks (R-Slofstra 2016, 2017, 2018):

- Versions of staircase diagrams exist for flag varieties in all finite types and affine type A.
- Haiman's enumeration (g.f.) of (rationally) smooth Schubert varieties generalizes to other types.

Theorem (Tenner 2007, Karuppuchamy 2013): Let $w \in S_n$. The following are equivalent (*Boolean permutations*):

- 1 $X(w)$ is a toric manifold.
- 2 w avoids 3412 and 321. (hence also avoids 4231)
- 3 $X(w)$ is an iterated fiber-bundle of \mathbb{CP}_1 's.

Remarks (R-Slofstra 2016, 2017, 2018):

- Versions of staircase diagrams exist for flag varieties in all finite types and affine type A.
- Haiman's enumeration (g.f.) of (rationally) smooth Schubert varieties generalizes to other types.

Theorem (Tenner 2007, Karuppuchamy 2013): Let $w \in S_n$. The following are equivalent (*Boolean permutations*):

- 1 $X(w)$ is a toric manifold.
- 2 w avoids 3412 and 321. (hence also avoids 4231)
- 3 $X(w)$ is an iterated fiber-bundle of \mathbb{CP}_1 's.

GF:
$$\sum_{n \geq 0} a_n x^n = \frac{1 - 2x}{1 - 3x + x^2}$$

Formula: $a_n = f_{2n-1}$ (Fibonacci)

Remarks (R-Slofstra 2016, 2017, 2018):

- Versions of staircase diagrams exist for flag varieties in all finite types and affine type A.
- Haiman's enumeration (g.f.) of (rationally) smooth Schubert varieties generalizes to other types.

Theorem (Tenner 2007, Karuppuchamy 2013): Let $w \in S_n$. The following are equivalent (*Boolean permutations*):

- 1 $X(w)$ is a toric manifold.
- 2 w avoids 3412 and 321. (hence also avoids 4231)
- 3 $X(w)$ is an iterated fiber-bundle of \mathbb{CP}_1 's.

GF:
$$\sum_{n \geq 0} a_n x^n = \frac{1 - 2x}{1 - 3x + x^2}$$
 Formula: $a_n = f_{2n-1}$ (Fibonacci)

Question: What are staircase diagrams for Boolean permutations?


Staircase diagram of a Boolean permutation:

$$\Lambda \left(\begin{array}{ccccccccc} & \text{2} & & & & & & & \\ \text{1} & & \text{3} & & \text{5} & & \text{7} & & \\ & & & \text{4} & & & & & \text{8} \end{array} \right) = (s_2)(s_1 s_3 s_5 s_7)(s_4 s_8) = 314625897.$$

Staircase diagram of a Boolean permutation:

$$\Lambda \left(\begin{array}{ccccccc} & \text{2} & & & & & \\ \text{1} & & \text{3} & & \text{5} & & \text{7} \\ & & & \text{4} & & & \\ & & & & & & \text{8} \end{array} \right) = (s_2)(s_1 s_3 s_5 s_7)(s_4 s_8) = 314625897.$$


Compare statements:

- A smooth permutation is Boolean if and only if it avoids 321.
- A staircase diagram is Boolean if and only if it avoids .


Staircase diagram of a Boolean permutation:

$$\Lambda \left(\begin{array}{ccccccc} & \text{2} & & & & & \\ \text{1} & & \text{3} & & \text{5} & & \text{7} \\ & & & \text{4} & & & \\ & & & & & & \text{8} \end{array} \right) = (s_2)(s_1 s_3 s_5 s_7)(s_4 s_8) = 314625897.$$

Compare statements:

- A smooth permutation is Boolean if and only if it avoids 321.
- A staircase diagram is Boolean if and only if it avoids .

TFAE:


- 1 The staircase diagram \mathcal{D} avoids .
- 2 The permutation $\Lambda(\mathcal{D})$ avoids 321.

Moreover $\Lambda \left(\begin{array}{cc} & \\ \text{1} & \text{2} \end{array} \right) = \text{"321"}$


Staircase diagram of a Boolean permutation:

$$\Lambda \left(\begin{array}{ccccccc} & \text{2} & & & & & \\ \text{1} & & \text{3} & & \text{5} & & \text{7} \\ & & & \text{4} & & & \\ & & & & & & \text{8} \end{array} \right) = (s_2)(s_1 s_3 s_5 s_7)(s_4 s_8) = 314625897.$$

Compare statements:

- A smooth permutation is Boolean if and only if it avoids 321.
- A staircase diagram is Boolean if and only if it avoids .



TFAE:

- 1 The staircase diagram \mathcal{D} avoids .
- 2 The permutation $\Lambda(\mathcal{D})$ avoids 321.


Moreover $\Lambda \left(\begin{array}{cc} & \\ \text{1} & \text{2} \end{array} \right) = \text{"321"}$

Question: Does this idea extend to other staircase shapes and patterns?


Staircase overlaps:



Staircase overlaps:






Theorem (Azam-R. 2025): The following are equivalent:

- 1 The staircase diagram \mathcal{D} avoids .

- 2 The permutation $\Lambda(\mathcal{D})$ avoids

$$(k+2)(k+3)(k+1)k \cdots 21 \quad \text{and} \quad (k+3)(k+2) \cdots 4312.$$



Theorem (Azam-R. 2025): Let $k = 2$ (avoids ) and


$$a_n = \#\{w \in S_n \mid w \text{ avoids } 3412, 4231, 45321, 54312\}.$$


Then

$$\sum_{n \geq 0} a_n x^n = \frac{1 - 4x + x^2}{1 - 5x + 4x^2}.$$

Proof:

Staircase overlaps are at most one:



Theorem (Azam-R. 2025): Let $k = 2$ (avoids ) and


$$a_n = \#\{w \in S_n \mid w \text{ avoids } 3412, 4231, 45321, 54312\}.$$

Then

$$\sum_{n \geq 0} a_n x^n = \frac{1 - 4x + x^2}{1 - 5x + 4x^2}.$$

Proof:

Staircase overlaps are at most one:



Curious connection:



Let

$$c_n = \#\{w \in S_n \mid w \text{ avoids } 4123 \text{ and } 4321\}.$$



Corollary: $a_n = c_n$ for all $n \geq 0$.

Proof: Same generating function (OEIS).



Strongly connected chains:





Strongly connected chains:



More strongly connected chains:





Weakly connected chains:



Theorem (Azam-R. 2025): The following are equivalent:

- ① The staircase diagram \mathcal{D} avoids strongly connected chains of length ℓ .
- ② The permutation $\Lambda(\mathcal{D})$ avoids



$$34 \cdots (\ell+1)(\ell+2)21 \quad \text{and} \quad (\ell+2)(\ell+1)12 \cdots \ell.$$



Theorem (Azam-R. 2025): The following are equivalent:

- ① The staircase diagram \mathcal{D} avoids strongly connected chains of length ℓ .
- ② The permutation $\Lambda(\mathcal{D})$ avoids

$$34 \cdots (\ell+1)(\ell+2)21 \quad \text{and} \quad (\ell+2)(\ell+1)12 \cdots \ell.$$




Theorem (Azam-R. 2025): Let $\ell = 3$ and

$$b_n = \#\{w \in S_n \mid w \text{ avoids } 3412, 4231, 34521, 54123\}.$$

Then


$$\sum_{n \geq 0} b_n x^n = \frac{1 - 5x + 6x^2 - 4x^3}{1 - 6x + 10x^2 - 8x^3 + 2x^4}.$$

Proof: Strongly connected chain length at most two:




Theorem (Gaetz-Gao 2020): Let $w \in S_n$. The following are equivalent (*polished permutations*):

- ① The Bruhat interval $[e, w]$ is self-dual.
- ② w is smooth and avoids 45321, 54312, 34521, 54123.




Theorem (Gaetz-Gao 2020): Let $w \in S_n$. The following are equivalent (*polished permutations*):


- 1 The Bruhat interval $[e, w]$ is self-dual.
- 2 w is smooth and avoids 45321, 54312, 34521, 54123.



Observation: Polished permutations correspond to staircase diagrams such that:

- 1 Overlaps are at most one ($k = 2$).
- 2 Strongly connected chain length at most two ($\ell = 3$).





Theorem (Azam-R. 2025): Let

$$a_n = \#\{w \in S_n \mid w \text{ is polished}\}.$$

Then

$$\sum_{n \geq 0} a_n x^n = \frac{1 - 2x + x^2 + x^4}{1 - 2x + x^3}.$$

Thank you!