

# MATH2130 FYTT 1

Ankush Madharha

January 22, 2023

## 1 Intermediate Value Theorem

Show that  $f(x) = \cos(x) - \sqrt{x}$  has a root in the interval  $[0, 1]$ .

$f(x)$  is continuous on the interval  $[0, 1]$  because  $\cos(x)$  is continuous everywhere and  $\sqrt{x}$  is continuous on the interval  $[0, \infty)$ .

$$f(0) = \cos(0) - \sqrt{0} = 1 - 0 = 1$$

$$f(1) = \cos(1) - \sqrt{1} \approx 0.5403 - 1 \approx -0.5403$$

Since  $-0.5403 < 0 < 1$ , by the IVT there exists a  $c \in (0, 1)$  such that  $f(c) = 0$ .

## 2 Intermediate Value Theorem

Find an interval on which  $f(x) = x^4 - 3x^3 + x^2 + 3x - 7$  has a root.

We must find a  $x_1$  and a  $x_2$  such that  $f(x_1) < 0$  and  $f(x_2) > 0$ .

$$x = 0: f(0) = (0)^4 - 3(0)^3 + (0)^2 + 3(0) - 7 = -7 = x_1$$

$$x = 3: f(3) = (3)^4 - 3(3)^3 + (3)^2 + 3(3) - 7 = 11 = x_2$$

Since  $-7 < 0 < 11$ , by the IVT there exists a  $c \in (0, 3)$  such that  $f(c) = 0$ .

### 3 Mean Value Theorem

Verify the Mean Value Theorem for  $f(x) = \sqrt{2x-1}$  on the interval  $x \in [1, 5]$ .

$f(x)$  is continuous and differentiable on the interval  $[\frac{1}{2}, \infty)$ . The slope of the secant between  $(1, f(1))$  and  $(5, f(5))$  is given by:

$$\begin{aligned} & \frac{f(5)-f(1)}{5-1} \\ &= \frac{\sqrt{2(5)-1}-\sqrt{2(1)-1}}{4} \\ &= \frac{\sqrt{9}-\sqrt{1}}{4} \\ &= \frac{2}{4} \\ &= \frac{1}{2} \end{aligned}$$

The slope of the tangent of  $f(x)$  is given by its derivative.  $f'(x) = \frac{1}{2}(2x-1)^{-\frac{1}{2}}$ .  
 $2 = \frac{1}{\sqrt{2x-1}}$ .

$$\begin{aligned} & \frac{1}{\sqrt{2x-1}} = \frac{1}{2} \\ \Rightarrow & \sqrt{2x-1} = 2 \\ \Rightarrow & 2x-1 = 4 \\ \Rightarrow & 2x = 5 \\ \Rightarrow & x = 5/2 \end{aligned}$$

Since  $1 < 5/2 < 5$ , this is our  $c$  in the Mean Value Theorem.

## 4 Mean Value Theorem

Verify the Mean Value Theorem for  $f(x) = \frac{1}{x}$  on the interval  $x \in [2, 3]$ .

$f(x)$  is continuous and differentiable on the interval  $(0, \infty)$ . The slope of the secant between  $(2, f(2))$  and  $(3, f(3))$  is given by:

$$\begin{aligned}\frac{f(3)-f(2)}{3-2} \\&= \frac{\frac{1}{3}-\frac{1}{2}}{1} \\&= -\frac{1}{6}\end{aligned}$$

The slope of the tangent of  $f(x)$  is given by its derivative.  $f'(x) = -\frac{1}{x^2}$

$$-\frac{1}{x^2} = -\frac{1}{6}$$

$$\Rightarrow x^2 = 6$$

$$\Rightarrow x = \sqrt{6} \approx 2.45$$

Since  $2 < \sqrt{6} < 3$ , this is our  $c$  value in the Mean Value Theorem.

## 5 Taylor's Theorem

Some commonly-used Taylor expansions are as follows:

1. About  $x = 0$ ,  $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$
2. About  $x = 0$ ,  $\cos(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$
3. About  $x = 0$ ,  $\sin(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$
4. About  $x = 0$ ,  $\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n$
5. About  $x = 1$ ,  $\ln(x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{(x-1)^n}{n}$

It is useful to have these on-hand or to have the ability to quickly derive them as needed! Research or recall how to derive each of these series using basic calculus.

1)

General form of a Taylor series expansion:

$$f(x) = f(a) + \frac{f'(a)}{1!}(x-a)^1 + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \dots$$

Choose to expand the series around  $a = 0$ :

$$f(x) = f(0) + \frac{f'(0)}{1!}(x-0)^1 + \frac{f''(0)}{2!}(x-0)^2 + \frac{f'''(0)}{3!}(x-0)^3 + \dots$$

Simplify:

$$f(x) = f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots$$

All order derivatives are equal to  $e^x$ , and plugging 0 into  $f(x)$  and into all the derivatives are equal to 1:

$$f(x) = 1 + \frac{1}{1!}x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \dots$$

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

2)

General form of a Taylor series expansion:

$$f(x) = f(a) + \frac{f'(a)}{1!}(x-a)^1 + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \dots$$

Choose to expand the series around  $a = 0$ :

$$f(x) = f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots$$

Take the derivatives at  $x = 0$ :

$$f(x) = \cos(x) \Rightarrow f(0) = \cos(0) = 1$$

$$f'(x) = -\sin(x) \Rightarrow f'(0) = -\sin(0) = 0$$

$$f''(x) = -\cos(x) \Rightarrow f''(0) = -\cos(0) = -1$$

$$f'''(x) = \sin(x) \Rightarrow f'''(0) = \sin(0) = 0$$

$$f^{(4)}(x) = \cos(x) \Rightarrow f^{(4)}(0) = \cos(0) = 1$$

This results in a cycle of  $\{1, 0, -1, 0\}$  repeats itself. Resulting in the following series.

$$\cos(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

3)

The Taylor series expansion around for  $\sin(x)$  is the same as  $\cos(x)$  except we add one to the  $2n$  terms:

$$\cos(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

$$\sin(x) = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$

4)

General form of a Taylor series expansion around  $a = 0$ :

$$f(x) = f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots$$

Find derivatives and evaluate at  $x = 0$ :

$$f(0) = 1$$

$$f'(x) = (1-x)^{-2} \Rightarrow f'(0) = 1$$

$$f''(x) = 2(1-x)^{-3} \Rightarrow f''(0) = 1$$

$$f'''(x) = 6(1-x)^{-4} \Rightarrow f'''(0) = 2$$

$$f^{(4)}(x) = 24(1-x)^{-5} \Rightarrow f^{(4)}(0) = 24$$

Plug into the general form:

$$f(x) = 1 + x + \frac{2}{2!}x^2 + \frac{6}{3!}x^3 + \frac{24}{4!}x^4$$

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + \dots$$

$$\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n$$

5)

General form of a Taylor series expansion:

$$f(x) = f(a) + \frac{f'(a)}{1!}(x-a)^1 + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \dots$$

Find derivatives and evaluate at  $x = 0$ :

$$f(x) = \ln(x) \Rightarrow f(1) = \ln(1) = 0$$

$$f'(x) = \frac{1}{x} \Rightarrow f'(1) = \frac{1}{1} = 1$$

$$f''(x) = -\frac{1}{x^2} \Rightarrow f''(1) = -\frac{1}{(1)^2} = -1$$

$$f'''(x) = \frac{2}{x^3} \Rightarrow f'''(1) = \frac{2}{(1)^3} = 2$$

$$f^{(4)}(x) = -\frac{(2)(3)}{x^4} \Rightarrow f^{(4)}(1) = -\frac{(2)(3)}{(1)^4} = -(2)(3)$$

We can see a pattern forming, so we can plug in this information into our formula:

$$0 + 1(x-1) - \frac{1(x-1)^2}{2!} + \frac{2(x-1)^3}{3!} - \frac{(2)(3)(x-1)^4}{4!} \dots$$

Writing this as a series:

$$\ln(x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{((n-1)!(x-1)^n}{n!}$$

$$\Rightarrow \ln(x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{(x-1)^n}{n}$$

## 6 Taylor's Theorem

Let  $P_n(x)$  be the  $n^{th}$ -order Taylor Polynomials for  $f(x) = \sin(x)$  about  $x = 0$ . Find  $n$  so that  $P_n(x)$  is within  $10^{-5}$  of  $\sin(x)$  for all  $x \in [0, \frac{\pi}{2}]$ .

Using Taylor's Theorem about  $x = 0$ :

$$\begin{aligned} e^x &= P_n(x) + R_n(x) \\ &= x - \frac{x^3}{3!} + \frac{x^5}{5!} + \dots + (-1)^n \frac{x^{2n+1}}{(2n+1)!} + \frac{f^{n+1}(c)}{(n+1)!} x^{n+1} \end{aligned}$$

$f(x) = \sin(x)$ 's derivative is cyclical:

$$f'(x) = \cos(x)$$

$$f''(x) = -\sin(x)$$

$$f'''(x) = -\cos(x)$$

$$f^{(4)}(x) = \sin(x)$$

We are considering the values of  $x \in [0, \frac{\pi}{2}]$ , we must determine for which values of  $n$  we have that:

$$\frac{f^{n+1}(c)}{(n+1)!} x^{n+1} < 10^{-5}$$

On this interval, we only consider the  $n$  values such that  $n \% 4 = 0$ .

$$n = 3 : \frac{f^4(c)}{4!} x^4 = \frac{1}{4!} \left(\frac{\pi}{2}\right)^4 = 0.253669507 \not< 10^{-5}$$

$$n = 7 : \frac{f^8(c)}{8!} x^8 = \frac{1}{8!} \left(\frac{\pi}{2}\right)^8 = 9.192602748 \times 10^{-4} \not< 10^{-5}$$

$$n = 11 : \frac{f^{12}(c)}{12!} x^{12} = \frac{1}{12!} \left(\frac{\pi}{2}\right)^{12} = 4.710874779 \times 10^{-7} < 10^{-5}$$