

# Adequate set of connectives

Lila Kari

The University of Western Ontario

# Adequate set of connectives

- A remarkable property of the standard set of connectives ( $\neg, \wedge, \vee, \rightarrow, \leftrightarrow$ ) is the fact that for every truth table

$p$	$q$	$\dots$	
1	1	$\dots$	*
			*
			*
0	0	$\dots$	*

there is a formula (depending on the variables  $p, q, \dots$  and using only the standard connectives) that has exactly this truth table.

- For every truth table there is Boolean function  $f(p, q, \dots)$  with exactly this truth table.

# Post's observation

- Any set of connectives with the capability to express all truth tables is said to be **adequate**.
- As Post (1921) observed that **the standard connectives are adequate**.



Emil Post, 1897-1954

# Proving adequacy of a connective set

We can show that a set  $\mathcal{S}$  of connectives is adequate if we can express all the standard connectives in terms of  $\mathcal{S}$ .

# Proving adequacy of a connective set

We can show that a set  $\mathcal{S}$  of connectives is adequate if we can express all the standard connectives in terms of  $\mathcal{S}$ .

- Formulas  $A \rightarrow B$  and  $\neg A \vee B$  are tautologically equivalent.

# Proving adequacy of a connective set

We can show that a set  $\mathcal{S}$  of connectives is adequate if we can express all the standard connectives in terms of  $\mathcal{S}$ .

- Formulas  $A \rightarrow B$  and  $\neg A \vee B$  are tautologically equivalent.
- Then  $\rightarrow$  is definable in terms of (or is reducible to or can be expressed in terms of)  $\neg$  and  $\vee$ .

# Proving adequacy of a connective set

We can show that a set  $\mathcal{S}$  of connectives is adequate if we can express all the standard connectives in terms of  $\mathcal{S}$ .

- Formulas  $A \rightarrow B$  and  $\neg A \vee B$  are tautologically equivalent.
- Then  $\rightarrow$  is definable in terms of (or is reducible to or can be expressed in terms of)  $\neg$  and  $\vee$ .
- Similarly,  $\vee$  is definable in terms of  $\neg$  and  $\rightarrow$  because  $A \vee B$  is tautologically equivalent to  $\neg A \rightarrow B$ .

**Theorem.**  $\{\neg, \wedge, \vee\}$  is an adequate set of connectives.



**Theorem.**  $\{\neg, \wedge, \vee\}$  is an adequate set of connectives.

**Proof.**

For any formulas  $A, B$ ,

$$\neg A \models \neg A$$

$$A \wedge B \models A \wedge B$$

$$A \vee B \models A \vee B$$

$$A \rightarrow B \models \neg A \vee B$$

$$A \leftrightarrow B \models (A \rightarrow B) \wedge (B \rightarrow A).$$

**Theorem.**  $\{\neg, \wedge, \vee\}$  is an adequate set of connectives.

**Proof.**

For any formulas  $A, B$ ,

$$\neg A \models \neg A$$

$$A \wedge B \models A \wedge B$$

$$A \vee B \models A \vee B$$

$$A \rightarrow B \models \neg A \vee B$$

$$A \leftrightarrow B \models (A \rightarrow B) \wedge (B \rightarrow A).$$

All the five standard connectives can be expressed in terms of  $\neg, \wedge, \vee$ , therefore  $\{\neg, \wedge, \vee\}$  is an adequate set of connectives.

**Theorem.**  $\{\neg, \wedge, \vee\}$  is an adequate set of connectives.

**Proof.**

For any formulas  $A, B$ ,

$$\neg A \models \neg A$$

$$A \wedge B \models A \wedge B$$

$$A \vee B \models A \vee B$$

$$A \rightarrow B \models \neg A \vee B$$

$$A \leftrightarrow B \models (A \rightarrow B) \wedge (B \rightarrow A).$$

All the five standard connectives can be expressed in terms of  $\neg, \wedge, \vee$ , therefore  $\{\neg, \wedge, \vee\}$  is an adequate set of connectives.

**Corollary**  $\{\neg, \wedge\}$ ,  $\{\neg, \vee\}$ , and  $\{\neg, \rightarrow\}$  are adequate.

# One connective is enough

Schroder showed in 1880 that each of the standard connectives is definable in terms of a single binary connective  $\downarrow$ , where the truth table associated with  $\downarrow$  is

$p$	$q$	$p \downarrow q$
1	1	0
1	0	0
0	1	0
0	0	1

# Proof of adequacy

We can express  $\downarrow$  in terms of the standard connectives by  $p \downarrow q \models \neg p \wedge \neg q$ , and also the standard connectives in terms of  $\downarrow$  by

# Proof of adequacy

We can express  $\downarrow$  in terms of the standard connectives by  $p \downarrow q \models \neg p \wedge \neg q$ , and also the standard connectives in terms of  $\downarrow$  by

$$\neg p \models p \downarrow p$$

# Proof of adequacy

We can express  $\downarrow$  in terms of the standard connectives by  $p \downarrow q \models \neg p \wedge \neg q$ , and also the standard connectives in terms of  $\downarrow$  by

$$\begin{array}{lcl} \neg p & \models & p \downarrow p \\ p \wedge q & \models & (p \downarrow p) \downarrow (q \downarrow q) \end{array}$$

# Proof of adequacy

We can express  $\downarrow$  in terms of the standard connectives by  $p \downarrow q \models \neg p \wedge \neg q$ , and also the standard connectives in terms of  $\downarrow$  by

$$\begin{array}{lcl} \neg p & \models & p \downarrow p \\ p \wedge q & \models & (p \downarrow p) \downarrow (q \downarrow q) \\ p \vee q & \models & (p \downarrow q) \downarrow (p \downarrow q) \end{array}$$



# Proof of adequacy

We can express  $\downarrow$  in terms of the standard connectives by  $p \downarrow q \models \neg p \wedge \neg q$ , and also the standard connectives in terms of  $\downarrow$  by

$$\begin{array}{ll} \neg p & \models p \downarrow p \\ p \wedge q & \models (p \downarrow p) \downarrow (q \downarrow q) \\ p \vee q & \models (p \downarrow q) \downarrow (p \downarrow q) \\ p \rightarrow q & \models ((p \downarrow p) \downarrow q) \downarrow ((p \downarrow p) \downarrow q) \end{array}$$

# Proof of adequacy

We can express  $\downarrow$  in terms of the standard connectives by  $p \downarrow q \models \neg p \wedge \neg q$ , and also the standard connectives in terms of  $\downarrow$  by

$$\begin{array}{ll} \neg p & \models p \downarrow p \\ p \wedge q & \models (p \downarrow p) \downarrow (q \downarrow q) \\ p \vee q & \models (p \downarrow q) \downarrow (p \downarrow q) \\ p \rightarrow q & \models ((p \downarrow p) \downarrow q) \downarrow ((p \downarrow p) \downarrow q) \\ p \leftrightarrow q & \models ((p \downarrow p) \downarrow q) \downarrow ((q \downarrow p) \downarrow p). \end{array}$$

Thus it follows that a single connective  $\downarrow$  is adequate.

Consequently, to test a given  $\mathcal{S}$  for being adequate it suffices to test if  $\downarrow$  can be expressed by  $\mathcal{S}$ .

# Sheffer stroke

In 1913 Sheffer showed that the **Sheffer stroke** | with associated truth table

$p$	$q$	$p q$
1	1	0
1	0	1
0	1	1
0	0	1

is also a single binary connective in terms of which the standard connectives can be expressed.

# Proving inadequacy

How do we show that a given set  $\mathcal{S}$  of connectives is not adequate?

# Proving inadequacy

How do we show that a given set  $\mathcal{S}$  of connectives is not adequate?

Show that some standard connective cannot be expressed by  $\mathcal{S}$ .

# Proving inadequacy

How do we show that a given set  $\mathcal{S}$  of connectives is **not adequate**?

Show that **some standard connective cannot be expressed by  $\mathcal{S}$** .

**Example.** The set  $\mathcal{S} = \{\wedge\}$  is not adequate.

# Proving inadequacy

How do we show that a given set  $\mathcal{S}$  of connectives is **not adequate**?

Show that **some standard connective cannot be expressed by  $\mathcal{S}$** .

**Example.** The set  $\mathcal{S} = \{\wedge\}$  is not adequate.

**Proof.** To see this, note that a formula depending on only one variable and which uses only the connective  $\wedge$  has the property that its truth value for a value assignment that makes  $p = 0$  is always 0.

# Proving inadequacy

How do we show that a given set  $\mathcal{S}$  of connectives is **not adequate**?

Show that **some standard connective cannot be expressed by  $\mathcal{S}$** .

**Example.** The set  $\mathcal{S} = \{\wedge\}$  is not adequate.

**Proof.** To see this, note that a formula depending on only one variable and which uses only the connective  $\wedge$  has the property that its truth value for a value assignment that makes  $p = 0$  is always 0.

In order to define the negation  $\neg p$  in terms of  $\wedge$ , there should exist a formula  $f$  depending on the variable  $p$  and using only the connective  $\wedge$  such that  $\neg p \models f$ .



# Proving inadequacy

How do we show that a given set  $\mathcal{S}$  of connectives is **not adequate**?

Show that **some standard connective cannot be expressed by  $\mathcal{S}$** .

**Example.** The set  $\mathcal{S} = \{\wedge\}$  is not adequate.

**Proof.** To see this, note that a formula depending on only one variable and which uses only the connective  $\wedge$  has the property that its truth value for a value assignment that makes  $p = 0$  is always 0.

In order to define the negation  $\neg p$  in terms of  $\wedge$ , there should exist a formula  $f$  depending on the variable  $p$  and using only the connective  $\wedge$  such that  $\neg p \models f$ .

However, for a value assignment  $v$  such that  $v(\neg p) = 1$ , we have  $v(p) = 0$  and therefore  $v(f) = 0$ , which shows that  $\neg p$  and  $f$  cannot be tautologically equivalent.

# A ternary connective

Let us use the symbol  $\tau$  for the **ternary connective** whose truth table is given by

$p$	$q$	$r$	$\tau$
1	1	1	1
1	1	0	1
1	0	1	0
1	0	0	0
0	1	1	1
0	1	0	0
0	0	1	1
0	0	0	0

# A ternary connective

It is easy to see that for any value assignment  $v$  we have

$$v(\tau(p, q, r)) = v(q) \text{ if } v(p) = 1 \text{ and } v(r) \text{ if } v(p) = 0.$$

# A ternary connective

It is easy to see that for any value assignment  $v$  we have

$$v(\tau(p, q, r)) = v(q) \text{ if } v(p) = 1 \text{ and } v(r) \text{ if } v(p) = 0.$$

This is the familiar **if-then-else** connective from computer science, namely

**if p then q else r**

We can consider now propositional logic based not upon the five common connectives, but upon any adequate set of connectives, for instance  $\{\neg, \wedge\}$ .

Let  $\mathcal{L}_0^P$  be a sublanguage of  $\mathcal{L}^P$  obtained by deleting from  $\mathcal{L}^P$  the three connectives  $\vee, \rightarrow, \leftrightarrow$ , and let  $\text{Form}(\mathcal{L}_0^P)$  be the set of formulas of  $\mathcal{L}_0^P$ .

**Theorem.**  $\text{Form}(\mathcal{L}_0^P) = \text{Form}(\mathcal{L}^P)$ .

# Proof

## Proof.

Obviously,  $\text{Form}(\mathcal{L}_0^p) \subseteq \text{Form}(\mathcal{L}^p)$ .

Conversely, for  $A \in \text{Form}(\mathcal{L}^p)$  we define (by recursion) its translation  $A_0$  into  $\mathcal{L}_0^p$  as follows:

$A_0 = A$  for atomic variable  $A$ ,

$(\neg A)_0 = \neg A_0$ ,

$(A \wedge B)_0 = A_0 \wedge B_0$ ,

$(A \vee B)_0 = \neg(\neg A_0 \wedge \neg B_0)$ ,

$(A \rightarrow B)_0 = \neg(A_0 \wedge \neg B_0)$ ,

$(A \leftrightarrow B)_0 = (A \rightarrow B)_0 \wedge (B \rightarrow A)_0 =$   
 $= \neg(A_0 \wedge \neg B_0) \wedge \neg(\neg A_0 \wedge B_0)$