

Propositional Language - Syntax

Lila Kari

The University of Western Ontario

Propositional language

- By using connectives, we can combine propositions, whether they are atomic themselves or compound.
- To prevent ambiguity, we introduce **fully parenthesized expressions** that can be parsed in a unique way.
- When possible, we will omit parentheses and use precedence rules.
- We construct the propositional language \mathcal{L}^P , which is the formal language of the propositional logic.

Propositional language \mathcal{L}^p : syntax

Propositional language \mathcal{L}^p is the formal language for propositional logic.

It consists of:

- **propositional variables**: p, q, r .
- **propositional constants**: 1 and 0 (true and false).
- **logical connectives**: $\neg, \wedge, \vee, \rightarrow, \leftrightarrow$.

Oral reading of logical connectives

\neg	not	negation
\wedge	and	conjunction
\vee	or (and/or)	(inclusive) disjunction
\rightarrow	if, then (imply)	implication
\leftrightarrow	iff (equivalent to)	equivalence

- **left and right parantheses** : $(,)$.

Expressions of \mathcal{L}^p

- **expressions** = finite strings of symbols
 - $p, pq, (r), p \wedge \rightarrow q$ and $\neg(p \vee q)$ are expressions in \mathcal{L}^p .
- **length** of expression: the number of occurrences of symbols in it
 - for expressions above, length is 1, 2, 3, 4 and 6
- **empty expression** = expression of length zero, \emptyset
- two expressions u and v are **equal** iff they are of the same length and have the same symbols in the same order.
- scanning of expressions proceeds from left to right

Expressions

- **concatenating** two expressions u , v in this order is denoted uv .
Note that $\emptyset u = u\emptyset = u$
- if $u = w_1vw_2$ where u , v , w_1 , w_2 are expressions, then v is a **segment of** u .
- If $v \neq u$ then v is a **proper segment** of u .
- If $u = vw$, where u , v , w are expressions, then v is an **initial segment** of u . Similarly, w is a **terminal segment** of u .

Atoms and formulas

- An **atom (atomic formula)** is an expression consisting of a proposition variable or constant only.
- An expression in \mathcal{L}^P is called a **formula (well-formed formula)** iff it can be constructed according to the following **formation rules**:
 1. Every atomic formula is formula;
 2. If A is a formula then so is $(\neg A)$;
 3. If A, B are formulas then so are $(A \vee B)$, $(A \wedge B)$, $(A \rightarrow B)$ and $(A \leftrightarrow B)$.
 4. No other expression is a formula.

Definition. (Negation, Conjunction, Disjunction, Implication, Equivalence).

- † $(\neg A)$ is called a **negation** (formula). It is the negation of A .
- † $(A \wedge B)$ is called a **conjunction** (formula). It is the conjunction of A and B . A and B are called the **conjuncts** of $(A \wedge B)$.
- † $(A \vee B)$ is called a **disjunction** (formula). It is the disjunction of A and B . A and B are called the **disjuncts** of $(A \vee B)$.
- † $(A \rightarrow B)$ is called an **implication** (formula). It is the implication of A and B . A and B are called the antecedent and consequent of $(A \rightarrow B)$.
- † $(A \leftrightarrow B)$ is called an **equivalence** (formula). It is the equivalence of A and B .

Formulas

- The set of atomic formulas in \mathcal{L}^P is denoted by $\text{Atom}(\mathcal{L}^P)$.
- The set of all formulas (well-formed formulas, wff) in \mathcal{L}^P is denoted by $\text{Form}(\mathcal{L}^P)$.
- $\text{Form}(\mathcal{L}^P)$ is the smallest class of expressions of \mathcal{L}^P closed under the formation rules of formulas.

- Every formula of \mathcal{L}^P has the same number of occurrences of left and right parentheses.
- Any proper initial segment of a formula in \mathcal{L}^P has more occurrences of left than right parentheses. Any proper terminal segment of a formula in \mathcal{L}^P has less occurrences of left than right parentheses.
Thus neither a proper initial segment nor a proper terminal segment of a formula can itself be a formula of \mathcal{L}^P .
- Every formula of \mathcal{L}^P is of exactly one of the six forms: an atom, $(\neg A)$, $(A \wedge B)$, $(A \vee B)$, $(A \rightarrow B)$, or $(A \leftrightarrow B)$, and in each case it is of the form in exactly one way.

Scope

- If $(\neg A)$ is a segment of a formula C , then A is called the **scope** of the negation \neg in the formula C .
- If $(A \wedge B)$ is a segment of a formula C , then A is called **the left scope** of the conjunction and B is called the **right scope** of the conjunction in the formula C .
- Similar definitions hold for left/right scopes of disjunction, implication and equivalence.

Example

Suppose $A = (\neg((p \wedge q) \vee ((\neg p) \rightarrow r)))$.

Example

Suppose $A = (\neg((p \wedge q) \vee ((\neg p) \rightarrow r)))$.

The scope of the first \neg is $((p \wedge q) \vee ((\neg p) \rightarrow r))$ and of the second \neg is p .

Example

Suppose $A = (\neg((p \wedge q) \vee ((\neg p) \rightarrow r)))$.

The scope of the first \neg is $((p \wedge q) \vee ((\neg p) \rightarrow r))$ and of the second \neg is p .

The left and right scopes of \wedge are p and q .

Example

Suppose $A = (\neg((p \wedge q) \vee ((\neg p) \rightarrow r)))$.

The scope of the first \neg is $((p \wedge q) \vee ((\neg p) \rightarrow r))$ and of the second \neg is p .

The left and right scopes of \wedge are p and q .

The left and right scopes of \vee are $(p \wedge q)$ and $((\neg p) \rightarrow r)$.

Example

Suppose $A = (\neg((p \wedge q) \vee ((\neg p) \rightarrow r)))$.

The scope of the first \neg is $((p \wedge q) \vee ((\neg p) \rightarrow r))$ and of the second \neg is p .

The left and right scopes of \wedge are p and q .

The left and right scopes of \vee are $(p \wedge q)$ and $((\neg p) \rightarrow r)$.

The scopes of \rightarrow are $(\neg p)$ and r .

Example 1 (generating formulas)

Example 1 (generating formulas).

The expression

$$((p \vee q) \rightarrow ((\neg p) \leftrightarrow (q \wedge r)))$$

is a formula. How is it generated?

Example 2 (parsing formulas)

If Michelle wins at the Olympics, everyone will admire her, and she will get rich, but if she does not win, all her effort was in vain.

p : Michelle wins at the olympics.

q : Everyone admires Michelle.

r : Michelle will get rich.

s : Michelle's effort was in vain.

Example 2 (parsing formulas)

If Michelle wins at the Olympics, everyone will admire her, and she will get rich, but if she does not win, all her effort was in vain.

p : Michelle wins at the olympics.

q : Everyone admires Michelle.

r : Michelle will get rich.

s : Michelle's effort was in vain.

The compound proposition becomes

$$((p \rightarrow (q \wedge r)) \wedge ((\neg p) \rightarrow s))$$

One can use parse trees to analyze formulas.

Precedence rules

- \neg has precedence over \wedge
- \wedge has precedence over \vee
- \vee has precedence over \rightarrow
- \rightarrow has precedence over \leftrightarrow

Precedence rules

- \neg has precedence over \wedge
- \wedge has precedence over \vee
- \vee has precedence over \rightarrow
- \rightarrow has precedence over \leftrightarrow

Examples:

- $\neg p \vee q$ is to be understood as $(\neg p) \vee q$.
- $p \wedge q \vee r$ is to be understood as $((p \wedge q) \vee r)$.
- $p \rightarrow q \vee r$ is to be understood as $p \rightarrow (q \vee r)$
- $p \leftrightarrow p \rightarrow q$ must be understood as
 $p \leftrightarrow (p \rightarrow q)$
- proposition about Michelle:
 $(p \rightarrow q \wedge r) \wedge (\neg p \rightarrow s)$

Examples of compound propositions

- No work, no pay.

Examples of compound propositions

- No work, no pay.

Reword: “If no work is done, then there is no pay.”

p = “work was done”,

q = “there is pay”.

Examples of compound propositions

- No work, no pay.

Reword: “If no work is done, then there is no pay.”

p = “work was done”,

q = “there is pay”.

Translation into propositional language: $\neg p \rightarrow \neg q$.

Examples of compound propositions

- Goods bought into this store can be returned only if they are in good condition, and only if the purchaser brings a receipt.

Examples of compound propositions

- Goods bought into this store can be returned only if they are in good condition, and only if the purchaser brings a receipt.

Atomic propositions:

p = “goods can be returned”,

q = “goods are in good condition”,

r = “the purchaser must bring a receipt”.

Examples of compound propositions

- Goods bought into this store can be returned only if they are in good condition, and only if the purchaser brings a receipt.

Atomic propositions:

p = “goods can be returned”,

q = “goods are in good condition”,

r = “the purchaser must bring a receipt”.

Translation into propositional language:

$$p \rightarrow (q \wedge r)$$

- If p is a prime number then, for even integers n , $n^p - n$ is divisible by p .

- If p is a prime number then, for even integers n , $n^p - n$ is divisible by p .

Atomic propositions:

$p =$ “ p is a prime”,

$q =$ “ n is an integer”,

$r =$ “ n is even”,

$s =$ “ $n^p - n$ is divisible by p .”

- If p is a prime number then, for even integers n , $n^p - n$ is divisible by p .

Atomic propositions:

$p =$ “ p is a prime”,

$q =$ “ n is an integer”,

$r =$ “ n is even”,

$s =$ “ $n^p - n$ is divisible by p .”

Translation into propositional language:

$$p \rightarrow (q \wedge r \rightarrow s)$$

Examples of compound propositions

- Set C is the intersection of set A and set B if and only if every element of set C is an element of both A and B .

Examples of compound propositions

- Set C is the intersection of set A and set B if and only if every element of set C is an element of both A and B .

Atomic propositions:

p = "Set C is an intersection of set A and set B ",

q = "Every element of C is an element of A ",

r = "Every element of C is an element of B ."

Examples of compound propositions

- Set C is the intersection of set A and set B if and only if every element of set C is an element of both A and B .

Atomic propositions:

p = "Set C is an intersection of set A and set B ",

q = "Every element of C is an element of A ",

r = "Every element of C is an element of B ."

Translation into propositional language:

$$p \leftrightarrow q \wedge r$$

Examples of compound propositions

- Nobody is allowed to turn on freeway. Police officers on duty are exempt from this rule.

Examples of compound propositions

- Nobody is allowed to turn on freeway. Police officers on duty are exempt from this rule.

Atomic propositions:

p = "X is allowed to turn on a freeway,"

q = "X is a police officer",

r = "X is on duty."

Examples of compound propositions

- Nobody is allowed to turn on freeway. Police officers on duty are exempt from this rule.

Atomic propositions:

p = "X is allowed to turn on a freeway,"

q = "X is a police officer",

r = "X is on duty."

Translation into propositional language:

$$\neg p \vee (q \wedge r)$$

Converse and contrapositive

Definition. Given the proposition $p \rightarrow q$,

the **converse** of $p \rightarrow q$ is the proposition $q \rightarrow p$

the **contrapositive** of $p \rightarrow q$ is the proposition $\neg q \rightarrow \neg p$.

Induction on the complexity of formulas

Theorem. Suppose R is a property. If

- 1 For any atomic formula $p \in \text{Atom}(\mathcal{L}^p)$, p has the property R .
- 2 For any formula A in $\text{Form}(\mathcal{L}^p)$, if A has property R , then $(\neg A)$ has the property R .
- 3 For any formulas A and B in $\text{Form}(\mathcal{L}^p)$, if A and B have property R then $(A * B)$ has property R , where $*$ $\in \{\wedge, \vee, \rightarrow, \leftrightarrow\}$.

then, any formula in $\text{Form}(\mathcal{L}^p)$ has property R .