

# TINA: Supplementary Proofs and Numerical Details

Scott Moeller

Bhaskar Krishnamachari

September 2026

Supplement to *A Theory of Information Architecture for Networked Decisions: Freshness, Locality, and Coordination*, revised v2-b draft. Unqualified references to main-paper results and equations use that draft's numbering.

## S1 Aggregate-coupling calculations

This section supplies the matrix arithmetic for Proposition 1 in the main paper. For  $Q = qI + \gamma \mathbf{1}\mathbf{1}^\top$ ,

$$Q^{-1} = \frac{1}{q} \left( I - \frac{\gamma}{q + \gamma n} \mathbf{1}\mathbf{1}^\top \right).$$

*Proof.* Recall the coupled normal equations (24):

$$Q_{ii}u_i + \sum_{j \neq i} Q_{ij}\mathbb{E}[u_j \mid \mathcal{G}_i] = \mathbb{E}[b_i(\theta_t) \mid \mathcal{G}_i].$$

Here  $Q_{ii} = q + \gamma$ ,  $Q_{ij} = \gamma$  for  $j \neq i$ , and  $b_i(\theta) = \theta_i$ . Let  $\mu_j = \mathbb{E}[u_j]$ . For  $j \neq i$ , the policy  $u_j$  is a function of  $\theta_{j,t}$ , which is independent of  $\theta_{i,t}$ , so

$$\mathbb{E}[u_j \mid \mathcal{G}_{i,t}^{\text{loc}}] = \mathbb{E}[u_j] = \mu_j,$$

while  $\theta_{i,t}$  is  $\mathcal{G}_{i,t}^{\text{loc}}$ -measurable. The conditional normal equation (24) therefore gives

$$(q + \gamma)u_i + \gamma \sum_{j \neq i} \mu_j - \theta_{i,t} = 0.$$

Taking expectations yields

$$q\mu_i + \gamma \sum_j \mu_j = 0.$$

Summing over  $i$  implies  $\sum_j \mu_j = 0$ , hence every  $\mu_i = 0$  and  $u_i = \theta_{i,t}/(q + \gamma)$ , which proves (45). This argument uses independence, zero means, square integrability, and strict convexity; Gaussianity is not required for the local-policy formula.

*Open-loop regret.* Here  $B = I$  and  $\Sigma_\infty = \sigma^2 I$ , so by (39),  $R_\infty = \frac{\sigma^2}{2} \text{tr}(Q^{-1})$ . From the Sherman–Morrison formula above,

$$\text{tr}(Q^{-1}) = \frac{1}{q} \left( n - \frac{\gamma n}{q + \gamma n} \right) = \frac{n(q + \gamma(n - 1))}{q(q + \gamma n)},$$

which gives (46).

*Local regret.* By the Pythagorean form of Lemma 1 (the orthogonality identity after Lemma 1, with both  $x^\star$  and  $u^{\text{loc}}$  mean zero),  $2R_{\text{loc}} = \|x^\star\|_Q^2 - \|u^{\text{loc}}\|_Q^2 = 2R_\infty - \|u^{\text{loc}}\|_Q^2$ . Since  $u^{\text{loc}} = \theta/(q+\gamma)$  and  $\text{tr}(Q) = n(q+\gamma)$ ,

$$\|u^{\text{loc}}\|_Q^2 = \frac{\sigma^2 \text{tr}(Q)}{(q+\gamma)^2} = \frac{n\sigma^2}{q+\gamma}.$$

Therefore

$$R_{\text{loc}} = \frac{n\sigma^2}{2} \left[ \frac{q+\gamma(n-1)}{q(q+\gamma n)} - \frac{1}{q+\gamma} \right] = \frac{n\sigma^2}{2} \cdot \frac{(q+\gamma(n-1))(q+\gamma) - q(q+\gamma n)}{q(q+\gamma)(q+\gamma n)}.$$

The numerator reduces to  $\gamma^2(n-1)$ , which gives (47).

*Share.* Dividing (47) by (46), the factors  $n\sigma^2/2$  and  $q(q+\gamma n)$  cancel, leaving (48).  $\square$

## S2 Canonical spatial omission: detailed calculation

We expand the proof of Lemma 3 in the main paper, including the equal-inverse-length case in the variance integral.

*Proof.* By spatial stationarity, take  $z = 0$  and suppress the time argument. Write  $\beta_c \triangleq 1/\ell_c$  and  $\beta_s \triangleq 1/\ell_s$  for the inverse lengths, so  $w(u) = \frac{\beta_c}{2} e^{-\beta_c|u|}$  and  $\text{Cov}(\theta(u), \theta(u')) = \sigma^2 e^{-\beta_s|u-u'|}$ .

**Step 0 (omission is a variance ratio).** With  $Q = qI$  the optimal policy given  $\mathcal{O}_r$  is  $\mathbb{E}[x^\star(0) \mid \mathcal{O}_r]$ . Its expected loss is  $\frac{q}{2} \mathbb{E}[\text{Var}(x^\star(0) \mid \mathcal{O}_r)]$ . For a Gaussian field the conditional variance is a deterministic number: it does not depend on the observed values (companion Section 1.3). Since  $R_\infty = \frac{q}{2} \text{Var}(x^\star(0))$ ,

$$\eta_S(r) = \frac{\text{Var}(x^\star(0) \mid \mathcal{O}_r)}{\text{Var}(x^\star(0))}.$$

**Step 1 (split into observed middle and two tails).** Write  $x^\star(0) = M_r + T_r^+ + T_r^-$ , where

$$M_r = \int_{-r}^r w(u)\theta(u) du, \quad T_r^+ = \int_r^\infty w(u)\theta(u) du, \quad T_r^- = \int_{-\infty}^{-r} w(u)\theta(u) du.$$

The middle  $M_r$  is computed from observed values, so it contributes no conditional variance.

**Step 2 (rewrite the right tail).** Substituting  $u = r + s$ ,

$$T_r^+ = \frac{\beta_c}{2} e^{-\beta_c r} Y_r, \quad Y_r \triangleq \int_0^\infty e^{-\beta_c s} \theta(r+s) ds.$$

**Step 3 (spatial Markov property).** By Section 4, the field is Markov in  $z$ . Given the observed segment  $[-r, r]$ , the right tail  $\{\theta(u) : u > r\}$  and the left tail  $\{\theta(u) : u < -r\}$  are conditionally independent, and each depends on the segment only through its adjacent endpoint  $\theta(r)$  or  $\theta(-r)$ . Therefore

$$\text{Var}(T_r^+ \mid \mathcal{O}_r) = \text{Var}(T_r^+ \mid \theta(r)), \quad \text{Var}(T_r^- \mid \mathcal{O}_r) = \text{Var}(T_r^- \mid \theta(-r)), \quad \text{Cov}(T_r^+, T_r^- \mid \mathcal{O}_r) = 0.$$

**Step 4 (the tail variance does not depend on  $r$ ).** Define  $C \triangleq \text{Var}(Y_0 \mid \theta(0))$ . Shifting the field by  $r$  does not change its law, so  $\text{Var}(Y_r \mid \theta(r)) = C$  for every  $r$ . By reflection symmetry (the covariance depends only on  $|u - u'|$ ), the left tail gives the same value. Hence

$$\text{Var}(x^*(0) \mid \mathcal{O}_r) = 2 \left( \frac{\beta_c}{2} \right)^2 e^{-2\beta_c r} C = \frac{\beta_c^2}{2} e^{-2\beta_c r} C.$$

At  $r = 0$  the same argument applies with  $\mathcal{O}_0 = \sigma(\theta(0))$  and  $M_0 = 0$ , giving  $\text{Var}(x^*(0) \mid \theta(0)) = \frac{\beta_c^2}{2} C$ . Dividing,  $\eta_S(r) = \eta_S(0) e^{-2\beta_c r} = \eta_S(0) e^{-2r/\ell_c}$ . The unknown constant  $C$  cancels, so it never has to be computed.

**Step 5 (the constant  $\eta_S(0)$ ).** Conditioning on the single Gaussian variable  $\theta(0)$  gives  $\text{Var}(x^*(0) \mid \theta(0)) = \text{Var}(x^*(0))(1 - \text{Corr}(x^*(0), \theta(0))^2)$  (companion Section 1.3), so  $\eta_S(0) = 1 - \text{Corr}^2$ . Two moments are needed.

*Covariance with the local value:*

$$\text{Cov}(x^*(0), \theta(0)) = \int_{\mathbb{R}} w(u) \sigma^2 e^{-\beta_s |u|} du = \frac{\sigma^2 \beta_c}{2} \int_{\mathbb{R}} e^{-(\beta_c + \beta_s)|u|} du = \frac{\sigma^2 \beta_c}{\beta_c + \beta_s}.$$

*Variance of the target:*  $\text{Var}(x^*(0)) = \frac{\sigma^2 \beta_c^2}{4} I$ , where  $I \triangleq \iint e^{-\beta_c(|u|+|v|) - \beta_s|u-v|} du dv$ .

To evaluate  $I$ , define the inner integral  $h(u) \triangleq \int_{\mathbb{R}} e^{-\beta_c|v|} e^{-\beta_s|u-v|} dv$ . Then  $h$  is even, so  $I = 2 \int_0^\infty e^{-\beta_c u} h(u) du$ . For  $u \geq 0$ , split the  $v$ -integral at 0 and  $u$ :

$$h(u) = \underbrace{\frac{e^{-\beta_s u}}{\beta_c + \beta_s}}_{v < 0} + \underbrace{\frac{e^{-\beta_c u} - e^{-\beta_s u}}{\beta_s - \beta_c}}_{0 \leq v \leq u} + \underbrace{\frac{e^{-\beta_c u}}{\beta_c + \beta_s}}_{v > u}.$$

(The middle term is read as  $u e^{-\beta_c u}$  when  $\beta_s = \beta_c$ .) Multiply by  $e^{-\beta_c u}$  and integrate over  $u \geq 0$  term by term:

$$\frac{1}{(\beta_c + \beta_s)^2}, \quad \frac{1}{\beta_s - \beta_c} \left( \frac{1}{2\beta_c} - \frac{1}{\beta_c + \beta_s} \right) = \frac{1}{2\beta_c(\beta_c + \beta_s)}, \quad \frac{1}{2\beta_c(\beta_c + \beta_s)}.$$

Summing and doubling,

$$I = 2 \left[ \frac{1}{(\beta_c + \beta_s)^2} + \frac{1}{\beta_c(\beta_c + \beta_s)} \right] = \frac{2(2\beta_c + \beta_s)}{\beta_c(\beta_c + \beta_s)^2}, \quad \text{Var}(x^*(0)) = \frac{\sigma^2 \beta_c (2\beta_c + \beta_s)}{2(\beta_c + \beta_s)^2}. \quad (\text{S1})$$

*Combine.* With  $\text{Var}(\theta(0)) = \sigma^2$ ,

$$\text{Corr}^2 = \frac{(\sigma^2 \beta_c / (\beta_c + \beta_s))^2}{\sigma^2 \cdot \sigma^2 \beta_c (2\beta_c + \beta_s) / (2(\beta_c + \beta_s)^2)} = \frac{2\beta_c}{2\beta_c + \beta_s}, \quad \eta_S(0) = \frac{\beta_s}{2\beta_c + \beta_s} = \frac{\ell_c}{\ell_c + 2\ell_s}.$$

□

### S3 Numerical settings and diagnostics

The six experiments and their principal findings appear in Section 6 of the main paper. The original detailed methodology and diagnostics follow; figure references point to that paper. No experiments have been regenerated.

### S3.1 Numerical Methodology and Synthetic Environments

The numerical objects are architecture regrets, so each calculation uses the same information structure as its analytical counterpart. We work with an exogenous common-rate Gaussian-affine environment: actions do not alter later states. For  $x^* = K\theta + k_0$ , with  $K = Q^{-1}B$ , we evaluate

$$R(\mathcal{A}) = \frac{1}{2} \|x^* - u_{\mathcal{A}}^*\|_Q^2$$

from conditional covariance and, where appropriate, from independently sampled quadratic losses. The solvers respect the information structure: the common delayed snapshot in Experiment 1 permits a joint conditional mean even for non-diagonal  $Q$ ; Experiment 2 uses the exact fresh-local rule; Experiments 3 and 4 evaluate the scalar composition; and Experiments 5 and 6 use  $Q = I$  so agentwise Gaussian conditional variances are exact. Experiment 5 holds the graph and covariance fixed while changing decision sensitivity. The graph covariances are scaled to average marginal variance one.

For an arbitrary graph, a convenient alternative is to define the covariance spectrally through the graph Laplacian  $L_G$ . The covariance in main-paper Eq. (95) is positive semidefinite and places more state energy in low graph-frequency modes. Such constructions are closely related to graph signal smoothness and Gaussian Markov random-field models [1, 2].

Pointwise Monte Carlo means use 95% normal confidence intervals. The Experiment 2 boundary intervals apply the delta method to the paired fresh-local/open-loop regret means, so their uncertainty reflects that the two regrets use the same draws.

Table 1 summarizes the reference settings. The repository stores fixed configurations, processed tables, selected raw arrays, and graph covariance/distance inputs where applicable, together with run metadata and diagnostics; it is not a complete archive of every random draw. The exact system construction, graph seeds, conditioning safeguards, boundary conventions, radius grids, crossover estimator, and interval calculation are specified in the reproduction documentation. Robustness beyond the common-rate Gaussian-affine model is discussed in Section 8.

Table 1: Reference numerical settings. The shared-resource sweep fixes  $q = 1$ ; graph covariances in Experiments 5–6 are scaled to average marginal variance one. Quantities not being swept retain the experiment-specific values in the configuration files.

| Experiment            | principal sweep                               | fixed setting/size                          | evaluation       |
|-----------------------|-----------------------------------------------|---------------------------------------------|------------------|
| 1. Temporal pairs     | $\tau/T \in [0, 3]$ (25 points)               | A–D; $n = 8, 20, 50, 32$                    | 20,000/point     |
| 2. Local–global       | 9 $\gamma/q$ values in $[0, 10]$              | $n = 5, 10, 25, 100$ ; $q = 1$              | 30,000/setting   |
| 3. Scalar radius      | $r/\ell_c \in [0, 2]$ (1,001 points)          | 3 regimes; $D/\ell_c = 2$                   | 50,000/point     |
| 4. Canonical map      | $(vT/\ell_c, \ell_s/\ell_c)$ ; $72 \times 68$ | $r/\ell_c \in [0, 4]$ ; $\Delta r = 0.0016$ | direct scalar    |
| 5. Decision relevance | $\ell_c = 0.7, 2, 6$                          | 64-node ring; $T = 6$                       | 12,000/check     |
| 6. General graphs     | path, ring, grid, geometric                   | 64 nodes; $T = 5$                           | exact covariance |

### S3.2 Temporal Scaling and the Local–Global Crossover

The first two experiments test two different consequences of common-rate temporal predictability: the exact stale-global law (38), and the endpoint crossover threshold (49).

Experiment 1 asks whether the normalized regret of a delayed global snapshot depends only on  $\rho = \tau/T$ , even when spatial covariance, objective coupling, and the decision operator change. We used the four systems in Table 1 and drew 20,000 exact stationary OU pairs  $(\theta_{t-\tau}, \theta_t)$  at each of 25 values of  $\rho$ . The first member has covariance  $\Sigma_S$  and the second is generated as  $e^{-\rho}\theta_{t-\tau} +$

$(1 - e^{-2\rho})^{1/2}\varepsilon$ , with an independent innovation  $\varepsilon$  of covariance  $\Sigma_S$ . For each pair, the optimal delayed-snapshot action was evaluated with the known common-rate coefficient and the quadratic loss was computed in the system’s  $Q$  metric.

Across the 100 system–parameter combinations, the Monte Carlo curves collapse onto  $1 - e^{-2\tau/T}$  (Fig. 1). The largest absolute normalized deviation was 0.0155 and the root-mean-square deviation was 0.00471. This supports the exact temporal scaling for these stationary common-rate pairs.

Experiment 2 asks whether the finite- $n$  threshold in Eq. (49) describes the point at which fresh local information becomes preferable to a stale global snapshot. We used the shared-resource model  $Q = qI + \gamma\mathbf{1}\mathbf{1}^\top$ ,  $q = 1$ , with independent OU components. For each of 36  $(n, \gamma/q)$  settings, the fresh-local and open-loop regrets were evaluated on the same 30,000 current-state draws. The empirical boundary was then inferred by inserting these two estimates into the OU temporal law,

$$\hat{\rho}^\star = \frac{1}{2} \log \left( \frac{\hat{R}_\infty}{\hat{R}_\infty - \hat{R}_{\text{loc}}} \right).$$

Accordingly, Fig. 2 reports the temporal-law boundary inferred from paired local/open-loop losses.

The mean absolute boundary error was 0.00111 and the maximum was 0.00557. Finite size matters most under strong coupling: at  $n = 5$  and  $\gamma/q = 10$ , the exact threshold is 0.109 below the large-system limit. As  $n$  grows, the inferred boundary approaches  $\frac{1}{2} \log(1 + \gamma/q)$ . Stronger decision coupling therefore makes older global information worth tolerating in this independent-component model.

### S3.3 Emergence of an Optimal Coordination Radius

Experiment 3 illustrates why expanding the information radius can help at first and hurt later: reduced spatial omission competes with increasing staleness. We used  $\eta_S(r) = \eta_0 e^{-2r/\ell_c}$  and  $\tau(r) = r/v$  on  $0 \leq r \leq D = 2$ , with local  $(\eta_0, vT) = (0.25, 0.2)$ , interior  $(0.7, 4)$ , and maximum allowed radius  $(0.9, 100)$  regimes (all with  $\ell_c = 1$ ). At each radius, independent standard-normal draws were scaled by the two prescribed component variances and added before squaring. This is a scalar loss check: no spatial field, graph, or neighborhood conditional distribution is sampled.

The grid minima were 0, 0.626, and 2; the corresponding analytical minima after imposing the cap were 0, 0.62638, and 2. At the interior grid minimum, the marginal-slope gap was  $|m_S(\hat{r}^\star) - m_T(\hat{r}^\star)| = 4.77 \times 10^{-4}$ , and the largest Monte Carlo deviation from the prescribed scalar curve was about 0.020. The maximum allowed radius case is not a global endpoint: its unconstrained minimizer lies beyond  $D$ , while  $\eta_S(D) = \eta_0 e^{-2D/\ell_c} > 0$ . Thus the cap still omits decision-relevant state. The three curves distinguish immediate dominance of staleness, an interior balance, and a benefit from expansion throughout the allowed range.

### S3.4 Scaling Laws and Regime Map

Experiment 4 asks whether the canonical closed form predicts both the zero-radius region and the interior radius over a two-parameter sweep, rather than only at selected examples. We swept 4,896 points over  $vT/\ell_c \in [0.1, 20]$  and  $\ell_s/\ell_c \in [0.05, 5]$ , using  $\ell_c = 1$  as the unit of length. Independently of the closed form, every point was minimized by direct scalar evaluation of Eq. (65) on the same fixed grid  $r_j = 0.0016j$ ,  $j = 0, \dots, 2500$ , so  $r \in [0, 4]$  for every parameter pair. No covariance simulation or adaptive radius range was used. Separate one-dimensional sweeps tested the four comparative statics.

The gray region in Fig. 4 selects fresh local information; above the threshold  $vT = 2\ell_s$ , positive scope becomes worthwhile. Moving upward increases the distance information can travel within a

coherence time and raises the preferred radius. Moving right increases spatial correlation, making remote observations more redundant and lowering the preferred radius. The slices show these changes as ordinary radius curves. The mean absolute grid-versus-closed-form error was  $2.25 \times 10^{-4}$  and the maximum was  $8.00 \times 10^{-4}$ , below the 0.0016 grid step. Ordinary grid error occurs throughout the positive-radius interior because the minimizer is discretized; it is not confined to the threshold. There were five zero/positive classification disagreements, all adjacent to the threshold where the continuous optimum fell below the first positive grid point. The one-dimensional sweeps were monotone in all four predicted directions:  $r^*$  increased with  $T$ ,  $v$ , and  $\ell_c$ , and decreased with  $\ell_s$ . This validates the scalar search and canonical scaling within the stated range.

### S3.5 State Predictability versus Decision-Relevant Predictability

Experiment 5 asks whether the architecture depends on which state modes affect the decision, rather than only on raw state correlation. We held a 64-node ring, its Laplacian covariance,  $T = 6$ , and the latency law (0.12 time units per hop) fixed, and changed only the decision-sensitivity operator  $K_{ij} = e^{-d_G(i,j)/\ell_c} / \|e^{-d_G(i,\cdot)/\ell_c}\|_2$ , using  $\ell_c \in \{0.7, 2, 6\}$  and  $Q = I$ . Exact Gaussian conditioning gives the decision-weighted omission  $\eta_S(r)$  at every graph radius. The raw state-correlation statistic was computed separately from the fixed covariance; it is not the same quantity as  $\eta_S(r)$ . Each curve is normalized by its own  $R_\infty = \frac{1}{2} \text{tr}(K\Sigma_S K^\top)$ , so normalized regrets compare unexplained fractions rather than equal absolute cost scales.

Increasing  $\ell_c$  from 0.7 to 2 to 6 increased  $\eta_S(0)$  from 0.0460 to 0.258 to 0.572 and moved the discrete optimum from 1 to 2 to 5 hops, even though the raw state correlation was identical. Independent 12,000-sample conditioning checks differed from the exact omission by at most  $2.39 \times 10^{-3}$  (the largest pointwise standard error was  $3.20 \times 10^{-3}$ ). Thus changing only decision sensitivity in a fixed environment produced substantially different preferred architectures.

### S3.6 Finite Graphs beyond the Canonical Geometry

Experiment 6 asks whether the conditional-variance construction and discrete radius comparison remain usable when the geometry is not the homogeneous line assumed by the canonical formula. We considered one 64-node graph of each of a path, ring, two-dimensional grid, and random geometric type; the geometric graph uses the fixed configured seed. For each instance we formed the covariance (95), scaled it to average marginal variance one, and computed every  $\eta_S(r)$  directly from the Gaussian conditional covariance of the desired action given the radius- $r$  neighborhood. The reference setting uses  $Q = I$ ,  $T = 5$ ,  $\ell_c = 2$ , and latency 0.12 per hop. No exponential omission fit or canonical line formula was used, and no graph ensemble average was taken.

Under the reference setting, Fig. 6 gives exact integer optima of 2, 2, 3, and 2 hops for the path, ring, grid, and geometric instance, respectively. In these four sampled instances, increasing  $T$  over the configured values 1.5, 2.5, 4, 6, 9, 14 changed the optimum from 1 to 3 hops on the path and ring, from 1 to 4 on the grid, and from 1 to 3 on the geometric graph. The sampled  $\ell_c$  sweep from 0.7 to 6 moved the optima from (0, 0, 1, 1) to (5, 5, 5, 3) in the same graph order. Increasing per-hop latency weakly decreased the optimum, as did decreasing  $\kappa_s$ , which increases spatial smoothness after variance normalization. These comparative statics describe the four specified graph instances. Since radii are integer-valued, the appropriate comparison is the one-hop regret difference in Eq. (68). For example, the geometric graph's three-hop regret exceeds its two-hop minimum by only  $0.00362R_\infty$ ; the archived neighbor differences quantify such shallow minima.

## References

- [1] D. I. Shuman, S. K. Narang, P. Frossard, A. Ortega, and P. Vandergheynst, “The emerging field of signal processing on graphs: Extending high-dimensional data analysis to networks and other irregular domains,” *IEEE Signal Processing Magazine*, vol. 30, no. 3, pp. 83–98, 2013.
- [2] F. Lindgren, H. Rue, and J. Lindström, “An explicit link between Gaussian fields and Gaussian Markov random fields: the stochastic partial differential equation approach,” *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, vol. 73, no. 4, pp. 423–498, 2011.