

Variants of Gödel's Ontological Proof in a Natural Deduction Calculus

the date of receipt and acceptance should be inserted later

Abstract This paper presents detailed formalizations of ontological arguments in a simple modal natural deduction calculus. The first formal proof closely follows the hints in Scott's manuscript about Gödel's argument and fills in the gaps, thus verifying its correctness. The second formal proof improves the first one, by relying on the weaker modal logic **KB** instead of the stronger modal logic **S5**; and by avoiding the introduction of the equality predicate. The second proof is also technically shorter than the first one, because it eliminates unnecessary detours and uses Axiom 1 for the positivity of properties only once. The third and fourth proofs formalize, respectively, Anderson's and Bjørdal's variants of the ontological argument, which are known to be immune to modal collapse.

Keywords: Ontological Argument, Higher-Order Logics, Modal Logics, Natural Deduction.

“There is a scientific (exact) philosophy and theology, which deals with concepts of the highest abstractness; and this is also most highly fruitful for science. [...] Religions are, for the most part, bad; but religion is not.”
- Kurt Gödel (*Wang*, 1996)[p. 316]

1 Introduction

Ontological arguments for the existence of God can be traced back at least to St. Anselm (1033-1109). His argument considers a greatest conceivable being, who must exist, because if it did not have the property of existence, then we could conceive of a greater being that, in addition to other properties, also had the property of existence. St. Anselm's argument was criticized and further developed by Descartes, Leibniz and Kant, among many others.

Leibniz identified the possible existence of God as a critical missing step in St. Anselm's argument. To fill this gap, he argued that the properties of God, the perfections, are compatible. This means that it is possible to satisfy all perfections at once, which implies that the existence of a greatest conceivable being with all these properties is possible.

Gödel built on Leibniz's work (*Adams*, 1995) and brought the ontological argument to a modern form using a modal logic with higher-order quantification over properties. In this setting, he gave precise axioms describing the notion of *positive* property and defined God as a being that has all positive properties. Gödel's work was saved in his own notes (*Gödel*, 1970) as well as in notes by Scott (*Scott*, 2001), in whom he confided his proof.

The increase in formality of the ontological argument has required a development of its basic notions. Gödel's notion of positive property and Leibniz's notion of perfection differ. A formal distinction is that Leibniz's perfections are atomic whereas Gödel's positive properties can consist of combinations of atomic properties (*Fitting*, 2002)[p.139]. In particular, one of Gödel's axioms states that any conjunction of positive properties is itself positive. From this

axiom, it is immediately deduced that the property of being God-like is positive. Intuitively, a (possibly infinite) conjunction of positive properties is deduced from the universal definition of God-likeness. This deductive inference is not formalizable in a finite first-order calculus. The interplay between universal quantification (in the definition of a God-like being) and infinite conjunctions (in Gödel’s axiom for positive properties) could explain why, starting with Scott (Scott, 2001), this axiom of Gödel has been replaced by another that simply assumes the positivity of the property of being god-like.

The aim of this paper is to present detailed formalizations of variants of ontological arguments in a natural deduction calculus. For a comprehensive introduction to natural deduction, the reader can consult (Prawitz, 2006) or (Gabbay, 1996). The natural deduction calculus proposed and used in this paper is an extension of a standard natural deduction calculus for (non-modal) higher-order logic (Benzmüller & Brown & Kohlhase, 2004) with new introduction and elimination rules for modal operators (as presented in Section 2). This calculus is sound and complete relative to the non-modal calculus extended with Axiom K and the Necessitation rule (as demonstrated in Subsection 2.1). There are various higher-order logics (Benzmüller & Brown & Kohlhase, 2004) and various higher-order modal logics (Muskens, 2006). The calculus proposed here can be regarded as a minimalistic calculus for a rigid higher-order modal logic K without extensionality principles. With additional axioms for stronger modal logics, such as KB and S5, this calculus is sufficient to achieve the stated aim.

The natural deduction style was chosen for four reasons. Firstly, presentations of Gödel’s proof are typically either informal or formalized in other styles of proof calculi (e.g. Fitting’s tableaux (Fitting, 2002) or Sobel’s sentential modal calculus (Sobel, 2001)). Therefore, a formalization in natural deduction is a valuable complement to the existing presentations. Secondly, it makes the ontological proof accessible to people who are familiarized with a natural deduction style. Thirdly, as natural deduction is the style used by proof assistants such as Coq (Paulin-Mohring, 2015) and Isabelle (Nipkow & Paulson & Wenzel, 2002), natural deduction formalizations can be verified step-by-step in such proof assistants, and we have in fact done this (Benzmüller & Woltzenlogel Paleo, 2015a). And finally, in comparison to other kinds of proof systems (e.g. resolution, tableaux, sequent calculi), natural deduction can be considered less technical, more human-readable and more intuitive, because it relies on natural principles of reasoning such as modus ponens (implication elimination) and reasoning by assumption (implication introduction); these qualities are conducive to rendering the formalizations of ontological arguments accessible to a wider audience.

The first main contribution of this paper is a detailed formalization of Scott’s version (Scott, 2001) of Gödel’s ontological argument (Gödel, 1970) (as shown in Section 4) in the proposed natural deduction calculus. The second main contribution of the paper is a new proof (shown in Section 5), also in natural deduction style. In contrast to Scott’s proof (Scott, 2001), which requires the modal logic **S5**, the new proof requires only the weaker modal logic **KB**. The new proof also does not rely on the equality predicate and is much shorter. Although there have been claims (as discussed in more detail in Section 5) that **KB** is sufficient and the equality predicate is superfluous, formal proofs substantiating these claims were missing up to now.

A major criticism against Gödel’s argument is an undesirable consequence of the stipulated axioms, called *modal collapse*. This is discussed in greater detail in Section 6, where a natural deduction derivation of the collapse is presented. Many recent works on the ontological argument have proposed modifications of the argument that do not entail modal collapse. In this paper we present natural deduction formalizations of two of these solutions: Anderson’s emendation (in Section 7) and Bjørdal’s variant (in Section 8).

An early unpublished draft of Section 5 of this paper, prepared by Woltzenlogel Paleo in March 2013, served as the starting point for two distinct but complementary research endeavours. Siders and Woltzenlogel Paleo transformed that early draft into this paper, aiming for human-readable and yet formal natural deduction proofs. Benzmüller and Woltzenlogel

Paleo, on the other hand, focused on the verification of variants of Gödel’s ontological arguments using higher-order automated reasoning tools, which quickly led to many publications. The main automatically obtained results about Gödel’s and Scott’s proofs appeared in the European Conference on Artificial Intelligence (*Benzmüller & Woltzenlogel Paleo*, 2014d) and, two years later, a detailed analysis of an inconsistency found in Gödel’s axioms was published in the International Joint Conference on Artificial Intelligence (*Benzmüller & Woltzenlogel Paleo*, 2016a). The automatic verification of variants by Anderson, Bjordal and Hájek, on the other hand were presented in the First World Congress on Logic and Religion (*Benzmüller & Weber & Woltzenlogel Paleo*, 2015) and published in (*Benzmüller & Weber & Woltzenlogel Paleo*, 2016) and results related to the modal collapse were reported in the 4th Word Congress on the Square of Opposition (*Benzmüller & Woltzenlogel Paleo*, 2014c) and later published in (*Benzmüller & Woltzenlogel Paleo*, 2016b). More detailed reconstructions of Scott’s variant within the interactive proof assistants Isabelle and Coq were released, respectively, in Isabelle’s Archive of Formal Proofs (*Benzmüller & Woltzenlogel Paleo*, 2013b) and in the Symposium of Computer Science in Russia (*Benzmüller & Woltzenlogel Paleo*, 2015a). The popularity of the topic led to invited talks and corresponding invited abstracts in the proceedings of workshops (*Benzmüller & Woltzenlogel Paleo*, 2014a,b, 2013a) and summer schools (*Benzmüller & Woltzenlogel Paleo*, 2015c,b) as well as articles targeting a more general audience (*Woltzenlogel Paleo*, 2013). More recently, this line of research ventured into other types of ontological arguments, and a computer-assisted reconstruction of an argument by Leibniz will appear as a chapter in a book dedicated to the 300th anniversary of Leibniz’s death (*Bentert & Benzmüller & Streit & Woltzenlogel Paleo*, 2016). The automatic verification of Gödel’s and Scott’s variants (*Benzmüller & Woltzenlogel Paleo*, 2014d) was preceded by an automatic verification of (a non-modal version of) Anselm’s ontological argument (*Oppenheimer & Zalta*, 2011). However, whereas the complexity of Gödel’s and Scott’s arguments required the embedding of higher-order modal logics into higher-order logics and the use of higher-order automated reasoners, Anselm’s argument greater simplicity enabled it to be verified with a first-order prover. Anselm’s argument has also been formalized within the interactive proof assistant PVS (*Rushby*, 2013) and there has been a so far failed attempt to formalize Avicenna’s argument in PVS as well.

2 Natural Deduction

The language of higher-order modal logic used here is inspired by that of Church’s simple type theory (*Church*, 1940).

Definition 1 *Simple types* are given by the following grammar:

$$\theta, \tau ::= \mu \mid o \mid \theta \rightarrow \tau$$

where μ is the atomic type for individuals, o is the atomic type for propositions and $\theta \rightarrow \tau$ is the type for functions taking an argument of type θ and returning something of type τ . ‘ \rightarrow ’ is assumed to be right associative.

Definition 2 *Terms and formulas* are given by the following grammar:

$$\begin{aligned} s, t ::= & p_\tau \mid X_\tau \mid (\lambda X_\theta. s_\tau)_{\theta \rightarrow \tau} \mid (s_{\theta \rightarrow \tau} t_\theta)_\tau \mid \\ & \perp_o \mid \rightarrow_{o \rightarrow o \rightarrow o} \mid \wedge_{o \rightarrow o \rightarrow o} \mid \\ & \forall_{(\tau \rightarrow o) \rightarrow o} \mid \exists_{(\tau \rightarrow o) \rightarrow o} \mid \Box_{o \rightarrow o} \mid \Diamond_{o \rightarrow o} \end{aligned}$$

where p_τ and X_τ range over, respectively, constants and variables of type τ . Parenthesis conventions, infix notation for propositional connectives and binding notation¹ for quantifiers are assumed. Furthermore, subscript types are omitted when they are clear from the context.

¹ $\forall x. F[x]$ and $\exists x. F[x]$ are assumed to be abbreviations for, respectively, $\forall(\lambda x. F[x])$ and $\exists(\lambda x. F[x])$.

Implicit alpha-conversion is assumed: two expressions are considered to be the same modulo renaming of their bound variables. Negation ($\neg_{o \rightarrow o}$), disjunction ($\vee_{o \rightarrow o \rightarrow o}$), equivalence ($\leftrightarrow_{o \rightarrow o \rightarrow o}$), equality ($=_{\tau \rightarrow \tau \rightarrow o}$), inequality ($\neq_{\tau \rightarrow \tau \rightarrow o}$), are defined by $\neg A \equiv A \rightarrow \perp$, $(A \vee B) \equiv \neg(\neg A \wedge \neg B)$, $(A \leftrightarrow B) \equiv (A \rightarrow B) \wedge (B \rightarrow A)$, $s = t \equiv \forall P.(P\ s) \leftrightarrow (P\ t)$ (Leibniz's equality) and $s \neq t \equiv \neg(s = t)$, .

The natural deduction calculus proposed and used here has standard rules for propositional connectives and quantifiers, as shown in Figures 1 and 3. The extension to classical logic is achieved by adding a rule for double negation elimination, shown in Figure 2, and β -reduction² is handled by the β rule shown in Figure 4. Finally, modal operators are handled by special new rules that insert or remove formulas from boxes, as shown in Figure 5. A *derivation* is a directed acyclic graph whose nodes are formulas and whose edges correspond to applications of the inference rules. A *proof* of a formula F is a derivation without open assumptions and having F as root not inside any box.

In every inference rule, (sub-)formulas in the conclusion correspond³ to formulas in the premises (even in the case of quantifier rules, where the formulas changes due to a substitution). For any expression occurring both in a formula in the premise and in its corresponding (sub-)formula in the conclusion, the occurrence in the premise is said to be an *ancestor* of the occurrence in the conclusion, and the occurrence in the conclusion is said to be a *descendant* of the occurrence in the premise.

Applications of the β rule are silently omitted. Double lines are used to abbreviate tedious propositional reasoning steps in the derivations where the reasoning is otherwise clear. Dashed lines are used to refer to an axiom or theorem with the proof shown elsewhere. When proof trees are too large to fit on the page, some branches may be displayed separately. In such cases, the conclusion of the branch and the location in the main proof tree where the branch belongs are annotated with the same symbol (a subscripted \star). Dotted lines are used to indicate folding and unfolding of definitions. Furthermore, as it is inconvenient to draw boxes around large derivations in L^AT_EX, formulas inside boxes are labeled with the names of the boxes surrounding them. Therefore, the boxes can be omitted without loss of information.

The calculus having only the rules shown in Figures 1, 2 and 3 is named **ND**. This calculus is essentially the same as the simplest natural deduction calculus in the family of calculi discussed in (Benzmüller & Brown & Kohlhase, 2004). The main difference is that they take disjunction as a primitive connective instead of conjunction, and they use a sequent notation, where the assumptions are listed in the left side of the sequent. The calculus with the new rules for modal operators shown in Figure 5 is named **ND_K**.

Fig. 1 Propositional rules

$$\begin{array}{cccc}
 \frac{}{A} \perp & \frac{B}{A \rightarrow B} \rightarrow_I & \frac{\overline{A}^n \vdots B}{A \rightarrow B} \rightarrow_I^n & \frac{A \quad A \rightarrow B}{B} \rightarrow_E \\
 \\
 \frac{A \quad B}{A \wedge B} \wedge_I & \frac{A \wedge B}{A} \wedge_{E1} & \frac{A \wedge B}{B} \wedge_{E2} &
 \end{array}$$

² Rules for η -expansion (Benzmüller & Brown & Kohlhase, 2004) are not included here, because they are not needed in the ontological arguments.

³ This correspondence could be defined formally, but it would be tedious to do so.

Fig. 2 Double negation elimination

$$\frac{\neg\neg A}{A} \neg\neg E$$

Fig. 3 Quantifier rules

$$\frac{A[\alpha]}{\forall x_{\tau}.A[x]} \forall_I \quad \frac{\forall x_{\tau}.A[x]}{A[t]} \forall_E \quad \frac{A[t]}{\exists x_{\tau}.A[x]} \exists_I \quad \frac{\exists x_{\tau}.A[x]}{A[\beta]} \exists_E$$

eigen-variable conditions:
 if ρ is a \forall_I inference eliminating a variable α , then any occurrence of α in the proof should be an ancestor of the occurrence of α eliminated by ρ and not a descendant of a variable introduced by a \exists_E inference;
 if ρ is a \exists_E inference introducing a variable β , then any occurrence of β in the proof should be a descendant of the occurrence of β introduced by ρ and not an ancestor of a variable eliminated by a \forall_I inference.

Fig. 4 Reduction rules

$$\frac{A}{B} \beta$$

only if B and A are β -equivalent
 (i.e. there is a C such that both B and A β -reduce to C)

Fig. 5 Rules for modal operators

$$\frac{\omega : \boxed{\begin{smallmatrix} \vdots \\ A \end{smallmatrix}}}{\Box A} \Box_I \quad \frac{\Box A}{w : \boxed{\begin{smallmatrix} A \\ \vdots \end{smallmatrix}}} \Box_E$$

$$\frac{w : \boxed{\begin{smallmatrix} \vdots \\ A \end{smallmatrix}}}{\Diamond A} \Diamond_I \quad \frac{\Diamond A}{\omega : \boxed{\begin{smallmatrix} A \\ \vdots \end{smallmatrix}}} \Diamond_E$$

eigen-box condition:
 An modal inference is said to *access* the box immediately above or below it.
 \Box_I and \Diamond_E are *strong* modal rules:
 ω must be a fresh name for the box they access
 (in analogy to the eigen-variable condition for strong quantifier rules).
 Every box must be accessed by *exactly one* strong modal inference.

boxed assumption condition:
 assumptions should be discharged within the box where they are created.

2.1 Suitability for Rigid Higher-Order Modal Logic K

ND is a calculus for a (non-modal) higher-order logic without functional and boolean extensionality principles (*Benzmüller & Brown & Kohlhase, 2004*). Adding the modal rules results in a calculus that is suitable for a basic rigid modal logic **K**, in the sense that **ND_K**

is sound and complete relative to **ND** extended with axiom K ($\Box(A \rightarrow B) \rightarrow (\Box A \rightarrow \Box B)$) and the necessitation rule (which establishes that $\Box A$ is a theorem if A is a theorem).

Theorem 1 **ND_K** is complete, relative to **ND** extended with axiom K and the necessitation rule.

Proof The necessitation rule can be immediately simulated with the \Box_I rule. Axiom K can be derived in **ND_K** as shown below:

$$\frac{\frac{\frac{\overline{\Box(A \rightarrow B)}}{\omega : A \rightarrow B} \Box_E \quad \frac{\frac{\overline{\Box A}}{\omega : A} \Box_E}{\omega : B} \Box_I}{\Box A \rightarrow \Box B} \rightarrow_I^1}{\Box(A \rightarrow B) \rightarrow (\Box A \rightarrow \Box B)} \rightarrow_I^2$$

Theorem 2 **ND_K** is sound, relative to **ND** extended with axiom K and the necessitation rule.

Proof It is necessary to show that **ND_K** proofs of the following form can be translated to proofs in **ND** extended with the axiom K and the necessitation rule.

$$\frac{\frac{\frac{\Box A_1}{\omega : A_1} \Box_E \quad \frac{\Box A_n}{\omega : A_n} \Box_E}{\vdots \quad \dots \quad \vdots} \Box_I}{\omega : B} \Box_I$$

A translation to **ND** extended with axiom K and necessitation is shown below for the case when $n = 1$:

$$\frac{\frac{\frac{\overline{A_1} \quad 1}{\vdots} \quad \frac{B}{A_1 \rightarrow B} \rightarrow_I^1}{\Box(A_1 \rightarrow B)} \text{ necessitation} \quad \frac{\text{Axiom K}}{\Box(A_1 \rightarrow B) \rightarrow (\Box A_1 \rightarrow \Box B)} \rightarrow_E}{\frac{\Box A_1 \rightarrow \Box B}{\Box B} \rightarrow_E} \Box A_1 \rightarrow_E$$

For $n > 1$, the translation is a straightforward generalization:

$$\frac{\frac{\frac{\overline{A_1} \quad 1}{\vdots} \quad \frac{\overline{A_n} \quad n}{\vdots} \quad \dots \quad \frac{B}{A_1 \rightarrow \dots \rightarrow A_n \rightarrow B} \rightarrow_I^*}{\Box(A_1 \rightarrow \dots \rightarrow A_n \rightarrow B)} \text{ nec.} \quad \frac{\text{Axiom K, iterated}}{\Box(A_1 \rightarrow \dots \rightarrow A_n \rightarrow B) \rightarrow (\Box A_1 \rightarrow \dots \rightarrow \Box A_n \rightarrow \Box B)} \rightarrow_E}{\Box A_1 \rightarrow \dots \rightarrow \Box A_n \rightarrow \Box B} \rightarrow_E$$

$$\frac{\Box A_1 \rightarrow \dots \rightarrow \Box A_n \rightarrow \Box B}{\Box B} \rightarrow_E$$

Without the restriction that every box must be accessed by exactly one strong modal inference, the calculus would be unsound for the modal logic **K**. For example, the formula $\forall \phi.(\Box \phi \rightarrow \Diamond \phi)$ is not valid in **K** but would be provable without this restriction:

$$\begin{array}{c}
\frac{}{\Box\phi} 1 \\
\frac{}{\omega:\phi} \Box_E \\
\frac{}{\Diamond\phi} \Diamond_I \\
\frac{}{\Box\phi \rightarrow \Diamond\phi} \rightarrow_I^1 \\
\frac{}{\forall\phi.(\Box\phi \rightarrow \Diamond\phi)} \forall_I
\end{array}$$

This example proof is unsound according to the **ND_K** calculus, because the eigen-box condition is violated: the box labelled by ω is not accessed by any strong inference.

The straightforward combinations of the quantifier rules of **ND** either with the modal rules of **ND_K** or with axiom K and the necessitation rule are suitable for a higher-order modal logic where constants and variables are *rigid*. From the point of view of a *possible world semantics*, rigidity means that their interpretation is independent of the world in which they are being interpreted. Rigidity is tacitly assumed by most works investigating the ontological argument, and is explicitly assumed here. Nevertheless, it should be noted that its adequacy for the ontological argument has already been contested (*Fitting*, 2002). Another assumption made here is that the quantification domains are constant (i.e. independent of the possible worlds). Neither Gödel’s manuscript nor Scott’s manuscript reveal whether they use constant or varying domains, and this is also the case for many variants (e.g. (*Hajek*, 1996)). Nevertheless, some authors of variants (*Anderson*, 1990, footnotes 11 and 14) of Gödel’s ontological argument do explicitly state a preference for varying domains. Our choice of constant domains is motivated by simplicity. As is well-known, varying domain quantifiers (also known as *actualistic* quantifiers) can be simulated by constant domain quantifiers (also known as *possibilistic* quantifiers) guarded by a primitive *existence* predicate⁴; therefore, the formalization of ontological arguments with varying domain quantification would be feasible in the natural deduction calculus presented here as well.

3 Some Useful Derivable Modal Principles

From an axiomatic point of view, modal logics differ with respect to which additional axiom schemas they admit. Some common axiom schemas of relevance to the ontological argument are T ($\Box A \rightarrow A$), B ($A \rightarrow \Box\Diamond A$), 4 ($\Box A \rightarrow \Box\Box A$) and 5 ($\Diamond A \rightarrow \Box\Diamond A$). From a semantic point of view, these axioms correspond to geometric frame conditions that must be satisfied by the accessibility relation in Kripke models (See for example (*Fitting & Mendelsohn*, 1998) or (*Blackburn & al.*, 2001, Section 3.1) on Frame definability): T corresponds to reflexivity, B corresponds to symmetry, 4 corresponds to transitivity and 5 corresponds to euclidianity. In the models of logic **S5**, all these axioms are satisfied; in the models of logic **KB**, K and B are satisfied; and in the models of logic **B**, K, T and B are satisfied.

In this section, natural deduction proofs of three convenient derived modal principles are presented. These principles are used in the ontological proofs formalized in the next sections. Although they are well-known among modal logicians, we include their proofs here in order to have a self-contained and fully formal presentation of the ontological proofs.

The *distribution principle* can be seen as a form of modus ponens within the scope of modalities: if $A \rightarrow B$ holds in all accessible worlds and A holds in an accessible world, then B holds in an accessible world. This principle is provable in the modal logic **K**.

Lemma (Distribution Principle)

$$\Box(A \rightarrow B) \rightarrow (\Diamond A \rightarrow \Diamond B)$$

Proof

⁴ Formally, $\forall^e x.F[x] \equiv \forall x.E(x) \rightarrow F[x]$ and $\exists^e x.F[x] \equiv \exists x.E(x) \wedge F[x]$

$$\begin{array}{c}
\frac{\overline{\Box(A \rightarrow B)}^2}{\omega : A \rightarrow B} \Box_E \quad \frac{\overline{\Diamond A}^1}{\omega : A} \Diamond_E \\
\hline
\frac{\omega : B}{\Diamond B} \Diamond_I \quad \frac{\omega : A}{\Diamond A \rightarrow \Diamond B} \rightarrow_I^1 \\
\hline
\frac{\omega : A \rightarrow B}{\Box(A \rightarrow B) \rightarrow (\Diamond A \rightarrow \Diamond B)} \rightarrow_I^2
\end{array}$$

Brouwer's reduction principle is derivable in modal logic **KB**, using axiom B. The proofs of Gödel and Scott (Section 4) do not make direct use of this principle, but the new proof presented in Section 5 does.

Lemma (Brouwer's Reduction Principle)

$$\Diamond \Box A \rightarrow A$$

Proof

$$\begin{array}{c}
\frac{\overline{\Diamond \Box A}^2}{\neg \Box \neg \Box A} \Box_E \quad \frac{\overline{\neg A}^1}{\Box \neg \Box \neg \neg A} \rightarrow_E \quad \frac{\text{Axiom B}}{\neg A \rightarrow \Box \Diamond \neg A} \\
\hline
\frac{\Box \neg \Box \neg \neg A}{\Box \neg \Box A} \rightarrow_E \quad \frac{\perp}{\neg \neg A} \rightarrow_I^1 \quad \frac{A}{\neg \neg A} \neg \neg_E \\
\hline
\frac{A}{\Diamond \Box A \rightarrow A} \rightarrow_I^2
\end{array}$$

In the modal logic **S5**, a sequential iteration of modalities can be collapsed to the last modality in the sequence. In other words, the following principles hold in **S5**:

- $\Diamond^n \Box A \rightarrow \Box A$
- $\Box^n \Diamond A \rightarrow \Diamond A$
- $\Diamond^n \Diamond A \rightarrow \Diamond A$
- $\Box^n \Box A \rightarrow \Box A$

Below we exhibit a natural deduction proof for the first principle listed above when $n = 1$. Informally, this particular case can be read as the claim that anything that is possibly necessary is in fact necessary, and it is sufficient for the ontological proofs of Gödel and Scott. The proof of this principle depends on the previously proven Brouwer's reduction principle, on the Modal Axiom 5, and on K, which was shown to be derivable in the proof of Theorem 1.

Lemma (Iteration Principle)

$$\Diamond \Box A \rightarrow \Box A$$

Proof

$$\begin{array}{c}
\frac{\text{Brouwer's Reduction}}{\omega : \Diamond \Box A \rightarrow A} \Box_I \quad \frac{\text{K}}{\Box(\Diamond \Box A \rightarrow A) \rightarrow (\Box \Diamond \Box A \rightarrow \Box A)} \rightarrow_E \quad \frac{\text{Axiom 5 for } \Box A}{\Diamond \Box A \rightarrow \Box \Diamond \Box A} \rightarrow_E \\
\hline
\frac{\Box \Diamond \Box A \rightarrow \Box A}{\Diamond \Box A \rightarrow \Box A} \rightarrow_I^1
\end{array}$$

Lemma (Reflexivity)

$$s = s$$

Proof

$$\begin{array}{c}
\frac{\overline{Ps}^1}{Ps \rightarrow Ps} \rightarrow_I^1 \quad \frac{\overline{Ps}^2}{Ps \rightarrow Ps} \rightarrow_I^2 \\
\hline
\frac{Ps \leftrightarrow Ps}{\forall P. Ps \leftrightarrow Ps} \forall_I \\
\hline
\frac{\dots}{s = s}
\end{array}$$

4 Scott's Proof in Natural Deduction

In this section we present a detailed formalization of Scott's proof in the natural deduction calculus defined in Section 2. The reasoning in Scott's manuscript has been reproduced step-by-step and all reasoning gaps are filled in by using the deduction considered most natural. All derived formulas that do appear in Scott's manuscript are marked with a † here. Unmarked formulas were derived in the process of filling the gaps between the marked formulas mentioned in Scott's manuscript.

Scott's version of Gödel's proof depends on five axioms that circumscribe the notion of *positive* property, with positivity being denoted by the undefined second-order predicate symbol P . Additionally, three definitions are used for establishing the notions of *God-like*, *essence* and *necessary existence*. Technically, these definitions only abbreviate certain complex formulas. The argument would still go through if all defined symbols were replaced by the complex formulas they define. This observation is particularly relevant in the case of *necessary existence*. Since this notion of "existence" is just an abbreviation, Gödel's argument is not susceptible to Kant's criticism against Anselm's argument (that existence should not be treated as a predicate). In Gödel's proof, existence is properly denoted by the existential quantifier. It is, therefore, unfortunately misleading to refer to the defined predicate symbol E as "necessary existence", when in fact it is just a convenient abbreviation.

In addition to the five axioms and three definitions that pertain specifically to the argument, the proof also uses **S5**'s iteration principle (in the proof of Lemma 2) and the lemma of reflexivity for equality (in the proof of Theorem 1). The use of reflexivity could be avoided, but we chose to use it because it allows the construction of a shorter proof.

Axiom 1 *Either a property or its negation is positive, but not both:*

$$\forall\varphi.[P(\neg\varphi) \leftrightarrow \neg P(\varphi)]$$

Axiom 2 *A property necessarily implied by a positive property is positive:*

$$\forall\varphi.\forall\psi.[(P(\varphi) \wedge \Box\forall x.(\varphi(x) \rightarrow \psi(x))) \rightarrow P(\psi)]$$

Theorem 1 *Positive properties are possibly exemplified:*

$$\forall\varphi.[P(\varphi) \rightarrow \Diamond\exists x.\varphi(x)]$$

Proof

$$\begin{array}{c}
\begin{array}{c}
\text{Reflexivity} \\
\frac{}{\omega : \bar{\gamma} = \bar{\gamma}} \rightarrow_I \\
\frac{}{\omega : \rho(\gamma) \rightarrow \gamma = \gamma} \forall_I \\
\frac{}{\omega : \forall x.(\rho(x) \rightarrow x = x)} \forall_I \\
\frac{}{P(\rho)} 5 \quad \frac{}{\Box\forall x.(\rho(x) \rightarrow x = x)^\dagger} \Box_I \\
\frac{}{P(\rho) \wedge \Box\forall x.(\rho(x) \rightarrow x = x)} \wedge_I
\end{array}
\quad
\begin{array}{c}
\text{Axiom 2 for } \rho \text{ and } \lambda x.x = x \\
\frac{}{P(\rho) \wedge \Box\forall x.(\rho(x) \rightarrow x = x) \rightarrow P(\lambda x.x = x)} \rightarrow_E \\
\frac{}{P(\lambda x.x = x)^\dagger} [\star_1]
\end{array}
\\
\\
\begin{array}{c}
\frac{}{\omega : \rho(\beta)} 1 \quad \frac{}{\omega : \neg\rho(\beta)} \rightarrow_E \\
\frac{}{\omega : \neg(\beta = \beta)} \perp_E \\
\frac{}{\omega : \beta \neq \beta} \rightarrow_I \\
\frac{}{\omega : \rho(\beta) \rightarrow \beta \neq \beta} \forall_I \\
\frac{}{\omega : \forall x.(\rho(x) \rightarrow x \neq x)} \forall_I \\
\frac{}{P(\rho)} 5 \quad \frac{}{\Box\forall x.(\rho(x) \rightarrow x \neq x)^\dagger} \Box_I \\
\frac{}{P(\rho) \wedge \Box\forall x.(\rho(x) \rightarrow x \neq x)} \wedge_I
\end{array}
\quad
\begin{array}{c}
\text{Axiom 2 for } \rho \text{ and } \lambda x.x \neq x \\
\frac{}{P(\rho) \wedge \Box\forall x.(\rho(x) \rightarrow x \neq x) \rightarrow P(\lambda x.x \neq x)} \rightarrow_E \\
\frac{}{P(\lambda x.x \neq x)^\dagger} [\star_2]
\end{array}
\end{array}$$

$$\begin{array}{c}
\frac{\overline{\omega : G(\delta)} \quad 2}{\dots \dots \dots} \quad \frac{\overline{\omega : \forall \varphi. P(\varphi) \rightarrow \varphi(\delta)} \quad \forall_E}{\omega : P(\rho) \rightarrow \rho(\delta)} \quad \frac{\omega : P(\rho)}{\omega : P(\rho)} \quad 3 \quad \frac{\omega : P(\rho) \rightarrow \rho(\delta)}{\omega : P(\rho) \rightarrow \rho(\delta)} \rightarrow_E \\
\frac{\omega : \rho(\delta)}{\omega : G(\delta) \rightarrow \rho(\delta)} \rightarrow_I^2 \quad \frac{\omega : G(\delta) \rightarrow \rho(\delta)}{\omega : \forall y. (G(y) \rightarrow \rho(y))} \forall_I \quad \frac{\omega : \forall y. (G(y) \rightarrow \rho(y))}{\omega : P(\rho) \rightarrow \forall y. (G(y) \rightarrow \rho(y))} \rightarrow_I^3 \\
\frac{\omega : P(\rho) \rightarrow \forall y. (G(y) \rightarrow \rho(y))}{\Box(P(\rho) \rightarrow \forall y. (G(y) \rightarrow \rho(y)))} \Box_I \quad \frac{\Box(P(\rho) \rightarrow \forall y. (G(y) \rightarrow \rho(y)))}{\Box P(\rho) \rightarrow \Box \forall y. (G(y) \rightarrow \rho(y))} \text{Axiom K} \\
\frac{\Box P(\rho)}{\Box P(\rho)^\dagger} \rightarrow_E \quad \frac{\Box \forall y. (G(y) \rightarrow \rho(y))}{\rho(\gamma) \rightarrow \Box \forall y. (G(y) \rightarrow \rho(y))} \rightarrow_I^4 \quad \frac{\rho(\gamma) \rightarrow \Box \forall y. (G(y) \rightarrow \rho(y))}{\forall \varphi. \varphi(\gamma) \rightarrow \Box \forall y. (G(y) \rightarrow \varphi(y))} \forall_I \\
\frac{G(\gamma)}{G(\gamma)} \quad 5 \quad \frac{G(\gamma) \wedge (\forall \varphi. \varphi(\gamma) \rightarrow \Box \forall y. (G(y) \rightarrow \varphi(y)))}{\dots \dots \dots} \wedge_I \\
\frac{G \text{ ess } \gamma}{G(\gamma) \rightarrow G \text{ ess } \gamma} \rightarrow_I^5 \quad \frac{G(\gamma) \rightarrow G \text{ ess } \gamma}{\forall x. G(x) \rightarrow G \text{ ess } x} \forall_I
\end{array}$$

Definition 3 *Necessary existence* of an individual is the necessary exemplification of all its essences:

$$E(x) \leftrightarrow \forall \varphi. [\varphi \text{ ess } x \rightarrow \Box \exists y. \varphi(y)]$$

Axiom 5 *Necessary existence is a positive property:*

$$P(E)$$

Lemma 1 *If there is a God-like being, then there is a God-like being necessarily:*

$$\exists z. G(z) \rightarrow \Box \exists x. G(x)$$

Proof

$$\begin{array}{c}
\frac{\overline{\exists z. G(z)} \quad 1}{G(\gamma)} \exists_E \quad \frac{\overline{\forall x. G(x) \rightarrow G \text{ ess } x}}{G(\gamma) \rightarrow G \text{ ess } \gamma} \forall_E \quad \frac{\overline{G \text{ ess } \gamma}^\dagger}{G \text{ ess } \gamma \rightarrow \Box \exists x. G(x)} \rightarrow_E \\
\frac{G \text{ ess } \gamma^\dagger}{\Box \exists x. G(x)} \rightarrow_E \quad \frac{\Box \exists x. G(x)}{\exists z. G(z) \rightarrow \Box \exists x. G(x)} \rightarrow_I^1 \\
\frac{\overline{\exists z. G(z)} \quad 1}{G(\gamma)} \exists_E \quad \frac{\overline{\forall \varphi. P(\varphi) \rightarrow \varphi(\gamma)}}{P(E) \rightarrow E(\gamma)} \forall_E \quad \frac{P(E)}{P(E)} \quad \text{Axiom 5} \\
\frac{P(E) \rightarrow E(\gamma)}{E(\gamma)^\dagger} \rightarrow_E \quad \frac{\overline{\forall \varphi. \varphi \text{ ess } \gamma \rightarrow \Box \exists x. \varphi(x)}}{G \text{ ess } \gamma \rightarrow \Box \exists x. G(x)} \forall_E \\
\frac{G \text{ ess } \gamma \rightarrow \Box \exists x. G(x)}{\Box \exists x. G(x)} \rightarrow_E
\end{array}$$

Lemma 2 *If the existence of a God-like being is possible, then it is necessary:*

$$\Diamond \exists z. G(z) \rightarrow \Box \exists x. G(x)$$

Proof

$$\begin{array}{c}
\frac{\overline{\omega : \exists x. G(x)} \quad 1}{\Diamond \exists x. G(x)} \quad \frac{\overline{\omega : \exists x. G(x) \rightarrow \Box \exists x. G(x)}}{\Box(\exists x. G(x) \rightarrow \Box \exists x. G(x))} \Box_I \quad \frac{\Box(\exists x. G(x) \rightarrow \Box \exists x. G(x))}{\Diamond \exists x. G(x) \rightarrow \Diamond \Box \exists x. G(x)} \text{Distribution Principle} \\
\frac{\Diamond \exists x. G(x)}{\Diamond \Box \exists x. G(x)^\dagger} \rightarrow_E \quad \frac{\Diamond \Box \exists x. G(x)^\dagger}{\Box \exists x. G(x)} \rightarrow_E \quad \frac{\Box \exists x. G(x)}{\Diamond \exists z. G(z) \rightarrow \Box \exists x. G(x)} \rightarrow_I^1 \\
\frac{\Box \exists x. G(x)}{\Diamond \exists z. G(z) \rightarrow \Box \exists x. G(x)} \rightarrow_I^1 \quad \frac{\overline{\Diamond \Box \exists x. G(x) \rightarrow \Box \exists x. G(x)}}{\Box \exists x. G(x)} \text{S5 Iteration Principle}
\end{array}$$

Theorem 3 *Necessarily, there exists a God-like being:*

$$\Box \exists x. G(x)$$

Proof

$$\frac{\frac{\text{Corollary 1}}{\diamond \exists x.G(x)} \quad \frac{\text{Lemma 2}}{\diamond \exists x.G(x) \rightarrow \Box \exists x.G(x)}}{\Box \exists x.G(x)} \rightarrow_E$$

5 A New Proof

In this section we present a new proof in **KB**. This proof uses exactly the same axioms and definitions of Scott’s proof shown in the previous sections, but it uses neither the axiom of reflexivity for equality nor **S5**’s modal iteration principle. Instead, it relies only on Brouwer’s reduction principle (for Theorem 3). The new proof is also shorter than Scott’s proof.

Axiom 1 *Either a property or its negation is positive, but not both.*

$$\forall \varphi. [P(\neg \varphi) \leftrightarrow \neg P(\varphi)]$$

Axiom 2 *A property necessarily implied by a positive property is positive:*

$$\forall\varphi.\forall\psi.[(P(\varphi) \wedge \Box\forall x.[\varphi(x) \rightarrow \psi(x)]) \rightarrow P(\psi)]$$

Theorem 1 *Positive properties are possibly exemplified:*

$$\forall \varphi. [P(\varphi) \rightarrow \Diamond \exists x. \varphi(x)]$$

Proof

$$\begin{array}{c}
\frac{\frac{\frac{}{\Box \neg \exists x. \rho(x)}{2} \quad \frac{\frac{}{\omega : \rho(x)}{1}}{\omega : \exists x. \rho(x)} \exists_I}{\omega : \neg \exists x. \rho(x)} \Box_E \quad \rightarrow_I}{\omega : \perp} \rightarrow_I^1 \\
\frac{\omega : \neg \rho(x)}{\omega : \rho(x) \rightarrow \neg \rho(x)} \rightarrow_I \\
\frac{\omega : \forall x. (\rho(x) \rightarrow \neg \rho(x))}{\Box \forall x. (\rho(x) \rightarrow \neg \rho(x))} \forall_I \\
\frac{P(\rho) \quad 3 \quad \Box \forall x. (\rho(x) \rightarrow \neg \rho(x))}{P(\rho) \wedge \Box \forall x. [\rho(x) \rightarrow \neg \rho(x)]} \Box_I \\
\frac{P(\rho) \wedge \Box \forall x. [\rho(x) \rightarrow \neg \rho(x)] \quad \wedge_I \quad \text{Axiom 2 for } \rho \text{ and } \lambda x. \neg \rho(x)}{P(\rho) \wedge \Box \forall x. [\rho(x) \rightarrow \neg \rho(x)] \rightarrow P(\lambda x. \neg \rho(x))} \rightarrow_E \\
\frac{[\star 5] \quad P(\lambda x. \neg \rho(x))}{\quad}
\end{array}$$

[illegible]

Comparing this new proof of Theorem 1 with the proof of Theorem 1 indicated by Gödel and Scott, it is noticeable that new proof is significantly shorter. The key idea to obtain a shorter proof is to instantiate the second universally quantified property of Axiom 2 by $\lambda x. \neg \rho(x)$ instead of using the predicate symbols $=$ or \neq as Gödel and Scott did. This eliminates the reliance of the ontological argument on the presence of $=$ and \neq in the logical language. Furthermore, Axiom 2 is then needed only once.

Although the simpler proof of Theorem 1 presented here was developed independently, it seems conceptually related to a possible simplification briefly described by Anderson (Anderson, 1990, footnote 2) and attributed by him to an anonymous referee. The simpler proof of Theorem 1 is not particularly special. In fact, it is probably the most likely proof that any well-trained logician would construct, if asked to derive Theorem 1 from Axioms 1 and 2 using methodical formal proof search techniques. Modern techniques usually rely on unification to compute or guess good instances for weakly quantified variables, and therefore they would never result in instantiations containing $=$ and \neq , which are symbols that do not appear anywhere in the axioms or in the theorems to be proved. We may wonder why Gödel preferred to use those unusual instantiations. It might be an interesting question from a historical point of view, as it might suggest that there was an evolution in the way logicians construct formal proofs; or from a philosophical point of view, if Gödel's preference was philosophically motivated.

Definition 1 A *God-like* being possesses all positive properties:

$$G(x) \leftrightarrow \forall \varphi. [P(\varphi) \rightarrow \varphi(x)]$$

Axiom 3 The property of being god-like is positive:

$$P(G)$$

Corollary 1 Possibly, a God-like being exists:

$$\Diamond \exists x. G(x)$$

Proof The proof of this corollary is identical to the proof of the same corollary shown in Section 4.

Axiom 4 Positive properties are necessarily positive:

$$\forall \varphi. [P(\varphi) \rightarrow \Box P(\varphi)]$$

Definition 2 An *essence* of an individual is a property possessed by it and necessarily implying any of its properties:

$$\varphi \text{ ess } x \leftrightarrow \varphi(x) \wedge \forall \psi. (\psi(x) \rightarrow \Box \forall x. (\varphi(x) \rightarrow \psi(x)))$$

At this point, Gödel and Scott proceed to prove Theorem 2, which is then used to prove Lemma 1. However, a close inspection of Scott's proof of Lemma 1 reveals that Theorem 2 is an unnecessary detour. Therefore, we do not include it in the new proof.

Definition 3 *Necessary existence* of an individual is the necessary exemplification of all its essences:

$$E(x) \leftrightarrow \forall \varphi. [\varphi \text{ ess } x \rightarrow \Box \exists y. \varphi(y)]$$

Axiom 5 *Necessary existence is a positive property:*

$$P(E)$$

This phenomenon is very intriguing: the new proof of Lemma 1 is technically simpler (because it has fewer inferences), but Scott’s proof of Lemma 1 with Theorem 2 can be considered intuitively simpler (because it is arguably easier to understand). This constitutes an interesting case for Hilbert’s 24th Problem (*Thiele*, 2003), which asks for criteria to properly compare the simplicity of proofs. While Hilbert had mathematical proofs in mind, insights into the 24th problem could be gained by analyzing philosophical proofs as well. As Gödel suggested, “a scientific (exact) philosophy and theology [...] is also most highly fruitful for science”.

$$\exists z.G(z) \rightarrow \Box \exists x.G(x)$$
[illegible]

Lemma 2 also turns out to be a superfluous detour and therefore the new proof does not include it. Inspecting Scott's proof of Lemma 2, we see that an important step is the derivation of $\Box\exists x.G(x)$ from $\Diamond\Box\exists x.G(x)$ and **S5**'s iteration principle. Instead, we can derive $\exists x.G(x)$ from $\Diamond\Box\exists x.G(x)$ and Brouwer's reduction theorem and then derive $\Box\exists x.G(x)$ from $\exists x.G(x)$ by the \Box_I (necessitation) inference rule. This is precisely what the new proof of Theorem 3 does.

Theorem 3 *Necessarily, there exists a God-like being:*

$$\Box\exists x.G(x)$$

$$\begin{array}{c}
\text{Proof} \quad \frac{\frac{\text{Distribution Principle}}{\frac{\Box[\exists x.G(x) \rightarrow \Box\exists x.G(x)] \rightarrow [\Diamond\exists x.G(x) \rightarrow \Diamond\Box\exists x.G(x)]}{\Diamond\exists x.G(x) \rightarrow \Diamond\Box\exists x.G(x)}} \quad \frac{\frac{\text{Lemma 1}}{\frac{\exists x.G(x) \rightarrow \Box\exists x.G(x)}{\Box[\exists x.G(x) \rightarrow \Box\exists x.G(x)]}} \Box_I}{\Box[\exists x.G(x) \rightarrow \Box\exists x.G(x)]} \rightarrow_E \\
\frac{\frac{\frac{\text{Corollary 1}}{\frac{[\Diamond\exists x.G(x) \rightarrow \Diamond\Box\exists x.G(x)]}{\Diamond\Box\exists x.G(x)}} \rightarrow_E \quad \frac{\text{Brouwer's Reduction Principle}}{\frac{\Diamond\Box\exists x.G(x) \rightarrow \exists x.G(x)}{\exists x.G(x)}} \rightarrow_E}{\frac{\exists x.G(x)}{\Box\exists x.G(x)}} \Box_I
\end{array}$$

Since the iteration principle requires the strong modal logic **S5** while Brouwer's reduction principle requires only the much weaker modal logic **KB**, the possibility of proving Theorem 3 using Brouwer's reduction principle instead of **S5**'s iteration principle is philosophically profound. There have been many discussions of what modal axioms are needed for the ontological argument. In their manuscripts, Gödel and Scott do not write explicitly which logic they use. However, a proof step in their notes clearly relies on **S5**'s *iteration principle*. Anderson (Anderson, 1990, footnote 5) cites several articles questioning the adequacy of **S5** for the ontological argument and conjectures that the ontological argument could be restricted to the weaker modal logic **B**. Sobel (Sobel, 2001)[p. 152] acknowledges Anderson's conjecture, claims that it is correct and tries to informally explain how his formal proofs could be modified to rely on **B** instead of **S5**. Nevertheless, neither Anderson nor Sobel presented any formal proof relying only on **B**. The new formal proof presented in this section relies on **KB**, which is weaker than **B**.

Curiously, in the new proof of Theorem 3, the actual existence of a God-like being ($\exists x.G(x)$) is proven *before* its necessity ($\Box\exists x.G(x)$). This contrasts with Gödel's proof, where the actual existence is never actually proven. It could, nevertheless, be derived as a trivial corollary by applying the modal axiom T.

6 Modal Collapse

A major criticism against Gödel's proof is that its axioms lead to the so-called *modal collapse* (Sobel, 1987): it is possible to prove that everything that is the case is so necessarily, and hence actuality, possibility and necessity coincide (Sobel, 2001)[Ch. 4, section 6, theorems 9 and 10]. That is: for all propositions A ,

$$A \leftrightarrow \Diamond A \leftrightarrow \Box A$$

Below we show natural deduction derivations of the modal collapse, thereby confirming that it holds⁵ for the axioms used in the previous sections. We show only the strongest implication above. The others are easy corollaries in modal logics where T holds.

⁵ It is well-known and uncontroversial that the modal collapse holds. We show a natural deduction proof here merely for the sake of self-containment and so that the proof becomes more accessible to readers who might be more familiarized with natural deduction than with Sobel's proof system.

$$\begin{array}{c}
A \rightarrow \Box A \\
\text{Theorem 2} \\
\frac{\forall y. [\overline{G(y)} \rightarrow \overline{G \text{ ess } y}]}{\forall y. [G(y) \rightarrow G(y) \wedge \forall \psi. (\psi(y) \rightarrow \Box \forall x. (G(x) \rightarrow \psi(x)))]} \text{ D2} \\
\frac{\forall y. [G(y) \rightarrow \forall \psi. (\psi(y) \rightarrow \Box \forall x. (G(x) \rightarrow \psi(x)))]}{\forall y. [G(y) \rightarrow (A \rightarrow \Box \forall x. (G(x) \rightarrow A))]} \\
\frac{\forall y. [G(y) \rightarrow (A \rightarrow \Box \forall x. (G(x) \rightarrow A))]}{[\star 8] \quad \exists y. G(y) \rightarrow (A \rightarrow \Box \forall x. (G(x) \rightarrow A))}
\end{array}$$

7 Anderson's Emendation in Natural Deduction

Axiom 1 (Emended) *If the negation of a property is positive, the property is not positive:*

$$\forall \varphi. [P(\neg \varphi) \rightarrow \neg P(\varphi)]$$

$$\forall\varphi.\forall\psi.[(P(\varphi) \wedge \Box\forall x.[\varphi(x) \rightarrow \psi(x)]) \rightarrow P(\psi)]$$
$$\forall \varphi. [P(\varphi) \rightarrow \Diamond \exists x. \varphi(x)]$$

16

Definition 1 (Emended) A *God-like* being necessarily possesses those and only those properties which are positive:

$$G(x) \leftrightarrow \forall \varphi. [P(\varphi) \leftrightarrow \Box \varphi(x)]$$

Axiom 3 (As in Scott's) *The property of being god-like is positive:*

$$P(G)$$

Corollary 1 *Possibly, a God-like being exists:*

$$\Diamond \exists x. G(x)$$

Proof This proof is omitted, because it is identical to the proof of the same corollary shown in Sections 4 and 5.

Axiom 4 (As in Scott's) *Positive properties are necessarily positive:*

$$\forall \varphi. [P(\varphi) \rightarrow \Box P(\varphi)]$$

Definition 2 (Emended) An *essence* of an individual is a property possessed by it and necessarily implying those and only those properties necessarily possessed by it:

$$\varphi \text{ ess } x \leftrightarrow \varphi(x) \wedge \forall \psi. (\Box \psi(x) \leftrightarrow \Box \forall x. (\varphi(x) \rightarrow \psi(x)))$$

Definition 3 (As in Scott's) *Necessary existence* of an individual is the necessary exemplification of all its essences:

$$E(x) \leftrightarrow \forall \varphi. [\varphi \text{ ess } x \rightarrow \Box \exists y. \varphi(y)]$$

Axiom 5 (As in Scott's) *Necessary existence is a positive property.*

$$P(E)$$

Because of Anderson's emendations to Axiom 1 and Definition 1, the proof of Lemma 1 has to be modified as well. The proof shown below uses the modal axiom T to eliminate the extra box modality introduced by Anderson in his emended Definition 1.

Lemma 1 *If there is a God-like being, then there is a God-like being necessarily:*

$$\exists z.G(z) \rightarrow \Box \exists x.G(x)$$

Proof

$$\frac{\frac{\frac{1}{\exists z.G(z)}}{[\star_9] \quad G(\gamma)} \exists_E}{\frac{\frac{\frac{\overline{\star_9} \quad \overline{G(\gamma)}}{\dots\dots\dots} D1}{\forall \varphi.[P(\lambda x.\varphi(x)) \leftrightarrow \Box \varphi(\gamma)]} \forall_E}{\frac{\frac{P(\lambda x.\rho(x)) \leftrightarrow \Box \rho(\gamma)}{P(\lambda x.\rho(x)) \rightarrow \Box \rho(\gamma)} \wedge_E}{\frac{}{P(\rho)} \rightarrow_E} \frac{\frac{\Box \rho(\gamma)}{}}{4} \frac{\text{Axiom 4}}{P(\rho) \rightarrow \Box P(\rho)} \forall_E}{[\star_{10}] \quad \Box P(\rho)} \rightarrow_E$$

which is the only part of the argument that depended on axioms 4 and 5 and definitions 2 and 3. Such a modified proof is shown below.

Lemma 1 *If there is a God-like being, then there is a God-like being necessarily:*

$$\exists z.G(z) \rightarrow \Box \exists x.G(x)$$

Proof

$$\frac{\frac{\frac{\overline{\exists z.G(z)}}{G(\gamma)} \exists_E \quad \frac{\overline{\forall \varphi.[P(\lambda x.\varphi(x)) \leftrightarrow \Box \varphi(\gamma)]}}{P(G) \leftrightarrow \Box G(\gamma)} \forall_E \quad \frac{\frac{P(G) \leftrightarrow \Box G(\gamma)}{P(G) \rightarrow \Box G(\gamma)} \wedge_E \quad \frac{\text{Axiom 3}}{P(G)} \rightarrow_E}{\Box G(\gamma)} \text{D1} \quad \frac{\frac{\frac{\overline{G(\gamma)}}{\exists x.G(x)} \exists_I \quad \frac{G(\gamma) \rightarrow \exists x.G(x)}{\Box[G(\gamma) \rightarrow \exists x.G(x)]} \rightarrow_I^1 \quad \frac{\Box[G(\gamma) \rightarrow \exists x.G(x)]}{\Box G(\gamma) \rightarrow \Box \exists x.G(x)} \Box_I \quad \frac{\Box G(\gamma) \rightarrow \Box \exists x.G(x)}{\Box \exists x.G(x)} \text{(Axiom K)} \rightarrow_E}{\frac{\Box \exists x.G(x)}{\exists z.G(z) \rightarrow \Box \exists x.G(x)} \rightarrow_I^2} \rightarrow_E$$

Proofs of other lemmas, theorems and corollaries remain the same as in the previous section. Theorem 1 is derived from axioms 1 and 2. Corollary 1 is derived from Axiom 3 and Theorem 1. Finally, Theorem 3 can be derived from Corollary 1 and Lemma 1, which, as shown above, depends only on Axiom 3 and Definition 1. Therefore, confirming Hájek's claim, axioms 4 and 5 and definitions 2 and 3 are superfluous: they are not needed to prove Theorem 3.

7.2 A Proof-Theoretical Explanation for the Immunity to Modal Collapse

Anderson's change in the definition of essence adds an extra box modality, which transforms the standard proof (shown in Section 6) of modal collapse (i.e. $\vdash A \rightarrow \Box A$) into a proof of a tautology ($\vdash \Box A \rightarrow \Box A$), as shown below.

$$\frac{\frac{\frac{\frac{\overline{\overline{\forall y.[G(y) \rightarrow G(y) \wedge \forall \psi.(\Box \psi(y) \leftrightarrow \Box \forall x.(G(x) \rightarrow \psi(x)))]}}{\overline{\forall y.[G(y) \rightarrow G(y) \wedge \forall \psi.(\Box \psi(y) \leftrightarrow \Box \forall x.(G(x) \rightarrow \psi(x)))]}} \text{D2} \quad \frac{\overline{\forall y.[G(y) \rightarrow G(y) \wedge \forall \psi.(\Box \psi(y) \leftrightarrow \Box \forall x.(G(x) \rightarrow \psi(x)))]}}{\overline{\forall y.[G(y) \rightarrow (\Box A \leftrightarrow \Box \forall x.(G(x) \rightarrow A))]}]} \text{Theorem 2} \quad \frac{\overline{\forall y.[G(y) \rightarrow (\Box A \leftrightarrow \Box \forall x.(G(x) \rightarrow A))]}]}{\overline{[\star 13] \quad \exists y.G(y) \rightarrow (\Box A \leftrightarrow \Box \forall x.(G(x) \rightarrow A))}} \text{Theorem 3} \quad \frac{\overline{\Box \exists y.G(y)}}{\overline{\exists y.G(y)}} \Box_E \quad \frac{\overline{\exists y.G(y) \rightarrow (\Box A \leftrightarrow \Box \forall x.(G(x) \rightarrow A))}}{\overline{\Box A \leftrightarrow \Box \forall x.(G(x) \rightarrow A)}} \star 13 \quad \frac{\overline{\Box A \leftrightarrow \Box \forall x.(G(x) \rightarrow A)}}{\overline{\Box A \leftrightarrow \Box (\exists x.G(x) \rightarrow A)}} \Box_E \quad \frac{\overline{\Box A \leftrightarrow \Box (\exists x.G(x) \rightarrow A)}}{\overline{\Box (\exists x.G(x) \rightarrow A)}} \Box_E \quad \frac{\overline{\Box (\exists x.G(x) \rightarrow A)}}{\overline{\omega : \exists x.G(x) \rightarrow A}} \Box_E \quad \frac{\overline{\omega : \exists x.G(x) \rightarrow A}}{\overline{\omega : A}} \Box_I \quad \frac{\overline{\omega : A}}{\overline{\Box A \rightarrow \Box A}} \rightarrow_I^1 \quad \frac{\overline{\Box A \rightarrow \Box A}}{\overline{\Box A \rightarrow \Box A}} \rightarrow_I^1$$

The exchange of the implication in Scott's definition of essence for an equivalence and the additional necessitation (box) modality before the property ψ (which is instantiated by $\lambda x.A$) prevents the implication of a necessary fact from its contingent counterpart, as seen in the second part of the proof above, because the hypothetical assumption of the contingent property A has to be exchanged for its necessitation $\Box A$ to preserve the proof structure.

8 Bjørdal's Variant in Natural Deduction

Bjørdal's variant (Bjørdal, 1999; Fuhrmann, 2005) is another solution to the Modal Collapse. In striking contrast to other variants, Bjørdal's axiomatization takes god-likeness (G_B) as a primitive notion, upon which the notion of positive property is defined. This variant also significantly reduces the number of necessary axioms. Scott's variants of Axioms 2, 3 and 4 become unnecessary to prove the final theorem from Bjørdal's modified definitions 1, 2 and 3 and axioms 1 and 5.

Definition 1 (Changed) A property is *positive* iff it is necessarily possessed by every God-like being.

$$P_B(\varphi) \equiv \Box \forall x. [G_B(x) \rightarrow \varphi(x)]$$

Bjørdal refines the notion of essence by restricting the attention to positive properties only. He calls this refined notion of essence a maximal composite.

Definition 2 (Changed) A *maximal composite* of an individual's positive properties is a positive property possessed by the individual and necessarily implying every positive property possessed by the individual.

$$MCP(\varphi, x) \equiv (\varphi(x) \wedge P_B(\varphi)) \wedge \forall \psi. ((\psi(x) \wedge P_B(\psi)) \rightarrow \Box \forall y. (\varphi(y) \rightarrow \psi(y)))$$

Definition 3 (Changed) *Necessary existence* of an individual is the necessary exemplification of all its maximal composites.

$$NE_B(x) \equiv \forall \varphi. (MCP(\varphi, x) \rightarrow \Box \exists y. \varphi(y))$$

Axiom 1 (Changed) If a property is positive, its negation is not positive:

$$\forall \varphi. [P_B(\varphi) \rightarrow \neg P_B(\neg \varphi)]$$

Axiom 5 *Necessary existence is a positive property.*

$$P_B(NE_B)$$

The proof of the possible instantiation of G_B is adjusted (as shown below) by proving Theorem 1 of section 5 directly for G_B and not for positive properties in general.

Theorem 1 *If God-likeness is a positive property, then it is possibly exemplified:*

$$P(G_B) \rightarrow \Diamond \exists x. G_B(x)$$

Proof

$$\begin{array}{c}
 \frac{\frac{\Box \neg \exists x. G_B(x)}{\omega : \neg \exists x. G_B(x)} \quad 2 \quad \frac{\omega : G_B(x)}{\omega : \exists x. G_B(x)} \quad 1}{\omega : \exists x. G_B(x)} \quad \Box_E \quad \exists_I \quad \rightarrow_E \\
 \frac{\omega : \perp}{\omega : \neg G_B(x)} \rightarrow_I^1 \\
 \frac{\omega : \neg G_B(x)}{\omega : G_B(x) \rightarrow \neg G_B(x)} \rightarrow_I \\
 \frac{\omega : G_B(x) \rightarrow \neg G_B(x)}{\omega : \forall x. (G_B(x) \rightarrow \neg G_B(x))} \quad \forall_I \\
 \frac{\omega : \forall x. (G_B(x) \rightarrow \neg G_B(x))}{\Box \forall x. (G_B(x) \rightarrow \neg G_B(x))} \quad \Box_I \\
 \frac{\Box \forall x. (G_B(x) \rightarrow \neg G_B(x))}{P(\neg G_B)} \quad D1 \\
 \frac{P(\neg G_B)}{\neg P(G_B)} \quad \rightarrow_E \\
 \frac{\neg P(G_B)}{P(G_B)} \quad 3 \quad \rightarrow_E \\
 \frac{\frac{\frac{\perp}{\neg \Box \neg \exists x. G_B(x)} \rightarrow_I^2}{\Diamond \exists x. G_B(x)} \rightarrow_I^3}{P(G_B) \rightarrow \Diamond \exists x. G_B(x)} \rightarrow_I^3
 \end{array}$$

Corollary 1 *Possibly, a God-like being exists:*

$$\Diamond \exists x. G(x)$$

$$\begin{array}{c}
\text{Proof} \quad \frac{\frac{\overline{\Box \forall x. [G_B(x) \rightarrow G_B(x)]}}{\dots\dots\dots} \text{D1} \quad \frac{\text{Theorem 1}}{P_B(G_B) \rightarrow \Diamond \exists x. G_B(x)} \wedge_I}{\frac{P_B(G_B)}{\Diamond \exists x. G_B(x)} \rightarrow_E}
\end{array}$$

Lemma 1 *If there is a God-like being, then there is a God-like being necessarily:*

$$\exists z.G_B(z) \rightarrow \Box \exists x.G_B(x)$$

Proof

[illegible]

9 Conclusions

The proof of section 5 shows explicitly that the weak modal system **KB** is sufficient for the ontological argument as opposed to the standard modal calculus **S5**. This weakening of the system is based on an explicit formalization and confirms the result briefly described by Anderson (*Anderson*, 1990, footnote 2). The weakening also applies to the emendation and variant of Anderson and Bjørdal respectively. The proofs are short and accessible to the general reader because of the choice of a natural deduction calculus.

References

- Adams, R. M. 1995. Introductory note to [Gödel 1970].
- Anderson, C. A. 1990. *Some Emendations of Gödel's Ontological Proof*. Faith and Philosophy, Vol. 7, Issue 3, pp. 291-303.
- Anderson, C. A. & Gettings, M. 1996. *Gödel's ontological proof revisited*. In: edited by Hajek P. *Gödel '96*, Springer.

- Bentert, M. & Benzmüller, C. & Streit, D. & Woltzenlogel Paleo, B. 2016. *Analysis of an Ontological Proof Proposed by Leibniz*. [to appear in] *Death and Anti-Death*, Volume 14: Four Decades after Michael Polanyi, Three Centuries after G. W. Leibniz. ed. Tandy, C. Ria University Press.
- Benzmüller, C. & Brown C. & Kohlhasse, M. In: *Journal of Symbolic Logic*, Vol. 69, N. 4, pp. 1027–1088.
- Benzmüller, C. & Weber, L. & Woltzenlogel Paleo, B. 2015. *Computer-Assisted Analysis of the Anderson-Hajek Ontological Controversy*. 1st World Congress on Logic and Religion.
- Benzmüller, C. & Weber, L. & Woltzenlogel Paleo, B. 2016. *Computer-Assisted Analysis of the Anderson-Hájek Ontological Controversy*. In: *Logica Universalis* 10 (2016)
- Benzmüller, C. & Woltzenlogel Paleo, B. 2013a. *Gödel's God on the Computer*. 10th International Workshop on the Implementation of Logics. pp. 1–2.
- Benzmüller, C. & Woltzenlogel Paleo, B. 2013b. *Gödel's God in Isabelle/HOL*. Archive of Formal Proofs.
- Benzmüller, C. & Woltzenlogel Paleo, B. 2014a. *On Logic Embeddings and Gödel's God*. Recent Trends in Algebraic Development Techniques.
- Benzmüller, C. & Woltzenlogel Paleo, B. 2014b. *On Logic Embeddings and Gödel's God*. 22nd International Workshop on Algebraic Development Techniques, pp. 8–9.
- Benzmüller, C. & Woltzenlogel Paleo, B. 2014c. *Formalization and Automated Verification of Gödel's Proof of God's Existence*. 4th World Congress on the Square of Opposition, pp. 24–25.
- Benzmüller, C. & Woltzenlogel Paleo, B. 2014d. *Automating Gödel's Ontological Proof of God's Existence with Higher-order Automated Theorem Provers*. ECAI 2014 - 21st European Conference on Artificial Intelligence, 18-22 August 2014, Prague, Czech Republic - Including Prestigious Applications of Intelligent Systems (PAIS 2014), pp. 93–98.
- Benzmüller, C. & Woltzenlogel Paleo, B. 2015a. *Interacting with Modal Logics in the Coq Proof Assistant*. Computer Science - Theory and Applications - 10th International Computer Science Symposium in Russia, CSR 2015, Listvyanka, Russia, July 13-17, 2015, Proceedings. pp. 398–411.
- Benzmüller, C. & Woltzenlogel Paleo, B. 2015b. *Higher-Order Modal Logics: Automation and Applications*. Reasoning Web. Web Logic Rules - 11th International Summer School 2015, Berlin, Germany, July 31 - August 4, 2015, Tutorial Lectures. pp. 32–74.
- Benzmüller, C. & Woltzenlogel Paleo, B. 2015c. *Experiments in Computational Metaphysics: Gödel's Proof of God's Existence*. Science and Spiritual Quest: Proceedings of the 9th All India Students' Conference on Science and Spiritual Quest (IIT Kharagpur, 30th October – 1st November). vol. 9. pp. 23–40.
- Benzmüller, C. & Woltzenlogel Paleo, B. 2016a. *The Inconsistency in Gödel's Ontological Argument: A Success Story for AI in Metaphysics*. International Joint Conference on Artificial Intelligence - IJCAI.
- Benzmüller, C. & Woltzenlogel Paleo, B. 2016b. *The Modal Collapse as a Collapse of the Modal Square of Opposition*. Studies in Universal Logic, Springer. pp. 1–7.
- Björda, F. 1999. Understanding Gödel's Ontological Argument, in Timothy Childers (ed.) *The Logica Yearbook 1998*, pp. 214-217, Filosofia.
- Blackburn, P. & M. de Rijke & Y. Venema. 2001. *Modal Logic*. Cambridge: Cambridge University Press.
- Church, A. 1940. *A Formulation of the Simple Theory of Types*, *Journal of Symbolic Logic*, 5: 56–68.
- Fitting, M. & Mendelsohn, R. 1998. *First-order Modal Logic*, Synthese Library vol. 277. Kluwer Academic Publishers.
- Fitting, M. 2002. *Types, Tableaus, and Gödel's God*, Kluwer Academic Publishers.
- Fuhrmann, A. 2005. Existenz und Notwendigkeit – Kurt Gödel's Axiomatische Theologie, in Olsson e.J., Schröder-Heister, P. and Spohn, W. (eds.) *Logik in der Philosophie*, pp. 349-374, Synchr.-Wissenschafts-Verlag.

- Gabbay, D. 1996. *Labelled Deductive Systems: volume I*. New York: Oxford Science publications.
- Gödel, K. Ontological Proof. In Kurt Gödel Collected Works vol. III. Ed. Feferman et al., pp. 403–404, 139, 145 or *Appendix A. Notes in Kurt Gödel's hand* in [Sobel 2001].
- Hajek, P. 1996. Magari and others on Gödel's Ontological Proof, in Ursini et alii (eds.) *Logic and Logical Algebra*, pp. 125–136, Marcel Dekker.
- Kant, I. original 1781. *Critique of Pure Reason*, J. M. Dent & Sons LTD, edition from 1959.
- Koons, R. 2006. Sobel on Gödel's Ontological Proof. *Philosophia Christi*, 2:235-248.
- Magari, R. 1988. *Logica e Teofilia*. In: *Notizie di Logica* VII.4.
- Muskens, R. 2006. *Higher Order Modal Logic*. In P. Blackburn, J.F.A.K. van Benthem, and F. Wolter, editors, *Handbook of Modal Logic*, Studies in Logic and Practical Reasoning, pages 621-653. Elsevier, Dordrecht, 2006.
- Nipkow, T. & Paulson, L.C. & Wenzel, M. 2002. *Isabelle/HOL: A Proof Assistant for Higher-Order Logic*. Number 2283 in LNCS, Springer.
- Oppenheimer, P. & Zalta, E. 2011. *A Computationally-Discovered Simplification of the Ontological Argument*. Australasian Journal of Philosophy, 89(2), 333-349.
- Paulin-Mohring, C. 2015. *Introduction to the calculus of inductive constructions*. In All about Proofs, Proofs for All, Mathematical Logic and Foundations. College Publications, London.
- Prawitz, D. 2006 (1st ed. 1965). *Natural deduction: a proof-theoretical study*. Mineola, New York: Dover publications.
- Rushby, J. 2013. *The Ontological Argument in PVS*. CAV Workshop "Fun with Formal Methods". St. Petersburg.
- Scott, D. *Appendix B. Notes in Dana Scott's hand* in *Logic and Theism: Arguments for and against Beliefs in God* by Sobel, J. H.
- Sobel, J. H. 1987. *Gödel's Ontological Proof*. In: J. J. Thompson (ed.). *On being and saying: essays for Richard Cartwright*, MIT Press.
- Sobel, J. H. 2001. *Logic and Theism: Arguments for and against Beliefs in God*, Cambridge University Press.
- Thiele, R. 2003. *Hilberts twenty-fourth problem*, American Mathematical Monthly, January.
- Wang, H. 1996. *A Logical Journey: From Gödel to Philosophy*, The MIT Press.
- Woltzenlogel Paleo, B. 2013. *Automated Verification and Reconstruction of Gödel's Proof of God's Existence*. Journal of the Austrian Computer Society. pp. 4–6.