

Variants of Gödel's Ontological Proof in a Natural Deduction Calculus

the date of receipt and acceptance should be inserted later

Abstract This paper presents detailed formalizations of ontological arguments in a simple modal natural deduction calculus. The first formal proof closely follows the hints in Scott's manuscript about Gödel's argument and fills in the gaps, thus verifying its correctness. The second formal proof improves the first one, by relying on the weaker modal logic **KB** instead of the stronger modal logic **S5**; and by avoiding the introduction of the equality predicate. The second proof is also technically shorter than the first one, because it eliminates unnecessary detours and uses Axiom 1 for the positivity of properties only once. The third and fourth proofs formalize, respectively, Anderson's and Bjørdal's variants of the ontological argument, which are known to be immune to modal collapse.

Keywords: Ontological Argument, Higher-Order Logics, Modal Logics, Natural Deduction.

“There is a scientific (exact) philosophy and theology, which deals with concepts of the highest abstractness; and this is also most highly fruitful for science. [...] Religions are, for the most part, bad; but religion is not.”
- Kurt Gödel [21][p. 316]

1 Introduction

Ontological arguments for the existence of God can be traced back at least to St. Anselm (1033-1109). His argument considers a greatest conceivable being, who must exist, because if it did not have the property of existence, then we could conceive of a greater being that, in addition to other properties, also had the property of existence. St. Anselm's argument was criticized and further developed by Descartes, Leibniz and Kant, among many others.

Leibniz identified the possible existence of God as a critical missing step in St. Anselm's argument. To fill this gap, he argued that the properties of God, the perfections, are compatible. This means that it is possible to satisfy all perfections at once, which implies that the existence of a greatest conceivable being with all these properties is possible.

Gödel built on Leibniz's work [1] and brought the ontological argument to a modern form using a modal logic with higher-order quantification over properties. In this setting, he gave precise axioms describing the notion of *positive* property and defined God as a being that has all positive properties. Gödel's work was saved in his own notes [9] as well as in notes by Scott [17], in whom he confided his proof.

The increase in formality of the ontological argument has required a development of its basic notions. Gödel's notion of positive property and Leibniz's notion of perfection differ. A formal distinction is that Leibniz's perfections are atomic whereas Gödel's positive properties can consist of combinations of atomic properties [7][p.139]. In particular, one of Gödel's axioms states that any conjunction of positive properties is itself positive. From this

axiom, it is immediately deduced that the property of being God-like is positive. Intuitively, a (possibly infinite) conjunction of positive properties is deduced from the universal definition of God-likeness. This deductive inference is not formalizable in a finite first-order calculus. The interplay between universal quantification (in the definition of a God-like being) and infinite conjunctions (in Gödel’s axiom for positive properties) could explain why, starting with Scott [17], this axiom of Gödel has been replaced by another that simply assumes the positivity of the property of being god-like.

The aim of this paper is to present two detailed formalizations of ontological arguments in a natural deduction calculus. For a comprehensive introduction to natural deduction, the reader can consult [16]. The higher-order modal natural deduction calculus proposed here has special introduction and elimination rules for modalities, as defined in Section 2, and is sound and complete relative to an axiomatic modal calculus, as proven in Section 2.1.

The natural deduction style was chosen for three reasons. Firstly, presentations of Gödel’s proof are typically either informal or formalized in other styles of proof calculi (e.g. Fitting’s tableaux [7] or Sobel’s sentential modal calculus [19]). Therefore, a formalization in natural deduction is a valuable complement to the existing presentations. Secondly, it makes the ontological proof accessible to people who are familiar with a natural deduction style. Thirdly, as natural deduction is the style used by proof assistants such as Coq [15] and Isabelle [14], natural deduction formalizations can be verified step-by-step in such proof assistants, and we have in fact done this.

The first main contribution of this paper is a detailed formalization of Scott’s version [17] of Gödel’s ontological argument [9] (as shown in Section 4) in the proposed natural deduction calculus. The second main contribution of the paper is a new proof (also in natural deduction style). In contrast to Scott’s proof [17], which requires the modal logic **S5**, the new proof requires only the weaker modal logic **KB**. The new proof also does not rely on the equality predicate and is much shorter.

A major criticism against Gödel’s argument is an undesirable consequence of the stipulated axioms, called *modal collapse*. This is discussed in greater detail in Section 6, where a natural deduction derivation of the collapse is presented. Many recent works on the ontological argument have proposed modifications of the argument that do not entail modal collapse. In this paper we present natural deduction formalizations of two of these solutions: Anderson’s emendation (in Section 7) and Bjørdal’s variant (in Section 8).

2 Natural Deduction

The language of higher-order modal logic used here is inspired by that of Church’s simple type theory [6].

Definition 1 *Simple types* are given by the following grammar:

$$\theta, \tau ::= \mu \mid o \mid \theta \rightarrow \tau$$

where μ is the atomic type for individuals, o is the atomic type for propositions and $\theta \rightarrow \tau$ is the type for functions taking an argument of type θ and returning something of type τ . ‘ \rightarrow ’ is assumed to be right associative.

Definition 2 *Terms and formulas* are given by the following grammar:

$$\begin{aligned} s, t ::= & p_\tau \mid X_\tau \mid (\lambda X_\theta. s_\tau)_{\theta \rightarrow \tau} \mid (s_{\theta \rightarrow \tau} t_\theta)_\tau \mid \\ & \perp_o \mid \rightarrow_{o \rightarrow o \rightarrow o} \mid \wedge_{o \rightarrow o \rightarrow o} \mid \vee_{o \rightarrow o \rightarrow o} \mid \\ & \forall_{(\tau \rightarrow o) \rightarrow o} \mid \exists_{(\tau \rightarrow o) \rightarrow o} \mid \Box_{o \rightarrow o} \mid \Diamond_{o \rightarrow o} \end{aligned}$$

where p_τ and X_τ range over, respectively, constants and variables of type τ . Parenthesis conventions, infix notation for propositional connectives and binding notation for quantifiers

are assumed. Furthermore, subscript types are omitted when they are clear from the context. Negation ($\neg_{o \rightarrow o}$) and equivalence ($\leftrightarrow_{o \rightarrow o \rightarrow o}$) are defined by $\neg A \equiv A \rightarrow \perp$ and $(A \leftrightarrow B) \equiv (A \rightarrow B) \wedge (B \rightarrow A)$.

The natural deduction calculus used here has standard rules for propositional connectives and quantifiers, as shown in Figures 1 and 3. The extension to classical logic is achieved by adding a rule for double negation elimination, shown in Figure 2. Finally, modal operators are handled by special rules that insert or remove formulas from boxes, as shown in Figure 4. Beta-reduction is implicit; all rules are assumed to operate modulo beta-reduction. A *derivation* is a directed acyclic graph whose nodes are formulas and whose edges correspond to applications of the inference rules. A *proof* of a formula F is a derivation without open assumptions and having F as root not inside any box.

Double lines are used to abbreviate tedious propositional reasoning steps in the derivations. Dashed lines are used to refer to an axiom or theorem with the proof shown elsewhere. When proof trees are too large to fit on the page, some branches may be displayed separately. In such cases, the conclusion of the branch and the location in the main proof tree where the branch belongs are annotated with the same symbol (a subscripted \star). Dotted lines are used to indicate folding and unfolding of definitions. Furthermore, as it is inconvenient to draw boxes around large derivations in L^AT_EX, formulas inside boxes are labeled with the names of the boxes surrounding them. Therefore, the boxes can be omitted without loss of information.

The calculus having only the rules shown in Figures 1, 2 and 3 is named **ND**, while the calculus with the additional rules shown in Figure 4 is named **ND_K**.

Fig. 1 Propositional rules

$$\begin{array}{c}
 \frac{\perp}{A} \perp_E \quad \frac{B}{A \rightarrow B} \rightarrow_I \quad \frac{\overline{A}^n \vdots B}{A \rightarrow B} \rightarrow_I^n \quad \frac{A \quad A \rightarrow B}{B} \rightarrow_E \\
 \\
 \frac{A \quad B}{A \wedge B} \wedge_I \quad \frac{A \wedge B}{A} \wedge_{E_1} \quad \frac{A \wedge B}{B} \wedge_{E_2}
 \end{array}$$

Fig. 2 Double negation elimination

$$\frac{\neg\neg A}{A} \neg\neg_E$$

2.1 Suitability for Rigid Higher-Order Modal Logic K

Adding the modal rules results in a calculus that is suitable for the basic modal logic **K**. In other words, **ND_K** is sound and complete relative to **ND** extended with axiom K ($\Box(A \rightarrow B) \rightarrow (\Box A \rightarrow \Box B)$) and the necessitation rule (which establishes that $\Box A$ is a theorem if A is a theorem).

Theorem 1 ***ND_K** is complete, relative to **ND** extended with axiom K and the necessitation rule.*

Fig. 3 Quantifier rules

$\frac{A[\alpha]}{\forall x_\tau. A[x]} \forall_I$	$\frac{\forall x_\tau. A[x]}{A[t]} \forall_E$	$\frac{A[t]}{\exists x_\tau. A[x]} \exists_I$	$\frac{\exists x_\tau. A[x]}{A[\beta]} \exists_E$
<p>eigen-variable conditions: if ρ is a \forall_I inference eliminating a variable α, then any occurrence of α in the proof should be an ancestor of the occurrence of α eliminated by ρ; if ρ is a \exists_E inference introducing a variable β, then any occurrence of β in the proof should be a descendant of the occurrence of β introduced by ρ.</p>			

Fig. 4 Rules for modal operators

$\frac{\omega : \boxed{\begin{smallmatrix} \vdots \\ A \end{smallmatrix}}}{\Box A} \Box_I$	$\frac{\Box A}{w : \boxed{\begin{smallmatrix} A \\ \vdots \end{smallmatrix}}} \Box_E$
$\frac{w : \boxed{\begin{smallmatrix} \vdots \\ A \end{smallmatrix}}}{\Diamond A} \Diamond_I$	$\frac{\Diamond A}{w : \boxed{\begin{smallmatrix} A \\ \vdots \end{smallmatrix}}} \Diamond_E$
<p>eigen-box condition: \Box_I and \Diamond_E are <i>strong</i> modal rules: ω must be a fresh name for the box they access (in analogy to the eigen-variable condition for strong quantifier rules). Every box must be accessed by <i>exactly one</i> strong modal inference.</p>	
<p>boxed assumption condition: assumptions should be discharged within the box where they are created.</p>	

Proof The necessitation rule can be immediately simulated with the \Box_I rule. Axiom K can be derived in **ND_K** as shown below:

$$\begin{array}{c}
 \frac{\frac{\Box(A \rightarrow B)}{\omega : A \rightarrow B} \Box_E \quad \frac{\Box A}{\omega : A} \Box_E}{\omega : B} \rightarrow_E \\
 \frac{\omega : B}{\Box B} \Box_I \\
 \frac{\Box B}{\Box A \rightarrow \Box B} \rightarrow_I^1 \\
 \frac{\Box A \rightarrow \Box B}{\Box(A \rightarrow B) \rightarrow (\Box A \rightarrow \Box B)} \rightarrow_I^2
 \end{array}$$

Theorem 2 **ND_K** is sound, relative to **ND** extended with axiom K and the necessitation rule.

Proof It is necessary to show that **ND_K** proofs of the following form can be translated to proofs in **ND** extended with the axiom K and the necessitation rule.

$$\frac{\frac{\Box A_1}{\omega : A_1} \Box_E \quad \frac{\Box A_n}{\omega : A_n} \Box_E \quad \vdots \quad \dots \quad \vdots}{\omega : B} \Box_I$$

A translation to **ND** extended with axiom K and necessitation is shown below for the case when $n = 1$:

$$\begin{array}{c}
\frac{}{A_1} 1 \\
\vdots \\
\frac{B}{A_1 \rightarrow B} \rightarrow_I^1 \\
\frac{}{\Box(A_1 \rightarrow B)} \text{ necessitation} \quad \frac{}{\Box(A_1 \rightarrow B) \rightarrow (\Box A_1 \rightarrow \Box B)} \text{ Axiom K} \\
\hline
\frac{\Box(A_1 \rightarrow B) \quad \Box(A_1 \rightarrow B) \rightarrow (\Box A_1 \rightarrow \Box B)}{\Box A_1 \rightarrow \Box B} \rightarrow_E \\
\hline
\frac{\Box A_1 \quad \Box A_1 \rightarrow \Box B}{\Box B} \rightarrow_E
\end{array}$$

For $n > 1$, the translation is a straightforward generalization:

$$\begin{array}{c}
\frac{}{A_1} 1 \quad \frac{}{A_n} n \\
\vdots \quad \dots \quad \vdots \\
\frac{B}{A_1 \rightarrow \dots \rightarrow A_n \rightarrow B} \rightarrow_I^* \\
\frac{}{\Box(A_1 \rightarrow \dots \rightarrow A_n \rightarrow B)} \text{ nec.} \quad \frac{}{\Box(A_1 \rightarrow \dots \rightarrow A_n \rightarrow B) \rightarrow (\Box A_1 \rightarrow \dots \rightarrow \Box A_n \rightarrow \Box B)} \text{ Axiom K, iterated} \\
\hline
\frac{\Box(A_1 \rightarrow \dots \rightarrow A_n \rightarrow B) \quad \Box(A_1 \rightarrow \dots \rightarrow A_n \rightarrow B) \rightarrow (\Box A_1 \rightarrow \dots \rightarrow \Box A_n \rightarrow \Box B)}{\Box A_1 \rightarrow \dots \rightarrow \Box A_n \rightarrow \Box B} \rightarrow_E \\
\hline
\frac{\Box A_1 \rightarrow \dots \rightarrow \Box A_n \rightarrow \Box B \quad \Box A_1 \quad \dots \quad \Box A_n}{\Box B} \rightarrow_E
\end{array}$$

Without the restriction that every box must be accessed by exactly one strong modal inference, the calculus would be unsound for the modal logic **K**. For example, the formula $\forall\phi.(\Box\phi \rightarrow \Diamond\phi)$ is not valid in **K** but would be provable without this restriction:

$$\begin{array}{c}
\frac{}{\Box\phi} 1 \\
\frac{}{\omega : \phi} \Box_E \\
\frac{}{\Diamond\phi} \Diamond_I \\
\frac{}{\Box\phi \rightarrow \Diamond\phi} \rightarrow_I^1 \\
\hline
\frac{}{\forall\phi.(\Box\phi \rightarrow \Diamond\phi)} \forall_I
\end{array}$$

This example proof is unsound according to the **ND_K** calculus, because the eigen-box condition is violated: the box labelled by ω is not accessed by any strong inference.

The straightforward combinations of the quantifier rules of **ND** either with the modal rules of **ND_K** or with axiom K and the necessitation rule are suitable for a higher-order modal logic where constants and variables are *rigid*. From the point of view of a *possible world semantics*, rigidity means that their interpretation is independent of the world in which they are being interpreted. Rigidity is tacitly assumed by most works investigating the ontological argument, and is explicitly assumed here. Nevertheless, it should be noted that its adequacy for the ontological argument has already been contested [7]. Another assumption made here is that the quantification domains are constant (i.e. independent of the possible worlds). Neither Gödel's manuscript nor Scott's manuscript reveal whether they use constant or varying domains, and this is also the case for many variants (e.g. [10]). Nevertheless, some authors of variants [2, footnotes 11 and 14] of Gödel's ontological argument do explicitly state a preference for varying domains. Our choice of assuming constant domains is motivated by simplicity.

3 Some Useful Derivable Modal Principles

From an axiomatic point of view, modal logics differ with respect to which additional axiom schemas they admit. Some common axiom schemas of relevance to the ontological argument are T ($\Box A \rightarrow A$), B ($A \rightarrow \Box\Diamond A$), 4 ($\Box A \rightarrow \Box\Box A$) and 5 ($\Diamond A \rightarrow \Box\Diamond A$). From a semantic point of view, these axioms correspond to geometric frame conditions that must be satisfied by the accessibility relation in Kripke models [13]: T corresponds to reflexivity, B corresponds to symmetry, 4 corresponds to transitivity and 5 corresponds to euclidianity. In the models

In this section, natural deduction proofs of three convenient derived modal principles are presented. These principles are used in the ontological proofs formalized in the next sections. Although they are well-known among modal logicians, we include their proofs here in order to have a self-contained and fully formal presentation of the ontological proofs, without any gaps.

Lemma (Distribution Principle)

Proof

Lemma (Brouwer's Reduction Principle)

Proof

6

Lemma (Iteration Principle)

$$\Diamond \Box A \rightarrow \Box A$$

Proof

$$\frac{\frac{\text{Brouwer's Reduction}}{\frac{\Diamond \Box A \rightarrow A}{\Box(\Diamond \Box A \rightarrow A)} \Box_I} \quad \frac{\frac{\text{K}}{\Box(\Diamond \Box A \rightarrow A) \rightarrow (\Box \Diamond \Box A \rightarrow \Box A)} \rightarrow_E \quad \frac{\frac{\text{Axiom 5 for } \Box A}{\Diamond \Box A \rightarrow \Box \Diamond \Box A} \quad \frac{1}{\Diamond \Box A} \rightarrow_E}{\Box \Diamond \Box A \rightarrow \Box A} \rightarrow_E \quad \frac{\Box A}{\Diamond \Box A \rightarrow \Box A} \rightarrow_I^1$$

4 Scott's Proof in Natural Deduction

In this section we present a detailed formalization of Scott's proof in the natural deduction calculus defined in Section 2. The reasoning in Scott's manuscript has been reproduced step-by-step and all reasoning gaps are filled in by using the deduction considered most natural. All derived formulas that do appear in Scott's manuscript are marked with a \dagger here. Unmarked formulas were derived in the process of filling the gaps between the marked formulas mentioned in Scott's manuscript.

Scott's version of Gödel's proof depends on five axioms that circumscribe the notion of *positive* property, with positivity being denoted by the undefined second-order predicate symbol P . Additionally, three definitions are used for establishing the notions of *God-like*, *essence* and *necessary existence*. Technically, these definitions only abbreviate certain complex formulas. The argument would still go through if all defined symbols were replaced by the complex formulas they define. This observation is particularly relevant in the case of *necessary existence*. Since this notion of "existence" is just an abbreviation, Gödel's argument is not susceptible to Kant's criticism against Anselm's argument (that existence should not be treated as a predicate). In Gödel's proof, existence is properly denoted by the existential quantifier. It is, therefore, unfortunately misleading to refer to the defined predicate symbol E as "necessary existence", when in fact it is just a convenient abbreviation.

In addition to the five axioms and three definitions that pertain specifically to the argument, the proof also uses **S5**'s iteration principle (in the proof of Lemma 2) and the equality axiom of reflexivity (in the proof of Theorem 1). The use of reflexivity could be avoided, but we chose to use it because it allows the construction of a shorter proof.

Axiom 1 *Either a property or its negation is positive, but not both:*

$$\forall \varphi. [P(\neg \varphi) \leftrightarrow \neg P(\varphi)]$$

Axiom 2 *A property necessarily implied by a positive property is positive:*

$$\forall \varphi. \forall \psi. [(P(\varphi) \wedge \Box \forall x. [\varphi(x) \rightarrow \psi(x)]) \rightarrow P(\psi)]$$

Theorem 1 *Positive properties are possibly exemplified:*

$$\forall \varphi. [P(\varphi) \rightarrow \Diamond \exists x. \varphi(x)]$$

Proof

$$\frac{\frac{\text{Reflexivity}}{\omega : \bar{\gamma} = \bar{\gamma}} \rightarrow_I \quad \frac{\omega : \rho(\gamma) \rightarrow \gamma = \gamma}{\omega : \forall x. \rho(x) \rightarrow x = x} \forall_I \quad \frac{P(\rho) \quad 5}{P(\rho) \wedge \Box(\forall x. \rho(x) \rightarrow x = x)} \Box_I \quad \frac{\Box(\forall x. \rho(x) \rightarrow x = x) \quad \wedge_I}{\frac{P(\rho) \wedge \Box(\forall x. \rho(x) \rightarrow x = x)}{P(\lambda x. x = x)} \wedge_I} \quad \frac{\text{Axiom 2 for } \rho \text{ and } \lambda x. x = x}{P(\rho) \wedge \Box(\forall x. \rho(x) \rightarrow x = x) \rightarrow P(\lambda x. x = x)} \rightarrow_E \quad \frac{P(\lambda x. x = x)}{P(\lambda x. x = x)} \dagger$$

Definition 2 An *essence* of an individual is a property possessed by it and necessarily implying any of its properties:

$$\varphi \text{ ess } x \leftrightarrow \varphi(x) \wedge \forall \psi. (\psi(x) \rightarrow \Box \forall x. (\varphi(x) \rightarrow \psi(x)))$$

Theorem 2 *Being god-like is an essence of any God-like being.*

$$\forall x.G(x) \rightarrow G \text{ ess } x$$

Proof

[illegible]

Definition 3 *Necessary existence* of an individual is the necessary exemplification of all its essences:

$$E(x) \leftrightarrow \forall \varphi. [\varphi \text{ ess } x \rightarrow \Box \exists y. \varphi(y)]$$

Axiom 5 *Necessary existence is a positive property:*

$$P(E)$$

Lemma 1 *If there is a God-like being, then there is a God-like being necessarily:*

$$\exists z.G(z) \rightarrow \Box \exists x.G(x)$$

Proof

$$\frac{\frac{\exists z.G(z)}{[\star_4]} 1}{G(\gamma)} \exists_E$$

$$\begin{array}{c}
\frac{\frac{\overline{\Box \neg \exists x. \rho(x)}}{2} \quad \frac{\overline{\omega : \rho(x)}}{1}}{\frac{\omega : \neg \exists x. \rho(x)}{\Box_E} \quad \frac{\omega : \exists x. \rho(x)}{\exists_I}} \rightarrow_E \\
\frac{\frac{\omega : \perp}{\rightarrow_I^1}}{\omega : \neg \rho(x)} \rightarrow_I \\
\frac{\omega : \rho(x) \rightarrow \neg \rho(x)}{\rightarrow_I} \forall_I \\
\frac{\omega : \forall x. (\rho(x) \rightarrow \neg \rho(x))}{\Box_I} \Box_I \\
\frac{P(\rho)}{3} \quad \frac{\Box \forall x. (\rho(x) \rightarrow \neg \rho(x))}{\wedge_I} \quad \frac{\text{Axiom 2 for } \rho \text{ and } \lambda x. \neg \rho(x)}{\overline{(P(\rho) \wedge \Box \forall x. [\rho(x) \rightarrow \neg \rho(x)]) \rightarrow P(\lambda x. \neg \rho(x))}} \rightarrow_E \\
\frac{P(\rho) \wedge \Box \forall x. [\rho(x) \rightarrow \neg \rho(x)]}{[\star_5] \quad P(\lambda x. \neg \rho(x))} \rightarrow_E \\
\frac{\frac{\text{Axiom 1}}{\forall \varphi. [P(\neg \varphi) \leftrightarrow \neg P(\varphi)]} \rightarrow_E}{\frac{P(\lambda x. \neg \rho(x)) \leftrightarrow \neg P(\rho)}{\wedge_E} \quad \frac{P(\lambda x. \neg \rho(x)) \rightarrow \neg P(\rho)}{\rightarrow_E} \quad \frac{P(\rho)}{3} \\
\frac{\neg P(\rho)}{\star_5} \quad \frac{\perp}{\neg \Box \neg \exists x. \rho(x)} \rightarrow_I^2 \\
\frac{\Diamond \exists x. \rho(x)}{P(\rho) \rightarrow \Diamond \exists x. \rho(x)} \rightarrow_I^3 \\
\frac{P(\rho) \rightarrow \Diamond \exists x. \rho(x)}{\forall \varphi. [P(\varphi) \rightarrow \Diamond \exists x. \varphi(x)]} \forall_I
\end{array}$$

Comparing this new proof of Theorem 1 with the proof of Theorem 1 indicated by Gödel and Scott, it is noticeable that new proof is significantly shorter. The key idea to obtain a shorter proof is to instantiate the second universally quantified property of Axiom 2 by $\lambda x. \neg \rho(x)$ instead of using the predicate symbols $=$ or \neq as Gödel and Scott did. This eliminates the reliance of the ontological argument on the presence of $=$ and \neq in the logical language. Furthermore, Axiom 2 is then needed only once.

Although the simpler proof of Theorem 1 presented here was developed independently, it seems conceptually related to a possible simplification briefly described by Anderson [2, footnote 2] and attributed by him to an anonymous referee. The simpler proof of Theorem 1 is not particularly special. In fact, it is probably the most likely proof that any well-trained logician would construct, if asked to derive Theorem 1 from Axioms 1 and 2 using methodical formal proof search techniques. Modern techniques usually rely on unification to compute or guess good instances for weakly quantified variables, and therefore they would never result in instantiations containing $=$ and \neq , which are symbols that do not appear anywhere in the axioms or in the theorems to be proved. We may wonder why Gödel preferred to use those unusual instantiations. It might be an interesting question from a historical point of view, as it might suggest that there was an evolution in the way logicians construct formal proofs; or from a philosophical point of view, if Gödel's preference was philosophically motivated.

Definition 1 A *God-like* being possesses all positive properties:

$$G(x) \leftrightarrow \forall \varphi. [P(\varphi) \rightarrow \varphi(x)]$$

Axiom 3 The property of being god-like is positive:

$$P(G)$$

Corollary 1 Possibly, a God-like being exists:

$$\Diamond \exists x. G(x)$$

[illegible]

Theorem 3 *Necessarily, there exists a God-like being:*

$$\begin{array}{c}
\text{Lemma 1} \\
\frac{\exists x. \bar{G}(x) \rightarrow \Box \exists x. \bar{G}(x)}{\Box [\exists x. \bar{G}(x) \rightarrow \Box \exists x. \bar{G}(x)]} \Box_I \\
\text{Distribution Principle} \\
\frac{\Box [\exists x. \bar{G}(x) \rightarrow \Box \exists x. \bar{G}(x)] \rightarrow [\Diamond \exists x. \bar{G}(x) \rightarrow \Diamond \Box \exists x. \bar{G}(x)]}{\Diamond \exists x. \bar{G}(x) \rightarrow \Diamond \Box \exists x. \bar{G}(x)} \rightarrow_E \\
\text{Corollary 1} \\
\frac{[\Diamond \exists x. \bar{G}(x) \rightarrow \Diamond \Box \exists x. \bar{G}(x)]}{\Diamond \Box \exists x. \bar{G}(x)} \rightarrow_E \\
\text{Brouwer's Reduction Principle} \\
\frac{\Diamond \Box \exists x. \bar{G}(x) \rightarrow \exists x. \bar{G}(x)}{\exists x. \bar{G}(x)} \rightarrow_E \\
\frac{\exists x. \bar{G}(x)}{\Box \exists x. \bar{G}(x)} \Box_I
\end{array}$$

13

Curiously, in the new proof of Theorem 3, the actual existence of a God-like being ($\exists x.G(x)$) is proven *before* its necessity ($\Box\exists x.G(x)$). This contrasts with Gödel's proof, where the actual existence is never actually proven. It could, nevertheless, be derived as a trivial corollary by applying the modal axiom T.

A major criticism against Gödel’s proof is that its axioms lead to the so-called *modal collapse* [18]: it is possible to prove that everything that is the case is so necessarily, and hence actuality, possibility and necessity coincide [19][Ch. 4, section 6, theorems 9 and 10]. That is: for all propositions A ,

Below we show natural deduction derivations of the modal collapse, thereby confirming that it holds¹ for the axioms used in the previous sections. We show only the strongest implication above. The others are easy corollaries in modal logics where T holds.

[illegible]

¹ It is well-known and uncontroversial that the modal collapse holds. We show a natural deduction proof here merely for the sake of self-containment and so that the proof becomes more accessible to readers who might be more familiarized with natural deduction than with Sobel's proof system.

property over which quantification is allowed might be too general and restrictions might be appropriate. Two alternative and more conservative solutions, requiring only modifications of the axioms and definitions but no change in the logic, are discussed in greater detail in the next section.

7 Anderson's Emendation in Natural Deduction

Anderson's solution [2] modifies the definitions of God-like being and essence, and eliminates half of Axiom 1 (replacing the equivalence by an implication).

Axiom 1 (Emended) *If the negation of a property is positive, the property is not positive:*

$$\forall\varphi.[P(\neg\varphi) \rightarrow \neg P(\varphi)]$$

Axiom 2 (As in Scott's) *A property necessarily implied by a positive property is positive:*

$$\forall\varphi.\forall\psi.[(P(\varphi) \wedge \Box\forall x.[\varphi(x) \rightarrow \psi(x)]) \rightarrow P(\psi)]$$

Theorem 1 *Positive properties are possibly exemplified:*

$$\forall\varphi.[P(\varphi) \rightarrow \Diamond\exists x.\varphi(x)]$$

Proof This proof is omitted, because any of the proofs of Theorem 1 in Sections 4 and 5 would suffice to prove this theorem with only small modifications: the equivalence of axiom 1 would not need to be reduced to an implication by conjunction elimination.

Definition 1 (Emended) A *God-like* being necessarily possesses those and only those properties which are positive:

$$G(x) \leftrightarrow \forall\varphi.[P(\varphi) \leftrightarrow \Box\varphi(x)]$$

Axiom 3 (As in Scott's) *The property of being god-like is positive:*

$$P(G)$$

Corollary 1 *Possibly, a God-like being exists:*

$$\Diamond\exists x.G(x)$$

Proof This proof is omitted, because it is identical to the proof of the same corollary shown in Sections 4 and 5.

Axiom 4 (As in Scott's) *Positive properties are necessarily positive:*

$$\forall\varphi.[P(\varphi) \rightarrow \Box P(\varphi)]$$

Definition 2 (Emended) An *essence* of an individual is a property possessed by it and necessarily implying those and only those properties necessarily possessed by it:

$$\varphi \text{ ess } x \leftrightarrow \varphi(x) \wedge \forall\psi.(\Box\psi(x) \leftrightarrow \Box\forall x.(\varphi(x) \rightarrow \psi(x)))$$

Definition 3 (As in Scott's) *Necessary existence* of an individual is the necessary exemplification of all its essences:

$$E(x) \leftrightarrow \forall\varphi.[\varphi \text{ ess } x \rightarrow \Box\exists y.\varphi(y)]$$

Axiom 5 (As in Scott's) *Necessary existence is a positive property:*

$$P(E)$$

ToDo: Proof of \star_7 is missing in the proof of lemma 1 below.

$$\exists z.G(z) \rightarrow \Box \exists x.G(x)$$
$$\frac{\overline{\exists z.G(z)}^1}{[\star_6] \quad G(\gamma)} \exists_E$$

Theorem 3 *Necessarily, there exists a God-like being:*

Proof This proof is identical to the proof of Theorem 3 shown in Section 5.

16

8 Bjørdal's Variant in Natural Deduction

Bjørdal's variant [5, 8] is another solution to the Modal Collapse. In striking contrast to other variants, Bjørdal's axiomatization takes god-likeness (G_B) as a primitive notion, upon which the notion of positive property is defined. This variant also significantly reduces the number of necessary axioms. Scott's variant of Axioms 2, 3 and 4 become unnecessary to prove the final theorem from Bjørdal's modified definitions 1, 2 and 3 and axioms 1 and 5.

Definition 1 (Changed) A property is *positive* iff it is necessarily possessed by every God-like being.

$$P_B(\varphi) \equiv \Box \forall x. [G_B(x) \rightarrow \varphi(x)]$$

Bjørdal refines the notion of essence (which he calls maximal composite) by restricting the attention to positive properties only. He calls this refined notion of essence a maximal composite.

Definition 2 (Changed) A *maximal composite* of an individual's positive properties is a positive property possessed by the individual and necessarily implying every positive property possessed by the individual.

$$MCP(\varphi, x) \equiv (\varphi(x) \wedge P_B(\varphi)) \wedge \forall \psi. ((\psi(x) \wedge P_B(\psi)) \rightarrow \Box \forall y. (\varphi(y) \rightarrow \psi(y)))$$

Definition 3 (Changed) *Necessary existence* of an individual is the necessary exemplification of all its maximal composites.

$$NE_B(x) \equiv \forall \varphi. (MCP(\varphi, x) \rightarrow \Box \exists y. \varphi(y))$$

Axiom 1 (Changed) *If a property is positive, its negation is not positive:*

$$\forall \varphi. [P_B(\varphi) \rightarrow \neg P_B(\neg \varphi)]$$

Axiom 5 *Necessary existence is a positive property.*

$$P_B(NE_B)$$

The proof of the possible instantiation of G_B is adjusted (as shown below) by proving Theorem 1 of section 5 directly for G_B and not for positive properties in general.

Theorem 1 *If God-likeness is a positive property, then it is possibly exemplified:*

$$P(G_B) \rightarrow \Diamond \exists x. G_B(x)$$

Proof

$$\begin{array}{c}
\frac{\frac{\overline{\Box \neg \exists x. G_B(x)}}{\omega : \neg \exists x. G_B(x)} 2 \quad \frac{\frac{\overline{\omega : G_B(x)}}{\omega : \exists x. G_B(x)} 1}{\omega : \exists x. G_B(x)} \exists_I}{\omega : \neg \exists x. G_B(x)} \Box_E \quad \frac{\omega : \neg \exists x. G_B(x)}{\omega : \neg G_B(x)} \rightarrow_I^1 \\
\frac{\omega : \neg G_B(x)}{\omega : G_B(x) \rightarrow \neg G_B(x)} \rightarrow_I \quad \frac{\omega : G_B(x) \rightarrow \neg G_B(x)}{\omega : \forall x. (G_B(x) \rightarrow \neg G_B(x))} \forall_I \\
\frac{\omega : \forall x. (G_B(x) \rightarrow \neg G_B(x))}{\Box \forall x. (G_B(x) \rightarrow \neg G_B(x))} \Box_I \quad \frac{\Box \forall x. (G_B(x) \rightarrow \neg G_B(x))}{P(\neg G_B)} \wedge_I \\
\frac{P(\neg G_B)}{\neg P(G_B)} \rightarrow_E \quad \frac{\text{Axiom 1}}{\forall \varphi. [P(\neg G_B) \rightarrow \neg P(G_B)]} \forall_E \\
\frac{\forall \varphi. [P(\neg G_B) \rightarrow \neg P(G_B)]}{P(\neg G_B(x)) \rightarrow \neg P(G_B)} \rightarrow_E \quad \frac{\neg P(G_B)}{P(G_B)} 3 \\
\frac{\frac{\frac{\overline{\perp}}{\neg \Box \neg \exists x. G_B(x)}}{\Diamond \exists x. G_B(x)} \rightarrow_I^2}{P(G_B) \rightarrow \Diamond \exists x. G_B(x)} \rightarrow_I^3
\end{array}$$

Corollary 1 *Possibly, a God-like being exists:*

$$\Diamond \exists x.G(x)$$

$$\text{Proof} \quad \frac{\frac{\overline{\overline{\Box \forall x.[G_B(x) \rightarrow G_B(x)]}}}{P_B(G_B)} \text{ D1} \quad \frac{\text{Theorem 1}}{P_B(G_B) \rightarrow \Diamond \exists x.G_B(x)} \wedge_I}{\Diamond \exists x.G_B(x)} \rightarrow_E$$

Lemma 1 *If there is a God-like being, then there is a God-like being necessarily:*

$$\exists z.G_B(z) \rightarrow \Box \exists x.G_B(x)$$

$$\begin{array}{c} \text{Proof} \quad \frac{\frac{\overline{\overline{\Box \forall x.[G_B(x) \rightarrow G_B(x)]}}}{P_B(G_B)} \text{ D1} \quad \frac{1}{(\psi(x) \wedge P_B(\psi))} \wedge_E}{\frac{G_B(x)}{G_B(x) \wedge P_B(G_B)} \wedge_I} \\ \frac{\frac{\frac{\frac{\frac{\overline{\overline{\Box \forall y.(G_B(y) \rightarrow \psi(y))}}}{P_B(\psi)} \text{ D1}}{(\psi(x) \wedge P_B(\psi)) \rightarrow \Box \forall y.(G_B(y) \rightarrow \psi(y))} \rightarrow_I^1}{\forall \psi.((\psi(x) \wedge P_B(\psi)) \rightarrow \Box \forall y.(G_B(y) \rightarrow \psi(y)))} \forall_I}{\frac{\overline{\overline{G_B(x) \wedge P_B(G_B)}}}{MCP(G_B, x)} \wedge_I} \\ \frac{\frac{\frac{\frac{\frac{\overline{\overline{\Box \forall x.[G_B(x) \rightarrow NE_B(x)]}}}{P_B(NE_B)} \text{ D1}}{\Box \forall x.[G_B(x) \rightarrow \forall \varphi.(MCP(\varphi, x) \rightarrow \Box \exists y.\varphi(y))]} \text{ D3}}{\frac{\forall x.[G_B(x) \rightarrow \forall \varphi.(MCP(\varphi, x) \rightarrow \Box \exists y.\varphi(y))]}{\Box_E} \quad \frac{2}{G_B(x)} \rightarrow_E}{\frac{\forall \varphi.(MCP(\varphi, x) \rightarrow \Box \exists y.\varphi(y))}{\forall_E} \quad \frac{\overline{\overline{MCP(G_B, x)}}}{\rightarrow_E} \\ \frac{\frac{\frac{\overline{\overline{\Box \exists y.G_B(y)}}}{G_B(x) \rightarrow \Box \exists y.G_B(y)} \rightarrow_I^2}{\frac{\overline{\overline{MCP(G_B, x)}}}{\rightarrow_E}} \end{array}$$

ToDo: Fix overfull hboxes

References

1. Adams, R. M. 1995. Introductory note to [Gödel 1970].
2. Anderson, C. A. 1990. *Some Emendations of Gödel's Ontological Proof*. Faith and Philosophy, Vol. 7, Issue 3, pp. 291-303.
3. Anderson, C. A. & Gettings, M. 1996. *Gödel's ontological proof revisited*. In: edited by Hajek P. *Gödel '96*, Springer.
4. Benz Müller, C. & Weber, L. & Woltzenlogel Paleo, B. 2016. *Computer-Assisted Analysis of the Anderson-Hájek Ontological Controversy*. In: *Logica Universalis* 10 (2016)
5. Bjørdal, F. 1999. Understanding Gödel's Ontological Argument, in Timothy Childers (ed.) *The Logica Yearbook 1998*, pp. 214-217, Filosofia.
6. Church, A. 1940. *A Formulation of the Simple Theory of Types*, Journal of Symbolic Logic, 5: 56-68.
7. Fitting, M. 2002. *Types, Tableaus, and Gödel's God*, Kluwer Academic Publishers.
8. Fuhrmann, A. 2005. Existenz und Notwendigkeit – Kurt Gödel's Axiomatische Theologie, in Olsson e.J., Schröder-Heister, P. and Spohn, W. (eds.) *Logik in der Philosophie*, pp. 349-374, Synchr.-Wissenschafts-Verlag.

9. Gödel, K. Ontological Proof. In Kurt Gödel Collected Works vol. III. Ed. Feferman et al., pp. 403–404, 139, 145 or *Appendix A. Notes in Kurt Gödel's hand* in [Sobel 2001].
10. Hajek, P. 1996. Magari and others on Gödel's Ontological Proof, in Ursini et alii (eds.) *Logic and Logical Algebra*, pp. 125–136, Marcel Dekker.
11. Kant, I. original 1781. *Critique of Pure Reason*, J. M. Dent & Sons LTD, edition from 1959.
12. Magari, R. 1988. *Logica e Teofilia*. In: *Notizie di Logica* VII.4.
13. Negri, S. 2005. *Proof analysis in modal logic*. J. Philosophical Logic, vol. 34, pp. 507–544.
14. Nipkow, T. & Paulson, L.C. & Wenzel, M. 2002. *Isabelle/HOL: A Proof Assistant for Higher-Order Logic*. Number 2283 in LNCS, Springer.
15. Paulin-Mohring, C. 2015. *Introduction to the calculus of inductive constructions*. In *All about Proofs, Proofs for All, Mathematical Logic and Foundations*. College Publications, London.
16. Prawitz, D. 2006 (1st ed. 1965). *Natural deduction: a proof-theoretical study*. Mineola, New York: Dover publications.
17. Scott, D. *Appendix B. Notes in Dana Scott's hand* in *Logic and Theism: Arguments for and against Beliefs in God* by Sobel, J. H.
18. Sobel, J. H. 1987. *Gödel's Ontological Proof*. In: J. J. Thompson (ed.). *On being and saying: essays for Richard Cartwright*, MIT Press.
19. Sobel, J. H. 2001. *Logic and Theism: Arguments for and against Beliefs in God*, Cambridge University Press.
20. Thiele, R. 2003. *Hilberts twenty-fourth problem*, American Mathematical Monthly, January.
21. Wang, H. 1996. *A Logical Journey: From Gödel to Philosophy*, The MIT Press.