
Patient name _____ Date _____

Lyme Disease

and other tickborne infections

Lyme disease is the most common illness spread by ticks in the U.S. An estimated **476,000 Americans** are diagnosed and treated for it each year. It is caused by *Borrelia* bacteria passed on by the bite of an infected blacklegged (deer) tick, and the same ticks can carry other infections at the same time. The best intervention is prevention, and the second best is early treatment.

HOW IT SPREADS



LYME DISEASE

Borrelia burgdorferi

IN GEORGIA


The lone star tick is the one people meet most. It can spread ehrlichiosis and trigger alpha-gal (red-meat) allergy.

OTHER INFECTIONS THE SAME TICKS CAN CARRY

Anaplasmosis · Babesiosis · Borrelia miyamotoi · Powassan virus

SYMPTOMS

EARLY: DAYS TO WEEKS **LATER: WEEKS TO MONTHS**



See a clinician promptly if a rash, fever, aches or joint swelling show up after a tick bite or time outdoors. Early antibiotic treatment works best.

Not everyone sees the tick. Nymphs are the size of a poppy seed, and many people never get the bull's-eye rash, which is why Lyme gets mistaken for flu, fibromyalgia or chronic fatigue.

After treatment, some people have fatigue, pain or brain fog that lingers for months. That recovery stretch is where food, sleep and the support on the back of this page come in.

KNOW YOUR TICKS

Blacklegged (deer) tick
Eastern U.S., Georgia included. The main Lyme carrier.

Western blacklegged
Pacific coast states. Also spreads Lyme.

Lone star tick
Common across the Southeast. Ehrlichiosis, alpha-gal allergy.

Lifestyle Recommendations

Avoiding tick bites

Clothing
Light colors, long sleeves, pants tucked into socks. Treat gear with permethrin.

Repellent
Use an EPA-registered repellent: DEET, picaridin, IR3535 or oil of lemon eucalyptus.

High heat
Tumble dry clothes on high for 10 minutes after outdoor time to kill ticks.

Shower and check
Shower within 2 hours. Check scalp, ears, armpits, waist and behind knees.

Pets
Check pets daily and ask your vet about tick prevention.

Remove it right
Fine-tipped tweezers, close to the skin, pull straight up. Note the date.

Nutrient Solutions

What your body needs, and the NutriDyn formula that delivers it

Protect your gut
probiotic + S. boulardii

START HERE



UltraBiotic Daily Multi-Strain
Practitioner exclusive

- S. boulardii keeps working during antibiotics
- Supports normal bowel function

IN EACH SERVING
30 billion CFU, seven strains: S. boulardii, L. reuteri, B. lactis, L. plantarum, L. acidophilus, L. gasseri, B. coagulans

1 capsule daily
2 hours from antibiotics

Immune readiness
beta-glucan



Immunity Pro
Practitioner exclusive

- Supports a healthy immune response
- One capsule a day

IN EACH SERVING
250 mg Wellmune beta-glucan, 85 mg FiberAid larch arabinogalactan

1 capsule daily

Inflammatory balance
curcumin



Curcumin 400X
Practitioner exclusive · softgels

- Supports a healthy inflammatory response
- Supports joint comfort

IN EACH SERVING
800 mg curcumin per 2 softgels, an enhanced-absorption form

2 softgels daily
with breakfast

Barrier support
serum-derived bovine immunoglobulins



GI Defend
Practitioner exclusive · powder

- Binds microbial toxins in the gut
- Supports immune balance

IN EACH SERVING
2.5 g ImmunoLin serum-derived bovine immunoglobulin per scoop

1 scoop daily
before breakfast

Cellular energy
mitochondrial support



Mito Recharge
90 capsules

- Supports energy production in every cell
- Antioxidant support

IN EACH SERVING
Per 3 capsules: 200 mg acetyl-L-carnitine, 100 mg NAC, 45 mg R-lipoic + 75 mg alpha lipoic acid, 100 mg resveratrol, 100 mg EGCG

3 capsules daily
with breakfast

YOUR DAILY RHYTHM

Morning
Immunity Pro 1 capsule
Curcumin 400X 2 softgels
GI Defend 1 scoop
Mito Recharge 3 capsules

Evening meal
nothing

Bedtime
UltraBiotic Daily Multi-Strain 1 capsule

The full plan is **7 pills and 1 scoop a day**. Overwhelmed? Start with the **Start here** pick, then add one product a week.

PHARMACIST CHECK

- **Doxycycline:** take it 2 to 3 hours apart from calcium, magnesium, iron and zinc, which block it.
- **During any antibiotic,** take the probiotic 2 hours apart from each dose, then keep going 2 weeks after.
- **Antibiotics can lower vitamin K and B vitamins.** Eat leafy greens and ask before adding herbs.

Special considerations for you:

Recommended by _____ Phone _____

◆ These statements have not been evaluated by the Food and Drug Administration. These products are not intended to diagnose, treat, cure, or prevent any disease. Nutrient support works alongside, never instead of, the care and

NutriDyn