

Your numbers caught this early.

That is the part working in your favor.

A 2026 systematic review and meta-analysis of 10 randomized trials in 599 adults (Viple et al., *Diabetologia*) tested time-restricted eating, meaning food kept inside a set number of hours each day. Fasting blood glucose came down. Then the researchers separated people by stage, and the result split: longer-term blood sugar control improved in the group still at the prediabetes stage, while in people who had already progressed to type 2 diabetes the same change did not reach statistical significance. Same approach, same four months. Being early is what made the difference.

WHY YOUR DOCTOR RECOMMENDED THIS

- **When you eat is its own lever.**

These trials did not hand people a new grocery list. They changed the hours during which food was eaten, and fasting blood glucose still came down across the pooled results. That makes this something you can start without overhauling your whole kitchen.

- **Being at this stage is an advantage you have right now.**

The improvement in longer-term blood sugar control showed up in the prediabetes group. In the group that had already progressed to type 2 diabetes, the same change did not reach statistical significance. Earlier is where the benefit appeared, and earlier is where you are.

- **The window is meant to fit your life.**

Across the ten trials the eating windows ranged from four to ten hours a day, and the studies ran about four months on average. That is a wide range on purpose. Pick the window you can actually keep, because the one you abandon in week three does nothing for you.

YOUR DAILY SUPPLEMENT

Gluco IR

NutriDyn | Practitioner Exclusive

Built around GlucoVantage® dihydroberberine, a form of berberine with up to five times greater bioavailability than the standard form, combined with Ceylon cinnamon, alpha lipoic acid, chromium, gymnema, and green tea extract. Formulated to promote healthy blood glucose levels already in the normal range and to provide antioxidant and metabolic support. ♦



HOW TO TAKE IT

Daily Dose: 5 capsules

Best Time: With a meal

With your window: Take it with food inside your eating hours

Duration: Daily, ongoing

Always: Follow your provider's guidance

KEY INGREDIENTS

- **Dihydroberberine 100 mg**
GlucoVantage®, the better-absorbed form of berberine, easier on the stomach than standard berberine.
- **Alpha Lipoic Acid 300 mg**
An antioxidant that supports how your cells handle glucose.
- **Ceylon Cinnamon 500 mg**
The true-cinnamon form, used traditionally for blood sugar support.
- **Chromium 600 mcg**
A mineral involved in how insulin moves sugar into your cells.
- **Gymnema sylvestre 200 mg**
A botanical long used to support healthy sugar metabolism.

WHAT TO EXPECT

Weeks 1 to 2

The window is the adjustment, not the food. The first days feel like a change of habit more than a diet. Keep water going through the fasting hours. Results vary by person.

Month 1 to 3

This is the stretch the research covered, about four months on average. Staying consistent here is exactly what the trials were measuring.

Long term

The goal is a pattern you keep without thinking about it, so your next visit reflects a habit and not a push before the appointment.

YOUR WINDOW AND WHAT GOES IN IT

Fewer Hours. Steadier Plate.

The eating window is one lever. What lands on the plate inside it is the other. A 2026 meta-analysis of 25 randomized trials in 1,973 people (Sun et al., *Current Diabetes Reviews*) found that low-glycemic-index eating reduced fasting blood glucose and HbA1c, with the effect strongest in metabolically high-risk groups and when people stayed with it. Low glycemic index simply means foods that raise blood sugar slowly instead of all at once. Use the window to decide *when*, and this page to decide *what*.

Food Group	Serving	Per Day	Best Choices
Non-Starchy Vegetables	1/2 to 1 cup	4 to 6	Leafy greens, broccoli, cauliflower, peppers, zucchini, green beans, asparagus. Fill half the plate here first. Almost no effect on blood sugar.
Protein	4 to 6 oz cooked	3 to 4	Chicken, turkey, fish, eggs, lean beef, Greek yogurt, cottage cheese, tofu, lentils. Protein blunts how fast a meal hits your blood sugar.
Healthy Fats	1 tsp to 1 Tbsp	2 to 3	Extra-virgin olive oil, avocado, nuts, seeds. Fat slows digestion, which flattens the spike. Avoid refined vegetable oils.
Slow Carbohydrates	1/2 cup cooked	1 to 2	Steel-cut oats, quinoa, barley, lentils, sweet potato, wild rice. Keep the portion to about a fist and always pair it with protein or fat.
Fruit	1 small or 1/2 cup	1 to 2	Berries, cherries, apple, pear, citrus. Whole fruit only. Juice delivers the sugar without the fiber that slows it down.
Dairy or Alternatives	6 to 8 oz	1 to 2	Plain Greek yogurt, kefir, unsweetened milk alternatives. Plain, not flavored. Flavored yogurt can carry as much sugar as dessert.
Fluids	8 oz	6 to 8+	Water, sparkling water, herbal tea, black coffee. These are fine during your fasting hours. Nothing with sugar in it.

SIMPLE SWAPS THAT LOWER THE SPIKE	
White bread	Sourdough or seeded rye
Instant oatmeal	Steel-cut oats
White rice	Quinoa or barley
White potato	Sweet potato
Fruit juice	Whole fruit
Flavored yogurt	Plain Greek yogurt
Crackers or chips	Nuts or olives
Cereal	Eggs

BUILDING YOUR EATING WINDOW	
Window length used in trials	4 to 10 hrs
A common starting point	10 hrs
Keep the same hours	most days
Water during fasting hours	always
Protein at your first meal	every day
Vegetables before starch	every meal

Window lengths describe what the trials tested. Your provider decides what is right for you, especially if you take blood sugar medication.

Talk to your provider first if: you take insulin or any blood-sugar-lowering medication, since changing when you eat can change how that medication affects you. Also avoid sugary drinks during fasting hours, which end the fast; skipping protein at your first meal; and stretching the window later and later on weekends until the pattern disappears.

Quality matters: Build every meal in the same order, vegetables and protein first, slow carbohydrate last. Take your Gluco IR with a meal inside your eating window, five capsules daily or as your practitioner directs.