

## **Week 3 Practice Problems**

**#1.** Write a program that takes the role of a store clerk. Ask a user to enter two floating point numbers: the cost of an item and the amount of money remitted to pay for the item. Then respond appropriately: calculate the change due to the customer or ask the customer for more money. For example:

Cost of the item: 3.56  
Amount tendered: 5.00  
Change: 1.44

or

Cost of the item: 3.56  
Amount tendered: 3.00  
Still due: 0.56

**#2.** Combine #1 above with #2 from Week 2: if in #1 you owe the customer some change, calculate the minimum number of \$100, \$50, \$20, \$10, \$5, toonies, loonies, quarters, dimes, nickels, and pennies needed to make up that amount. Modify the function from Week 2 Q#2 to print out the way to make the change.

**#3.** Your code for #2 doesn't work in Canada since we no longer have pennies. Fix it to work in Canada.

Hint: In stores when the amount isn't a multiple of 5, the store rounds to the nearest 5 cents. E.g., \$1.53 is rounded to \$1.55, \$2.47 is rounded to \$2.50, \$4.42 is rounded to \$4.40.

**#4.** Redo Week 2 #5 to use an if-statement. Write a function that takes a positive integer input less than 100000 and returns an integer corresponding to the number of digits in the number. Use this function in a program that prompts the user for an integer and prints the number of digits in the user's input.

**#5.** Water exists in three states- solid, liquid, and gas. Write a function that takes in the temperature in Celsius and returns a string "solid", "liquid", or "gas" depending on the

## APS106 - Practice Problems

2 of 2

temperature. Write a program using this function that prompts the user for a temperature and prints out the resulting state of water.

#6. In fluid mechanics the dimensionless Reynolds Number is defined below, where  $v$  is the speed of the fluid (m/s),  $L$  is the characteristic length of the flow situation (m), and  $n$  is the kinematic viscosity ( $\text{m}^2/\text{s}$ ).

$$\text{Reynolds Number} = vL/n$$

The magnitude of the Reynolds Number tells if a flowing fluid is moving in a laminar, transitional, or turbulent mode. Two flow situations are of interest, flow in a pipe and flow on a flat plate.

For a pipe:	<ul style="list-style-type: none"><li>• When the Reynolds number is less than 2000, then the flow is laminar.</li><li>• When the Reynolds number is greater than or equal to 2000 and less than or equal to 4000, then the flow is transitional.</li><li>• When the Reynolds number is greater than 4000, then the flow is turbulent.</li></ul>
For a plate:	<ul style="list-style-type: none"><li>• When the Reynolds number is less than <math>5 \times 10^5</math>, then the flow is laminar.</li><li>• When the Reynolds number is greater than or equal to <math>5 \times 10^5</math>, then the flow is turbulent.</li></ul>

Write a function that takes the speed of the fluid, the characteristic length, the kinematic viscosity, and the flow situation as arguments and returns a string indicating if the flow is "laminar", "transitional", or "turbulent".

You may assume that:

- the first three arguments are positive numbers in the correct units.
- the fourth argument is a string equal to "pipe" or "plate"