Course notes

Vector calculus

Chasles's relation $\overrightarrow{AB} + \overrightarrow{BC} = \overrightarrow{AC}$, the sum of vectors and the product by a real number extend to space. Two non-zero vectors are collinear if one is a multiple of the other. The zero vector is collinear with every vector. Three vectors are coplanar if they belong to the same vector plane. If $\vec{u}, \vec{v}$ are not collinear, $\vec{u}, \vec{v}, \vec{w}$ are coplanar if and only if $\vec{w} = \alpha\vec{u} + \beta\vec{v}$ for some real numbers α, β.

> **Key point · Line and plane.** The line through A with non-zero direction vector $\vec{u}$ is the set of points M such that $\overrightarrow{AM} = t\vec{u}$ for a real number t. The plane through A spanned by two non-collinear vectors $\vec{u}, \vec{v}$ is the set of points M such that $\overrightarrow{AM} = s\vec{u} + t\vec{v}$ for real numbers s, t.

Three non-collinear points define a plane. Four points A, B, C, D, with A, B, C not collinear, are coplanar if and only if $\overrightarrow{AD}$ is a linear combination of $\overrightarrow{AB}$ and $\overrightarrow{AC}$.

Relative positions

Two lines are coincident, strictly parallel, intersecting or non-coplanar (skew). Collinear direction vectors characterise parallel lines in the broad sense. Non-collinear direction vectors do not guarantee an intersection. A line and a plane: line contained in the plane, parallel and disjoint, or meeting it in a single point. Two planes: coincident, strictly parallel, or intersecting along a line.

Example in a cube. In $ABCDEFGH$, the faces $ABCD$ and $EFGH$ are parallel and the edges AE, BF, CG, DH correspond to one another. The lines (AB) and (CG) are neither parallel nor intersecting: they are non-coplanar.

Basis and coordinates

A basis of space is a family of three non-coplanar vectors. Every vector can be written uniquely in it as $x\vec{\imath} + y\vec{\jmath} + z\vec{k}$. A frame of reference consists of an origin O and a basis.

$$\overrightarrow{AB} = (x_B - x_A, y_B - y_A, z_B - z_A), \qquad I\left(\frac{x_A + x_B}{2}, \frac{y_A + y_B}{2}, \frac{z_A + z_B}{2}\right),$$

where I is the midpoint of $[AB]$. In an orthonormal frame, $\|\vec{u}\| = \sqrt{x^2 + y^2 + z^2}$ and $AB = \sqrt{(x_B - x_A)^2 + (y_B - y_A)^2 + (z_B - z_A)^2}$.

Example. In an orthonormal frame, for $A(1, 0, 2)$, $B(3, -1, 4)$, $\overrightarrow{AB} = (2, -1, 2)$, $AB = 3$ and the midpoint is $(2, -\frac{1}{2}, 3)$.

Method · **Testing a linear combination.** Write the three coordinate equalities. Determine the coefficients from two equations when possible, then check the third. A contradiction proves non-membership; do not discard the awkward equation.