CHAPTER 9

The exponential function and continuous models

Key point · Chapter path. Objectives. Use the algebraic properties; differentiate; study growth or decay; link the exponential to geometric sequences.

Prerequisites. Powers, geometric sequences, differentiation and sign.

Yearly reference point. 2–11 February: 8 hours

Diagnostic: six questions, 10 minutes

Answer without consulting the lesson. Mark the questions for which help is needed. This diagnostic is not graded.

1. Simplify $2^3 \times 2^4$.
2. Express 3^{-2}.
3. Solve $2x + 1 = 5$.
4. Differentiate $x^2 + 2x$.
5. What multiplicative factor corresponds to a 20% decrease?
6. For a geometric sequence $u_0 = 5, q = 2$, calculate u_n.

After the diagnostic. Revisit the flagged topics with the teacher; redo a similar question, then explain the rule used.

Speaking and logic. Explain the difference between a continuous rate and a discrete multiplicative factor; present an inequality proved by finding the minimum of a difference.

Investigate before formalising

Compare the increments of e^x and the values of the function around 0, with several positive steps. Conjecture a relation between $\dfrac{e^h - 1}{h}$ and 1 when h becomes small.


Organisation. Individual investigation 5 min, comparison in pairs 5 min, class discussion 5 min. Keep a record of one attempt, even an unfinished one. A conjecture must then be justified.

9.1 Defining a new function

Key point · Definition and results. The exponential function is the unique function f differentiable on $\mathbb{R}$ satisfying $f' = f$ and $f(0) = 1$. Its existence and uniqueness are accepted. It is written exp, then e^x with $e = \exp(1) \approx 2.71828$. The number e is a constant; x is the variable. The exponential is strictly positive.

Status and justification. Existence and uniqueness accepted in accordance with the programme. Positivity is deduced, after the algebraic properties accepted below, from $\exp(x) = \exp(\frac{x}{2})^2$ and from $\exp(x)\exp(-x) = 1$.

Example 1. $\exp(0) = 1$, $\exp(1) = e$, $\exp'(0) = 1$. The tangent at 0 is therefore $y = 1 + x$. The notation e^x extends the properties of powers to every real exponent.



e^x (blue) and its tangent $1 + x$ at 0 (green).

Watch out. $e^0 = 1$, not 0. The function never takes the value 0, even for negative x.

Application. What is the equation of the tangent to e^x at 1?

9.2 Algebraic properties and comparisons

Key point · Definition and results. For all real x,y, $e^{x+y} = e^x e^y$, $e^{-x} = \frac{1}{e^x}$, $e^{x-y} = \frac{e^x}{e^y}$. For integer n, $(e^x)^n = e^{nx}$. The function is strictly increasing on $\mathbb{R}$, so $e^u = e^v \iff u = v$ and $e^u < e^v \iff u < v$.

Status and justification. The addition relation is accepted in the common pathway; it can be explored further via uniqueness. Positivity and $\exp' = \exp > 0$ give strict increase. This justifies the equivalences for equations and inequalities.

Example 2. $e^{2x}e^{1-x} = e^{x+1}$. $e^{3x-2} = e^{x+4}$ is equivalent to $3x - 2 = x + 4$, so x=3. $e^{2x} \leq 1 = e^0$ is equivalent to $x \leq 0$.

Watch out. e^{x+y} is a product, not in general $e^x + e^y$. We do not solve $e^x = 2$ using a logarithm, which is not studied in Première.

Application. Simplify $\dfrac{e^{3x}}{e^{x-1}}$; solve $e^{x+2} > e^{2x}$.

9.3 Differentiating and studying

Key point · Definition and results. For real a, $t \mapsto e^{at}$ is differentiable and its derivative is ae^{at}. For $e^{at+b} = e^b e^{at}$, the derivative is ae^{at+b}. The product and quotient rules apply with the derivatives already known. The sign of the exponential often simplifies the study of the sign of the derivative.

Status and justification. The formula for differentiating e^{at} is accepted; the case with b follows by taking out the constant factor e^b. The general chain rule is not needed here.

Example 3. For $f(x) = (x+1)e^{-x}$, $f\prime(x) = e^{-x} - (x+1)e^{-x} = -xe^{-x}$. Since $e^{-x} > 0$, the derivative has the sign of $-x$: global maximum 1 at 0.

Watch out. Do not forget the factor a or the negative sign in e^{-x}. The exponential factor is positive, not necessarily increasing.


Application. Differentiate $g(t) = 3e^{-2t}$ and study its monotonicity.

9.4 From discrete to continuous

Key point · Definition and results. A model $Q(t) = Q_0 e^{kt}$, with $Q_0 > 0$, represents growth if $k > 0$, decay if $k < 0$, constancy if $k = 0$. For a fixed step h, $\dfrac{Q(t+h)}{Q(t)} = e^{kh}$. The values at integer times form a geometric sequence with common ratio e^k. The unit of k is the inverse of the unit of time.

Status and justification. The ratio follows from $e^{k(t+h)} = e^{kt}e^{kh}$. Monotonicity follows from $Q\prime(t) = kQ_0e^{kt}$ and the positivity of the remaining factor.

Example 4. $Q(t) = 100e^{-0.2t}$, t in hours: $Q(0) = 100$, $\dfrac{Q(t+1)}{Q(t)} = e^{-0.2}$. A simulation can look for the first whole number of hours at which Q falls below 50; the exact equation is not solved with a logarithm in this chapter.



The curves for $k = \frac{1}{2}$ and $k = -\frac{1}{2}$ are symmetric about the vertical axis. The line styles also distinguish them without colour.

Watch out. The relative rate per discrete unit of time is $e^k - 1$, not exactly k; they are close only for small k.

Application. For $Q(t) = 5e^{0.3t}$, express $Q(n)$ and $\dfrac{Q(t+2)}{Q(t)}$.

9.5 Proving an inequality

Key point · Definition and results. The inequality $e^x \geq 1 + x$, true for every real x, is obtained by studying the function $g(x) = e^x - 1 - x$: the curve of the exponential lies above its tangent at 0.

Status and justification. For $g(x) = e^x - 1 - x$, $g'(x) = e^x - 1$, negative if x<0 and positive if x>0. The minimum is $g(0) = 0$, which proves the inequality; equality only at 0.

Example 5. With $x = 0.1$, the inequality gives $e^{0.1} \geq 1.1$; the calculator displays $e^{0.1} \approx 1.10517$. With $x = -1$, it gives $e^{-1} \geq 0$, which is true but not very precise.

Beyond the syllabus · Euler's method. To approximate the solution of $y' = y$, $y(0) = 1$ with a step h, we replace the increment by hy: $y_{k+1} = (1 + h)y_k$. With $h = \dfrac{1}{n}$ on $[0, 1]$, we obtain $\left(1 + \dfrac{1}{n}\right)^n$, that is 2.25 for n=2 and about 2.59374 for n=10, to be compared with $e \approx 2.71828$. This method is not part of the première syllabus and is not assessed.

Watch out. A graph showing a curve above a line does not prove the inequality for all real numbers.

Application. Show that $e^{2x} \geq 1 + 2x$ for every real number x.

Exercise path

Start with exercises 1 and 2. Then choose one of exercises 3 or 4 and do it without help. Exercises 5 to 7 are a reserve for transfer, investigation or programming: the teacher designates one. Exercise 8 is revisited after a delay.

Exercise 9.1 · Transforming

For every real number x, consider the expressions

$$A = e^{x}e^{2-x}, \qquad B = \frac{(e^{2x})^{3}}{e^{x}}, \qquad C = e^{-x}(e^{x}+1).$$

1. Simplify A and B, justifying each grouping of exponents.
2. Expand then simplify C. Explain why a sum does not simplify in the same way as a product.

Exercise 9.2 · Equations and inequalities

Consider an equation and an inequality in the real unknown x:

$$e^{2x-1} = e^{x+3} \qquad \text{and} \qquad e^{-3x} \geq e^{x-2}.$$

1. Solve each in $\mathbb{R}$, citing the property of the exponential that allows the exponents to be compared.
2. While solving the inequality, explain at which step the direction of the inequality changes. Give the solution set as an interval.

Exercise 9.3 · Product and maximum

The function f is defined on $[0,4]$ by $f(x) = xe^{-x}$.

1. Calculate $f\prime(x)$, then study its sign on the interval.
2. Draw the monotonicity table of f. Deduce its maximum and the x-coordinate where it is reached.
3. Check that the values at the endpoints do not give a larger maximum. Keep the exact values.

Exercise 9.4 · A modelled population

The following two parts are independent.

Part A: A population. A population is modelled by $P(t) = 200e^{0.1t}$, where $t \geq 0$ is the time elapsed in years. $P(t)$ is a theoretical count, which is not necessarily an integer.

1. Determine the initial count, then the multiplicative factor between times t and $t+1$. Deduce the exact annual rate and express it as a percentage rounded to two decimal places.
2. Explain why this model cannot necessarily describe a real population indefinitely.
3. **Part B: Two curves.** In the same coordinate system, sketch the curves of $t \mapsto e^{\frac{t}{2}}$ and $t \mapsto e^{-\frac{t}{2}}$. State where they increase or decrease, their common point and their symmetry. Identify each curve clearly.

Exercise 9.5 · Searching for a threshold

The quantity of a substance, in milligrams, is modelled at time $t \geq 0$ by $Q(t) = 80e^{-\frac{t}{2}}$, where t is expressed in hours. Checks take place at integer times $n = 0, 1, 2, \ldots$. We look for the first check at which the quantity is strictly less than 10 mg.

1. Write an algorithm that searches for this first integer. Provide an iteration bound and state what the program returns if the bound is reached.
2. Run the algorithm, then check numerically the quantity at the index found and at the previous index.
3. Justify, using the decrease of Q, that this double check establishes that it is indeed the first index. Do not use a logarithm.

Exercise 9.6 · A symmetry

Consider the function f defined on $\mathbb{R}$ by $f(x) = e^x + e^{-x}$.

1. Study the parity of f and interpret the result for its curve.
2. Prove that $f(x) \geq 2$ for every real number x, choosing a method. State the equality case.
3. Suggest a check of the result using a second method, based on a square or a study of monotonicity.

Exercise 9.7 · Uniqueness under a hypothesis

Extension. Assume that the exponential function has already been constructed. Let f be a function differentiable on $\mathbb{R}$ that satisfies $f\prime(x) = f(x)$ for every real number x and $f(0) = 1$. Let $g(x) = f(x)e^{-x}$.

1. Calculate $g\prime$ and explain why g is constant on $\mathbb{R}$.
2. Determine this constant, then deduce the expression of f.
3. Does this proof establish the existence or the uniqueness of the function sought? Justify.

Exercise 9.8 · Delayed review

Define the sequence (u_n) by $u_n = e^{2n+1}$ for every natural number n.

1. Prove that this sequence is geometric, then state its first term and common ratio.
2. Calculate $u_0 + u_1 + u_2$ in exact form. Compare the direct calculation with the use of a sum formula.

Homework. Do only the task designated at the end of the session, in the dated progression: 10–20 min at most. The exercises depend on the topics already studied, so some questions are deferred. Revisit exercise 8 one week after the chapter (10 min). If the allotted time is exceeded, note the difficulty and bring your attempts; do not pile up the whole reserve.

To do in the CQFD textbook

These exercises are done in your textbook. The progression indicates the session; they replace the homework planned for that day (10 to 20 minutes in total).

CQFD, exercise 18 p. 187, questions a, b and c. Simplify the products and quotients, citing the property of exponentials used.

CQFD, exercise 30 p. 187, questions a, b and c. Solve the three inequalities and write the solution sets as intervals.

MCQ 1: understand and choose

Five single-answer questions for practice. Choose one answer per question; then explain one distractor. This chapter format does not reproduce the official EAM format.

1. $e^0 =$
   - A. 0
   - B. 1
   - C. e
   - D. -1
2. $e^x e^y =$
   - A. e^{x+y}
   - B. e^{xy}
   - C. $e^x + e^y$
   - D. e^{x-y}
3. Derivative of e^{-2x}:
   - A. e^{-2x}
   - B. $2e^{-2x}$
   - C. $-e^{-2x}$
   - D. $-2e^{-2x}$
4. $e^x = 1$ is equivalent to
   - A. $x = 1$
   - B. $x = e$
   - C. $x = 0$
   - D. no solution
5. The sign of e^{-x} is
   - A. negative if x>0
   - B. always strictly positive
   - C. zero at 0
   - D. variable

MCQ 2: understand and choose

Five single-answer questions for practice. Choose one answer per question; then explain one distractor. This chapter format does not reproduce the official EAM format.

1. $e^{2x} < e^3$ is equivalent to
   - A. $x < 3$
   - B. $x > \frac{3}{2}$
   - C. $x < \frac{3}{2}$
   - D. $x < \sqrt{3}$
2. $u_n = e^{3n}$ has common ratio
   - A. e^3

B. 3
C. e
D. $3e$

3. Tangent to e^x at 0:
   A. $y = x$
   B. $y = ex$
   C. $y = 1$
   D. $y = x + 1$
4. Derivative of xe^x:
   A. e^x
   B. $(x+1)e^x$
   C. xe^x
   D. xe^{x-1}
5. $e^x \geq 1 + x$ holds with equality
   A. for all x
   B. at 1 only
   C. at 0 only
   D. never

True or false? Justify.

1. $e^{2x} = 2e^x$ for all x.
2. The function $t \mapsto e^{-3t}$ is decreasing.
3. A continuous rate k=0.1 corresponds exactly to a 10% increase in one unit of time.

> **Key point · Self-assessment.** I keep the rules for powers; I know the positivity; I differentiate with the factor; I separate model, exact calculation and simulation.
>
> For each objective, indicate: achieved without help; achieved with help; to be reworked. Write one example and one mistake to avoid.