CHAPTER 8

Random variables and dispersion

Key point · Chapter path. Objectives. Build a distribution; interpret events; calculate expectation, variance, standard deviation; use linearity; simulate samples.

Prerequisites. Probability, weighted mean, squares, trees and lists.

Yearly reference point. 19–28 January: 8 hours

Diagnostic: six questions, 10 minutes

Answer without consulting the lesson. Mark the questions for which help is needed. This diagnostic is not graded.

1. Calculate the mean of 2, 4, 6.
2. Check whether $(\frac{1}{2}, \frac{1}{3}, \frac{1}{6})$ can be a distribution.
3. Calculate $(2-5)^2$.
4. If $P(A) = 0.6$, calculate $P(\overline{A})$.
5. A gross gain of 8 units and a stake of 3 give what net gain?
6. Calculate $\sqrt{0.25}$.

After the diagnostic. Revisit the flagged topics with the teacher; redo a similar question, then explain the rule used.

Speaking and logic. Explain why a theoretical mean can be an impossible gain. Present the difference between a distribution, a simulation and a data set.

Investigate before formalising

Create two games with three equally likely outcomes, both with expectation 0 but different standard deviations. Check their distributions and explain the difference.


Organisation. Individual investigation 5 min, comparison in pairs 5 min, class discussion 5 min. Keep a record of one attempt, even an unfinished one. A conjecture must then be justified.

8.1 Variable, event and distribution

Key point · Definition and results. On a finite sample space Ω, a real random variable X is a function that associates a real number with each outcome. The event $\{X = a\}$ groups the outcomes giving the value a. Its distribution associates with each distinct value x_i the probability $p_i = P(X = x_i)$, with $p_i \geq 0$ and $\sum p_i = 1$.

Status and justification. Definitions. The events associated with distinct values form a partition of the sample space; this justifies the sum of the probabilities.

Example 1. A fair die gives a gain X equal to 2 for 1 or 2, and to -1 otherwise. The distribution is $P(X = 2) = \frac{1}{3}$, $P(X = -1) = \frac{2}{3}$. $\{X \leq 0\} = \{3, 4, 5, 6\}$: probability $\frac{2}{3}$.



The height shows the probability: $P(X = -1) = \frac{2}{3}$ and $P(X = 2) = \frac{1}{3}$. The exact probabilities are written on the two bars.

Watch out. The values are not always equally likely even if the outcomes are. Group the outcomes that give the same gain.

Application. Two fair coins are tossed. X counts the heads. Write its distribution and $P(X \leq 1)$.

8.2 Expectation and linearity

Key point · Definition and results. The expectation is $E(X) = \sum_i x_i p_i$. It describes the long-term theoretical mean, not a result that is necessarily possible on a single trial. For variables X and Y on the same finite sample space and real numbers a,b: $E(aX + b) = aE(X) + b$ and $E(X + Y) = E(X) + E(Y)$, with no assumption of independence.

Status and justification. Summing over the outcomes ω, $E(X + Y) = \sum_\omega (X(\omega) + Y(\omega))P(\omega)$; distributing the sum gives the result. The constant b has expectation b because the probabilities sum to 1.

Example 2. For the previous distribution $P(X = 2) = \frac{1}{3}$, $P(X = -1) = \frac{2}{3}$, $E(X) = 0$: the game is fair if X is the net gain. For $Y = 3X - 2$, $E(Y) = 3 \times 0 - 2 = -2$.

Watch out. An expectation is not an individual guarantee; a game can be losing at every observed horizon despite a positive expectation.

Application. A gross gain is 10 with probability $\frac{1}{5}$ and 0 otherwise. What stake makes the game fair?

8.3 Variance, standard deviation and König–Huygens

Key point · Definition and results. Writing $\mu = E(X)$, $V(X) = E((X-\mu)^2) = \sum_i (x_i - \mu)^2 p_i$. The standard deviation is $\sigma(X) = \sqrt{V(X)}$. The König–Huygens formula gives $V(X) = E(X^2) - E(X)^2$. A variance is always positive or zero; the standard deviation has the same unit as X, the variance a squared unit.

Status and justification. Expand $(X-\mu)^2 = X^2 - 2\mu X + \mu^2$, apply linearity and $E(X) = \mu$: $E(X^2) - 2\mu^2 + \mu^2$. We obtain the formula.

Example 3. For X with values 0 and 4 with probabilities $\frac{3}{4}$ and $\frac{1}{4}$, $E(X) = 1$, $E(X^2) = 4$, $V(X) = 3$ and $\sigma = \sqrt{3}$. Direct calculation: $(0-1)^2(\frac{3}{4}) + (4-1)^2(\frac{1}{4}) = 3$.

Watch out. $E(X^2)$ is not $E(X)^2$. A variance calculated as negative signals an error or a very small rounding error to be dealt with explicitly.

Application. For X with values $-1, 1$ equally likely, calculate the expectation, variance and standard deviation.

8.4 Transforming and comparing gains

Key point · Definition and results. For real a,b, $V(aX+b) = a^2 V(X)$ and $\sigma(aX+b) = |a|\sigma(X)$. A translation moves the mean without changing the dispersion; a multiplication changes the scale. Two games with the same expectation can have very different dispersions.

Status and justification. $aX + b - E(aX+b) = a(X - E(X))$. Squaring and then taking the expectation gives $a^2 V(X)$. The square root of a^2 is $|a|$.

Example 4. A game A always pays 2. A game B pays 0 or 4 with equal probability. Both have expectation 2; variances 0 and 4. Their distributions are therefore not interchangeable despite the same mean.

Watch out. The variance of a sum is not, in general, the sum of the variances; no rule of this type is used without hypotheses.

Application. $E(X) = 3$, $V(X) = 4$. Give the characteristics of $Y = -2X + 5$.

8.5 Samples and observed mean

Key point · Definition and results. A sample of size n is here a list of n independent realisations of the same variable X. Its observed mean m varies from one sample to another. We compare m with $\mu = E(X)$; repetitions allow us to observe the proportion of samples satisfying $|m - \mu| \leq \dfrac{2\sigma}{\sqrt{n}}$. This threshold is an experimental task of the programme, not a claim of an exact coverage rate.

Status and justification. Experimentation: neither a convergence theorem nor the normal distribution is proved here. We check the size of the lists, the calculation of the mean and the distinction between n and the number N of samples.

Example 5. For X with values 0 and 2 equally likely, $\mu = 1$, $\sigma = 1$. For n=100, threshold $\dfrac{2}{10} = 0.2$: we count the means in $[0.8, 1.2]$. A result of 956 out of 1000 means an observed frequency of 0.956 for this simulation only.

Watch out. Do not copy a single draw n times; do not announce that the proportion is necessarily 95%.

Application. For $\mu = 5$, $\sigma = 3$ and n=36, give the experimental interval asked for.

Exercise path

Start with exercises 1 and 2. Then choose one of exercises 3 or 4 and do it without help. Exercises 5 to 7 are a reserve for transfer, investigation or programming: the teacher designates one. Exercise 8 is revisited after a delay.

Exercise 8.1 · A distribution to build

A fair six-sided die, with faces numbered 1 to 6, is rolled. The random variable X associates with the result the remainder of its Euclidean division by 3. For example, a result divisible by 3 gives $X = 0$.

1. For each of the six faces, state the value taken by X.
2. Present the probability distribution of X in a table and check that the probabilities sum to 1.
3. Calculate $P(X < 2)$, stating which values of X are retained.

Exercise 8.2 · Two variance calculations


The random variable X takes the values -2, 0 and 4 with respective probabilities $\frac{1}{4}$, $\frac{1}{2}$ and $\frac{1}{4}$.

1. Calculate its expectation $\mu = E(X)$ and the second moment $E(X^2)$.
2. Calculate $V(X)$ using the König–Huygens formula, then deduce the exact standard deviation.
3. Recover the variance from the definition $V(X) = E((X - \mu)^2)$. Present the three contributions to the calculation.

Exercise 8.3 · A fair game

A wheel has five sectors of equal angle, bearing the numbers 0, 0, 0, 5 and 10. The five sectors are assumed equally likely. The number obtained represents a gross gain G, in monetary units. To play, one pays a stake m; the net gain is therefore $X = G - m$.

1. Establish the distribution of G, then determine the stake m that makes the game fair, that is, such that $E(X) = 0$.
2. After studying variance, calculate the standard deviation of the gross gain. Deduce that of the net gain when the stake is fair.
3. Explain why changing the stake changes the expectation of the net gain but not its standard deviation.



The five sectors have the same angle. The three sectors marked 0 are distinct and must all be counted.

Exercise 8.4 · Same mean, different risk

Two fictitious investments are offered. The first yields a certain gain $X = 1$. The second yields a gain $Y = -2$ with probability $\frac{1}{2}$ and a gain $Y = 4$ with probability $\frac{1}{2}$. The gains are expressed in the same monetary unit.

1. Compare the expectations of the two gains, then their variances.
2. Someone states: "These investments are equivalent since they have the same mean." Explain what this sentence omits and what the variance makes it possible to distinguish here.

Exercise 8.5 · Linearity without independence

A fair die numbered 1 to 6 is rolled. Let X be the result of the roll and define $Y = 7 - X$. The two variables are therefore obtained from the same roll.

1. Calculate $E(X)$ and $E(Y)$, then $E(X + Y)$. Check the linearity of expectation on this example.
2. Are the variables X and Y independent? Justify using, for example, the events $\{X = 1\}$ and $\{Y = 6\}$.
3. Does the linearity of expectation require independence? Conclude using your results.

Exercise 8.6 · A minimal loss

A random variable X takes finitely many values and has an expectation μ. For a real number a, the mean squared error associated with the constant forecast a is measured by $E((X - a)^2)$.

1. Prove that $E((X - a)^2) = V(X) + (a - \mu)^2$. You may write $X - a = (X - \mu) + (\mu - a)$.
2. Deduce the value of a that minimises this error, then the minimum value. Justify the uniqueness of the optimal choice.

Exercise 8.7 · Letter frequencies

We wish to measure the frequency of the letter "a" in a text. The denominator is the number of alphabetic characters: spaces, digits and punctuation marks are excluded. Capital letters are treated as lower case; accented letters remain distinct from "a".

1. Write a Python function that returns this frequency. State the result chosen when the text contains no letters.
2. Test the function with `Aba !`, then with `123 !` and with the empty string. Explain the expected results before running the program.

Exercise 8.8 · Delayed review

An experiment whose probability of success is $\frac{1}{2}$ is repeated twice, independently. The random variable X counts the number of successes obtained.

1. Recover the distribution of X using the four possible sequences of results.
2. Calculate $E(X)$ and $V(X)$ starting from this distribution.

Homework. Do only the task designated at the end of the session, in the dated progression: 10–20 min at most. The exercises depend on the topics already studied, so some questions are deferred. Revisit exercise 8 one week after the chapter (10 min). If the allotted time is exceeded, note the difficulty and bring your attempts; do not pile up the whole reserve.

To do in the CQFD textbook

These exercises are done in your textbook. The progression indicates the session; they replace the homework planned for that day (10 to 20 minutes in total).

CQFD, exercise 28 p. 306, all questions. Calculate the expectation from the distribution table; keep the probabilities associated with each value.

CQFD, exercise 31 p. 306, all questions; use the distribution of exercise 28. Calculate the variance then the standard deviation, respecting the rounding to the nearest tenth asked for by the textbook.

MCQ 1: understand and choose

Five single-answer questions for practice. Choose one answer per question; then explain one distractor. This chapter format does not reproduce the official EAM format.

1. A probability distribution has a sum of probabilities equal to
   - A. 0
   - B. 1
   - C. n
   - D. the mean
2. X is 0 or 4 with probabilities $\frac{3}{4}$ and $\frac{1}{4}$; $E(X) =$
   - A. 1
   - B. 2
   - C. 3
   - D. 4
3. $V(X)$ equals
   - A. $E(X^2)$
   - B. $E(X)^2$
   - C. $E(X^2) + E(X)^2$
   - D. $E(X^2) - E(X)^2$
4. If $V(X) = 9$, $\sigma(X) =$
   - A. 81
   - B. 9
   - C. 3
   - D. $\frac{1}{3}$
5. $E(2X + 1)$ equals
   - A. $2E(X)$
   - B. $2E(X) + 1$
   - C. $E(X) + 1$
   - D. $E(X)^2 + 1$

MCQ 2: understand and choose

Five single-answer questions for practice. Choose one answer per question; then explain one distractor. This chapter format does not reproduce the official EAM format.

1. $V(X+5)$ equals
   A. $V(X)+5$
   B. $V(X)+25$
   C. $V(X)$
   D. $5V(X)$
2. $\sigma(-3X)$ equals
   A. $3\sigma(X)$
   B. $-3\sigma(X)$
   C. $9\sigma(X)$
   D. $\sigma(X)$
3. The expectation is necessarily
   A. a possible value
   B. an integer
   C. positive
   D. none of these three statements
4. $E(X+Y)=E(X)+E(Y)$ requires
   A. independence
   B. no independence
   C. X=Y
   D. X positive
5. For $\sigma=2$, n=100, $\frac{2\sigma}{\sqrt{n}}=$
   A. 4
   B. 0.04
   C. 0.4
   D. 2

True or false? Justify.

1. A variance can be negative.
2. Two variables with the same expectation have the same distribution.
3. Adding a constant to a variable does not change its standard deviation.

> **Key point · Self-assessment.** I build a complete distribution; I distinguish gross from net; I check both variance formulas; I do not confuse frequency with probability.
>
> For each objective, indicate: achieved without help; achieved with help; to be reworked. Write one example and one mistake to avoid.

Solutions and teaching notes for chapter 8

Teacher use. The answers below concern the tasks not worked in the lesson. First have students state their method; give graded help before the solution. The durations are estimates to be calibrated in class.

Diagnostic and remediation

1. 4. *Action:* Add, then divide by the number of values.
2. Yes: positive numbers with sum 1. *Action:* Put over a common denominator.
3. 9. *Action:* Keep the brackets before squaring.
4. 0.4. *Action:* Use the complement.
5. 5. *Action:* Write net gain = received minus paid.
6. 0.5. *Action:* Check the square of the answer.

Investigation activity

Gains $(-1, 0, 1)$ and $(-3, 0, 3)$: expectation 0 in both cases, variances $\frac{2}{3}$ and 6. In standard deviation, the second is three times more dispersed.

Immediate applications

1. Values 0, 1, 2; probabilities $\frac{1}{4}$, $\frac{1}{2}$, $\frac{1}{4}$. $P(X \leq 1) = \frac{3}{4}$.
2. $E(G) = \frac{10}{5} = 2$. For a stake m, $E(G - m) = 2 - m$; fair stake 2.
3. $E(X) = 0$, $E(X^2) = 1$, variance 1 and standard deviation 1.
4. $E(Y) = -1$, $V(Y) = 16$, $\sigma(Y) = 4$.
5. $\frac{2\sigma}{\sqrt{n}} = 1$; interval $[4, 6]$.

Exercises: method and results

Exercise 8.1 – A distribution to build. Values 0, 1, 2; each obtained from two faces, probabilities $\frac{1}{3}$. $P(X < 2) = \frac{2}{3}$. Beware the remainder 0 for 3 and 6.

Exercise 8.2 – Two variance calculations. $E(X) = -\frac{1}{2} + 1 = \frac{1}{2}$. $E(X^2) = 1 + 4 = 5$; $V = 5 - \frac{1}{4} = \frac{19}{4}$, $\sigma = \frac{\sqrt{19}}{2}$. Direct: $(\frac{25}{4})(\frac{1}{4}) + (\frac{1}{4})(\frac{1}{2}) + (\frac{49}{4})(\frac{1}{4}) = \frac{76}{16} = \frac{19}{4}$.

Exercise 8.3 – A fair game. $E(G) = \frac{15}{5} = 3$, so m=3. $E(G^2) = \frac{125}{5} = 25$; $V(G) = 25 - 9 = 16$, standard deviation 4, unchanged by the stake.

Exercise 8.4 – Same mean, different risk. $E(X) = E(Y) = 1$; $V(X) = 0$; $E(Y^2) = 10$, $V(Y) = 9$. X is certain, Y is dispersed. A player's preferences cannot be ranked from the expectation alone.

Exercise 8.5 – Linearity without independence. $E(X) = \dfrac{1+2+3+4+5+6}{6} = \dfrac{7}{2}$; $E(Y) = 7 - E(X) = \dfrac{7}{2}$. Since $X + Y = 7$, its expectation is 7, the sum of the two expectations. The events $\{X = 1\}$ and $\{Y = 6\}$ are identical: their intersection has probability $\dfrac{1}{6}$, whereas the product of their probabilities is $\dfrac{1}{36}$. The variables are not independent. Linearity remains valid without this hypothesis.

Exercise 8.6 – A minimal loss. Write $X - a = (X - \mu) + (\mu - a)$, expand, then use $E(X - \mu) = 0$. The square $(a - \mu)^2$ is minimal, zero only for $a = \mu$. The mean minimises the mean squared error.

Exercise 8.7 – Letter frequencies. Convert the text to lower case, keep the characters for which `isalpha()` is true, then count the "a"s. Divide this number by the number of letters if it is nonzero. For a text with no letter, return `None`, because the frequency is not defined. For `Aba !`, two "a"s out of three letters give $\dfrac{2}{3}$; the other two tests give `None`. The supplied program `frequence_a` follows this convention.

Exercise 8.8 – Delayed review. Distribution: $0, 1, 2$ with probabilities $\dfrac{1}{4}, \dfrac{1}{2}, \dfrac{1}{4}$. Expectation 1; second moment: $\dfrac{1}{2} + 1 = \dfrac{3}{2}$, variance $\dfrac{1}{2}$. The four paths are counted explicitly.

CQFD, exercise 28 p. 306, all questions. $E(X) = (-10) \times 0.35 + (-5) \times 0.15 + 0 \times 0.4 + 20 \times 0.1 = -2.25 = -\dfrac{9}{4}$. A negative expectation is compatible with a possible positive value.

CQFD, exercise 31 p. 306, all questions; use the distribution of exercise 28. $E(X^2) = 100 \times 0.35 + 25 \times 0.15 + 400 \times 0.1 = 78.75$. Thus $V(X) = 78.75 - (-2.25)^2 = 73.6875 = \dfrac{1179}{16}$, then $\sigma(X) = \dfrac{\sqrt{1179}}{4} \approx 8.6$. Do not round the expectation before calculating the variance.

MCQ: answers and reasons

MCQ 1.

1. B. Partition of the sample space.
2. A. $0 + \dfrac{4}{4}$.
3. D. König–Huygens.
4. C. Positive root.
5. B. Linearity.

MCQ 2.

1. C. Translation with no new dispersion.

2. A. Absolute value of the factor.
3. D. A variable can be negative or have a mean that is never attained.
4. B. A common finite sample space is enough.
5. C. $\frac{4}{10} = 0.4$.

True or false

1. False: it is a mean of squares with positive weights.
2. False: the constant 1 and an equally likely variable taking $-2, 4$ have the same mean.
3. True: the deviations from the new mean stay the same.

Differentiated teaching. For a technical block, offer only the first step (domain, factor or reference sample space), then withdraw the help. For an independent student, ask for a second method and the treatment of a limiting case. Orally, allow an initial reformulation in the language the student has mastered, then require a precise mathematical sentence in French.