CHAPTER 13

Conditional probability and the binomial distribution

S36 (revision), S37, S38, S39, S41

Key point · Objectives. Build a tree; choose a binomial distribution; calculate a cumulative probability and a threshold; interpret expected value and variance.

Course notes

Revision: conditioning and independence

If $P(A) > 0$, $P_A(B) = \frac{P(A \cap B)}{P(A)}$. A branch of a tree carries a conditional probability; the probability of a path is the product of its branches. For a partition $A_1, \ldots, A_r$, $P(B) = \sum_i P(A_i \cap B)$; if all the $P(A_i) > 0$, it is written $\sum_i P(A_i) P_{A_i}(B)$. Two events are independent when $P(A \cap B) = P(A)P(B)$. For $P(A) > 0$, this is equivalent to $P_A(B) = P(B)$. Do not confuse $P_A(B)$ and $P_B(A)$.

Example. $P(A) = 0.4$, $P_A(B) = 0.7$, $P_{\overline{A}}(B) = 0.2$. Then $P(B) = 0.4 \times 0.7 + 0.6 \times 0.2 = 0.4$ and $P_B(A) = \frac{0.28}{0.4} = 0.7$.

Bernoulli and the binomial distribution

A Bernoulli trial has two outcomes, success with probability p and failure with probability $1 - p$. The indicator variable B equals 1 in case of success and 0 otherwise: $E(B) = p$, $V(B) = p(1-p)$. A Bernoulli scheme is a repetition of n identical and independent trials. The number X of successes follows the binomial distribution $\mathcal{B}(n, p)$.

Key point · Distribution and parameters. For $k \in \{0, \ldots, n\}$,

$$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k}.$$

$$E(X) = np, \qquad V(X) = np(1-p), \qquad \sigma(X) = \sqrt{np(1-p)}.$$

Proof of the distribution. Each path containing k successes has probability $p^k(1-p)^{n-k}$ by independence. There are $\binom{n}{k}$ choices for the positions of the successes. These paths are mutually exclusive; add their probabilities.

Cumulative events

$$P(X \leq k) = \sum_{j=0}^{k} P(X = j), \qquad P(X \geq k) = 1 - P(X \leq k-1).$$

$$P(a \leq X \leq b) = P(X \leq b) - P(X \leq a-1).$$

For at least one success, $P(X \geq 1) = 1 - (1-p)^n$. The complement of "at least k" is "strictly fewer than k".

Example. For $X \sim \mathcal{B}(10, 0.2)$, $E(X) = 2$ and $V(X) = 1.6$. $P(X = 2) = 45(0.2)^2(0.8)^8 \simeq 0.3020$ and $P(X \geq 1) = 1 - 0.8^{10} \simeq 0.8926$. The expected value is neither a certain result nor necessarily the most likely one.

Threshold and interval

To find the smallest n such that $1 - (1 - p)^n \geq q$, a loop on n is enough. After taking logarithms, one can solve $(1 - p)^n \leq 1 - q$ then round in the right direction. To find the smallest integer k such that $P(X > k) \leq \alpha$, increase k until $P(X \leq k) \geq 1 - \alpha$ is obtained, then check the previous value. One can also look for an interval $[a, b]$ containing at least a prescribed probability, without assuming that it is unique.

Independent trials, beyond two outcomes

For independent experiments, the probability of a list of results is the product of the probabilities of its components. The sample space is the Cartesian product of the sample spaces of each experiment. For example, for two independent throws of a fair die, the sample space contains the 36 pairs (i, j), $1 \leq i, j \leq 6$, each with probability $\frac{1}{36}$. The probability that the sum is 5 is $\frac{4}{36} = \frac{1}{9}$, because the favourable pairs are $(1, 4), (2, 3), (3, 2), (4, 1)$. In a Bernoulli scheme, success is coded by 1 and failure by 0: the sample space is $\{0, 1\}^n$. Independence concerns the whole succession; a mere equality between two observed probabilities is not enough to prove it.

Research activity

Checking components — 40 min.

A component is chosen at random from a production coming from two workshops. Denote by A the event "the component comes from the first workshop", $B = \overline{A}$ and D the event "the component is defective". We are given $P(A) = 0.6$, $P_A(D) = 0.02$ and $P_B(D) = 0.05$. Approximate results will be rounded to 10^{-4}.

1. Represent the situation by a weighted tree, then calculate $P(D)$.
2. The component chosen is defective. Calculate the probability that it comes from the first workshop.
3. We now choose twenty components independently, each following the previous model. Denote by X the number of defective components among the twenty. Justify the distribution followed by X and state its parameters.
4. Calculate $P(X = 0)$, $P(X \geq 2)$ and $E(X)$.
5. If the twenty components were taken from a single batch, which assumption of the model would need to be re-examined? Explain why.

Progressive exercises

E1 · Practice. Consider two events A and B such that $P(A) = 0.3$, $P_A(B) = 0.8$ and $P_{\overline{A}}(B) = 0.1$. Calculate $P(B)$, then $P_B(A)$.

E2 · Practice. The random variable X follows the binomial distribution with parameters 8 and $\frac{1}{4}$. Give the exact values of $P(X = 3)$, $E(X)$ and $V(X)$.

E3 · Practice. Let X be a random variable taking integer values. For every integer k, set $F(k) = P(X \leq k)$. Express $P(X \geq 4)$ and $P(2 \leq X \leq 5)$ in terms of F.

E4 · Practice. Five cards are drawn in succession, without replacement, from a pack of 52 cards containing four aces. Let X be the number of aces obtained. Does the variable X follow a binomial distribution? Justify.

E5 · Practice. A trial whose probability of success is $\frac{1}{2}$ is repeated independently. Determine the smallest number of trials for which the probability of obtaining at least one success is greater than or equal to 0.99.

Textbook or past-paper work

Baccalauréat, Metropolitan France, 17 June 2025: exercise 1, question 5 (p. 3). In class after S41. Read the context of questions 1 to 4; for this work, accept the result $p = 0.0714$ from question 4. The paper is available here: official past paper. Use the data and instructions of this past paper.

Homework

Homework 13 — Reservations and overbooking — 60 min.

A hall has eighteen seats. The organiser accepts twenty reservations, paid in advance. Each person who has reserved turns up with probability 0.8, independently of the others. Denote by X the number of people who turn up. If more than eighteen people come, the surplus people are turned away.

1. Justify the distribution of X, then calculate its expected value and variance.
2. Calculate the probability that the organiser has to turn away at least one person. Round to 10^{-6}.
3. Each reservation brings in 10 monetary units, even if the person does not come. Each person turned away receives a total payment of 30 monetary units, refund and compensation included. Denote by $R = \max(X - 18, 0)$ the number of people turned away. Express the profit G in terms of R, then calculate $E(G)$.
4. Calculate the profit if the organiser accepts only eighteen reservations, with the same payment rules. Compare the two strategies in terms of mean profit.
5. State one assumption of the model that may be questionable in practice, then a criterion other than monetary profit to take into account in this decision.