Session 15 — Definition of the common logarithm

Common logarithm function

17 Nov 2026 • B: 2 h – C: 1 h

Session objectives

- Link powers of 10 and logarithms.
- Calculate common values.
- Use the calculator.

Fluency – 6 to 8 minutes, no calculator

1. Calculate 20 % of 80. .....
2. Give the domain of log. .....
3. Calculate $\log(10^4)$. .....
4. Solve $10^x = 0.01$. .....
5. Expand $\log(3b)$, $b > 0$. .....

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: link powers of 10 and logarithms and calculate common values.

Investigation

Complete the pairs $10^x = b$ and $x = \log(b)$ for $b = 0.001; 0.1; 1; 10; 1\,000$. Then use the calculator for $b = 7$.

Written record

For $b > 0$, $\log(b)$ is the unique solution of $10^x = b$. In particular $\log(1) = 0$, $\log(10) = 1$ and $\log(100) = 2$.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Practice – essential

1. Calculate $\log(10^2)$ and $\log(10^{-2})$.
2. Expand $\log\left(\dfrac{100a^2}{b}\right)$ for $a, b > 0$.
3. Solve $2^x = 22$ and give an approximate value to the nearest hundredth.

Textbook references

Further practice: Calao, course pp. 64–69; exercises p. 70, no. 1–5. The application exercise below is original; these references point to textbook exercises for extra practice and for its figures.

Application exercise

Solve the following equations in $\mathbb{R}$. When a solution exists, give its exact value and then its value rounded to the nearest thousandth.

a. $4 \times 10^x = 9.2$

b. $7 \times 10^x - 1.5 = 3 \times 10^x + 0.9$

c. $2.5 \times 10^x + 12 = 6 - 0.5 \times 10^x$

d. $10^{x+1} = 50$ (you may write $10^{x+1} = 10 \times 10^x$)

Homework

Redo practice exercise 2 with the following data: replace the initial value by 195 and the rate or common ratio by 5. Write it up in four lines: data, formula, calculation, conclusion.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1, 2 and 3.

Session 16 — Variations and graph

Common logarithm function

19 Nov 2026 • B: 1 h – C: 2 h

Session objectives

- Know the domain.
- Use the fact that the function is increasing.
- Read the curve.

Fluency – 6 to 8 minutes, no calculator


1. Write the multiplier for an increase of 25 %. .....
2. Calculate $\log(10^0)$. .....
3. Solve $10^x = 0.01$. .....
4. Expand $\log(3b)$, $b > 0$. .....
5. Give the domain of log. .....

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: know the domain and use the fact that the function is increasing.

Investigation

Plot $y = 10^x$ and the line $y = x$, then obtain the curve of the logarithm by symmetry. Deduce its domain, its direction of variation and its asymptote.



Written record

The function log is defined on $]0, +\infty[$, is strictly increasing, and its curve has $x = 0$ as a vertical asymptote.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Practice – essential

1. Calculate $\log(10^3)$ and $\log(10^{-3})$.
2. Expand $\log\left(\dfrac{100a^2}{b}\right)$ for $a, b > 0$.
3. Solve $3^x = 23$ and give an approximate value to the nearest hundredth.

Textbook references

Further practice: Calao, course pp. 64–69; exercises p. 70, no. 6–10. The application exercise below is original; these references point to textbook exercises for extra practice and for its figures.

Application exercise

Recall that the function log is strictly increasing on $]0;+\infty[$ and that $\log(1) = 0$.

1. Without a calculator, give the sign of each of the following numbers, with a justification:
   a. $\log(0.9)$;
   b. $\log\left(\dfrac{4}{3}\right)$;
   c. $\log(CM)$, where CM is the multiplier associated with an increase of $0.2\,\%$.
2. Without a calculator, compare $\log(0.35)$ and $\log(0.53)$.

Homework

Redo practice exercise 3 with the following data: replace the initial value by 200 and the rate or common ratio by 6. Write it up in four lines: data, formula, calculation, conclusion.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1 and 2.

Session 17 — Algebraic properties

Common logarithm function

24 Nov 2026 • B: 2 h – C: 1 h

Session objectives

- Expand or condense logarithms.
- Check the conditions.
- Avoid false properties.

Fluency – 6 to 8 minutes, no calculator

1. Simplify $3x + 6 - x + 2$. .....
2. Solve $10^x = 0.01$. .....
3. Expand $\log(3b)$, $b > 0$. .....
4. Give the domain of log. .....
5. Calculate $\log(10^1)$. .....

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: expand or condense logarithms and check the conditions.

Investigation

Starting from $10^2 \times 10^3 = 10^5$, conjecture $\log(ab) = \log a + \log b$. Test numerically with $a = 2, b = 5$, then explain why a, b must be positive.

Written record

For $a, b > 0$, $\log(ab) = \log a + \log b$ and $\log\left(\frac{a}{b}\right) = \log a - \log b$. Moreover $\log(a^n) = n \log a$.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Practice – essential

1. Calculate $\log(10^4)$ and $\log(10^{-4})$.
2. Expand $\log\left(\frac{100a^2}{b}\right)$ for $a, b > 0$.
3. Solve $4^x = 24$ and give an approximate value to the nearest hundredth.

Textbook references

Further practice: Calao, course pp. 66–69; exercises p. 72, no. 20–23. The application exercise below is original; these references point to textbook exercises for extra practice and for its figures.

Application exercise

Write each of the following numbers in the form $\log(A)$, where A is a strictly positive real number to be given as an integer or a fraction. When possible, give the exact value of the number.

a. $\log(6) + \log(5) - \log(3)$

b. $2\log(4) - \log(8)$

c. $\log(5) + \log(11)$

d. $2\log(3) - 3\log(2)$

e. $\log(40) - \log(8) + 2\log(2)$

Homework

Redo practice exercise 2 with the following data: replace the initial value by 205 and the rate or common ratio by 7. Write it up in four lines: data, formula, calculation, conclusion.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1, 2 and 3.

Session 18 — Solving $10^x = b$

Common logarithm function

26 Nov 2026 • B: 1 h – C: 2 h

Session objectives

- Solve $10^x = b$.
- Bracket a solution.
- Interpret an order of magnitude.

Fluency – 6 to 8 minutes, no calculator

1. Calculate 5 % of 140. .....
2. Expand $\log(3b)$, $b > 0$. .....
3. Give the domain of log. .....
4. Calculate $\log(10^2)$. .....
5. Solve $10^x = 0.01$. .....

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: solve $10^x = b$ and bracket a solution.

Investigation

A device measures intensities $b = 0.04; 3.2; 850$. For each value, solve $10^x = b$, give x to the nearest hundredth and check by calculating 10^x.

Written record

The equation $10^x = b$ has a solution only if $b > 0$; this solution is $x = \log(b)$.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Practice – essential

1. Calculate $\log(10^5)$ and $\log(10^{-5})$.
2. Expand $\log\left(\dfrac{100a^2}{b}\right)$ for $a, b > 0$.
3. Solve $5^x = 25$ and give an approximate value to the nearest hundredth.

Textbook references

Further practice: Calao, p. 73, no. 30–32. The application exercise below is original; these references point to textbook exercises for extra practice and for its figures.

Application exercise

A delivery company buys a van for 38 000 €. Its value is estimated to fall by 15 % each year. The company will sell the van as soon as its value becomes strictly less than 10 000 €. Let V be the value of the van (in euros) and N the number of years since the purchase.

```
V ← 38 000
N ← 0
While V .....
    V ← .....
    N ← N + 1
End While
```

1. Complete the algorithm so that, at the end of its execution, the variable N contains the number of years after which the van will be sold.
2. Using the calculator (table of values), run the algorithm and find the value of N obtained. Give the value of the van when it is sold, rounded to the nearest euro.
3. Write this program in Python, run it and compare with the result of question 2.

Homework

Redo practice exercise 3 with the following data: replace the initial value by 210 and the rate or common ratio by 2. Write it up in four lines: data, formula, calculation, conclusion.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1 and 2.

Session 19 — Solving $a^x = b$

Common logarithm function

1 Dec 2026 • B: 2 h – C: 1 h

Session objectives

- Solve $a^x = b$.
- Use a quotient of logarithms.
- Round sensibly.

Fluency – 6 to 8 minutes, no calculator

1. Write the multiplier for an increase of 10 %. .....
2. Give the domain of log. .....
3. Calculate $\log(10^3)$. .....
4. Solve $10^x = 0.01$. .....
5. Expand $\log(3b)$, $b > 0$. .....

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: solve $a^x = b$ and use a quotient of logarithms.

Investigation

The number of subscribers is multiplied by 1.15 each month. Determine after how many months it will have tripled. Set up the equation, solve it using logarithms and interpret the rounding.

Written record

For $a > 0$, $a \neq 1$ and $b > 0$, $a^x = b$ is equivalent to $x = \dfrac{\log(b)}{\log(a)}$.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Practice – essential

1. Calculate $\log(10^6)$ and $\log(10^{-6})$.
2. Expand $\log\left(\dfrac{100a^2}{b}\right)$ for $a, b > 0$.
3. Solve $6^x = 26$ and give an approximate value to the nearest hundredth.

Textbook references

Further practice: Calao, p. 74, no. 33–39. The application exercise below is original; these references point to textbook exercises for extra practice and for its figures.

Application exercise

An online shop A received 2 500 orders in January 2026. This number is assumed to increase by 7 % per month. Let a_n be the number of orders n months after January 2026.

1. a. Express a_n in terms of n.

   b. After how many months will the monthly number of orders have doubled?

   c. After how many months will it have quadrupled? Is this result twice the previous one? Explain.

2. A competing shop B received 4 000 orders in January 2026, with an increase of 3 % per month. From how many months on will shop A have more monthly orders than shop B? In which month will this happen?

Homework

Redo practice exercise 2 with the following data: replace the initial value by 215 and the rate or common ratio by 3. Write it up in four lines: data, formula, calculation, conclusion.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1, 2 and 3.

Session 20 — Equations $x^a = b$ and simple inequalities

Common logarithm function

3 Dec 2026 • B: 1 h – C: 2 h

Session objectives

- Solve equations where the variable is the base.
- Deal with an inequality.
- Take the domain into account.

Fluency – 6 to 8 minutes, no calculator

1. Simplify $3x + 9 - x + 2$. .....
2. Calculate $\log(10^4)$. .....
3. Solve $10^x = 0.01$. .....
4. Expand $\log(3b)$, $b > 0$. .....
5. Give the domain of log. .....

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: solve equations where the variable is the base and deal with an inequality.

Investigation

Solve on $[0, +\infty[$: $x^3 = 40$, then $x^4 = 16$. Then solve $1.08^x < 1.5$ and explain the role of the function being increasing.

Written record

To solve $x^n = b$, take the domain and the parity of n into account. For an exponential inequality, the direction depends on whether the function is increasing or decreasing.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Practice – essential

1. Calculate $\log(10^2)$ and $\log(10^{-2})$.
2. Expand $\log\left(\dfrac{100a^2}{b}\right)$ for $a, b > 0$.
3. Solve $2^x = 22$ and give an approximate value to the nearest hundredth.

Textbook references

Further practice: Calao, p. 75, no. 40–42. The application exercise below is original; these references point to textbook exercises for extra practice and for its figures.

Application exercise

The retail price of a solar panel model went from 420 € in March 2021 to 260 € in September 2025, a period of 4 years and 6 months. The price is assumed to have changed at a constant annual rate t.

1. Justify that $(1+t)^{4.5} = \dfrac{260}{420}$.
2. Deduce t, rounded to 0.1 %. Interpret the sign of the result.
3. A second model costs 395 € today. If its price changes at the same annual rate, what will its price be in 2 years and 3 months? Round to the nearest euro.
4. The manufacturer of this second model would rather reach a price of 300 € in 2 years and 3 months. What constant annual rate should it apply? Round to 0.1 %.

Homework

Redo practice exercise 3 with the following data: replace the initial value by 220 and the rate or common ratio by 4. Write it up in four lines: data, formula, calculation, conclusion.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1 and 2.

Session 21 — Problems in context

Common logarithm function

8 Dec 2026 • B: 2 h – C: 1 h

Session objectives

- Model a situation.
- Choose an equation.
- Conclude in context.

Fluency – 6 to 8 minutes, no calculator

1. Calculate 20 % of 60. .....
2. Solve $10^x = 0.01$. .....
3. Expand $\log(3b)$, $b > 0$. .....
4. Give the domain of log. .....
5. Calculate $\log(10^3)$. .....

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: model a situation and choose an equation.

Investigation

The sound level is $L = 10\log\left(\frac{I}{I_0}\right)$. Calculate L when $I = 10^6 I_0$. Then determine the ratio $\frac{I}{I_0}$ for $L = 85$ dB.

Written record

In a problem, define the variable, set up the equation, solve, check the domain and conclude with the unit. A numerical solution without interpretation is incomplete.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Practice – essential

1. Calculate $\log(10^3)$ and $\log(10^{-3})$.
2. Expand $\log\left(\frac{100a^2}{b}\right)$ for $a, b > 0$.
3. Solve $3^x = 23$ and give an approximate value to the nearest hundredth.

Textbook references

Further practice: Calao, p. 74, no. 38–39, then p. 75, no. 40–42. The application exercise below is original; these references point to textbook exercises for extra practice and for its figures.

Application exercise

In a laboratory, a cell culture grows by successive divisions: at each generation, the number of cells doubles. When the number of cells goes from N_0 to N, the number of generations that have elapsed is

$$g = \frac{\log\left(\dfrac{N}{N_0}\right)}{\log(2)}.$$

For example, going from 1 to 8 cells corresponds to $g = \dfrac{\log(8)}{\log(2)} = 3$ generations.

1. A culture goes from 1 000 to 64 000 cells. How many generations have elapsed?
2. Another culture goes from 500 to 128 000 cells. Same question.
3. A culture of 1 000 cells develops for 9.5 generations. About how many cells does it contain?
4. Starting from a single cell, how many cells are obtained after 20 generations?
5. A laboratory starts from 1 000 cells and wants to obtain 1 000 000 000. How many generations are needed? Given that one generation lasts 30 minutes, how much time should be planned?

Homework

Redo practice exercise 2 with the following data: replace the initial value by 225 and the rate or common ratio by 5. Write it up in four lines: data, formula, calculation, conclusion.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1, 2 and 3.

Session 22 — Review and assessment

Common logarithm function

10 Dec 2026 • B: 1 h – C: 2 h

Session objectives

- Summarise the methods.
- Identify the conditions.
- Correct mistakes.

Fluency – 6 to 8 minutes, no calculator

1. Write the multiplier for an increase of 25 %. .....
2. Expand $\log(3b)$, $b > 0$. .....
3. Give the domain of log. .....
4. Calculate $\log(10^4)$. .....
5. Solve $10^x = 0.01$. .....

Reminder and prerequisites

Reread the last written record and be able to carry out the basic calculations needed today. Before the investigation, have students rephrase the key words: summarise the methods and identify the conditions.

Investigation

Make a map of methods: domain; properties; equations $10^x = b$; equations $a^x = b$; inequalities. Attach a condition and an example to each branch.

Written record

Before calculating a logarithm: check positivity. Before a comparison: identify the direction of variation. Before a conclusion: check that the result is plausible.

Worked example

Revisit the data from the investigation and write up the solution in order: definition of the objects, property used, exact or approximate calculation, then a concluding sentence. The model answer appears in the teacher edition.

Practice – essential

1. Calculate $\log(10^4)$ and $\log(10^{-4})$.
2. Expand $\log\left(\dfrac{100a^2}{b}\right)$ for $a, b > 0$.
3. Solve $4^x = 24$ and give an approximate value to the nearest hundredth.

Textbook references

Further practice: Calao, pp. 76–79. The application exercise below is original; these references point to textbook exercises for extra practice and for its figures.

Application exercise

We look for a method to find the number of digits of a natural number $N \geqslant 1$ without writing it out in full. The integer part of a real number $y \geqslant 0$ is the greatest integer less than or equal to y; it is denoted $E(y)$.

1. Let $N = 4\,718$.
   a. Bound N between two consecutive powers of 10. Deduce bounds for $\log(N)$ between two consecutive integers.
   b. Compare $E(\log N)$ with the number of digits of N.
2. An integer N is written with 15 digits.
   a. Bound N between two consecutive powers of 10, then $\log(N)$ between two consecutive integers.
   b. What is $E(\log N)$?
3. State a rule giving the number of digits of N from $\log(N)$. Test it on $N = 1\,000$.
4. Use this rule to find the number of digits of: a. 3^{100}; b. 2^{64}; c. 365×10^{15}.
5. A password is made of 12 characters, each chosen from 62 symbols (26 lower-case letters, 26 upper-case letters, 10 digits). There are 62^{12} possible passwords. How many digits does this number have?

Homework

Redo practice exercise 3 with the following data: replace the initial value by 230 and the rate or common ratio by 6. Write it up in four lines: data, formula, calculation, conclusion.

Exit ticket – 3 minutes

On a half-sheet: write the main rule of the session, work through a very short example and state your confidence level from 1 to 4. Key question(s) of the day: 1 and 2.