## Course notes

**Definitions**

$u_n \to \ell \in \mathbb{R}$ means that any open interval containing $\ell$ contains all terms from a certain index. We say that $(u_n)$ converges to $\ell$. $u_n \to +\infty$ means that for every real number $A$, there is an index $N$ such that $n \geq N \Rightarrow u_n \geq A$. $u_n \to -\infty$ means that for every real number $A$, there is $N$ such that $n \geq N \Rightarrow u_n \leq A$.

A sequence may have no limit, not even an infinite one: $(-1)^n$ oscillates between $-1$ and 1. Every convergent sequence is bounded, but the converse is false.

## Operations and comparisons

For two convergent sequences with limits $a, b$, the sum tends to $a+b$, the product tends to $ab$ and the quotient to $\frac{a}{b}$ if $b \neq 0$. Rules involving infinity are used while checking signs. The forms $+\infty - \infty$, $0 \times \infty$, $\frac{\infty}{\infty}$, $\frac{0}{0}$ do not allow a direct conclusion.

**Example.** $\frac{3n^2-n+1}{2n^2+4} = \frac{3-\frac{1}{n}+\frac{1}{n^2}}{2+\frac{4}{n^2}} \to \frac{3}{2}$. For $\sqrt{n^2+n}-n$, multiplying by the conjugate gives $\frac{1}{\sqrt{1+\frac{1}{n}}+1} \to \frac{1}{2}$.

**Theorems**

If $u_n \leq v_n$ from a certain index onwards and $u_n \to +\infty$, then $v_n \to +\infty$. If $a_n \leq u_n \leq b_n$ from a certain index and $a_n, b_n \to \ell$, then $u_n \to \ell$ (squeeze theorem). Every increasing sequence bounded above converges. Every decreasing sequence that is bounded below converges. This theorem is assumed.

**Proof by comparison.** For every $A$, the terms of $u$ exceed $A$ from some index onwards $N_1$. The inequality $v_n \geq u_n$ holds beyond some index $N_2$. For $n \geq \max(N_1, N_2)$, $v_n \geq A$.

**Proof: increasing and not bounded above.** For every $A$, the absence of an upper bound gives $N$ such that $u_N > A$. Since it is increasing, $n \geq N \Rightarrow u_n \geq u_N > A$, hence $u_n \to +\infty$. Not being bounded above is not enough on its own.

## Geometric sequences

$$q^n \longrightarrow \begin{cases} 0 & \text{if } -1 < q < 1, \\ 1 & \text{if } q = 1, \\ +\infty & \text{if } q > 1. \end{cases}$$

If $q \leq -1$, $(q^n)$ has no limit: alternation for $q = -1$; even and odd subsequences of opposite signs and with unbounded moduli for $q < -1$.


**Bernoulli and geometric limits.** Fix $a \geq 0$. Let us prove $P(n)$ : "$(1+a)^n \geq 1+na$ ". At index 0, both sides equal 1. Fix $n \geq 0$ and suppose $P(n)$ true. Since $1+a > 0$,

$$(1+a)^{n+1} \geq (1+na)(1+a) = 1+(n+1)a+na^2 \geq 1+(n+1)a.$$

This is $P(n+1)$ since $na^2 \geq 0$. By induction, the inequality holds for every $n \geq 0$. If $q > 1$, take $a = q-1 > 0$ and compare to $1+na$. If $0 < q < 1$, apply this result to $\frac{1}{q}$, then take reciprocals. For $-1 < q < 0$, $|q^n| = |q|^n \to 0$ and squeezing concludes. The case $q = 0$ is handled from index 1.

**Recursively defined sequence**

For $u_{n+1} = f(u_n)$, establish a bound and monotonicity to prove convergence. If $f$ is continuous at the limit $\ell$, taking the limit gives $\ell = f(\ell)$ (continuity discussed in Chapter 4). Solving this equation never proves, on its own, that the sequence converges.




The points get closer to 2 without reaching it: $u_n = 2 + \frac{3}{n+1}$.

## Give meaning to the limits

### 2.1 A band and all subsequent indices

First use: S14.

**Definition · Converging to $\ell$**

We say $(u_n)$ converges to $\ell$ if, for any precision $\varepsilon > 0$, there exists an index $N$ such that $n \geq N$ implies $|u_n - \ell| < \varepsilon$. The index depends on the precision, but once chosen, it works for all the following terms.

**Worked example · Calculating a valid index**

For $u_n = 2 + \frac{3}{n+1}$, $|u_n - 2| = \frac{3}{n+1}$. The precision $0{,}1$ requires $n + 1 > 30$, so $n \geq 30$. Index 29 does not work: the gap there is exactly $0{,}1$. For any precision, every $N$ such that $N + 1 > 3/\varepsilon$ works.





**Figure 2.1.** Band $]1{,}9;2{,}1[$ : all terms from index $n \geq 30$ remain there.

**A1 · Immediate application** For $v_n = 5 - \frac{2}{n+1}$, determine the first index from which $4{,}95 < v_n < 5.05$. Check the previous index and explain why all later indices work.

**A2 · Counterexample** "There is a term in each band around 0, so the sequence converges to 0." Refute with a sequence whose even terms are zero and whose odd terms are 1.

### 2.2 Transform before concluding

First use: S16 and S19.

**Worked example · A form does not determine a limit**

The three quotients $n/n$, $n/n^2$ & $n^2/n$ all have the form $\infty/\infty$. Yet their limits are $1$, $0$ and $+\infty$. For $\frac{3n^2-n+1}{2n^2+4}$, dividing numerator and denominator by $n^2$ gives the limit $\frac{3}{2}$. The transformation is valid for $n \geq 1$, which is enough for a limit.

**A3 · Choose before calculating** Determine the limits of $\frac{2n+3}{n^2+1}$, $\sqrt{n^2+n}-n$ and $\frac{(-1)^n}{n+1}$. Name, for each, the decisive transformation or theorem.

A first step: In the second expression, multiply by the conjugate. In the third, bound the absolute value.

### 2.3 Converge, then identify

First use: S20; identification by continuity after S32.

**Property · Monotone convergence · accepted without proof**

An increasing sequence that is bounded above converges to a real number. The theorem says neither that it reaches its limit, nor that its limit is any one of its upper bounds.

**Worked example · A complete chain**

Let $u_0 = 1$ and $u_{n+1} = \sqrt{2+u_n}$. The interval $[1;2]$ is stable because $1 \leq u_n \leq 2$ implies $\sqrt{3} \leq u_{n+1} \leq 2$. On this interval, $2+x-x^2 = (2-x)(x+1) \geq 0$, therefore $u_{n+1} \geq u_n$ after comparison of positive squares. The sequence converges to $\ell \in [1;2]$. After continuity, $\ell = \sqrt{2+\ell}$ implies $(\ell-2)(\ell+1) = 0$; only $\ell = 2$ lies in $[1;2]$.

**A4 · Fix a line of reasoning** Seq. $a_0 = 1$, $a_{n+1} = 2a_n$ satisfies a relation whose only fixed point is 0. A student concludes $a_n \to 0$. Show the mistake and determine the actual behaviour.

**A5 · Independent proof** Let $b_0 = 0$, $b_{n+1} = \frac{1}{3}b_n + 2$. Prove $0 \leq b_n < 3$, then that it is increasing. Deduce convergence and determine the limit by the explicit formula or, after S32, by continuity.

### 2.4 Do not forget negative ratios

First use: S19 and S21.

**Property · Behaviour of $q^n$**

If $|q| < 1$, $q^n \to 0$. If $q > 1$, $q^n \to +\infty$. For $q = 1$, the sequence is constant. For $q \leq -1$, it has no limit, not even an infinite one: the signs alternate, and the two subsequences of even and odd indices have incompatible behaviours.

**Proof.** For $q > 1$, write $q = 1+a$ with $a > 0$. Induction gives $(1+a)^n \geq 1+na$, which tends to $+\infty$ ; comparison gives the result. For $0 < q < 1$, apply this result to $1/q$ and take the reciprocal. For $-1 < q < 0$, $|q^n| = |q|^n \to 0$ implies $q^n \to 0$ by the squeeze theorem. The case $q = 0$ is immediate from index 1. □

**Worked example · Common ratio and behaviour**

For $q = -2$, the even-index terms $4^k$ tend to $+\infty$ and the odd ones $-2 \times 4^k$ to $-\infty$ : no common limit. For $q = -1/2$, the signs also alternate, but the absolute value $2^{-n}$ tends to 0; the sequence converges.

**A6 · Compare properties** For $(-1)^n$, $-\frac{1}{n+1}$ & $n$, state: bounded? monotonic? convergent ? Give justifications, not just a table of answers.

**One minute, spoken aloud.** Explain why "increasing" means neither "positive" nor "tends to $+\infty$ ".

**Delayed review · S33 then S66.** Go back to identifying the limit by continuity; for a sequence of integrals, rebuild the same chain: bounds, monotonicity, convergence, identification.

## Investigation

**A limit is not a last term - 30 min.**

Consider the sequence $(u_n)$ defined on $\mathbb{N}$ by $u_n = 2 + \frac{3}{n+1}$.

1. Calculate $u_0$, $u_9$ and $u_{99}$. Is there an integer $n$ such that $u_n = 2$? Justify.
2. Determine the smallest integer $N$ such that, for every integer $n \geq N$, $|u_n - 2| < 0{,}01$.
3. Let $\varepsilon > 0$. Give, in terms of $\varepsilon$, an integer $N$ such that $|u_n - 2| < \varepsilon$ for all $n \geq N$.
4. On $\mathbb{N}$ define $(v_n)$ by $v_n = 2 + \frac{(-1)^n}{n+1}$. Prove that it converges to 2 and that it is not monotonic. What does this example show?