CHAPTER 10

Trigonometry: circle, radians and angles associated

Chapter path

Objectives. Convert degrees to radians; plot points; calculate sine and cosine standard values; use the symmetries of the circle.
Prerequisites. Circle, Pythagoras, cosine and sine in a right triangle, symmetries, fractions.
Yearly reference point. 2·11 March: 8 a.m.

Diagnostic check: six questions, 10 minutes

Answer without looking at your notes. Indicate the questions for which help is needed. This diagnostic test is not graded.

1. Circumference of a circle of radius 1?

2. How many degrees is a quarter turn?

3. Calculate $\sqrt{\frac{1}{2}}$.

4. Coordinates of the reflection of ($a; b$) with respect to the x-axis?

5. Simplify $\frac{5\pi}{3} - 2\pi$.

6. In a right triangle, opposite side 3 and hypotenuse 5: sine?

After the diagnostic check. Go over the flagged concepts with the teacher; redo a question similar, then explain the rule used.

Oral work and logic. Present an equality between associated angles, naming the symmetry that justifies it; explain what a radian measures.

Investigate before formalising

Construct, without a protractor, a point at angle $\frac{\pi}{3}$ then a corner point $\frac{\pi}{4}$. Justify the exact values of their coordinates from the figures.

Organisation. Individual work 5 min, discussion in pairs 5 min, whole-class sharing 5 min. Keep a record of an attempt, even an unfinished one. A conjecture must then be justified.

10.1 Measuring an angle in radians

Definition and results

The unit circle is the circle with centre O and radius 1, oriented anticlockwise clockwise. A radian corresponds to an arc of length 1 on this circle. One turn is 2 π radians, half a turn π. On a circle of radius r, an arc length is $r\theta$ for a positive angle θ expressed in radians.

Status and justification. Definitions and proportionality between angle at the centre and arc length. Conversion is 180 $^\circ = \pi$ radians; the degrees are multiplied by $\frac{\pi}{180}$.

Worked example. $60^\circ = \frac{\pi}{3}$; $\frac{3\pi}{2} = 270^\circ$. An angle of $\frac{\pi}{3}$ on a circle of radius 6 cm gives an arc of 2 π cm.

Watch out

The length of an arc is not that of the chord. In trigonometric calculations, check the mode degree or radian mode of the tool.

Immediate application. Convert 45 $^\circ$, 120° ; give the arc with radius 2 m and angle $\frac{3\pi}{4}$.

10.2 Wrapping and the image of a real number

Definition and results

To each real number x corresponds a point M on the circle obtained by a directed path of length x from I(1;0). Two real numbers have the same image if their difference is an integer multiple of 2 π. A negative number indicates travel in the clockwise direction.

Status and justification. Adding or removing a whole number of turns keeps the end point. Conversely, returning to the same point requires an integer number of turns.

Worked example. $\frac{13\pi}{3} = 4\pi + \frac{\pi}{3}$ has the same image as $\frac{\pi}{3}$. $-\frac{\pi}{2}$ has the same image as $\frac{3\pi}{2}$: point $(0; -1)$.

Watch out

Add π gives the opposite point, not the same point.

Immediate application. Plot the images of $\frac{7\pi}{2}$ and $-\frac{5\pi}{3}$ reducing them to one turn.

10.3 Sine, cosine and standard values


Definition and results

If M is the image of x, $M = (\cos x; \sin x)$. Hence $-1 \leq \cos x, \sin x \leq 1$ and $\cos^2 x + \sin^2 x = 1$.

For $x = 0, \frac{\pi}{6}, \frac{\pi}{4}, \frac{\pi}{3}, \frac{\pi}{2}$, the cosines are 1 $, \frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{2}, \frac{1}{2}$, 0 and the sines 0 $, \frac{1}{2}, \frac{\sqrt{2}}{2}, \frac{\sqrt{3}}{2}, 1$.

Status and justification. À $\frac{\pi}{4}$, the two coordinates are equal and positive ; Pythagoras gives $2a^2 = 1$, therefore $a = \frac{\sqrt{2}}{2}$. An equilateral triangle with side 1, cut by its altitude, has half-base $\frac{1}{2}$ and height $\frac{\sqrt{3}}{2}$: $\cos(\frac{\pi}{3}) = \frac{1}{2}$, $\sin(\frac{\pi}{3}) = \frac{\sqrt{3}}{2}$. The complementary angle gives $\frac{\pi}{6}$.

Worked example. The corner point $\frac{\pi}{3}$ is $(\frac{1}{2}; \frac{\sqrt{3}}{2})$. The scalar product of two unit vectors making this angle equals its cosine, so $\frac{1}{2}$. The geometry chapters thus connect to the circle.



For the angle $\frac{\pi}{3}$, the projections of M give its cosine and sine directly. The fractions are placed outside the lines of the drawing.

Watch out

The cosine is the x-coordinate, the sine the y-coordinate. The identity gives an absolute value: the quadrant determines the sign.

Immediate application. If $\cos x = \frac{3}{5}$ and the image of x is in the fourth quadrant, give $\sin x$.

10.4 Related angles and symmetries

Definition and results

The symmetries of the circle give $\cos(-x) = \cos x$, $\sin(-x) = -\sin x$; $\cos(\pi - x) = -\cos x$, $\sin(\pi - x) = \sin x$; $\cos(\pi + x) = -\cos x$, $\sin(\pi + x) = -\sin x$. Add 2 $k\pi$, k integer, does not change neither sine nor cosine.

Status and justification. Each equality is read on the coordinates of symmetric points: axis horizontal for $-x$, vertical axis for $\pi - x$, centre O for $\pi + x$. This is not a catalogue without justification.

Worked example. $\cos(\frac{5\pi}{6}) = -\frac{\sqrt{3}}{2}$, $\sin(\frac{5\pi}{6}) = \frac{1}{2}$ by symmetry of $\frac{\pi}{6}$. $\cos(-\frac{\pi}{4}) = \frac{\sqrt{2}}{2}$, $\sin(-\frac{\pi}{4}) = -\frac{\sqrt{2}}{2}$.

Watch out

Two angles with the same cosine do not necessarily have the same sine. Reduce the angle, then read the signs.

Immediate application. Calculate the two coordinates of the images of $\frac{4\pi}{3}$ and $-\frac{7\pi}{4}$.

10.5 Approximating pi and reusing ideas

Definition and results

A regular n-sided polygon inscribed in the unit circle has semi-perimeter $n\sin(\frac{\pi}{n})$; a polygon circumscribed has semi-perimeter $n\tan(\frac{\pi}{n})$, with $\tan\theta = \frac{\sin\theta}{\cos}\theta$ here. Archimedes' method doubles the number of sides and calculates the new lengths by Pythagoras, without knowing pi in advance.

Status and justification. The perimeter of the inscribed polygon is less than that of the circle, and that of the circumscribed upper: geometric bounding assumed. If c is the chord of one side, set $h = \sqrt{1 - \frac{c^2}{4}}$; the chord of the half-angle is $\frac{c}{\sqrt{2(1+h)}}$. This formula comes from Pythagoras and avoids subtracting real numbers close.

Worked example. For the inscribed hexagon, c=1 and semi-perimeter 3. The circumscribed one gives 2 $\sqrt{3} \approx$ 3,4641. Doubling the number of sides tightens the bounds. The given program starts from n=6 and c=1.

Watch out

Using a sine function directly with pi already known would check a calculation, but would not would not construct an approximation of pi independently.

Immediate application. For n=6 and c=1, calculate the new chord after doubling.

Exercise path

Start with exercises 1 and 2. Then choose exercise 3 or 4 without help. Exercises 5 to 7 are a reserve for transfer, investigation or programming: the teacher chooses one. Exercise 8 is revisited later.

Exercise 10.1 · Convert and Measure

The unit circle is used to measure angles in radians.

1. Convert 150 $^\circ$ and -90° in radians, showing the conversion factor.
2. A circular sector has a radius of 4 cm and an angle of $\frac{\pi}{3}$ radians. Calculate the exact length of its arc.
3. Explain why an angle measure can be negative while a geometric arc length is positive.

Exercise 10.2 · Reduce angles

Consider the real numbers $\frac{17\pi}{6}$, $-\frac{7\pi}{4}$ and 5π.

1. For each, determine a representative in [0; 2 π[having the same image on the circle trigonometric. Show the number of turns added or removed.
2. Place the three images on a circle and label them with the original real numbers.

Exercise 10.3 · Exact values

We look for the exact values of the cosine and sine of $\frac{2\pi}{3}$, $-\frac{\pi}{6}$ and $\frac{5\pi}{4}$.

1. Place the three images on the unit circle. For each, state the expected sign of its two coordinates.
2. Calculate the exact values using a standard angle and a symmetry. Present the answer in a table: angle, cosine, sine.

Exercise 10.4 · Find the sign

A real number x satisfies $\sin x = \frac{5}{13}$. Its image on the unit circle lies in the second quadrant, hence to the left of the y-axis and above the x-axis.

1. Determine the exact value of cos x. Justify both its absolute value and its sign.

2. Explain why knowing only sin x would not have been enough to choose the sign of cos x.

Exercise 10.5 · A wheel

A wheel of radius 0 ,30 m rolls on horizontal ground without slipping. It turns through an angle $\frac{5\pi}{3}$ radians. We want the distance travelled by its axle in the direction of motion.

1. Calculate this distance exactly, explaining the link with the unrolled arc length.

2. Explain why the length of a chord of the circle must not be used. The path of a fixed point on the rim of the wheel, seen from the ground, the same as that of the axle?

Exercise 10.6 · Chord and scalar product

On the trigonometric circle with centre O, the point A is the image of 0 and the point B that of $\frac{2\pi}{3}$.

1. Determine the exact coordinates of A and B, then calculate the length AB.

2. Find this length again by applying Al-Kashi's formula to the triangle OAB. State the lengths and the angle used.

Exercise 10.7 · Read an equation on the circle

We look for the real numbers x of the interval [0; 2 π[whose image on the trigonometric circle has x-coordinate $\frac{1}{2}$.

1. Express the condition on the x-coordinate as an equation involving cos x.

2. Draw the corresponding vertical line on a circle, locate its points of intersection and give the two solutions. Check that they belong to the required interval.

Exercise 10.8 · Delayed review

Consider the angle with measure $\frac{25\pi}{4}$ radians. Answers must be exact, without approximation decimal.

1. Reduce this angle by removing a whole number of complete turns.

2. Deduce its cosine and sine. Justify the values using the unit circle.

Independent work. Do only the task assigned at the end of the session, in the scheme of work Date: 10·20 min maximum. The exercises depend on the concepts already studied; some questions are therefore postponed. Revisit exercise 8 one week after the chapter (10 min). If the planned time is exceeded, note the difficulty and bring your attempts; do not use up the whole reserve.

To be done in the CQFD textbook

These exercises are done in your textbook. The scheme of work gives the session; they replace the personal work planned that day (10 to 20 minutes in total).

CQFD, exercise 32 p. 220, questions a and b. Use the exact value given and the angles associated, without a calculator.

CQFD, Exercise 39 p. 220, Questions 1, 2, 3 and 5. Do the four calculation and reading the circle; justify the answers on your script.

MCQ 1: understand and choose

Five teaching questions with a single answer. Choose one answer per question; then explain one distractor. This chapter format does not reproduce the official format of the EAM.

1. $180°$ is
   - A. 2π
   - B. π
   - C. $\frac{\pi}{2}$
   - D. 1
2. $\cos(\frac{\pi}{3}) =$
   - A. $\frac{1}{2}$
   - B. $\frac{\sqrt{3}}{2}$
   - C. 0
   - D. 1
3. $\sin(-\frac{\pi}{2}) =$
   - A. 1
   - B. 0
   - C. $\frac{1}{2}$
   - D. -1
4. $\cos^2 x + \sin^2 x =$
   - A. 0
   - B. 2
   - C. 1
   - D. x
5. The same point corresponds to angles differing by
   - A. π
   - B. $2k\pi$, k an integer
   - C. $\frac{\pi}{2}$
   - D. 1

MCQ 2: understand and choose

Five teaching questions with a single answer. Choose one answer per question; then explain one distractor. This chapter format does not reproduce the official format of the EAM.

1. $\cos(\pi - x) =$
   - A. $\cos x$
   - B. $\sin x$
   - C. $-\cos x$
   - D. $-\sin x$
2. $\sin(\frac{\pi}{4}) =$
   - A. $\frac{\sqrt{2}}{2}$
   - B. $\frac{1}{2}$
   - C. $\frac{\sqrt{3}}{2}$
   - D. $\sqrt{2}$
3. Image of $\frac{3\pi}{2}$:
   - A. $(1;0)$
   - B. $(-1;0)$
   - C. $(0;1)$
   - D. $(0;-1)$
4. Arc with radius 3 and angle 2 radians: length
   - A. $\frac{2}{3}$
   - B. 6
   - C. $\frac{3}{2}$
   - D. 6π
5. Inside the second quadrant: sign of the cosine
   - A. positive
   - B. zero everywhere
   - C. Negative
   - D. cannot be determined

True or false? Justify.

1. $\cos(-x) = -\cos x$ for all x.
2. $\sin(x + 2\pi) = \sin x$.
3. Knowing sin x always determines cos x.

Personal review

I reduce the angles by complete turns; I read cosine and sine in the right order; I justify the signs; I keep the exact values.
For each objective, indicate: mastered without help; achieved with help; needs more work. Write an example and a mistake to avoid.