Meeting 47 — Expectance of a random variable

Discrete random variables

06/04/2027 • B : 2 h – C : 1 h

Objectives of the session

· Define a law.

· Calculate a expectation.

· Interpret a theoretical average.

Fluency · 6 to 8 minutes, without calculator

1. Reduce 3 $x + 8 - x + 2$. .....
2. Provide $E(X)$ for $X \sim \mathcal{B}(10, 0{,}3)$. .....
3. Calculate $\binom{5}{2}$. .....
4. Calculate the expectation of X constant of 8 .....
5. Translate $X \geq 1$ by an additional .....

Reminder and Prerequisites

Read the last written record and know how to perform the elementary calculations mobilized today. Before the activity, reformulate the keywords: define a law and calculate a expectation.

Research activity

A game costs 2 euros. The gross gain is 0, 3 or 10 euros with probability 0 ,6, 0,3, 0,1. Building the law of net gain X, calculate $E(X)$ and decide if the game is favorable to the player.

Trace written

The law of X associated with each value x_i probability p_i. Expectation $E(X) = \sum x_i p_i$ is an average long-term theory.

Worked example

Retrieve activity data and write the solution in order: definition of objects, property used, exact calculation or approach, then response phrase. The model response figure in the teaching edition.

Training · essential

1. Justify that the number of successes among 8 independent trials, with $p = 0{,}3$, follows a law binomial.
2. Calculate $P(X = 2)$ and $P(X = 0)$.
3. Deduct from it $P(X \geq 1)$ and calculate $E(X)$.

Reference in the manual

Supplements: Calao, course pp. 154-159; exercises p. 160, no. 1-5.
session is identified below by its page and number; the reference also allows you to find
the figures and digital resources of the manual.

Manuel Calao · p. 160, exercise 2

A game consists of launching a well balanced dice. If the result is:

· 1, 2 or 3, then you lose 2 points;

· 3 or 4, so you don't lose or gain anything;

· 6, then we win 3 points.

Set G the random variable that, at each throw, combines the player's (positive or negative) gain.

1. Complete the table that represents the probability distribution of the random variable G.
2. Calculate $E(G)$. What can we notice?

Personal work

Resuming Exercise 2 with the following data:
355 and rate or common difference / ratio by 7. Write in four lines: data, formula, calculation, conclusion.

Exit ticket · 3 minutes

On half a sheet: write the main rule of the session, deal with a very short example and
indicate confidence level of 1 to 4. Key question(s): 1, 2 and 3.

Meeting 48 — Binomial distribution: recognizing a situation

Discrete random variables

08/04/2027 • B : 1 h – C : 2 h

Objectives of the session

· Recognize a Bernoulli scheme.

· Identify n and p.

· Define X.

Fluency · 6 to 8 minutes, without calculator

1. Calculate 5 per cent of 180 .....

2. Calculate $\binom{5}{2}$. .....

3. Calculate the expectation of X constant of 9 .....

4. Translate $X \geq 1$ by an additional .....

5. Provide $E(X)$ for $X \sim \mathcal{B}(10,0{,}3)$. .....

Reminder and Prerequisites

Read the last written record and know how to perform the elementary calculations mobilized today. Before the activity, have the keywords reworded: recognize a bernoulli scheme and identify n and p.

Research activity

A customer accepts an offer with probability 0 ,25. An independent adviser contacts eight Clients. Define X, justify the binomial distribution and identify n and p.

Trace written

$X \sim \mathcal{B}(n, p)$ if X counts the successes in n independent tests of the same probability p.

Worked example

Retrieve activity data and write the solution in order: definition of objects, property used, exact calculation or approach, then response phrase. The model response figure in the teaching edition.

Training · essential

1. Justify that the number of successes among 9 independent trials, with $p = 0{,}3$, follows a law binomial.

2. Calculate $P(X = 2)$ and $P(X = 0)$.

3. Deduct from it $P(X \geq 1)$ and calculate $E(X)$.

Reference in the manual

Supplements: Calao, courses p. 154-159; exercises p. 160, n° 5, and p. 162, n° 14-20. reproduced in this session is identified below by its page and number; the reference allows

also find the figures and digital resources of the manual.

Manuel Calao · p. 163, exercise 22

The mayor of a city has decided to hold a referendum on the possible development of a park.
The results are as follows: 63% for and 37% against.
The number of participants is large enough to be able to assimilate this
draw at random and with two ballot papers.
The random variable is considered X which, in any draw of two ballots, combines the number "For" bulletins.

1. What is the law followed by X ? Determine its parameters.
2. Calculate the probability that, in such a sampling, one has: none "For"; exactly one " For."
3. Deduce from question 2 the probability that, in such a sampling, there is at most one "For ».

Personal work

Resuming Exercise 3 with the following data:
360 and rate or common difference / ratio by 2. Write in four lines: data, formula, calculation, conclusion.

Exit ticket · 3 minutes

On half a sheet: write the main rule of the session, deal with a very short example and indicate the degree of confidence from 1 to 4. Key question(s): 1 and 2.

Meeting 49 — Pascal triangle and binomial coefficients

Discrete random variables

27/04/2027 • B : 2 h – C : 1 h

Objectives of the session

· Build Pascal's triangle.

· Read $\binom{n}{k}$.

· Interpret a coefficient.

Fluency · 6 to 8 minutes, without calculator

1. Write the multiplier coefficient of a 10 per cent increase . . . . . .
2. Calculate the expectation of X constant equal to 3 . . . . . .
3. Translate $X \geq 1$ by an additional . . . . . .
4. Provide $E(X)$ for $X \sim \mathcal{B}(10,0{,}3)$. . . . . .
5. Calculate $\binom{5}{2}$. . . . . .

Reminder and Prerequisites

Read the last written record and know how to perform the elementary calculations mobilized today. Before the activity, reformulate the keywords: build the pascal triangle and read $\binom{n}{k}$.

Research activity

Build Lines $n = 0$ à $n = 6$ of Pascal's triangle. Explain the calculation of the number 15 of the line 6, then read $\binom{6}{2}$ and $\binom{6}{4}$.

			1			
		1		1		
	1		2		1	
	1		3		3		1
1		4		6		4		1

Trace written

Pascal's triangle checks $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$. The edges are worth 1.

Worked example

Retrieve activity data and write the solution in order: definition of objects, property used, exact calculation or approach, then response phrase. The model response figure in the teaching edition.

Training · essential

1. Justify that the number of successes among 10 independent trials, with $p = 0{,}3$, follows a law binomial.
2. Calculate $P(X = 2)$ and $P(X = 0)$.

3. Deduct from it $P(X \geq 1)$ and calculate $E(X)$.

Reference in the manual

Supplements: Calao, p. 164, Nos. 24 to 29, then p. 166, No. 36.
is identified below by its page and number; the reference also allows to find the figures and numerical resources of the manual.

Manuel Calao · p. 161, exercise 13

We throw a well-balanced coin five times in a row. F the random variable which, at each experience thus defined, combines the number of "faces" obtained.

1. F follow a binomial distribution, specify its parameters.
2. Realize the triangle of Pascal up to $n = 5$.
3. Calculate $P(F = 3)$. Interpret this result.
4. Calculate the probability that the "Face" side will appear at least three times.

Personal work

Resuming Exercise 2 with the following data:
365 and rate or common difference / ratio by 3. Write in four lines: data, formula, calculation, conclusion.

Exit ticket · 3 minutes

On half a sheet: write the main rule of the session, deal with a very short example and indicate confidence level of 1 to 4. Key question(s): 1, 2 and 3.

Meeting 50 — Binomial probabilities

Discrete random variables

29/04/2027 • B : 1 h – C : 2 h

Objectives of the session

· Calculate $P(X = k)$.

· Translate at least/at most.

· Use the supplement.

Fluency · 6 to 8 minutes, without calculator


1. Reduce 3 $x + 4 - x + 2$. .....
2. Translate $X \geq 1$ by an additional .....
3. Provide $E(X)$ for $X \sim \mathcal{B}(10, 0{,}3)$. .....
4. Calculate $\binom{5}{2}$. .....
5. Calculating the Expectation of X constant of 4 .....

Reminder and Prerequisites

Read the last written record and know how to perform the elementary calculations mobilized today. Before the activity, reformulate the keywords: calculate $p(x = k)$ and translate at least/at most.

Research activity

For $X \sim \mathcal{B}(8, 0{,}25)$, calculate $P(X = 2)$, $P(X = 0)$ then $P(X \geq 1)$. Translate each event into sentence.



Trace written

If $X \sim \mathcal{B}(n, p)$, then $P(X = k) = \binom{n}{k} p^k (1-p)^{n-k}$. Carefully translate "at least", "at most and "none".

Worked example

Retrieve activity data and write the solution in order: definition of objects, property used, exact calculation or approach, then response phrase. The model response figure in the teaching edition.

Training · essential

1. Justify that the number of successes among 6 independent trials, with $p = 0{,}3$, follows a law binomial.
2. Calculate $P(X = 2)$ and $P(X = 0)$.
3. Deduct from it $P(X \geq 1)$ and calculate $E(X)$.

Reference in the manual

Supplements: Calao, p. 164, Nos. 24 to 29, and p. 168, Nos. 41 to 45.
session is identified below by its page and number; the reference also allows you to find the figures and digital resources of the manual.

Manuel Calao · p. 167, exercise 38

A parcel delivery company succeeds, for packages taken care of before 10 a.m., in delivering them as soon as possible the next day with a probability of 0 ,75. A merchant shall deposit before 10:07 a.m. parcels for his Clients. X the random variable which, in each batch of seven packages, combines the number of packages The next day, we admit that X follows a binomial distribution.

1. Determine parameters n and p of the law X.
2. Calculate the probability of the following events: A: "All packages will arrive the next day." ; B: "At least one package will not arrive the next day."
3. Complete Pascal's triangle up to the line $n = 7$. Deduct $\binom{7}{2}$. Calculate probability Event C: " Exactly two packages will arrive the next day."
4. Interpret the event $\{X = 4\}$. Calculate probability $P(X = 4)$.

Personal work

Resuming Exercise 3 with the following data:
370 and the rate or common difference / ratio by 4. Write in four lines: data, formula, calculation, conclusion.

Exit ticket · 3 minutes

On half a sheet: write the main rule of the session, deal with a very short example and indicate the degree of confidence from 1 to 4. Key question(s): 1 and 2.

Meeting 51 — Chapter review and evaluation

Discrete random variables

04/05/2027 • B : 2 h – C : 1 h

Objectives of the session

· Calculate and interpret $E(X)$.

· Solve a binomial problem.

· Analyze errors.

Fluency · 6 to 8 minutes, without calculator

1. Calculate 20 per cent of 100 .....
2. Provide $E(X)$ for $X \sim \mathcal{B}(10, 0{,}3)$. .....
3. Calculate $\binom{5}{2}$. .....
4. Calculate the expectation of X constant equal to 5 .....
5. Translate $X \geq 1$ by an additional .....

Reminder and Prerequisites

Read the last written record and know how to perform the elementary calculations mobilized today. Before the activity, reformulate the keywords: calculate and interpret $e(x)$ and solve a problem binomial.

Research activity

One centre receives 120 groups of 10 clients, each client probably claiming 0 ,08. Estimate the average number of complaints per group, then on all groups.
Expectation may or may not predict.

Trace written

For $X \sim \mathcal{B}(n, p)$, $E(X) = np$. The value gives the average number of successes over a large number of series of n tests.

Worked example

Retrieve activity data and write the solution in order: definition of objects, property used, exact calculation or approach, then response phrase. The model response figure in the teaching edition.

Training · essential

1. Justify that the number of successes among 7 independent trials, with $p = 0{,}3$, follows a law binomial.
2. Calculate $P(X = 2)$ and $P(X = 0)$.
3. Deduct from it $P(X \geq 1)$ and calculate $E(X)$.

Reference in the manual

Supplements: Calao, pp. 160-169. The exercise reproduced in this session is identified below. by its page and number; the reference also makes it possible to find figures and resources Numbers of the manual.

Manuel Calao · p. 166, exercise 34

A company markets, in large quantities, low consumption and long-lasting LED bulbs 98% of these bulbs have a running time of at least 20,000 hours. bulbs are sold in lots of 10. X the random variable which, in each batch of 10 ampoules, combines the number of bulbs whose operating time is at least equal to 20 000 h. The quantity of ampoules is large enough to assimilate the choice of a batch at random sampling and with 10 ampoules delivered.

1. What is the law followed by X ? Specify its parameters.
2. Copy and complete: $\binom{10}{9} = \cdots$; $\binom{10}{10} = \cdots$.
3. Deduce probabilities (round to 0 ,001) : $P(X = 9)$ and $P(X = 10)$. Interpret results previous.
4. Deduce the probability (at 0 ,001) that a batch of 10 ampoules contains at least 9 bulbs with a running time of at least 20,000 hours.

Personal work

Resuming Exercise 2 with the following data:
Write in four lines: data, formula, calculation, conclusion.

Exit ticket · 3 minutes

On half a sheet: write the main rule of the session, deal with a very short example and indicate confidence level of 1 to 4. Key question(s): 1, 2 and 3.